

International Journal of Modern Physics D
© World Scientific Publishing Company

REVISITING THE CONFRONTATION OF THE ENERGY CONDITIONS WITH SUPERNOVAE DATA

MARIANA P. LIMA, SANDRO D. P. VITENTI and MARCELO J. REBOUÇAS

*Centro Brasileiro de Pesquisas Físicas
Rua Dr. Xavier Sigaud 150
22290-180 Rio de Janeiro – RJ, Brazil
penna@cbpf.br, vitenti@cbpf.br, reboucas@cbpf.br*

Received Day Month Year
Revised Day Month Year
Communicated by Managing Editor

In the standard Friedmann–Lemaître–Robertson–Walker (FLRW) approach to model the Universe the violation of the so-called energy conditions is related to some important properties of the Universe as, for example, the current and the inflationary accelerating expansion phases. The energy conditions are also necessary in the formulation and proofs of Hawking–Penrose singularity theorems. In two recent articles we have derived bounds from energy conditions and made confrontations of these bounds with supernovae data. Here, we extend these results in following way: first, by using our most recent statistical procedure for calculating new $q(z)$ estimates from the *gold* and *combined* type Ia supernovae samples; second, we use these estimates to obtain a new picture of the energy conditions fulfillment and violation for the recent past ($z \leq 1$) in the context of the standard cosmology.

Keywords: Energy conditions; energy condition confrontation with supernovae data.

1. Introduction

In the absence of constraints on the energy-momentum tensor $T_{\mu\nu}$ any metric satisfies Einstein's equations since they can be regarded as a definition of $T_{\mu\nu}$, i.e., a set of equations determining $T_{\mu\nu}$ for any given metric $g_{\mu\nu}$. However, if one wishes to explore general properties that hold for a variety of different physical sources it is convenient to impose the so-called *energy conditions* that limit the arbitrariness of $T_{\mu\nu}$ on physical grounds.¹

On scales relevant for cosmology, an important point in the study of the energy conditions is the confrontation of their predictions with the observational data. By using model-independent energy-condition *integrated* bounds on the cosmological observables as, for example, the distance modulus and lookback time, this confrontation has been made in some recent articles^{2–8} (see also the pioneering Refs. 9 by Visser). In Ref. 10, however, it was shown that the fulfillment (or the violation) of these *integrated* bounds at a specific redshift z is not a sufficient (nor a necessary)

2 Mariana P. Lima, Sandro D.P. Vitenti, Marcelo J. Rebouças

local condition for the fulfillment (or respectively the violation) of the associated energy condition at z .^a In this way, the confrontation between the prediction of these *integrated* bounds and observational data cannot be used to draw conclusions on the fulfillment (or violation) of the energy conditions at z . In Ref. 10 this problem was overcome by deriving new *non-integrated* energy-condition bounds, and confrontations between the new bounds with type Ia supernovae (SNe Ia) data of the *gold*¹⁴ and *combined*¹⁵ samples were performed by using the upper and lower limits of confidence regions on $E(z) - q(z)$ plane. More recently, in Ref. 16 a new statistical way for estimating the deceleration parameter $q(z)$ was carried out, and a new picture of the energy conditions fulfillment and violation for recent past ($z \leq 1$) was calculated by using the recently compiled *Union* sample.¹⁷

In this work, we use the most recent statistical procedure introduced in Ref. 16 along with the *gold*¹⁴ and *combined*¹⁵ samples to obtain estimates of $q(z)$ in order to build up a new picture of the confrontation between the energy condition *integrated* bounds and these SNe Ia data sets, completing therefore the cycle of this type of analysis which involves these three samples and the two statistical procedures to $q(z)$ estimates of Refs.10 and 16.

2. Preliminaries

It is known that the energy conditions can be stated in a coordinate-invariant way in terms of $T_{\mu\nu}$ and vector fields of fixed character (timelike, null and spacelike). However, within the framework of the standard Friedmann-Lemaître-Robertson-Walker (FLRW) model, we only need to consider the energy-momentum tensor of a perfect fluid with density ρ and pressure p , i.e., $T_{\mu\nu} = (\rho + p)u_\mu u_\nu - pg_{\mu\nu}$. In this context, the energy conditions take the following forms:¹

$$\begin{aligned} \text{NEC} : \quad & \rho + p \geq 0, \\ \text{WEC} : \quad & \rho \geq 0 \quad \text{and} \quad \rho + p \geq 0, \\ \text{SEC} : \quad & \rho + 3p \geq 0 \quad \text{and} \quad \rho + p \geq 0, \\ \text{DEC} : \quad & \rho \geq 0 \quad \text{and} \quad -\rho \leq p \leq \rho, \end{aligned} \tag{1}$$

where NEC, WEC, SEC and DEC correspond, respectively, to the null, weak, strong, and dominant energy conditions. For a FLRW metric with a scale factor $a(t)$, the density ρ and pressure p of the cosmological fluid are given by

$$\rho = \frac{3}{8\pi G} \left[\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right] \quad \text{and} \quad p = -\frac{1}{8\pi G} \left[2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right], \tag{2}$$

where overdots denote the derivative with respect to the time t and G is Newton's gravitational constant.

^aEnergy conditions constraints on the so-called $f(R)$ -gravity have also been investigated in Ref. 11 and more recently in Refs. 12 and 13.

The *non-integrated* bounds from energy conditions derived in Ref. 10 can be obtained in terms of the deceleration parameter $q(z) = -\ddot{a}/aH^2$, the normalized Hubble function $E(z) = H(z)/H_0$, and the curvature density parameter $\Omega_{k0} = -k/(a_0H_0)^2$, simply by substituting Eqs. (2) into Eqs. (1). This gives^b

$$\mathbf{NEC} \Leftrightarrow q(z) - \Omega_{k0} \frac{(1+z)^2}{E^2(z)} \geq -1, \quad (3)$$

$$\mathbf{WEC} \Leftrightarrow \frac{E^2(z)}{(1+z)^2} \geq \Omega_{k0}, \quad (4)$$

$$\mathbf{SEC} \Leftrightarrow q(z) \geq 0, \quad (5)$$

$$\mathbf{DEC} \Leftrightarrow q(z) + 2\Omega_{k0} \frac{(1+z)^2}{E^2(z)} \leq 2, \quad (6)$$

where $z = (a_0/a) - 1$ is the redshift, $H(z) = \dot{a}/a$, and the subscript 0 stands for present-day quantities.

In this work, we focus on the FLRW flat ($\Omega_{k0} = 0$) universe. In this case the **NEC**, **SEC** and **DEC** bounds reduce, respectively, to $q(z) \geq -1$, $q(z) \geq 0$ and $q(z) \leq 2$, while the **WEC** bound is fulfilled identically. Thus, having estimates of $q(z_*)$ for different redshifts z_* , one can test the fulfillment or violation of the energy conditions at each z_* .

Now, the $q(z_*)$ estimates are obtained by using a SNe Ia data set, by approximating the deceleration parameter $q(z)$ function as the following linear piecewise continuous function:

$$q(z) = q_l + q'_l \Delta z_l, \quad z \in (z_l, z_{l+1}), \quad (7)$$

where the subscript l means that the quantity is taken at z_l , $\Delta z_l \equiv (z - z_l)$, and the prime denotes the derivative with respect to z . The supernovae observations provide the redshifts and distance modulus

$$\mu(z) = 5 \log_{10} \left[\frac{c(1+z)}{H_0 \text{ 1Mpc}} \int_0^z \frac{dz'}{E(z')} \right] + 25. \quad (8)$$

Then, by using the following well known relation between $q(z)$ and $E(z)$:

$$E(z) = \exp \int_0^z \frac{1+q(z')}{1+z'} dz', \quad (9)$$

along with Eq. (8), we fitted the parameters of the $q(z)$, as given by (7), by using the SNe Ia redshift–distance modulus data from the *gold*¹⁴ and *combined*¹⁵ samples.

3. Results and Conclusions

Since in the flat case the energy condition bounds given by Eqs. (3), (5) and (6) depend only on $q(z)$, we have obtained the $q(z_*)$ estimates at $1\sigma - 3\sigma$ confidence

^bThrough out this paper we use the notation of Ref. 10 in which **NEC**, **WEC**, **SEC** and **DEC** correspond, respectively, to $\rho + p \geq 0$, $\rho \geq 0$, $\rho + 3p \geq 0$ and $\rho - p \geq 0$.

4 Mariana P. Lima, Sandro D.P. Vitenti, Marcelo J. Rebouças

levels from *gold* and *combined* SNe Ia samples by marginalizing over $E(z_*)$ and the other parameters (q'_i 's) of the $q(z)$ function [Eq.(7)].^c

A global picture of the breakdown and fulfillment of the energy conditions in the recent past has been built up with the $q(z_*)$ estimates at 200 equally spaced redshifts in the interval $(0, 1]$. Fig. 1(a) shows the **NEC**, **SEC**, and **DEC** bounds along with the best-fit values and the 1σ , 2σ and 3σ limits of $q(z_*)$ in the $q(z) - z$ plane. We recall that **WEC** bound [$E^2(z) \geq 0$] is fulfilled identically.




Fig. 1. The best-fit, the upper and lower 1σ , 2σ and 3σ limits of $q(z_*)$ estimates, obtained with the *gold* [panel (a)] and the *combined* [panel (b)] samples, for 200 equally spaced redshifts. The **NEC** and **SEC** lower bounds, and also the **DEC** upper bound are shown. This figure shows that the **SEC** is violated with 1σ confidence level from $\simeq 0$ until $z \simeq 0.31$ [*gold* sample, panel (a)], and until $z \simeq 0.52$ [*combined* sample, panel (b)]. It also shows that the **DEC** and **NEC** is violated within 3σ confidence level for high redshifts for both supernovae samples, and that the **NEC** is violated for $z \lesssim 0.105$ [panel (a)] and $z \lesssim 0.085$ [panel (b)].

In Fig. 1 it is showed that the **SEC** bound is violated with 1σ confidence level until $z = 0.31$ for *gold* and $z = 0.52$ for *combined* sample, while in the redshift intervals $(0.09, 0.17)$ [panel (a)] and $(0.08, 0.18)$ [panel (b)] this violation occurs with more than 3σ confidence level, where the highest evidence is found at $z = 0.135$ with $\simeq 3.86\sigma$ [*gold*, panel (a)] and $\simeq 4.28\sigma$ [*combined*, panel (b)]. Unlike the result of Ref. 10, wherein these analyses have been performed by computing the confidence regions on the $E(z_*) - q(z_*)$ plane, revealing no redshift value for the **SEC** fulfillment with at least 1σ , we note here the **SEC** is fulfilled with more than 1σ for $z \gtrsim 0.615$ [*gold*, panel (a)] and $z \gtrsim 0.855$ [*combined*, panel (b)]. According to the present **SEC** analysis, with 1σ confidence level, the universe crosses over from a decelerated expansion phase to an accelerated expansion during the redshift interval ($\simeq 0.31, \simeq 0.615$) for *gold* and ($\simeq 0.52, \simeq 0.855$) for *combined* sample.^d

^cWe note that this statistical approach has been previously used in Ref. 16 but for the SNe Ia *Union* sample.¹⁷

^dWe recall that in a similar **SEC** analysis of Ref. 16 performed by using the *Union* sample, the

Regarding the **NEC**, Fig. 1 indicates its breakdown within 3σ confidence level for low redshift, $z \lesssim 0.105$ [panel (a)] and $z \lesssim 0.085$ [panel (b)]. For higher values of redshift, **NEC** is violated within 3σ at $z \gtrsim 0.94$ for *gold* and $z \gtrsim 0.96$ for *combined* sample.

Concerning the **DEC**, Fig. 1 indicates that it is violated within 3σ for $z \gtrsim 0.795$ [*gold*, panel (a)] and $z \gtrsim 0.83$ [*combined*, panel (b)], which are intervals where the error in the estimates of $q(z)$ grow significantly, though. Finally, we note that the **DEC** violation of the present analysis is weaker than that obtained in Ref. 10 in the sense that, differently from that analysis, now the **DEC** is fulfilled with 1σ confidence level in the entire redshift interval for both samples.

Acknowledgments

This work is supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) - Brasil, under grant No. 472436/2007-4. M.P.L., S.V. and M.J.R. thank CNPq for the grants under which this work was carried out. We also thank A.F.F. Teixeira for the indication of the relevant misprints and omissions.

References

1. S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Spacetime*, (Cambridge University Press, England, 1973); S. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*, (Addison Wesley, New York, 2004); R. M. Wald, *General Relativity*, (University of Chicago Press, Chicago, 1984).
2. J. Santos, J. S. Alcaniz and M. J. Rebouças, *Phys. Rev. D* **74** (2006) 067301.
3. J. Santos, J. S. Alcaniz, N. Pires and M. J. Rebouças, *Phys. Rev. D* **75** (2007) 083523.
4. Y. Gong, A. Wang, Q. Wu and Y. Z. Zhang, *J. Cosmol. Astropart. Phys.* **08** (2007) 018.
5. Y. Gong and A. Wang, *Phys. Lett. B* **652** (2007) 63.
6. J. Santos, J. S. Alcaniz, M. J. Rebouças and N. Pires, *Phys. Rev. D* **76** (2007) 043519.
7. A. A. Sen and R. J. Scherrer, *Phys. Lett. B* **659** (2008) 457.
8. C. Cattoën and M. Visser, arXiv:0712.1619v1 [gr-qc].
9. M. Visser, *Science* **276** (1997) 88; *Phys. Rev. D* **56** (1997) 7578.
10. M. P. Lima, S. Vitenti and M. J. Rebouças, *Phys. Rev. D* **77** (2008) 083518.
11. S. E. Perez Bergliaffa, *Phys. Lett. B* **642** (2006) 311.
12. J. Santos, J. S. Alcaniz, M. J. Rebouças and F. C. Carvalho, *Phys. Rev. D* **76** (2007) 083513.
13. J. Santos, M. J. Rebouças and J.S. Alcaniz, arXiv:0807.2443v2 [astro-ph].
14. A. G. Riess *et al.*, *Astrophys. J.* **659** (2007) 98.
15. W. M. Wood-Vasey *et al.*, *Astrophys. J.* **666** (2007) 694; A. G. Riess *et al.*, *Astrophys. J.* **659** (2007) 98; P. Astier *et al.*, *Astron. Astrophys.* **447** (2006) 31; T. M. Davis *et al.*, *Astrophys. J.* **666** (2007) 716.
16. M. P. Lima, S. Vitenti and M. J. Rebouças, *Phys. Lett. B* **668** (2008) 83.
17. M. Kowalski *et al.*, arXiv:0804.4142 [astro-ph].