DOUBLE AFFINE HECKE ALGEBRAS AND BISPECTRAL QUANTUM KNIZHNIK-ZAMOLODCHIKOV EQUATIONS

MICHEL VAN MEER AND JASPER V. STOKMAN

ABSTRACT. We use the double affine Hecke algebra of type GL_N to construct an explicit consistent system of q-difference equations, which we call the bispectral quantum Knizhnik-Zamolodchikov (BqKZ) equations. BqKZ includes, besides Cherednik's quantum affine KZ equations associated to principal series representations of the underlying affine Hecke algebra, a compatible system of q-difference equations acting on the central character of the principal series representations. We construct a meromorphic self-dual solution Φ of BqKZ which, upon suitable specializations of the central character, reduces to symmetric self-dual Laurent polynomial solutions of quantum KZ equations. We give an explicit correspondence between solutions of BqKZ and solutions of a particular bispectral problem for Ruijsenaars' commuting trigonometric qdifference operators. Under this correspondence Φ becomes a self-dual Harish-Chandra series solution Φ^+ of the bispectral problem. Specializing the central character as above, we recover from Φ^+ the symmetric self-dual Macdonald polynomials.

Contents

1.	Introduction	1
2.	The double affine Hecke algebra	4
3.	The bispectral quantum KZ equations	8
4.	The explicit form of the bispectral quantum KZ equations	12
5.	Solutions of the bispectral quantum KZ equations	19
6.	The correspondence with bispectral problems	28
7.	Polynomial theory	38
8.	Appendix on holonomic systems of q -difference equations	45
References		51

1. INTRODUCTION

Let M be an N!-dimensional complex vector space and write $T = (\mathbb{C} \setminus \{0\})^N$. We derive an explicit holonomic system of q-difference equations on M-valued meromorphic functions on $T \times T$, which we call the bispectral quantum Knizhnik-Zamolodchikov (BqKZ) equations. It is related to Cherednik's [9] quantum affine KZ equations in the following way.

²⁰⁰⁰ Mathematics Subject Classification. 33D80, 33D52.

Key words and phrases. Double affine Hecke algebras, quantum Knizhnik-Zamolodchikov equations, Macdonald polynomials, bispectral problems.

Let H be the extended affine Hecke algebra of type GL_N . For fixed $\zeta \in T$ the vector space M admits an H-module structure, turning M into the principal series module $M(\zeta)$ of H with central character ζ . The BqKZ naturally splits up into two subsystems of q-difference equations. The first subsystem acts only on the first T-component of $T \times T$ and as such it realizes, for any fixed $\zeta \in T$, Cherednik's [9] quantum affine KZ equation qKZ_{\zeta} acting on $M(\zeta)$ -valued meromorphic functions on $T \times \{\zeta\}$. The second, dual subsystem is obtained from the first by replacing the role of $(t, \zeta) \in T \times T$ by (ζ^{-1}, t^{-1}) and conjugating the q-connection matrices by an explicit complex linear automorphism C_t of M. Hence, it acts only on the second T-component of $T \times T$ and it essentially realizes qKZ_{t-1} for fixed $t \in T$. In particular, this provides a quantum isomonodromic interpretation of qKZ. This should be compared with the interpretation of rational KZ equations as quantizations of Schlesinger equations, see [38] and [22].

The BqKZ is constructed using Cherednik's [9] double affine Hecke algebra \mathbb{H} of type GL_N. As a vector space \mathbb{H} is isomorphic to $\mathbb{C}[T] \otimes H \simeq \mathbb{C}[T] \otimes H_0 \otimes \mathbb{C}[T] \simeq$ $\mathbb{C}[T \times T] \otimes H_0$ with H_0 the finite Hecke algebra of type A_{N-1} . Cherednik's antialgebra involution $* \colon \mathbb{H} \to \mathbb{H}$ essentially interchanges, under the above vector space identification, the role of the two copies of $\mathbb{C}[T]$. For $w, w' \in W = S_N \ltimes \mathbb{Z}^N$ we consider the map $h \mapsto \widetilde{S}_w h \widetilde{S}_{w'}^*$ ($h \in \mathbb{H}$), where the $\widetilde{S}_w \in \mathbb{H}$ are Cherednik's nonnormalized (X-)intertwiners. Restricted to $w, w' \in \mathbb{Z}^N$, suitable renormalizations of these maps become the q-connection matrices of BqKZ, with H_0 playing the role of M. The anti-involution * of \mathbb{H} gives rise to the automorphism C_ι interchanging the qKZ subsystem of BqKZ with its dual subsystem in BqKZ.

 qKZ_{ζ} is gauge equivalent to Frenkel and Reshetikhin's [20] quantum KZ equations associated with the N-fold tensor product of the vector representation of quantum \mathfrak{sl}_N (see [9, §1.3.2]). A special case of qKZ_{ζ} was considered earlier by Smirnov [42]. Etingof and Varchenko [18] used quantum group methods to construct systems of q-difference equations (so-called dynamical q-difference equations) that are compatible with Frenkel and Reshetikhin's quantum KZ equations associated to evaluation representations of quantum affine algebras. It is likely that the system of dynamical q-difference equations associated with qKZ_{ζ} is equivalent to the dual qKZ subsystem in BqKZ.

Preceding the above mentioned work [18] of Etingof and Varchenko, dynamical equations for various degenerations of quantum KZ equations have been analyzed in detail; see, e.g., [19], [47], [44], [46], [45] and [29]. An interesting aspect in, e.g., [47] and [46], is the observation that various degenerations of quantum KZ equations are the duals of their associated dynamical equations with respect to $(\mathfrak{gl}_r, \mathfrak{gl}_s)$ duality. In the present set-up (which corresponds to r = s = N), this duality is incorporated by the automorphism C_{ι} , which reflects Cherednik's duality anti-involution of the double affine Hecke algebra \mathbb{H} on the level of BqKZ.

We investigate the space SOL of *M*-valued meromorphic solutions of BqKZ in detail. We first analyze BqKZ in a suitable asymptotic region. It leads to a solution Φ of BqKZ which is self-dual, in the sense that $\Phi(t, \gamma) = C_{\iota} \Phi(\gamma^{-1}, t^{-1})$ as *M*-valued meromorphic functions in $(t, \gamma) \in T \times T$. We construct a basis of solutions of SOL in terms of Φ , and we give an explicit formula for the leading term of $\Phi(t, \gamma)$ as function of *t*.

Cherednik [6, Thm. 3.4] constructed for arbitrary root systems a correspondence between solutions of quantum KZ equations and solutions of a system of q-difference equations. In [6, Thm. 4.4], Cherednik made the correspondence precise for GL_N . It yields an explicit map χ_+ from solutions of qKZ_{ζ} to solutions of the spectral problem of Ruijsenaars' [39] commuting trigonometric *q*-difference operators with spectral parameter ζ^{-1} (the Ruijsenaars operators are also frequently referred to as Macdonald-Ruijsenaars operators). The latter result has been generalized to arbitrary root systems in [26, Thm. 4.6] and [7]. We analyze the map χ_+ in the present bispectral setting. It leads to the interpretation of χ_+ as an embedding of SOL into the space of meromorphic solutions of a bispectral problem involving the above Ruijsenaars operators as well as Ruijsenaars operators acting on the spectral parameter.

The application of the correspondence to the self-dual solution Φ of BqZK leads to a self-dual Harish-Chandra series solution Φ^+ of the bispectral problem. Harish-Chandra series solutions of the Ruijsenaars operators with fixed spectral parameter were investigated before in, e.g., [15], [16], [27] and [31]. The present approach to Harish-Chandra series, which uses quantum KZ equations in an essential way, has the advantage that it leads to new results on the convergence and singularities of the Harish-Chandra series. These results, together with Cherednik's recent work [10], form important building blocks in deriving the *c*-function expansion of Cherednik's global (q, t)-spherical function (this will be detailed in a forthcoming paper of the second author).

The q-connection matrices of the dual qKZ subsystem of BqKZ can be used to map solutions of qKZ_{ζ} for a fixed central character $\zeta \in T$ to solutions of qKZ_{ζ'} with respect to a suitably shifted central character ζ' . Applied to a constant solution we obtain a symmetric Laurent polynomial solution Q_{λ} for any non-increasing Ntuple λ of integers, which is automatically self-dual. We express Q_{λ} in terms of a suitable specialization of the solution Φ of BqKZ. Applying χ_{+} to Q_{λ} we obtain a symmetric self-dual Laurent polynomial eigenfunction of the Ruijsenaars operators, which is the normalized Macdonald [32] polynomial of degree λ . We show that the well-known duality and evaluation formula for the Macdonald polynomials (see [32, Chpt. VI]) are direct consequences of the properties of the Laurent polynomials Q_{λ} .

For various other approaches to the construction of solutions of quantum KZ equations see, e.g., [20], [14], [24], [27], [36], [11] and [25].

Many results in the present paper, in particular the BqZK equations themselves, can be extended to arbitrary root systems. These extensions use the root system generalizations of \mathbb{H} and their fundamental properties (see [9]). We have chosen to restrict the present exposition to the GL_N theory, not only in order to present the basic ideas while limiting the technicalities, but also because the GL_N theory deserves extra attention by its particularly strong ties with quantum groups and quantum integrable lattice models (see, e.g., [14], [24]). The theory for arbitrary root systems is detailed in a forthcoming paper of the first author.

Conventions

 $-\otimes$ always stands for tensor product over \mathbb{C} and $\operatorname{End}(M)$, for a module M over \mathbb{C} , stands for \mathbb{C} -linear endomorphisms.

 $-\mathbb{N} = \{1, 2, \ldots\}.$

– For a module M over a commutative ring R and a ring extension $R \subset S$, we write $M^S = S \otimes_R M$.

Acknowledgments

Both authors are supported by the Netherlands Organization for Scientific Research (NWO) in the VIDI-project "Symmetry and modularity in exactly solvable models". We thank the referee for valuable comments.

2. The double affine Hecke Algebra

2.1. The extended affine Weyl group. Let $N \ge 2$ and let $D = D_N$ be the affine Dynkin diagram of affine type \widehat{A}_{N-1} (the cyclic graph with N vertices if $N \ge 3$). The N vertices are labeled by the numbers $0, 1, \ldots, N-1$ (anticlockwise if $N \ge 3$). We identify occasionally the set of labels by the group \mathbb{Z}_N of integers modulo N.

Write W_Q for the affine Weyl group of affine type \widehat{A}_{N-1} . In terms of its Coxeter generators s_i $(i \in \mathbb{Z}_N)$, the characterizing group relations are the quadratic relations $s_i^2 = 1$ and, if $N \geq 3$, the braid relations

(2.1)
$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \\ s_i s_j = s_j s_i, \qquad i - j \neq 0, \pm 1.$$

The subgroup generated by s_1, \ldots, s_{N-1} is isomorphic to the symmetric group S_N in N letters, where s_i is identified with the simple transposition $i \leftrightarrow i + 1$.

Let $\operatorname{Aut}(D)$ be the group of automorphisms of the affine Dynkin diagram of type \widehat{A}_{N-1} . Let $c \in \operatorname{Aut}(D)$ the element of order N, acting on the label set \mathbb{Z}_N of the vertices of D by c(i) = i + 1. We view c as automorphism of W_Q by $c(s_i) = s_{i+1}$.

Let $\Omega = \langle \pi \rangle$ be the infinite cyclic group with cyclic generator π . It acts by group automorphisms on W_Q by $\pi \mapsto c$. Accordingly we can define the semi-direct product group $W = \Omega \ltimes W_Q$, which is called the extended affine Weyl group (associated to GL_N). We denote *e* for the identity element of *W*.

Since $s_0 = \pi s_{N-1} \pi^{-1}$, the subgroups S_N and Ω already generate W as a group. Furthermore we have $W \simeq S_N \ltimes \mathbb{Z}^N$. The cyclic generator π of Ω corresponds to $\pi = \sigma \epsilon_N$, where $\{\epsilon_i\}_{i=1}^N$ denotes the standard \mathbb{Z} -basis of \mathbb{Z}^N and $\sigma = s_1 s_2 \cdots s_{N-1} \in S_N$ is the "clockwise rotation" which maps N to 1 and all other i to i+1. Conversely,

(2.2)
$$\epsilon_j = s_{j-1} \cdots s_2 s_1 \pi s_{N-1} s_{N-2} \cdots s_j$$

for j = 1, ..., N.

Remark 2.1. Under the identification $W \simeq S_N \ltimes \mathbb{Z}^N$, we have $W_Q = S_N \ltimes Q$ with $Q \subset \mathbb{Z}^N$ the sublattice of rank N-1 consisting of N-tuples of integers that sum up to zero (this is the (co)root lattice of the root system $R = \{\epsilon_i - \epsilon_j\}_{1 \le i \ne j \le N}$ of type A_{N-1}).

For $w \in W$ let $w' \in W_Q$ and $\omega \in \Omega$ be the unique group elements such that $w = w'\omega$, then we define the length $\ell(w)$ of w to be the length of $w' \in W_Q$, i.e., it is the minimal number r such that w' can be expressed as

$$w' = s_{i_1} \cdots s_{i_r}$$

for some $i_k \in \mathbb{Z}_N$ (such an expression of w', as well as the resulting expression for $w = w'\omega$, is called a reduced expression). Thus Ω consists of the elements of W of length zero.

A central role in this paper is played by an action of the extended affine Weyl group W by q-difference reflection operators on suitable function spaces on $T := (\mathbb{C}^{\times})^N$, where $\mathbb{C}^{\times} := \mathbb{C} \setminus \{0\}$. Here q is taken to be real and strictly between zero and one (with minor technical adjustments the condition on q may be relaxed to 0 < |q| < 1, and a parallel theory can be developed for |q| > 1). Since q is fixed once and for all, we will in general suppress the dependence on q in notations. We start with an action of W on T by

(2.3)
$$wt = (t_{w^{-1}(1)}, \dots, t_{w^{-1}(N)}), \qquad w \in S_N,$$
$$\lambda t = (q^{\lambda_1} t_1, \dots, q^{\lambda_N} t_N), \qquad \lambda = (\lambda_1, \dots, \lambda_N) \in \mathbb{Z}^N,$$

for $t = (t_1, \ldots, t_N) \in T$. It is convenient to introduce $\kappa^{\lambda} := (\kappa^{\lambda_1}, \ldots, \kappa^{\lambda_N})$ for $\kappa \in \mathbb{C}^{\times}$ and $\lambda \in \mathbb{Z}^N$, so that the action of $\lambda \in \mathbb{Z}^N$ on $t \in T$ can simply be written as

$$\lambda t = q^{\lambda} t$$

in standard vector notation. Note that the action of $\pi \in \Omega$ is given by

$$\pi(t_1,\ldots,t_N)=(qt_N,t_1,\ldots,t_{N-1}).$$

Consider the algebra $\mathbb{C}[T] = \mathbb{C}[x_1^{\pm 1}, \ldots, x_N^{\pm 1}]$ of complex-valued, regular functions on T, where $x_i(t) := t_i$ for $t = (t_1, \ldots, t_N) \in T$ are the standard coordinate functions. We write

$$x^{\lambda} := x_1^{\lambda_1} \cdots x_N^{\lambda_N} \in \mathbb{C}[T]$$

for $\lambda = (\lambda_1, \ldots, \lambda_N) \in \mathbb{Z}^N$, which form the monomial basis of $\mathbb{C}[T]$.

Let $\mathbb{C}(T)$ be the field of rational functions on T, $\mathcal{O}(T)$ be the ring of analytic functions on T, and $\mathcal{M}(T)$ be the field of meromorphic functions on T. Note that $\mathcal{M}(T)$ is the quotient field of $\mathcal{O}(T)$ (cf. [23, Thm. 7.4.6]). The W-action on Tgives rise to a left W-action by algebra automorphisms on $\mathbb{C}[T]$, $\mathbb{C}(T)$, $\mathcal{O}(T)$ and $\mathcal{M}(T)$, via

$$(wf)(t) = f(w^{-1}t)$$

for $w \in W$, $t \in T$. We can, in particular, form the smashed product algebra $\mathbb{C}(T) \# W$. Recall that if G is a group and A is a G-algebra over \mathbb{C} (that is, a unital associative algebra over \mathbb{C} endowed with a left G-action by algebra automorphisms), then the smashed product algebra A # G is the unique complex unital associative algebra such that

- (i) $A \# G = A \otimes \mathbb{C}[G]$ as a complex vector space;
- (ii) the canonical linear embeddings $A \hookrightarrow A \# G$, $\mathbb{C}[G] \hookrightarrow A \# G$ are algebra homomorphisms; and
- (iii) the cross relations

$$(a \otimes g) \cdot (b \otimes h) = ag(b) \otimes gh,$$

are satisfied for $a, b \in A$ and $g, h \in G$.

We will always write $ag := a \otimes g \in A \# G$ $(a \in A, g \in G)$. Observe that A # G canonically acts on any G-algebra B containing A as a G-subalgebra.

Note that the smashed product algebra $\mathbb{C}(T) \# W$ depends on q, since the W-action on $\mathbb{C}(T)$ depends on q (see (2.3)). Sometimes it is convenient to emphasize its q-dependence, in which case we write $\mathbb{C}(T) \#_q W$ instead of $\mathbb{C}(T) \# W$.

The canonical left $\mathbb{C}(T) \# W$ -action on $\mathbb{C}(T)$ (and $\mathcal{M}(T)$) is faithful and realizes $\mathbb{C}(T) \# W$ as the algebra of q-difference S_N -reflection operators with coefficients in $\mathbb{C}(T)$. If $f \in \mathbb{C}(T)$ then we write f(X) for the associated element in $\mathbb{C}(T) \# W$ (it is the operator defined as multiplication by f). In particular, X_i is multiplication by the coordinate function x_i .

2.2. The extended affine Hecke algebra and Cherednik's basic representation. In this subsection we recall some constructions and results due to Cherednik (see, e.g., [9, Chpt. 1] and references therein).

Fix a nonzero complex number k.

Definition 2.2. The affine Hecke algebra $H_Q = H_Q(k)$ is the complex, unital, associative algebra generated by T_i $(i \in \mathbb{Z}_N)$ and satisfying

(1) if $N \geq 3$, the braid relations

$$\begin{split} T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, \\ T_i T_j &= T_j T_i, \qquad i-j \neq 0, \pm 1; \end{split}$$

(2) the quadratic relations $(T_i - k)(T_i + k^{-1}) = 0$.

The Dynkin diagram automorphism $c \in \operatorname{Aut}(D)$ can also be viewed as automorphism of H_Q by $c(T_i) = T_{i+1}$. Accordingly, Ω acts by algebra automorphisms on H_Q by $\pi \mapsto c$. The extended affine Hecke algebra H = H(k) is the associated smashed product algebra $H_Q \# \Omega$.

For a reduced expression $w = s_{i_1} \cdots s_{i_r} \omega \in W$ $(i_k \in \mathbb{Z}_N, \omega \in \Omega)$, the element

$$T_w := T_{i_1} \cdots T_{i_r} \omega \in H$$

is well defined. The T_w ($w \in W$) form a linear basis of H. For k = 1, the extended affine Hecke algebra H is isomorphic to the group algebra $\mathbb{C}[W]$ of W via the identification $T_w \leftrightarrow w$ ($w \in W$).

The finite Hecke algebra is the subalgebra H_0 of H generated by T_1, \ldots, T_{N-1} . The elements T_w ($w \in S_N$) form a linear basis of H_0 . Note that H is already generated as algebra by H_0 and $\pi^{\pm 1}$, since $T_0 = \pi T_{N-1}\pi^{-1}$.

Put

(2.4)
$$Y_i := T_{i-1}^{-1} \cdots T_2^{-1} T_1^{-1} \pi T_{N-1} T_{N-2} \cdots T_i \in H$$

for i = 1, ..., N. Note that Y_i becomes the translation element ϵ_i in W if k = 1. We furthermore write $Y^{\lambda} := Y_1^{\lambda_1} \cdots Y_N^{\lambda_N}$ for $\lambda = (\lambda_1, ..., \lambda_N) \in \mathbb{Z}^N$. We have the following characterization of H, due to Bernstein. For details we refer to Lusztig [30] or Macdonald [34, §4.2].

Theorem 2.3. *H* is the unique unital complex associative algebra, such that

- (i) H₀ ⊗ ℂ[T] ≃ H as complex vector spaces, via h ⊗ f ↦ hf(Y) for h ∈ H₀, f ∈ ℂ[T], where f(Y) = ∑_λ c_λY^λ if f = ∑_λ c_λx^λ ∈ ℂ[T];
 (ii) the canonical maps H₀, ℂ[T] ↔ H are algebra embeddings; we write ℂ_Y[T] =
- (ii) the canonical maps $H_0, \mathbb{C}[T] \hookrightarrow H$ are algebra embeddings; we write $\mathbb{C}_Y[T] = \operatorname{span}_{\mathbb{C}} \{Y^{\lambda}\}_{\lambda \in \mathbb{Z}^N}$ for the image of $\mathbb{C}[T]$ in H; and
- (iii) the following cross relations

$$\begin{split} T_i^{-1} Y_i T_i^{-1} &= Y_{i+1}, \\ Y_j T_i &= T_i Y_j, \qquad i \neq j-1, j \end{split}$$

are satisfied for $1 \leq i < N$ and $1 \leq j \leq N$.

Cherednik realized the affine Hecke algebra H inside the algebra $\mathbb{C}(T) \# W$ of q-difference reflection operators as follows.

Theorem 2.4. There is a unique injective algebra homomorphism $\rho = \rho_{k,q} \colon H(k) \to \mathbb{C}(T) \#_q W$ satisfying

$$\rho(T_i) = k + c_k (X_i / X_{i+1}) (s_i - 1),
\rho(\pi) = \pi,$$

for i = 1, ..., N - 1, where

(2.5)
$$c_k(z) := \frac{k^{-1} - kz}{1 - z}.$$

Note that for the affine Hecke algebra H = H(k) with fixed parameter k, Theorem 2.4 yields a one-parameter family of realizations of H (the additional parameter being q).

Remark 2.5. The image $\rho(H)$ preserves $\mathbb{C}[T]$, viewed as a subspace of the canonical $\mathbb{C}(T) \# W$ -module $\mathbb{C}(T)$. The resulting representation of H on $\mathbb{C}[T]$ is faithful and is called the basic representation of H.

We frequently identify H with its image under ρ in $\mathbb{C}(T) \# W$.

We now come to the definition of Cherednik's double affine Hecke algebra which depends, besides on k, on the additional parameter q.

Definition 2.6. The double affine Hecke algebra $\mathbb{H} = \mathbb{H}(k,q)$ is the subalgebra of $\mathbb{C}(T) \#_q W$ generated by $\rho_{k,q}(H)$ and by the multiplication operators f(X) $(f \in \mathbb{C}[T])$.

Let $\mathbb{L} = \mathbb{C}[T] \otimes \mathbb{C}[T] \simeq \mathbb{C}[T \times T]$ denote the complex-valued regular functions on $T \times T$. We view \mathbb{H} as \mathbb{L} -module by

(2.6)
$$(f \otimes g) \cdot h := f(X)hg(Y)$$

for $f, g \in \mathbb{C}[T]$ and $h \in \mathbb{H}$. The following theorem is the so-called Poincaré-Birkhoff-Witt (PBW) property of the double affine Hecke algebra.

Theorem 2.7. We have $\mathbb{H} \simeq H_0^{\mathbb{L}} = \mathbb{L} \otimes H_0$ as \mathbb{L} -modules.

The PBW property is an essential ingredient in deriving the characterizing relations for the double affine Hecke algebra \mathbb{H} in terms of its algebraic generators T_i $(1 \leq i < N)$, $\pi^{\pm 1}$ and $X_j^{\pm 1}$ $(1 \leq j \leq N)$. Since we are not going to use this presentation explicitly in this paper, we refer the reader to [9] for further details. We use though one of its direct consequences, namely the existence of the duality anti-isomorphism (see Cherednik [9, Thm. 1.4.8]):

Theorem 2.8. There exists a unique \mathbb{C} -linear anti-algebra involution $*: \mathbb{H} \to \mathbb{H}$ determined by

$$T_w^* = T_{w^{-1}}, \qquad w \in S_N,$$

$$(Y^{\lambda})^* = X^{-\lambda}, \qquad \lambda \in \mathbb{Z}^N,$$

$$(X^{\lambda})^* = Y^{-\lambda}, \qquad \lambda \in \mathbb{Z}^N.$$

2.3. Intertwiners. In this subsection we recall the construction of the (nonnormalized) affine intertwiners associated to the double affine Hecke algebra \mathbb{H} . The intertwiners play an important role in the construction of a nontrivial $W \times W$ -cocycle in the next section. Consider the elements

$$\widetilde{S}_{i} = (k - k^{-1} X_{i+1} / X_{i}) s_{i}, \qquad 1 \le i < N,$$

$$\widetilde{S}_{0} = (k - k^{-1} q^{-1} X_{1} / X_{N}) s_{0},$$

$$\widetilde{S}_{\pi} = \pi$$

in $\mathbb{C}(T) \#_q W$. The following facts are well known (cf., e.g., [9, §1.3]). For the convenience of the reader, we give a short sketch of the proof.

Proposition 2.9. Let $w \in W$ and let $w = s_{j_1} \cdots s_{j_r} \pi^m$ be a reduced expression $(j_l \in \mathbb{Z}_N, m \in \mathbb{Z}).$

(i) S_w := S_{j1} ··· S_{jr} S_π^m is a well-defined element of C(T)#W;
(ii) S_w ∈ H;
(iii) the S_i (i ∈ Z_N) satisfy the Â_{N-1}-type braid relations;
(iv) S_wf(X) = (wf)(X)S_w in C(T)#W for all f ∈ C(T); and
(v) S_iS_i = (k - k⁻¹X_{i+1}/X_i)(k - k⁻¹X_i/X_{i+1}) for 1 ≤ i < N.

Proof. (i) Set $d_i := (k - k^{-1}X_{i+1}/X_i)$ $(1 \le i < N)$ and $d_0 := (k - k^{-1}q^{-1}X_1/X_N)$. We have

$$S_w = d_{j_1}(s_{j_1}d_{j_2})\cdots(s_{j_1}\cdots s_{j_{r-1}}d_{j_r})w$$

in $\mathbb{C}(T) \# W$. By, e.g., Macdonald [34, (2.2.9)], we know that

(2.7)
$$d_w := d_{j_1}(s_{j_1}d_{j_2})\cdots(s_{j_1}\cdots s_{j_{r-1}}d_{j_r})$$

is independent of the reduced expression of w. Hence $\widetilde{S}_w \in \mathbb{C}(T) \# W$ is well defined. (ii) Note that \widetilde{S}_i can be written as

(2.8)
$$\widetilde{S}_i = (1 - X_{i+1}/X_i)(T_i - k) + k - k^{-1}X_{i+1}/X_i$$

for $1 \leq i < N$, which shows that it lies in \mathbb{H} . Furthermore, $\pi^{\pm 1} \in \mathbb{H}$, hence $\widetilde{S}_{\pi}^{\pm 1} \in \mathbb{H}$ and $\widetilde{S}_0 = \pi \widetilde{S}_{N-1} \pi^{-1} \in \mathbb{H}$. Consequently, $\widetilde{S}_w \in \mathbb{H} \subset \mathbb{C}(T) \# W$.

(iii) is immediate from (i), while (iv) and (v) are clear from the definition of the \tilde{S}_i and \tilde{S}_{π} .

Definition 2.10. The elements \widetilde{S}_w ($w \in W$) are called the affine intertwiners of \mathbb{H} .

3. The bispectral quantum KZ equations

3.1. Construction of the cocycle. Let ι denote the nontrivial element of the two group \mathbb{Z}_2 . We define the group \mathbb{W} as the semidirect product

$$\mathbb{W} := \mathbb{Z}_2 \ltimes (W \times W),$$

where $\iota \in \mathbb{Z}_2$ acts on $W \times W$ by switching the components: $\iota(w, w') = (w', w)\iota$ for $w, w' \in W$. We first use the double affine Hecke algebra, its affine intertwiners, and its duality anti-isomorphism to construct a group homomorphism $\tau = \tau_k : \mathbb{W} \to \operatorname{GL}_{\mathbb{C}}(H_0^{\mathbb{K}})$ depending on the Hecke algebra parameter k, where $\mathbb{K} := \mathcal{M}(T \times T)$ is the field of meromorphic functions on $T \times T$.

The representation τ will be constructed from the complex linear endomorphisms $\widetilde{\sigma}_{(w,w')}$ $(w,w' \in W)$ and $\widetilde{\sigma}_{\iota}$ of the double affine Hecke algebra \mathbb{H} , defined by

$$\widetilde{\sigma}_{(w,w')}(h) = \widetilde{S}_w h \widetilde{S}_{w'}^*$$
$$\widetilde{\sigma}_\iota(h) = h^*$$

for $h \in \mathbb{H}$. In the following lemma we collect some elementary properties of the maps $\widetilde{\sigma}_{(w,w')}$ and $\widetilde{\sigma}_{\iota}$. First we introduce some auxiliary notations.

For a regular function $q \in \mathbb{C}[T]$, we write $q(x) \in \mathbb{C}[T \times T]$ (respectively $q(y) \in$ $\mathbb{C}[T \times T]$ for the corresponding regular function on $T \times T$ constant with respect to the second (respectively first) T-component. In particular, the x_i (respectively y_i) are the standard coordinate functions of the first (respectively second) copy of T in $T \times T$. Recall the regular function $d_w \in \mathbb{C}[T]$ $(w \in W)$ such that

$$S_w = d_w(X)w$$

in $\mathbb{C}(T) \# W$; see (2.7).

Lemma 3.1. The complex linear endomorphisms $\widetilde{\sigma}_{(w,w')}$ and $\widetilde{\sigma}_{\iota}$ of \mathbb{H} satisfy the following properties:

- (i) the $\tilde{\sigma}_{(s_i,e)}$ $(i \in \mathbb{Z}_N)$ satisfy the \hat{A}_{N-1} braid relations;

- (ii) $\widetilde{\sigma}_{(s_i,e)}^2 = d_{s_i}(x)(s_i d_{s_i})(x) \cdot \text{id for } i \in \mathbb{Z}_N;$ (iii) $\widetilde{\sigma}_{(\pi,e)} \widetilde{\sigma}_{(s_i,e)} \widetilde{\sigma}_{(\pi^{-1},e)} = \widetilde{\sigma}_{(s_{i+1},e)} \text{ for } i \in \mathbb{Z}_N;$ (iv) $\widetilde{\sigma}_{\iota}^2 = \text{id and } \widetilde{\sigma}_{(e,w)} = \widetilde{\sigma}_{\iota} \widetilde{\sigma}_{(w,e)} \widetilde{\sigma}_{\iota} \text{ for } w \in W; \text{ and}$
- (v) $\widetilde{\sigma}_{(w,e)}\widetilde{\sigma}_{(e,w')} = \widetilde{\sigma}_{(w,w')} = \widetilde{\sigma}_{(e,w')}\widetilde{\sigma}_{(w,e)}$ for $w, w' \in W$.

Proof. These are direct consequences of Proposition 2.9 and Theorem 2.8.

To construct a W-action from the maps $\widetilde{\sigma}_{(w,w')}$ and $\widetilde{\sigma}_{\iota}$, we need to renormalize the maps appropriately. To do so, we describe as a first step the behavior of the maps $\widetilde{\sigma}_{(w,w')}$ and $\widetilde{\sigma}_{\iota}$ with respect to the L-module structure (2.6) on H. This will allow us to extend the maps $\widetilde{\sigma}_{(w,w')}$ and $\widetilde{\sigma}_{\iota}$ to endomorphisms of $H_0^{\mathbb{K}} \simeq \mathbb{K} \otimes_{\mathbb{L}} \mathbb{H}$, which is a suitably flexible surrounding for the normalizations of the maps to take place in.

Consider the group involution $\diamond : W \to W$ defined by $w^{\diamond} = w$ for $w \in S_N$ and $\lambda^{\diamondsuit} = -\lambda$ for $\lambda \in \mathbb{Z}^N$. Then \mathbb{W} acts on $T \times T$ by

$$(w, w')(t, \gamma) = (wt, w'^{\diamond} \gamma),$$
$$\iota(t, \gamma) = (\gamma^{-1}, t^{-1})$$

for $w, w' \in W$, where $t^{-1} := (t_1^{-1}, \dots, t_N^{-1}) \in T$ and the action of W on T is by q-dilations and permutations; see (2.3). By transposition, this defines an action of \mathbb{W} on $\mathbb{K} = \mathcal{M}(T \times T)$ by field automorphisms,

(3.1)
$$(\mathbf{w}f)(t,\gamma) = f(\mathbf{w}^{-1}(t,\gamma)), \qquad \mathbf{w} \in \mathbb{W}.$$

Note that $\mathbb{L} = \mathbb{C}[T \times T]$ is a W-subalgebra of K.

Lemma 3.2. For $h \in \mathbb{H}$ and $f \in \mathbb{L}$ we have

(3.2)
$$\widetilde{\sigma}_{(w,w')}(f \cdot h) = ((w,w')f) \cdot \widetilde{\sigma}_{(w,w')}(h),$$
$$\widetilde{\sigma}_{\iota}(f \cdot h) = (\iota f) \cdot \widetilde{\sigma}_{\iota}(h)$$

for $w, w' \in W$.

Proof. From Proposition 2.9 we know that $\widetilde{S}_w p(X) = (wp)(X)\widetilde{S}_w$ in \mathbb{H} for $p \in \mathbb{C}[T]$ and $w \in W$. For $p \in \mathbb{C}[T]$ let $p^{\diamondsuit} \in \mathbb{C}[T]$ be defined by $p^{\diamondsuit}(t) = p(t^{-1})$, then we also have

$$p(Y)\widetilde{S}_{w'}^* = (\widetilde{S}_{w'}p^{\diamondsuit}(X))^* = ((w'p^{\diamondsuit})(X)\widetilde{S}_{w'})^* = \widetilde{S}_{w'}^*(w'p^{\diamondsuit})^{\diamondsuit}(Y)$$

in \mathbb{H} . Hence for $p, r \in \mathbb{C}[T]$,

$$\widetilde{\sigma}_{(w,w')}(p(X)hr(Y)) = (wp)(X)\widetilde{S}_wh\widetilde{S}_{w'}^*(w'r^{\diamondsuit})^{\diamondsuit}(Y).$$

The first formula of (3.2) now follows since $(w'r^{\diamond})^{\diamond} = w'^{\diamond}r$. The second is immediate from the definition of the duality anti-involution.

As a direct consequence of Lemma 3.2 the maps $\tilde{\sigma}_{(w,w')}$ $(w,w' \in W)$ and $\tilde{\sigma}_{\iota}$ uniquely extend to complex linear endomorphisms of $H_0^{\mathbb{K}} \simeq \mathbb{K} \otimes_{\mathbb{L}} \mathbb{H}$ such that (3.2) is valid for all $f \in \mathbb{K}$ and $h \in H_0^{\mathbb{K}}$. We keep the same notations $\tilde{\sigma}_{(w,w')}$ and $\tilde{\sigma}_{\iota}$ for these maps. Note that the properties of $\tilde{\sigma}_{(w,w')}$ and $\tilde{\sigma}_{\iota}$ as described in Lemma 3.1 also hold true as identities between endomorphisms of $H_0^{\mathbb{K}}$.

Theorem 3.3. There is a unique group homomorphism

$$\tau \colon \mathbb{W} \to \mathrm{GL}_{\mathbb{C}}(H_0^{\mathbb{K}})$$

satisfying

(3.3)
$$\tau(w,w')(f) = d_w(x)^{-1} d_{w'}^{\diamond}(y)^{-1} \cdot \widetilde{\sigma}_{(w,w')}(f),$$
$$\tau(\iota)(f) = \widetilde{\sigma}_{\iota}(f)$$

for $w, w' \in W$ and $f \in H_0^{\mathbb{K}}$. It satisfies $\tau(w)(g \cdot f) = wg \cdot \tau(w)(f)$ for $g \in \mathbb{K}$, $f \in H_0^{\mathbb{K}}$ and $w \in \mathbb{W}$.

Proof. The last statement is clear.

The action τ of $W \times \{e\}$ arises naturally from left multiplication by normalized affine intertwiners on a suitable localization of the double affine Hecke algebra (see Cherednik [9, §1.3]). In the present set-up, one observes that Lemma 3.2 and Lemma 3.1(i)-(ii) imply that the $\tau(s_i, e)$ $(i \in \mathbb{Z}_N)$ satisfy the \widehat{A}_{N-1} braid relations and the quadratic relations $\tau(s_i, e)^2 = \operatorname{id}_{H_0^{\mathbb{K}}}$. Since furthermore $\tau(\pi, e)$ is a complex linear automorphism of $H_0^{\mathbb{K}}$ with inverse $\tau(\pi^{-1}, e)$, and $\tau(\pi, e)\tau(s_i, e)\tau(\pi^{-1}, e) =$ $\tau(s_{i+1}, e)$ for $i \in \mathbb{Z}_N$ by Lemma 3.2 and Lemma 3.1(iii), we conclude that the formulas (3.3) for the maps $\tau(s_i, e)$ $(i \in \mathbb{Z}_N)$ and $\tau(\pi, e)$ uniquely extend to a group homomorphism $\tau : W \times \{e\} \to \operatorname{GL}_{\mathbb{C}}(H_0^{\mathbb{K}})$. It follows from Proposition 2.9 and its proof that the resulting group homomorphism satisfies

$$\sigma(w,e)f = d_w(x)^{-1} \cdot \widetilde{\sigma}_{(w,e)}f$$

for $w \in W$. This is in accordance with formula (3.3).

Combining Lemma 3.1(iv) with Lemma 3.2 we can relate the complex endomorphism $\tau(e, w)$ (see (3.3)) of $H_0^{\mathbb{K}}$ to $\tau(w, e)$ by the formula

(3.4)
$$\tau(e,w) = \tau(\iota)\tau(w,e)\tau(\iota), \qquad w \in W,$$

where $\tau(\iota)$ is given by the second formula of (3.3). Since $\tau(\iota)^2 = \tilde{\sigma}_{\iota}^2 = \operatorname{id}_{H_0^{\mathbb{K}}}$ we conclude that $W \ni w \mapsto \tau(e, w)$ (see (3.3)) defines a left W-action on $H_0^{\mathbb{K}}$.

By Lemma 3.1 and Lemma 3.2 (v) we have

$$\tau(w, e)\tau(e, w') = \tau(w, w') = \tau(e, w')\tau(w, e)$$

for all $w, w' \in W$. Thus $\tau \colon W \times W \to \operatorname{GL}_{\mathbb{C}}(H_0^{\mathbb{K}})$, defined by the first formula of (3.3), is a group homomorphism. Combined with (3.4) and $\tau(\iota)^2 = \operatorname{id}_{H_0^{\mathbb{K}}}$ we conclude that τ (3.3) indeed defines a complex linear action of \mathbb{W} on $H_0^{\mathbb{K}}$. \Box

For $w \in W$ and $f \in H_0^{\mathbb{K}} = \mathbb{K} \otimes H_0$ we write wf for the action of w on the \mathbb{K} -coefficients of f in its expansion along a basis of H_0 . In other words, viewing $f(t,\gamma)$ as H_0 -valued meromorphic function in $(t,\gamma) \in T \times T$, the action is given by $(wf)(t,\gamma) = f(w^{-1}(t,\gamma))$. Consider $\operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$ as a \mathbb{W} -group by the corresponding conjugation action

(3.5)
$$(\mathbf{w}, A) \mapsto \mathbf{w}A\mathbf{w}^{-1}, \qquad \mathbf{w} \in \mathbb{W}, \ A \in \mathrm{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$$

by group automorphisms. We have the following direct consequence of the previous theorem.

Corollary 3.4. The map $w \mapsto C_w := \tau(w)w^{-1}$ is a cocycle of \mathbb{W} with values in the \mathbb{W} -group $\operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$. In other words, $C_w \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$ and

$$C_{\mathbf{w}\mathbf{w}'} = C_{\mathbf{w}}\mathbf{w}C_{\mathbf{w}'}\mathbf{w}^{-1}$$

for all $w, w' \in W$.

For more details on non-abelian group cohomology, see, e.g., the appendix in [43].

Remark 3.5. Interpreting $A \in \operatorname{End}_{\mathbb{K}}(H_0^{\mathbb{K}})$ as $\operatorname{End}(H_0)$ -valued meromorphic function $A(t,\gamma)$ in $(t,\gamma) \in T \times T$, the action (3.5) becomes $(wA)(t,\gamma) = A(w^{-1}(t,\gamma))$. In particular, $wAw^{-1} = A$ for all $w \in \mathbb{W}$ if $A \in \operatorname{End}_{\mathbb{K}}(H_0^{\mathbb{K}})$ is the \mathbb{K} -linear extension of a complex linear endomorphism of H_0 . This is, for instance, the case for the cocycle value C_{ι} (see Subsection 4.2).

Remark 3.6. One may replace in this subsection \mathbb{K} by the field $\mathbb{C}(T \times T)$ of rational functions on $T \times T$. Consequently, the cocycle value $C_{w}(t, \gamma)$ for $w \in \mathbb{W}$ is a rational $\operatorname{End}(H_{0})$ -valued function in $(t, \gamma) \in T \times T$. We presented the results with respect to $\mathbb{K} = \mathcal{M}(T \times T)$ since this is the natural setting for the applications of the cocycle C in the analytic theory of the quantum KZ equations (to which we come at a later stage).

3.2. **Bispectral quantum KZ equations.** In this subsection, we use the cocycle $C_{\mathbf{w}} \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$ ($\mathbf{w} \in \mathbb{W}$) to define a holonomic system of *q*-difference equations on the space $H_0^{\mathbb{K}}$ of H_0 -valued meromorphic functions on $T \times T$.

The constructions thus far have led to a \mathbb{C} -linear action τ of \mathbb{W} on $H_0^{\mathbb{K}}$. In terms of the cocycle C_{w} (w $\in \mathbb{W}$), it is given by

(3.6)
$$(\tau(\mathbf{w})f)(t,\gamma) = C_{\mathbf{w}}(t,\gamma)f(\mathbf{w}^{-1}(t,\gamma))$$

for $\mathbf{w} \in \mathbb{W}$ and $f \in H_0^{\mathbb{K}}$, where (3.6) should be read as identities between H_0 -valued meromorphic functions in $(t, \gamma) \in T \times T$. It follows that $f \in H_0^{\mathbb{K}}$ is $\tau(\mathbb{Z}^N \times \mathbb{Z}^N)$ -invariant if and only if

(3.7)
$$C_{(\lambda,\mu)}(t,\gamma)f(q^{-\lambda}t,q^{\mu}\gamma) = f(t,\gamma) \quad \forall \lambda,\mu \in \mathbb{Z}^N$$

viewed as identities between H_0 -valued meromorphic functions on $T \times T$.

Definition 3.7. We call the q-difference equations (3.7) the bispectral quantum KZ (BqKZ) equations. We write SOL for the set of functions $f \in H_0^{\mathbb{K}}$ satisfying the BqKZ equations (3.7).

Let $\mathbb{F} \subset \mathbb{K}$ denote the subfield consisting of $f \in \mathbb{K}$ satisfying $(\lambda, \mu)f = f$ for all $\lambda, \mu \in \mathbb{Z}^N$. Let furthermore \mathbb{S}_N denote the subgroup $\mathbb{Z}_2 \ltimes (S_N \times S_N)$ of \mathbb{W} .

Corollary 3.8. (i) The BqKZ equations (3.7) form a holonomic system of qdifference equations. In other words, the connection matrices $C_{(\lambda,\mu)}$ $(\lambda, \mu \in \mathbb{Z}^N)$ satisfy the compatibility conditions

(3.8)
$$C_{(\lambda,\mu)}(t,\gamma)C_{(\nu,\xi)}(q^{-\lambda}t,q^{\mu}\gamma) = C_{(\nu,\xi)}(t,\gamma)C_{(\lambda,\mu)}(q^{-\nu}t,q^{\xi}\gamma)$$

for $\lambda, \mu, \nu, \xi \in \mathbb{Z}^N$, as End(H_0)-valued meromorphic functions in $(t, \gamma) \in T \times T$. (ii) The solution space SOL of BqKZ is a $\tau(\mathbb{S}_N)$ -invariant \mathbb{F} -subspace of $H_0^{\mathbb{K}}$.

Proof. (i) The cocycle condition implies that both sides of (3.8) are equal to $C_{(\lambda+\nu,\mu+\xi)}(t,\gamma)$.

(ii) Clearly, SOL is a \mathbb{F} -subspace of $H_0^{\mathbb{K}}$. Note, furthermore, that $\mathbb{Z}^N \times \mathbb{Z}^N$ is a normal subgroup of \mathbb{W} with quotient group isomorphic to \mathbb{S}_N . Hence the \mathbb{F} subspace SOL of $\tau(\mathbb{Z}^N \times \mathbb{Z}^N)$ -invariant elements in the $\tau(\mathbb{W})$ -module $H_0^{\mathbb{K}}$ is $\tau(\mathbb{S}_N)$ invariant.

4. The explicit form of the bispectral quantum KZ equations

In this section we derive explicit expressions for the cocycle values $C_{\mathbf{w}}$ ($\mathbf{w} \in \mathbb{W}$) and, in particular, for the *q*-connection matrices $C_{(\lambda,\mu)}$ ($\lambda, \mu \in \mathbb{Z}^N$) of the BqKZ. It will become apparent that the $C_{(\lambda,e)}(\cdot,\zeta)$ ($\lambda \in \mathbb{Z}^N$) with $\zeta \in T$ fixed coincide with the *q*-connection matrices of Cherednik's quantum affine KZ equation associated to the principal series module of H(k) with central character ζ . They also turn up as gauged *q*-connection matrices for a Frenkel-Reshetikhin [20] type quantum KZ equation associated to the quantum affine algebra $\mathcal{U}_k(\widehat{sl}_N)$.

4.1. Formal principal series. View the commutative subalgebra $\mathbb{C}_Y[T]$ of H as left $\mathbb{C}_Y[T]$ -module by left multiplication. Let $M = \operatorname{Ind}_{\mathbb{C}_Y[T]}^H(\mathbb{C}_Y[T])$ be the corresponding induced left H-module. With respect to the $\mathbb{C}[T] \simeq \mathbb{C}[\{1\} \times T]$ -module structure

$$f \cdot (h \otimes_{\mathbb{C}_Y[T]} g(Y)) = h \otimes_{\mathbb{C}_Y[T]} (fg)(Y) \qquad f, g \in \mathbb{C}[T], \ h \in H$$

on M we have $M \simeq H_0^{\mathbb{C}[\{1\} \times T]} = \mathbb{C}[\{1\} \times T] \otimes H_0$ as $\mathbb{C}[\{1\} \times T]$ -modules. The left H-action on M is $\mathbb{C}[\{1\} \times T]$ -linear, hence we obtain an algebra homomorphism

$$\eta: H \to \operatorname{End}_{\mathbb{C}[\{1\} \times T]}(H_0^{\mathbb{C}[\{1\} \times T]}).$$

We occasionally view $\eta(h)$ as $\operatorname{End}(H_0)$ -valued regular function in $\gamma \in T$, in which case we write it as $T \ni \gamma \mapsto \eta(h)(\gamma)$. Extending the ground ring $\mathbb{C}[\{1\} \times T]$ to $\mathbb{K} = \mathcal{M}(T \times T)$ we obtain an algebra homomorphism

$$H \to \operatorname{End}_{\mathbb{K}}(H_0^{\mathbb{K}}),$$

which we shall also denote by η . From this viewpoint, $\eta(h)(\gamma)$ is the regular $\operatorname{End}(H_0)$ -valued function in $(t, \gamma) \in T \times T$ which is constant in t. Note that $\eta(h)$ for $h \in H_0$ is constant as $\operatorname{End}(H_0)$ -valued function on $T \times T$ (see Remark 3.5).

Lemma 4.1. For $w \in S_N$ and $1 \le i < N$ we have

(4.1)
$$\eta(T_i)T_w = \begin{cases} T_{s_iw} & \text{if } \ell(s_iw) = \ell(w) + 1, \\ (k - k^{-1})T_w + T_{s_iw} & \text{if } \ell(s_iw) = \ell(w) - 1, \end{cases}$$

and

(4.2)
$$\eta(\pi)(\gamma)T_w = \gamma_{w^{-1}(N)}T_{\sigma w}$$

as regular H_0 -valued functions in $\gamma \in T$.

Proof. The first formula follows directly from the definitions. For the second formula it suffices to verify that $\pi T_w = T_{\sigma w} Y_{w^{-1}(N)}$ in H.

If w = 1 then $\pi = T_{\sigma}Y_N$ since $\sigma = s_1s_2\cdots s_{N-1}$ is a reduced expression. If $w = s_{N-1}$ then

$$\pi T_w = \pi T_{N-1} = T_1 \cdots T_{N-2} Y_{N-1} = T_{s_1 \cdots s_{N-2}} Y_{w^{-1}(N)} = T_{\sigma w} Y_{w^{-1}(N)}.$$

Next, we prove that $\pi T_w = T_{\sigma w} Y_{w^{-1}(N)}$ in H if $w \neq 1$ and $\ell(s_i w) = \ell(w) + 1$ for all $1 \leq i \leq N-2$. Then $w = s_{N-1}s_{N-2}\cdots s_j$ for some $1 \leq j < N$ (and this is a reduced expression of w). We find, making repetitive use of the cross relation $T_r Y_{r+1} T_r = Y_r$ $(1 \leq r < N)$ in H,

$$\pi T_w = \pi T_{N-1} T_{N-2} \cdots T_j = T_1 \cdots T_{N-2} Y_{N-1} T_{N-2} \cdots T_j$$

= $T_1 \cdots T_{N-3} Y_{N-2} T_{N-3} \cdots T_j$
:
= $T_1 \cdots T_{j-1} Y_j = T_{s_1 \cdots s_{j-1}} Y_j$
= $T_{\sigma w} Y_j = T_{\sigma w} Y_{w^{-1}(N)},$

which is the desired relation in H.

The general case is now proved by induction on $\ell(w)$. Let $w \neq 1$ and decompose it as $w = s_i u$ with $1 \leq i < N$ and $u \in S_N$ such that $\ell(s_i u) = \ell(u) + 1$. Suppose that $\pi T_u = T_{\sigma u} Y_{u^{-1}(N)}$ in H. In order to prove that $\pi T_w = T_{\sigma w} Y_{w^{-1}(N)}$ we may, in view of the previous paragraph, assume without loss of generality that $1 \leq i \leq N - 2$. Then $s_i(N) = N$ and $\ell(s_{i+1}\sigma u) = \ell(\sigma u) + 1$. (The latter equality is equivalent to $(\sigma u)^{-1}(i+1) < (\sigma u)^{-1}(i+2)$, which follows from $u^{-1}(i) < u^{-1}(i+1)$, which again is equivalent to the assumption $\ell(s_i u) = \ell(u) + 1$.) Then

$$\pi T_w = \pi T_i T_u = T_{i+1} \pi T_u = T_{i+1} T_{\sigma u} Y_{u^{-1}(N)}$$

= $T_{s_{i+1}\sigma u} Y_{u^{-1}s_i(N)} = T_{\sigma s_i u} Y_{w^{-1}(N)}$
= $T_{\sigma w} Y_{w^{-1}(N)}$

in H, which completes the proof.

In view of the explicit expression (2.8) for the intertwiner $\widetilde{S}_i \in \mathbb{H}$ $(1 \leq i < N)$ and the definition of the duality anti-involution, we have $\widetilde{S}_w^* \in H$ for all $w \in S_N$. We now set

$$\xi_w := \eta(\widetilde{S}_{w^{-1}}^*) T_e \in H_0^{\mathbb{K}}, \qquad w \in S_N.$$

Note that $\xi_w \in H_0^{\mathbb{C}[\{1\} \times T]} \subset H_0^{\mathbb{K}}$ for $w \in S_N$. We view ξ_w as regular H_0 -valued function in $\gamma \in T$, as well as meromorphic H_0 -valued function in $(t, \gamma) \in T \times T$ constant in $t \in T$.

Lemma 4.2. $\{\xi_w\}_{w\in S_N}$ is a \mathbb{K} -basis of $H_0^{\mathbb{K}}$ consisting of common eigenfunctions for the η -action of $\mathbb{C}_Y[T]$ on $H_0^{\mathbb{K}}$. For $p \in \mathbb{C}[T]$ and $w \in S_N$ we have

(4.3)
$$\eta(p(Y))(\gamma)\xi_w(\gamma) = (w^{-1}p)(\gamma)\xi_w(\gamma)$$

as H_0 -valued regular functions in $\gamma \in T$.

Proof. For $p \in \mathbb{C}[T]$ we have $\eta(p(Y))(\gamma)T_e = p(\gamma)T_e$. Note, furthermore, that $p(Y)\widetilde{S}_{w^{-1}}^* = \widetilde{S}_{w^{-1}}^*(w^{-1}p)(Y)$ in H for $p \in \mathbb{C}[T]$ and $w \in S_N$; see the proof of Lemma 3.2. Combining the two observations gives (4.3). It follows from Proposition 2.9(iv)-(v) that the ξ_w ($w \in S_N$) are nonzero in $H_0^{\mathbb{K}}$. The eigenvalue equations (4.3) then show that the ξ_w ($w \in S_N$) are \mathbb{K} -linearly independent in $H_0^{\mathbb{K}}$.

4.2. The cocycle values. We define

$$R_i(z) = c_k(z)^{-1}(\eta(T_i) - k) + 1, \qquad 1 \le i < N,$$

viewed as a rational $\operatorname{End}(H_0)$ -valued function in z. The results of the previous subsection implies that the $R_i(z)$ satisfy the following Yang-Baxter type equations (see Cherednik [9, §1.3.2]).

Lemma 4.3. We have

$$C_{(s_i,e)}(t,\gamma) = R_i(t_i/t_{i+1}), \qquad 1 \le i < N,$$

as rational $\operatorname{End}(H_0)$ -valued functions in $(t, \gamma) \in T \times T$. In particular, the $R_i(z)$ satisfy

(4.4)
$$\begin{aligned} R_i(z)R_i(z^{-1}) &= \mathrm{id}, \\ R_j(z)R_{j+1}(zz')R_j(z') &= R_{j+1}(z')R_j(zz')R_{j+1}(z), \end{aligned}$$

for $1 \le i < N$ and $1 \le j < N - 1$ as $End(H_0)$ -valued rational functions.

Proof. For $1 \leq i < N$ and $h \in H_0$ we have, as H_0 -valued meromorphic functions in $(t, \gamma) \in T \times T$,

$$C_{(s_i,e)}(t,\gamma)h = (\tau(s_i,e)h)(t,\gamma)$$

= $d_{s_i}(t)^{-1}(\tilde{S}_ih)(t,\gamma)$
= $c_k(t_i/t_{i+1})^{-1}(\eta(T_i) - k)h + h$

in view of the explicit expression (2.8) for \tilde{S}_i , c_k (2.5) and d_{s_i} (see the proof of Lemma 2.9). For the second statement of the lemma, note that the cocycle property of C implies for $1 \leq i < N$ and $1 \leq j < N - 1$ that

$$\begin{split} C_{(s_i,e)}(t,\gamma) C_{(s_i,e)}(s_i t,\gamma) &= \mathrm{id}, \\ C_{(s_j,e)}(t,\gamma) C_{(s_{j+1},e)}(s_j t,\gamma) C_{(s_j,e)}(s_{j+1} s_j t,\gamma) \\ &= C_{(s_{j+1},e)}(t,\gamma) C_{(s_j,e)}(s_{j+1} t,\gamma) C_{(s_{j+1},e)}(s_j s_{j+1} t,\gamma), \end{split}$$

as rational End(H_0)-valued functions in $(t, \gamma) \in T \times T$. Using $C_{(s_i, e)}(t, \gamma) = R_i(t_i/t_{i+1})$ these formulas imply (4.4).

Observe that C_{ι} is the K-linear extension of the anti-algebra involution of H_0 mapping T_w to $T_{w^{-1}}$ for all $w \in S_N$. Note furthermore that

Together with the explicit description of $C_{(s_i,e)}$ $(1 \leq i < N)$ from the previous lemma, these formulas determine the values C_{w} ($w \in W$) uniquely (cf. Corollary 3.4). In particular, the cocycle property of C implies that

$$C_{(e,w)}(t,\gamma) = C_{\iota}C_{(w,e)}(\gamma^{-1},t^{-1})C_{\iota}, \qquad w \in W,$$

as $\operatorname{End}(H_0)$ -valued rational functions in $(t, \gamma) \in T \times T$.

Lemma 4.4. Let $w \in W$.

(i) $C_{(w,e)} \in (\mathbb{C}(T) \otimes \mathbb{C}[T]) \otimes \operatorname{End}(H_0)$. (ii) $The \mathbb{C}[T] \otimes \operatorname{End}(H_0)$ -valued rational function $t \mapsto C_{(w,e)}(t, \cdot)$ in $t \in T$ is regular at $t \in T \setminus S$, where

(4.6)
$$\mathcal{S} = \{ t \in T \mid t^{\alpha} \in k^{-2}q^{\mathbb{Z}} \text{ for some } \alpha \in R \}.$$

Proof. By the cocycle condition, $C_{(w,e)}(t,\gamma)$ can be written as a product of factors $C_{(s_i,e)}(ut,\gamma)$ $(1 \le i < N, u \in W)$ and $\eta(\pi^{\pm 1})(\gamma)$. By Lemma 4.1 and the fact that $R_i(z)$ has a single pole at $z = k^{-2}$, we conclude (i) and (ii).

4.3. The q-connection matrices. Besides the standard \mathbb{Z} -basis $\{\epsilon_i\}_{i=1}^N$ of \mathbb{Z}^N , we also have the \mathbb{Z} -basis $\{\varpi_i\}_{i=1}^N$ consisting of the fundamental weights $\varpi_i := \sum_{i=1}^i \epsilon_j \in \mathbb{Z}^N$. Note that

(4.7)
$$\varpi_i = \pi^i \sigma^{-i}$$

in W for all $1 \leq i \leq N$. In the following lemma, we compute the q-connection matrices $C_{(\lambda,e)}$ for $\lambda \in \mathbb{Z}^N$ of BqKZ explicitly in case λ is one of these two types of basis elements of \mathbb{Z}^N .

Lemma 4.5. (i) For $1 \le j \le N$ we have

$$C_{(\epsilon_j,e)}(t,\gamma) = R_{j-1}(t_{j-1}/t_j)R_{j-2}(t_{j-2}/t_j)\cdots R_1(t_1/t_j)$$

$$\times \eta(\pi)(\gamma)R_{N-1}(qt_N/t_j)\cdots R_{j+1}(qt_{j+2}/t_j)R_j(qt_{j+1}/t_j)$$

as rational $\operatorname{End}(H_0)$ -valued functions in $(t, \gamma) \in T \times T$. (ii) For $1 \leq i < N$ we have

$$C_{(\varpi_i,e)}(t,\gamma) = (\eta(\pi)(\gamma))^i (R_{N-i}(qt_N/t_1) \cdots R_2(qt_{i+2}/t_1)R_1(qt_{i+1}/t_1))$$

$$\times \cdots \times (R_{N-2}(qt_N/t_{i-1}) \cdots R_i(qt_{i+2}/t_{i-1})R_{i-1}(qt_{i+1}/t_{i-1}))$$

$$\times (R_{N-1}(qt_N/t_i) \cdots R_{i+1}(qt_{i+2}/t_i)R_i(qt_{i+1}/t_i))$$

as rational $\operatorname{End}(H_0)$ -valued functions in $(t, \gamma) \in T \times T$. (iii) We have

$$C_{(\varpi_N,e)}(t,\gamma) = \gamma^{\varpi_N} \mathrm{id}$$

as rational $\operatorname{End}(H_0)$ -valued functions in $(t, \gamma) \in T \times T$.

Proof. (i) By the cocycle property of C and by the expression (2.2) for $\epsilon_j \in W$, we obtain an explicit expression for $C_{(\epsilon_j,e)}$ in terms of the $C_{(s_i,e)}$ $(1 \le i < N)$ and $C_{(\pi,e)}$. Combining (4.5) and the previous lemma then gives the desired expression for $C_{(\epsilon_j,e)}(t,\gamma)$.

(ii) σ^i is the permutation

so we find a reduced expression

(4.8) $\sigma^{i} = (s_{i} \cdots s_{N-1})(s_{i-1} \cdots s_{N-2}) \cdots (s_{2} \cdots s_{N-i+1})(s_{1} \cdots s_{N-i}).$

Combined with (4.7) we get a reduced expression for ϖ_i . Using the cocycle condition for C_w repeatedly, we get the desired result.

(iii) Since $\sigma^N = 1$, we get $C_{(\varpi_N, e)}(t, \gamma) = (\eta(\pi)(\gamma))^N$, which maps T_w to $\gamma^{\varpi_N} T_w$ for all $w \in S_N$ in view of Lemma 4.1.

We end this subsection by computing the asymptotic leading terms of the *q*-connection matrices $C_{(\lambda,e)}(t,\gamma)$ ($\lambda \in \mathbb{Z}^N$) as $|t^{-\alpha_i}| \to 0$ ($1 \le i < N$), where we take $\alpha_i := \epsilon_i - \epsilon_{i+1}$ ($1 \le i < N$) as a base of the root system $R = \{\epsilon_i - \epsilon_j\}_{1 \le i \ne j \le N}$ of type A_{N-1} . Let $R_+ = \{\epsilon_i - \epsilon_j\}_{1 \le i < j \le N}$ denote the associated set of positive roots and $Q_+ = \bigoplus_{i=1}^{N-1} \mathbb{Z}_{\ge 0} \alpha_i$ the corresponding cone in the root lattice Q of R. Let furthermore $\delta := (N - 1, N - 3, \dots, 1 - N) \in \mathbb{Z}^N$ and write $w_0 \in S_N$ for the longest Weyl group element (mapping i to N - i + 1 for $1 \le i \le N$).

Consider the subring $\mathcal{A} := \mathbb{C}[x^{-\alpha_1}, \ldots, x^{-\alpha_{N-1}}]$ of $\mathbb{C}[T \times \{1\}] = \mathbb{C}[x_1^{\pm 1}, \ldots, x_N^{\pm 1}] \subset \mathbb{C}[T \times T]$. We write $Q(\mathcal{A})$ for its quotient field and $Q_0(\mathcal{A})$ for the subring of $Q(\mathcal{A})$ consisting of rational functions which are analytic at the point $x^{-\alpha_i} = 0$ $(1 \leq i < N)$. We consider $Q_0(\mathcal{A}) \otimes \mathbb{C}[T]$ as a subring of $\mathbb{C}(T \times T)$ in the natural way. The first part of the following corollary is a refinement of Lemma 4.4(i) in case $w \in \mathbb{Z}^N$.

Corollary 4.6. Let $\lambda \in \mathbb{Z}^N$. We have

(4.9)
$$C_{(\lambda,e)} \in (Q_0(\mathcal{A}) \otimes \mathbb{C}[T]) \otimes \operatorname{End}(H_0)).$$

Writing

$$C_{(\lambda,e)}^{(0)} = C_{(\lambda,e)}\big|_{x^{-\alpha_1}=0,\dots,x^{-\alpha_{N-1}}=0} \in \mathbb{C}[T] \otimes \operatorname{End}(H_0),$$

we have $C_{(\lambda,e)}^{(0)} = k^{\langle \delta,\lambda \rangle} \eta(T_{w_0} Y^{w_0(\lambda)} T_{w_0}^{-1})$, where $\langle \cdot, \cdot \rangle$ is the standard scalar product on \mathbb{R}^N .

Proof. To prove (4.9) it suffices, in view of the cocyle property of C, to verify (4.9) for $\lambda = \epsilon_i$. The statement then follows from Lemma 4.5(i), Lemma 4.1 and the explicit expression of $R_i(z)$.

Observe that $\lim_{z\to 0} R_i(z) = k\eta(T_i^{-1})$ for $1 \le i < N$. Combined with Lemma 4.5(i) and the explicit expression for Y_j (see (2.4)) we obtain for $1 \le j \le N$,

$$C_{(\epsilon_j,e)}(t,\gamma) \to k^{2j-N-1}\eta(Y_j)(\gamma)$$

as $|t^{\alpha_i}| \to 0$ for all $1 \le i < N$, hence

$$C_{(\lambda,e)}(t,\gamma) \to k^{-\langle \delta,\lambda \rangle} \eta(Y^{\lambda})(\gamma)$$

as $|t^{\alpha_i}| \to 0$ for all $1 \le i < N$. To derive the asymptotics of $C_{(\lambda,e)}(t,\gamma)$ as $|t^{-\alpha_i}| \to 0$ for $1 \le i < N$ we use the cocycle property to write

$$C_{(\lambda,e)}(t,\gamma) = C_{(w_0,e)}(t,\gamma)C_{(w_0(\lambda),e)}(w_0t,\gamma)C_{(w_0,e)}(q^{-w_0(\lambda)}w_0t,\gamma)$$

Note that $C_{(w_0,e)}(t,\gamma) \to k^{\ell(w_0)}\eta(T_{w_0}^{-1})$ if $|t^{\alpha_i}| \to 0$ for all $1 \le i < N$. Hence

$$C_{(\lambda,e)}^{(0)} = k^{\langle \delta,\lambda \rangle} \eta(T_{w_0} Y^{w_0(\lambda)} T_{w_0}^{-1})$$

as desired.

4.4. Relation to quantum KZ equations. Fix $\zeta \in T$ and let $\chi_{\zeta} : \mathbb{C}[\{1\} \times T] \to \mathbb{C}$ denote the corresponding evaluation character $\chi_{\zeta}(f) = f(\zeta)$. Recall the formal principal series $\eta : H \to \operatorname{End}_{\mathbb{C}[\{1\} \times T]}(M)$. The corresponding complex *H*representation $M(\zeta) = \mathbb{C} \otimes_{\chi_{\zeta}} M$ of dimension N! is the principal series module of *H* with central character ζ . We identify $M(\zeta)$ with H_0 as a complex vector space, and push the *H*-action on $M(\zeta)$ through the linear isomorphism to H_0 . We denote the corresponding representation map by

$$\eta_{\zeta}: H \to \operatorname{End}(H_0).$$

As in Subsection 4.1 we have as identities in H_0 ,

$$\eta_{\zeta}(T_i)T_w = \begin{cases} T_{s_iw} & \text{if } \ell(s_iw) = \ell(w) + 1, \\ (k - k^{-1})T_w + T_{s_iw} & \text{if } \ell(s_iw) = \ell(w) - 1, \end{cases}$$

for $1 \leq i < N$ and $w \in S_N$,

$$\eta_{\zeta}(\pi)T_w = \zeta_{w^{-1}(N)}T_{\sigma w},$$

for $w \in S_N$, as well as

$$\eta_{\zeta}(f(Y))\xi_w(\zeta) = (w^{-1}f)(\zeta)\xi_w(\zeta),$$

for $w \in S_N$, where $\xi_w(\zeta) \in H_0$ is the regular H_0 -valued function $\xi_w(\gamma)$ in $\gamma \in T$ specialized at $\gamma = \zeta$. Extending the base field to $\mathcal{M}(T)$ we get an algebra homomorphism $H \to \operatorname{End}_{\mathcal{M}(T)}(H_0^{\mathcal{M}(T)})$, which is also denoted by η_{ζ} .

In this subsection we consider the BqKZ for specialized values of γ . In view of Lemma 4.4(i) we may specialize $C_{(w,e)}(t,\gamma)$ ($w \in W$) at $\gamma = \zeta$. We write

$$C_w^{\zeta}(t) := C_{(w,e)}(t,\zeta), \qquad w \in W,$$

for the resulting specialized cocycle values, viewed as $\operatorname{End}(H_0)$ -valued rational functions in $t \in T$. For $\zeta \in T$ the map $W \ni w \mapsto C_w^{\zeta}$ defines a cocycle of W with values in the W-group $\operatorname{GL}_{\mathbb{C}(T)}(H_0^{\mathbb{C}(T)})$. In other words,

$$C_{ww'}^{\zeta}(t) = C_w^{\zeta}(t)C_{w'}^{\zeta}(w^{-1}t), \qquad w, w' \in W,$$

as rational H_0 -valued functions in $t \in T$. Comparing the cocycle values C_{λ}^{ζ} ($\lambda \in \mathbb{Z}^N$) to the ones in [9, §1.3] we obtain the following result.

Corollary 4.7. Fix $\zeta \in T$. The holonomic system of q-difference equations

(4.10)
$$C_{\lambda}^{\zeta}(t)f(q^{-\lambda}t) = f(t), \quad \forall \lambda \in \mathbb{Z}^{N}$$

for $f \in H_0^{\mathcal{M}(T)}$ is Cherednik's quantum affine KZ equation associated to the principal H-module $M(\zeta)$ with central character ζ .

Let $\operatorname{SOL}_{\zeta} \subset H_0^{\mathcal{M}(T)}$ denote the set of solutions of the quantum KZ equations (4.10). Write $\mathcal{E}(T) \subset \mathcal{M}(T)$ for the subfield of meromorphic functions f satisfying $f(q^{\lambda}t) = f(t)$ for all $\lambda \in \mathbb{Z}^N$ as meromorphic functions in $t \in T$. The set $\operatorname{SOL}_{\zeta}$ of solutions is a $\mathcal{E}(T)$ -subspace of $H_0^{\mathcal{M}(T)}$. Furthermore, $\operatorname{SOL}_{\zeta}$ is invariant for the S_N -action

(4.11)
$$(\varsigma(w)f)(t) := C_w^{\zeta}(t)f(w^{-1}t), \quad w \in S_N$$

on $H_0^{\mathcal{M}(T)}$ (note that ς does not depend on ζ since $C_w^{\zeta}(t) = C_{(w,e)}(t,\zeta)$ is independent of ζ for $w \in S_N$).

Remark 4.8. The quantum KZ equations (4.10) are gauge equivalent to Frenkel and Reshetikhin's [20] quantum KZ equations associated with the N-fold tensor product representation $\mathbb{C}^N(t_1) \otimes \cdots \otimes \mathbb{C}^N(t_N)$ of the quantum affine algebra $\mathcal{U}_k(\widehat{sl}_N)$, where $\mathbb{C}^N(t_i)$ is the evaluation representation of the vector representation \mathbb{C}^N of $\mathcal{U}_k(sl_N)$ (see [9, §1.3.2] and [14] for the details). In view of Corollary 4.7 the BqKZ equations (3.7) are a holonomic extension of the quantum KZ equations (4.10) by q-difference equations in the central character ζ of $M(\zeta)$. These may be thought of as analogs of isomonodromy transformations; in fact, in view of Lemma 4.2 and Corollary 4.6 the q-difference equations in ζ (which are essentially the quantum KZ equations again!) are reminiscent of Schlesinger transformations. This should be compared with the quantum isomonodromic interpretation of (rational) KZ equations as quantizations of Schlesinger equations, see [38] and [22].

From a different perspective we may think of the cocycle values $C_{(e,w)}$ $(w \in W)$ as shift operators, in the sense that they map solutions of quantum KZ equations to solutions of quantum KZ equations with respect to shifted central characters. To formulate the precise result, we view in the following proposition $\gamma \mapsto C_{(e,w)}(\cdot, \gamma)$ as $\mathbb{C}[T] \otimes \operatorname{End}(H_0)$ -valued rational function in $\gamma \in T$.

Proposition 4.9. Let $w \in W$ and $\zeta \in T$ such that $\gamma \mapsto C_{(e,w)}(\cdot, \gamma)$ is regular at $\gamma = \zeta$. Then $f \mapsto C_{(e,w)}(\cdot, \zeta)f$ defines an S_N -equivariant linear map $\operatorname{SOL}_{w^{\Diamond -1}\zeta} \to \operatorname{SOL}_{\zeta}$.

Proof. By the cocycle property we have for $f \in \text{SOL}_{w^{\Diamond -1}\zeta}$ and $\lambda \in \mathbb{Z}^N$,

$$C_{\lambda}^{\zeta}(t) \left(C_{(e,w)}(q^{-\lambda}t,\zeta) f(q^{-\lambda}t) \right) = C_{(\lambda,w)}(t,\zeta) f(q^{-\lambda}t)$$
$$= C_{(e,w)}(t,\zeta) C_{\lambda}^{w^{\Diamond - 1}\zeta}(t) f(q^{-\lambda}t)$$
$$= C_{(e,w)}(t,\zeta) f(t).$$

Hence $C_{(e,w)}(\cdot,\zeta)f \in \text{SOL}_{\zeta}$. The S_N -equivariance of the map is again a consequence of the cocycle property of C_w ($w \in W$); indeed, for $v \in S_N$ and $f \in \text{SOL}_{w^{\Diamond -1}\zeta}$ we have

$$C_{v}^{\zeta}(t) \big(C_{(e,w)}(v^{-1}t,\zeta)f(v^{-1}t) \big) = C_{(v,w)}(t,\zeta)f(v^{-1}t) = C_{(e,w)}(t,\zeta) \big(C_{v}^{w^{\diamond -1}\zeta}(t)f(v^{-1}t) \big),$$

which is the desired result.

From the quantum group perspective (see Remark 4.8), Proposition 4.9 resembles the action of the dynamical Weyl group on solutions of quantum KZ equations from [18]. We expect that the second half of the BqKZ is closely related to the Varchenko-Etingof dynamical difference equations [18, §9]; see also [19], [47], [44], [46], [45] and [29] for detailed studies of various degenerate cases. An interesting aspect, e.g., in [47] and [46], is the observation that KZ equations are dual to the associated dynamical equations using $(\mathfrak{gl}_r, \mathfrak{gl}_s)$ duality (our set-up relates to r = s = N). In the present theory this duality is incorporated by the cocycle value C_ι , which relates the *q*-connection matrices $C_{(\lambda,e)}$ ($\lambda \in \mathbb{Z}^N$) of the quantum KZ equation to the dual *q*-connection matrices $C_{(e,\lambda)}$ by conjugation,

$$C_{(e,\lambda)}(t,\gamma) = C_{\iota}C_{(\lambda,e)}(\gamma^{-1},t^{-1})C_{\iota}$$

as End (H_0) -valued meromorphic functions in $(t, \gamma) \in T \times T$. In turn, C_{ι} is a direct reflection of (the existence of) Cherednik's duality anti-isomorphism of the double affine Hecke algebra (see Theorem 2.8).

5. Solutions of the bispectral quantum KZ equations

In this section we use asymptotic analysis to construct a ι -invariant solution Φ_{κ} of BqKZ, which we call the basic asymptotically free solution. It depends in a mild way on an auxiliary parameter $\kappa \in \mathbb{C}^{\times}$ (in fact, $\mathbb{F}\Phi_{\kappa}$ is independent of κ). The orbit of Φ_{κ} under the action of $\{e\} \times S_N \subset \mathbb{S}_N$ turns out to be an \mathbb{F} -basis of SOL consisting of asymptotically free solutions. Along the way we derive various additional properties of Φ_{κ} .

5.1. The leading term. Let $\theta \in \mathcal{M}(T)$ denote the renormalized Jacobi theta function

(5.1)
$$\theta(z) := \prod_{m \ge 0} (1 - q^m z)(1 - q^{m+1}/z)$$

for $z \in \mathbb{C}^{\times}$. It satisfies

(5.2)
$$\theta(q^m z) = (-z)^{-m} q^{-\frac{1}{2}m(m-1)} \theta(z), \qquad m \in \mathbb{Z}$$

For $\kappa \in \mathbb{C}^{\times}$ we define $W_{\kappa} \in \mathbb{K}$ by

(5.3)
$$W_{\kappa}(t,\gamma) := \prod_{i=1}^{N} \frac{\theta(\kappa t_i \gamma_{N-i+1}^{-1})}{\theta(\kappa k^{\langle \delta, \epsilon_i \rangle} t_i) \theta(\kappa k^{-\langle \delta, \epsilon_i \rangle} \gamma_{N-i+1}^{-1})}$$

By Corollary 4.6, the formal asymptotic form of the quantum KZ equations

$$\mathcal{L}_{(\lambda,e)}(t,\gamma)f(q^{-\lambda}t,\gamma) = f(t,\gamma), \qquad \lambda \in \mathbb{Z}^N$$

in the asymptotic region $|t^{\alpha_i}| \gg 0$ $(1 \le i < N)$ is

C

(5.4)
$$k^{\langle \delta, \lambda \rangle} \eta(T_{w_0} Y^{w_0(\lambda)} T_{w_0}^{-1})(\gamma) f(q^{-\lambda} t, \gamma) = f(t, \gamma), \qquad \lambda \in \mathbb{Z}^N.$$

Lemma 5.1. $W_{\kappa} \in \mathbb{K}$ enjoys the following properties.

- (i) f_κ⁽⁰⁾(t, γ) := W_κ(t, γ)T_{w0} is a solution of (5.4).
 (ii) ι(W_κ) = W_κ and τ(ι)f_κ⁽⁰⁾ = f_κ⁽⁰⁾.

Proof. (i) Since $\eta(T_{w_0}Y^{w_0(\lambda)}T_{w_0}^{-1})(\gamma)T_{w_0} = \gamma^{w_0(\lambda)}T_{w_0}$ for all $\lambda \in \mathbb{Z}^N$, it suffices to show that

$$W_{\kappa}(q^{-\lambda}t,\gamma) = k^{-\langle \delta,\lambda\rangle} \gamma^{-w_0(\lambda)} W_{\kappa}(t,\gamma), \qquad \lambda \in \mathbb{Z}^N,$$

which follows from (5.2).

(ii) Clearly $\iota(W_{\kappa}) = W_{\kappa}$, i.e. $W_{\kappa}(\gamma^{-1}, t^{-1}) = W_{\kappa}(t, \gamma)$. Since $C_{\iota}(T_{w_0}) = T_{w_0}$, it follows that $\tau(\iota)f_{\kappa}^{(0)} = f_{\kappa}^{(0)}$.

Observe that, more generally, $W_{\kappa} \in \mathbb{K}$ satisfies the q-difference equations

(5.5)
$$W_{\kappa}(q^{-\lambda}t,q^{\mu}\gamma) = k^{-\langle\delta,\lambda+\mu\rangle} t^{w_0(\mu)} \gamma^{-w_0(\lambda)} q^{-\langle w_0(\lambda),\mu\rangle} W_{\kappa}(t,\gamma), \qquad \lambda,\mu\in\mathbb{Z}^N.$$

5.2. The basic asymptotically free solution Φ_{κ} . We now gauge BqKZ by $W_{\kappa} \in$ K. Concretely, for $\lambda, \mu \in \mathbb{Z}^N$ we write

$$D_{(\lambda,\mu)}(t,\gamma) = W_{\kappa}(t,\gamma)^{-1}C_{(\lambda,\mu)}(t,\gamma)W_{\kappa}(q^{-\lambda}t,q^{\mu}\gamma)$$

as $\operatorname{End}(H_0)$ -valued meromorphic functions in $(t, \gamma) \in T \times T$. It is independent of κ in view of (5.5). For $f \in H_0^{\mathbb{K}}$ we have $f \in \text{SOL}$ if and only if $g := W_{\kappa}^{-1} f \in H_0^{\mathbb{K}}$ satisfies the holonomic system of q-difference equations

(5.6)
$$D_{(\lambda,\mu)}(t,\gamma)g(q^{-\lambda}t,q^{\mu}\gamma) = g(t,\gamma), \qquad \lambda,\mu \in \mathbb{Z}^N$$

as H_0 -valued rational functions in $(t, \gamma) \in T \times T$.

The existence of a solution $\Psi \in H_0^{\mathbb{K}}$ of (5.6) admitting a convergent H_0 -valued power series expansion

(5.7)
$$\Psi(t,\gamma) = \sum_{\alpha,\beta \in Q_+} K_{\alpha,\beta} t^{-\alpha} \gamma^{\beta}, \qquad K_{0,0} = T_{w_0}$$

in the asymptotic region $|t^{\alpha_i}| \gg 0$ and $|\gamma^{-\alpha_i}| \gg 0$ $(1 \le i < N)$ is guaranteed by the following properties of the gauged q-connection matrices $D_{(\lambda,\mu)}$.

Consider the subring $\mathcal{B} := \mathbb{C}[y^{\alpha_1}, \dots, y^{\alpha_{N-1}}]$ of $\mathbb{C}[\{1\} \times T] = \mathbb{C}[y_1^{\pm 1}, \dots, y_N^{\pm 1}].$ Write $Q(\mathcal{B})$ for its quotient field and $Q_0(\mathcal{B})$ for the subring of $Q(\mathcal{B})$ consisting of rational functions which are analytic at the point $y^{\alpha_j} = 0$ $(1 \le j < N)$. We consider $Q_0(\mathcal{A}) \otimes \mathcal{B}$ and $\mathcal{A} \otimes Q_0(\mathcal{B})$ as subrings of $\mathbb{C}(T \times T)$ in the natural way.

Lemma 5.2. Set $A_i = D_{(\varpi_i, e)}$ and $B_i = D_{(e, \varpi_i)}$ for $1 \le i \le N$.

- (i) $A_N = B_N = \text{id } on H_0^{\mathbb{K}}$.
- (i) A_i ∈ (Q₀(A) ⊗ B) ⊗ End(H₀) and B_j ∈ (A ⊗ Q₀(B)) ⊗ End(H₀).
 (iii) Set A_i^(0,0) ∈ End(H₀) and B_j^(0,0) ∈ End(H₀) for the value of A_i and B_j at x^{-α_r} = 0 = y^{α_s} (1 ≤ r, s < N). For w ∈ S_N we have

(5.8)
$$A_i^{(0,0)}(T_{w_0}T_w) = \begin{cases} 0 & \text{if } w^{-1}w_0(\varpi_i) \neq w_0(\varpi_i), \\ T_{w_0}T_w & \text{if } w^{-1}w_0(\varpi_i) = w_0(\varpi_i) \end{cases}$$

and

(5.9)
$$B_{i}^{(0,0)}(T_{w_{0}}T_{w}) = \begin{cases} 0 & \text{if } w(\varpi_{i}) \neq \varpi_{i}, \\ T_{w_{0}}T_{w} & \text{if } w(\varpi_{i}) = \varpi_{i}. \end{cases}$$

Proof. (i) We only give the proof of $A_N = id$. Since $\varpi_N = \pi^N$ in W, we have

$$A_N(t,\gamma) = W_\kappa(t,\gamma)^{-1} C_{(\varpi_N,e)}(t,\gamma) W_\kappa(q^{-\varpi_N}t,\gamma) = \gamma^{-\varpi_N}(\eta(\pi)(\gamma))^N = \mathrm{id}$$

where we use (5.5) and (4.5) for the second equality, and Lemma 4.1 and $\sigma^N = e$ for the third equality.

(ii) Note that

$$A_i(t,\gamma) = k^{-\langle \delta, \varpi_i \rangle} \gamma^{-w_0(\varpi_i)} C_{(\varpi_i,e)}(t,\gamma)$$

by (5.5). Since $\varpi_i = \pi^i \sigma^{-i}$ the cocycle property of C gives

$$A_i(t,\gamma) = k^{-\langle \delta, \varpi_i \rangle} \gamma^{-w_0(\varpi_i)} \big(\eta(\pi)(\gamma) \big)^i C_{(\sigma^{-i},e)}(\pi^{-i}t,\gamma)$$

It follows from the explicit expressions for the cocycle values $C_{(s_i,e)}$ $(1 \le i < N)$ that the End(H_0)-valued rational function $C_{(\sigma^{-i},e)}(\pi^{-i}t,\gamma)$ in $(t,\gamma) \in T \times T$ lies in $Q_0(\mathcal{A}) \otimes \operatorname{End}(H_0)$ (in particular, it is independent of γ). Furthermore, for $w \in S_N$

$$\gamma^{-w_0(\varpi_i)} (\eta(\pi)(\gamma))^i (T_w) = \gamma^{w^{-1}w_0(\varpi_i) - w_0(\varpi_i)} T_{\sigma^i w}$$

by Lemma 4.1, hence the End(H_0)-valued regular function $\gamma^{-w_0(\varpi_i)}(\eta(\pi)(\gamma))^i$ in $\gamma \in T$ lies in $\mathcal{B} \otimes \operatorname{End}(H_0)$. Consequently, $A_i \in (Q_0(\mathcal{A}) \otimes \mathcal{B}) \otimes \operatorname{End}(H_0)$. The statement for B_i follows from this using the cocycle property $C_{(e,\varpi_i)}(t,\gamma) =$ $C_{\iota}C_{(\varpi_i,e)}(\gamma^{-1},t^{-1})C_{\iota}.$

(iii) Recall that $\xi_w = \eta(\widetilde{S}_{w^{-1}}^*)T_e$ with \widetilde{S}_w the intertwiners of \mathbb{H} (see Proposition 2.9). By induction on $\ell(w)$, using the explicit expression (2.8) of the intertwiners \widetilde{S}_i , it follows that $\xi_w \in \mathcal{B} \otimes H_0$ and that the value of ξ_w at $y^{\alpha_i} = 0$ $(1 \le i < N)$ is $T_w \in H_0$. Set

$$A_i^{(0)} = A_i|_{x^{-\alpha_1} = 0, \dots, x^{-\alpha_{N-1}} = 0} \in \mathcal{B} \otimes \operatorname{End}(H_0).$$

By Corollary 4.6 and (5.5),

$$A_i^{(0)} = y^{-w_0(\varpi_i)} \eta(T_{w_0} Y^{w_0(\varpi_i)} T_{w_0}^{-1}).$$

Lemma 4.2 then gives

(5.10)
$$A_i^{(0)}(\eta(T_{w_0})\xi_w) = y^{w^{-1}w_0(\varpi_i) - w_0(\varpi_i)}\eta(T_{w_0})\xi_w, \quad \forall w \in S_N$$

as identities in $\mathcal{B} \otimes H_0$. Specializing (5.10) at $y^{\alpha_j} = 0$ $(1 \le j < N)$ yields (5.8). To prove (5.9) we consider

$$\widetilde{B}_j^{(0)} = B_j|_{y^{\alpha_1} = 0, \dots, y^{\alpha_{N-1}} = 0} \in \mathcal{A} \otimes \operatorname{End}(H_0).$$

It is the rational $End(H_0)$ -valued function

$$\widetilde{B}_{j}^{(0)}(t) = t^{w_{0}(\varpi_{j})} C_{\iota} \big(\eta(T_{w_{0}} Y^{w_{0}(\varpi_{j})} T_{w_{0}}^{-1})(t^{-1}) \big) C_{\iota}$$

in $t \in T$. Denoting $\tilde{\xi}_w \in \mathcal{A} \otimes H_0$ for the rational H_0 -valued function $\xi_w(t^{-1})$ in $t \in T$, it follows that

(5.11)
$$\widetilde{B}_{j}^{(0)}\left(C_{\iota}\eta(T_{w_{0}})\widetilde{\xi}_{w}\right) = x^{-w^{-1}w_{0}(\varpi_{j})+w_{0}(\varpi_{j})}C_{\iota}\eta(T_{w_{0}})\widetilde{\xi}_{w}$$

for all $w \in S_N$. The value of $C_{\iota}\eta(T_{w_0})\xi_w$ at $x^{-\alpha_i} = 0$ $(1 \leq i < N)$ is $C_{\iota}(T_{w_0}T_w)$. In addition, C_{ι} restricts to the anti-algebra involution on H_0 mapping T_w to $T_{w^{-1}}$ for $w \in S_N$, hence

$$C_{\iota}(T_{w_0}T_w) = T_{w^{-1}}T_{w_0} = T_{w_0}T_{w_0w^{-1}w_0}.$$

Formula (5.9) then follows from specializing (5.11) at $x^{-\alpha_i} = 0$ $(1 \le i < N)$ and replacing w by $w_0 w^{-1} w_0$ in the resulting formula.

For $\epsilon > 0$, put $B_{\epsilon} := \{t \in T \mid |t^{\alpha_i}| < \epsilon, \forall i\}$ and $B_{\epsilon}^{-1} := \{t \in T \mid t^{-1} \in B_{\epsilon}\}.$

Theorem 5.3. There exists a unique solution $\Psi \in H_0^{\mathbb{K}}$ of the gauged equations (5.6) satisfying, for some $\epsilon > 0$,

(i) $\Psi(t,\gamma)$ admits an H₀-valued power series expansion

(5.12)
$$\Psi(t,\gamma) = \sum_{\alpha,\beta \in Q_+} K_{\alpha,\beta} t^{-\alpha} \gamma^{\beta}, \qquad (K_{\alpha,\beta} \in H_0)$$

for $(t, \gamma) \in B_{\epsilon}^{-1} \times B_{\epsilon}$ which is normally convergent on compact of $B_{\epsilon}^{-1} \times B_{\epsilon}$. In particular, $\Psi(t, \gamma)$ is analytic at $(t, \gamma) \in B_{\epsilon}^{-1} \times B_{\epsilon}$; (ii) $K_{0,0} = T_{w_0}$.

Proof. It follows from the previous lemma that the commuting endomorphisms $A_i^{(0,0)}, B_j^{(0,0)} \in \text{End}(H_0) \ (1 \leq i, j < N)$ are semisimple. For $a, b \in \mathbb{C}^{N-1}$ set

$$H_0[(a,b)] = \{ v \in H_0 \mid A_i^{(0,0)}v = a_iv \text{ and } B_j^{(0,0)}v = b_jv \ (1 \le i, j < N) \},\$$

so that $H_0 = \bigoplus_{(a,b)\in S} H_0[(a,b)]$ with S the finite set of $(a,b) \in \mathbb{C}^{N-1} \times \mathbb{C}^{N-1}$ for which $H_0[(a,b)] \neq 0$. By the previous lemma, $(1^{N-1}, 1^{N-1}) \in S$ and we have $H_0[(1^{N-1}, 1^{N-1})] = \operatorname{span}_{\mathbb{C}} \{T_{w_0}\}$. Furthermore, $a_i, b_i \notin q^{-\mathbb{N}}$ for all $(a,b) \in S$ and i. Under these conditions, the holonomic system of q-difference equations (5.6) admits a unique solution Ψ satisfying the desired properties; see Theorem 8.6 in the appendix (to show that the gauged BqKZ falls in the class of holonomic systems of *q*-difference equations to which Theorem 8.6 applies, one should take M = 2(N-1), $q_i = q$ for $1 \le i < N$ and variables $z_i = x^{-\alpha_i}$ and $z_{N-1+j} = y^{\alpha_j}$ for $1 \le i, j < N$ in the appendix).

Remark 5.4. In a small neighborhood of a fixed $(t', \gamma') \in T \times T$, the meromorphic solution Ψ of (5.6) can be expressed in terms of the power series expansion (5.12) by the formula

$$\Psi(t,\gamma) = D_{(\lambda,\mu)}(t,\gamma)\Psi(q^{-\lambda}t,q^{\mu}\gamma)$$

= $D_{(\lambda,\mu)}(t,\gamma)\sum_{\alpha,\beta\in Q_{+}}K_{\alpha,\beta}(q^{-\lambda}t)^{-\alpha}(q^{\mu}\gamma)^{\beta},$

where $\lambda, \mu \in \mathbb{Z}^N$ are such that $(q^{-\lambda}t', q^{\mu}\gamma') \in B_{\epsilon}^{-1} \times B_{\epsilon}$.

Definition 5.5. We call $\Phi_{\kappa} := W_{\kappa} \Psi \in \text{SOL}$ the basic asymptotically free solution of BqKZ.

Note that $\Phi_{\kappa} \in \mathbb{F}^{\times} \Phi_{\kappa'}$ for $\kappa, \kappa' \in \mathbb{C}^{\times}$. The κ -flexibility will come in handy when we consider specializations of Φ_{κ} . In the following subsections, we derive various properties of the basic asymptotically free solution Φ_{κ} .

5.3. Duality.

Theorem 5.6. The basic asymptotically free solution Φ_{κ} of BqKZ is self-dual, in the sense that

$$\tau(\iota)\Phi_{\kappa} = \Phi_{\kappa}$$

Proof. SOL is \mathbb{S}_N -invariant, hence $\tau(\iota)\Phi_{\kappa} \in SOL$. In addition,

$$\tau(\iota)\Phi_{\kappa} = W_{\kappa}(\tau(\iota)\Psi)$$

because $\iota(W_{\kappa}) = W_{\kappa}$. Hence $\tau(\iota)\Psi$ is a solution of the gauged equations (5.6) having a convergent H_0 -valued power series expansion

$$(\tau(\iota)\Psi)(t,\gamma) = C_{\iota}\Psi(\gamma^{-1},t^{-1}) = \sum_{\alpha,\beta\in Q_{+}} C_{\iota}(K_{\alpha,\beta})\gamma^{\alpha}t^{-\beta}$$

for $(t, \gamma) \in B_{\epsilon}^{-1} \times B_{\epsilon}$. Since $C_{\iota}(K_{0,0}) = C_{\iota}(T_{w_0}) = T_{w_0}$, we conclude from Theorem 5.3 that $\tau(\iota)\Psi = \Psi$, hence $\tau(\iota)\Phi_{\kappa} = \Phi_{\kappa}$.

5.4. Singularities. Define

$$\Lambda = \{\lambda \in \mathbb{Z}^N \mid \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_N\} = \bigoplus_{i=1}^{N-1} \mathbb{Z}_{\ge 0} \varpi_i \oplus \mathbb{Z} \varpi_N,$$

i.e., Λ consists of the $\lambda \in \mathbb{Z}^N$ such that $\langle \lambda, \alpha \rangle \in \mathbb{Z}_{\geq 0}$ for all $\alpha \in R_+$. Set

$$\mathcal{S}_+ := \{ t \in T \mid t^\alpha \in k^{-2}q^{-\mathbb{N}} \text{ for some } \alpha \in R_+ \}.$$

Write $\Psi(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma_{\alpha}(\gamma) t^{-\alpha}$ for $(t,\gamma) \in B_{\epsilon}^{-1} \times B_{\epsilon}$, where Γ_{α} is the H_0 -valued analytic function on B_{ϵ} defined by the H_0 -valued power series

$$\Gamma_{\alpha}(\gamma) := \sum_{\beta \in Q_+} K_{\alpha,\beta} \gamma^{\beta}.$$

Lemma 5.7. The Γ_{α} ($\alpha \in Q_+$) extend uniquely to a meromorphic H_0 -valued function on T, analytic at $T \setminus S_+$, such that $\Psi(t, \gamma)$ admits an H_0 -valued power series expansion

$$\Psi(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma_{\alpha}(\gamma) t^{-\alpha}$$

for $(t, \gamma) \in B_{\epsilon}^{-1} \times T \setminus S_+$, converging normally on compact of $B_{\epsilon}^{-1} \times T \setminus S_+$.

Proof. Using Lemma 5.2 we write for $\mu \in \Lambda$,

Ð

$$D_{(e,\mu)}(t,\gamma) = \sum_{\beta \in Q_+} F^{\mu}_{\beta}(\gamma) t^{-\beta}$$

with $F^{\mu}_{\beta} \in Q_0(\mathcal{B}) \otimes \operatorname{End}(H_0)$ for all $\beta \in Q^{\vee}_+$. Note that $F^{\mu}_{\beta} \equiv 0$ for all but finitely many $\beta \in Q_+$.

We first show that $F^{\mu}_{\beta}(\gamma)$ is regular at $\gamma \in T \setminus S_+$. By (5.5) and by the cocycle property, $F^{\mu}_{\beta}(\gamma)$ is regular at $\gamma = \zeta$ if $C_{(e,\varpi_j)}(\cdot, q^{\nu}\gamma) \in \mathbb{C}[T] \otimes \operatorname{End}(H_0)$ is regular at $\gamma = \zeta$ for all $1 \leq j \leq N$ and $\nu \in \Lambda$. The latter statement follows from the fact that $R_i(z)$ has only a (simple) pole at $z = k^{-2}$ and from the explicit expression (5.13)

$$C_{(e,\varpi_j)}(t,\gamma) = C_{\iota} (\eta(\pi)(t^{-1}))^{j} (R_{N-j}(q\gamma_1/\gamma_N) \cdots R_2(q\gamma_1/\gamma_{j+2})R_1(q\gamma_1/\gamma_{j+1}))$$

$$\times \cdots \times (R_{N-2}(q\gamma_{j-1}/\gamma_N) \cdots R_j(q\gamma_{j-1}/\gamma_{j+2})R_{j-1}(q\gamma_{j-1}/\gamma_{j+1})))$$

$$\times (R_{N-1}(q\gamma_j/\gamma_N) \cdots R_{j+1}(q\gamma_j/\gamma_{j+2})R_j(q\gamma_j/\gamma_{j+1}))C_{\iota},$$

which follows from Lemma 4.5(ii) and the cocycle property of C.

Let $U \subset T \setminus S_+$ be a relatively compact open subset. Choose $\mu \in \Lambda$ such that the closure of $q^{\mu}U$ is contained in B_{ϵ} , where $q^{\mu}U := \{q^{\mu}\gamma \mid \gamma \in U\}$. As meromorphic H_0 -valued function in $(t, \gamma) \in B_{\epsilon}^{-1} \times U$, we have

$$\begin{split} f(t,\gamma) &= D_{(e,\mu)}(t,\gamma)\Psi(t,q^{\mu}\gamma) \\ &= \sum_{\alpha,\beta\in Q_{+}} F^{\mu}_{\beta}(\gamma) \big(\Gamma_{\alpha}(q^{\mu}\gamma)\big)t^{-\alpha-\beta} \\ &= \sum_{\alpha\in Q_{+}} \Big(\sum_{\beta\in Q_{+}:\alpha-\beta\in Q_{+}} F^{\mu}_{\beta}(\gamma) \big(\Gamma_{\alpha-\beta}(q^{\mu}\gamma)\big)\Big)t^{-\alpha}, \end{split}$$

with the sums converging normally on compact of $B_{\epsilon}^{-1} \times U$ (note that the sums over β are finite). It follows that Γ_{α} ($\alpha \in Q_{+}$) has a unique H_{0} -valued meromorphic extension to T which, on U, is given by

(5.14)
$$\Gamma_{\alpha}(\gamma) = \sum_{\beta \in Q_{+}: \alpha - \beta \in Q_{+}} F^{\mu}_{\beta}(\gamma) \big(\Gamma_{\alpha - \beta}(q^{\mu}\gamma) \big),$$

such that Ψ on $B_{\epsilon}^{-1} \times U$ admits the power series expansion

$$\Psi(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma_{\alpha}(\gamma) t^{-\alpha}$$

which converges normally on compact of $B_{\epsilon}^{-1} \times U$. It follows from (5.14) and the previous paragraph that Γ_{α} is analytic on $T \setminus S_+$.

The arguments from the proof of Lemma 5.7, applied to both torus variables of $\Psi(t, \gamma)$ at the same time, directly lead to the following result.

Proposition 5.8. The H_0 -valued meromorphic function $\Psi(t, \gamma)$ is analytic at $(t, \gamma) \in T \setminus S_+^{-1} \times T \setminus S_+$.

For specialized spectral parameter, we obtain the following result.

Proposition 5.9. Let $\zeta \in T \setminus S_+$.

(i) The H_0 -valued meromorphic function $\Psi(t, \gamma)$ in $(t, \gamma) \in T \times T$ can be specialized at $\gamma = \zeta$, giving rise to a meromorphic H_0 -valued function $\Psi(t, \zeta)$ in $t \in T$. It has the power series expansion

$$\Psi(t,\zeta) = \sum_{\alpha \in Q_+} \Gamma_\alpha(\zeta) t^{-\alpha}$$

for $t \in B_{\epsilon}^{-1}$, normally converging on compact of B_{ϵ}^{-1} . (ii) $\Psi(t,\zeta)$ satisfies the gauged q-difference equations

(5.15)
$$D_{(\lambda,e)}(t,\zeta)\Psi(q^{-\lambda}t,\zeta) = \Psi(t,\zeta), \quad \forall \lambda \in \mathbb{Z}^N$$

Proof. (i) Restricting to $t \in B_{\epsilon}^{-1}$ for $\epsilon > 0$ small enough, the statement is correct by Lemma 5.7. If $t' \in T$ is arbitrary then there exists a $\lambda \in \Lambda$ such that $q^{-\lambda}t' \in B_{\epsilon}^{-1}$. For $t \in T$ in a small neighborhood of t' we then have

$$\Psi(t,\gamma) = D_{(\lambda,e)}(t,\gamma)\Psi(q^{-\lambda}t,\gamma).$$

Since $D_{(\lambda,e)} \in (Q_0(\mathcal{A}) \otimes \mathcal{B}) \otimes \text{End}(H_0)$ by Lemma 5.2(ii) the statement now follows in a small open neighborhood of t'.

(ii) Specializing the gauged q-difference equations $D_{(\lambda,e)}(t,\gamma)\Psi(q^{-\lambda}t,\gamma) = \Psi(t,\gamma)$ $(\lambda \in \mathbb{Z}^N)$ to $\gamma = \zeta$ yields the desired result.

5.5. **Evaluation formula.** We write $(z;q)_{\infty} = \prod_{m=0}^{\infty} (1-q^m z)$ for the *q*-shifted factorial. Recall the power series expansion $\Psi(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma_{\alpha}(\gamma)t^{-\alpha}$ for $|t^{\alpha_i}| \gg 0$ $(1 \leq i < N)$ from Subsection 5.4. We call the following result the evaluation formula for the basic asymptotically free solution $\Phi_{\kappa} = W_{\kappa}\Psi$ of BqKZ, since it implies the celebrated evaluation formula for the Macdonald polynomials (see Subsection 7.4).

Theorem 5.10. We have

$$\Gamma_0(\gamma) = K(\gamma)T_{w_0}$$

with $K \in \mathcal{M}(T)$ explicitly given by

(5.16)
$$K(\gamma) := \prod_{1 \le i < j \le N} \frac{(q\gamma_i/\gamma_j; q)_{\infty}}{(qk^2\gamma_i/\gamma_j; q)_{\infty}}.$$

Proof. We use the notations of Lemma 5.2. Recall that Ψ satisfies the gauged q-difference equations

$$A_i(t,\gamma)\Psi(q^{-\varpi_i}t,\gamma) = \Psi(t,\gamma)$$

for $1 \le i \le N$. In view of the proof of Lemma 5.2 and Lemma 5.7, it reduces in the limit $|t^{-\alpha_i}| \to 0$ $(1 \le i < N)$ to

$$\gamma^{-w_0(\varpi_i)} \eta(T_{w_0} Y^{w_0(\varpi_i)} T_{w_0}^{-1})(\gamma) \Gamma_0(\gamma) = \Gamma_0(\gamma)$$

for $1 \leq i \leq N$, as H_0 -valued meromorphic functions in $\gamma \in T$. This forces $\Gamma_0(\gamma) = K(\gamma)\eta(T_{w_0})\xi_e(\gamma) = K(\gamma)T_{w_0}$ for some $K \in \mathcal{M}(T)$; see Lemma 4.2.

It remains to show that K is explicitly given by (5.16). Write $L(\gamma)$ for the right hand side of (5.16). Then $L \in \mathcal{M}(T)$ is characterized by the following three properties:

(i) for some $\epsilon > 0$ we have a power series expansion

$$L(\gamma) = \sum_{\alpha \in Q_+} l_\alpha \gamma^\alpha$$

for $\gamma \in B_{\epsilon}$, converging normally on compact of B_{ϵ} ; (ii) $l_0 = 1$; and

(iii) $L(\gamma)$ satisfies the q-difference equations

$$\prod_{\substack{1 \le r \le j \\ +1 \le s \le N}} \frac{1 - q\gamma_r / \gamma_s}{1 - qk^2 \gamma_r / \gamma_s} L(q^{\varpi_j} \gamma) = L(\gamma), \qquad 1 \le j \le N.$$

It thus suffices to show that $K(\gamma)$ satisfies the three properties (i)–(iii). It is clear that $K \in \mathcal{M}(T)$ satisfies (i); see Subsection 5.4. Theorem 5.3(ii) implies (ii) for K. What remains is the verification of the q-difference equations (iii) for K. Using the notations of Lemma 5.2, we write

$$B_j^{(0)} := B_j|_{x^{-\alpha_1} = 0, \dots, x^{-\alpha_{N-1}} = 0} \in Q_0(\mathcal{B}) \otimes \operatorname{End}(H_0).$$

We view $B_j^{(0)}(\gamma)$ as an End (H_0) -valued meromorphic function in $\gamma \in T$. Taking the limit $|t^{-\alpha_i}| \to 0$ $(1 \le i < N)$ in the gauged q-difference equations

$$B_j(t,\gamma)\Psi(t,q^{\varpi_j}\gamma) = \Psi(t,\gamma), \qquad 1 \le j \le N$$

and using $\Gamma_0(\gamma) = K(\gamma)T_{w_0}$ we obtain

j -

$$K(q^{\varpi_j}\gamma)B_j^{(0)}(\gamma)T_{w_0} = K(\gamma)T_{w_0}$$

for $1 \leq j \leq N$, as meromorphic H_0 -valued functions in $\gamma \in T$. Writing $B_j^{(0)}(\gamma)T_{w_0} = \sum_{w \in S_N} a_w^j(\gamma)T_w$ with $a_w^j \in \mathcal{M}(T)$ it thus suffices to show that

(5.17)
$$a_{w_0}^j(\gamma) = \prod_{\substack{1 \le r \le j\\ j+1 \le s \le N}} \frac{1 - q\gamma_r/\gamma_s}{1 - qk^2\gamma_r/\gamma_s} = k^{-\langle \delta, \varpi_j \rangle} \prod_{\substack{1 \le r \le j\\ j+1 \le s \le N}} c_k (q\gamma_r/\gamma_s)^{-1}$$

for $1 \leq j \leq N$, where the second equality follows from a direct computation using the explicit expression (2.5) of c_k .

By (5.5) we have

$$B_j(t,\gamma) = k^{-\langle \delta, \varpi_j \rangle} t^{w_0(\varpi_j)} C_{(e,\varpi_j)}(t,\gamma)$$

and $C_{(e,\varpi_j)}(t,\gamma)$ is given explicitly by (5.13). Since $R_i(z) = c_k(z)^{-1}(\eta(T_i) - k) + 1$, Lemma 4.1 and the reduced expression (4.8) for σ^i imply that

$$B_j(t,\gamma)T_{w_0} = k^{-\langle \delta, \varpi_j \rangle} t^{w_0(\varpi_j)} C_\iota(\eta(\pi)(t^{-1}))^j \Big(\sum_{w < \sigma^{-j}} b_w^j(\gamma) T_{ww_0}\Big)$$

with \leq the Bruhat order on S_N and with

$$b_{\sigma^{-j}}^{j}(\gamma) = \prod_{\substack{1 \le r \le j \\ j+1 \le s \le N}} c_k (q\gamma_r / \gamma_s)^{-1}.$$

By Lemma 4.1 we have

$$t^{w_0(\varpi_j)}C_{\iota}(\eta(\pi)(t^{-1}))^j T_{ww_0} = t^{w_0(\varpi_j) - w_0 w^{-1} w_0(\varpi_j)} T_{w_0 w^{-1} \sigma^{-j}}$$

Hence

$$B_j^{(0)}(\gamma)T_{w_0} = k^{-\langle \delta, \varpi_j \rangle} \sum_w b_w^j(\gamma)T_{w_0w^{-1}\sigma^{-j}},$$

with the sum running over $w \in S_N$ satisfying $w \leq \sigma^{-j}$ and $w(\varpi_j) = w_0(\varpi_j)$. In particular, $a_{w_0}^j(\gamma) = k^{-\langle \delta, \varpi_j \rangle} b_{\sigma^{-j}}^j(\gamma)$. This completes the proof of (5.17). \Box

5.6. Consistency of the bispectral quantum KZ equations. In this subsection, we show that BqKZ is a consistent system of q-difference equations, i.e., $\dim_{\mathbb{F}}(\text{SOL}) = \dim_{\mathbb{C}}(H_0)$, by explicitly constructing an \mathbb{F} -basis of SOL. Since the q-connection matrices $C_{(\lambda,\mu)}(t,\gamma)$ $(\lambda, \mu \in \mathbb{Z}^N)$ depend rationally on $(t,\gamma) \in T \times T$, the consistency of BqKZ follows also from the abstract arguments in [13, §5].

We start with a preliminary lemma on the cocycle values $C_{(e,w)}$ for $w \in S_N$.

Lemma 5.11. Let $w \in S_N$. We have $C_{(e,w)} \in Q_0(\mathcal{B}) \otimes \operatorname{End}(H_0)$ and

$$C_{(e,w)}^{(0)}(h) = k^{-\ell(w)} h T_{w^{-1}}, \qquad h \in H_0,$$

where

$$C_{(e,w)}^{(0)} = C_{(e,w)}|_{y^{\alpha_1}=0,\dots,y^{\alpha_{N-1}}=0} \in \text{End}(H_0)$$

Proof. Let $w = s_{i_1}s_{i_2}\cdots s_{i_r}$ be a reduced expression for $w \in S_N$ $(1 \le i_j < N)$ and write $\beta_j := s_{i_1}\cdots s_{i_{j-1}}(\alpha_{i_j}) \in R_+$ for $1 \le j \le r$, where β_1 should be read as α_{i_1} . By Subsection 4.2 and the cocycle property, we have

$$C_{(e,w)}(t,\gamma) = C_{\iota}C_{(w,e)}(\gamma^{-1},t^{-1})C_{\iota} = C_{\iota}\left(R_{i_{r}}(\gamma^{\beta_{r}})\cdots R_{i_{2}}(\gamma^{\beta_{2}})R_{i_{1}}(\gamma^{\beta_{1}})\right)^{-1}C_{\iota}.$$

From the expression for $R_i(z)$ it now follows that $C_{(e,w)} \in Q_0(\mathcal{B}) \otimes \operatorname{End}(H_0)$. Since $\lim_{z\to 0} R_i(z) = k\eta(T_i^{-1})$ we furthermore have

$$C_{(e,w)}^{(0)} = k^{-\ell(w)} C_{\iota} \eta(T_w) C_{\iota}$$

The map C_{ι} is the K-linear extension of the anti-algebra involution of H_0 mapping T_w to $T_{w^{-1}}$. Hence $C_{(e,w)}^{(0)}(h) = k^{-\ell(w)}hT_{w^{-1}}$ for $h \in H_0$.

Define $U \in \operatorname{End}(H_0)^{\mathbb{K}} := \mathbb{K} \otimes \operatorname{End}(H_0)$ by

(5.18)
$$U(k^{-\ell(w)}T_{w_0}T_{w^{-1}}) = \tau(e,w)\Phi_{\kappa}, \qquad w \in S_N.$$

Since SOL is \mathbb{S}_N -invariant, U is an End (H_0) -valued solution of the BqKZ, i.e.

$$C_{(\lambda,\mu)}(t,\gamma)U(q^{-\lambda}t,q^{\mu}\gamma) = U(t,\gamma), \qquad \lambda,\mu \in \mathbb{Z}^{N}$$

as $\operatorname{End}(H_0)$ -valued meromorphic functions in $(t, \gamma) \in T \times T$.

Lemma 5.12. $U \in \text{End}(H_0)^{\mathbb{K}}$ is invertible.

Proof. Using the natural identification $\operatorname{End}(H_0)^{\mathbb{K}} \simeq \operatorname{End}_{\mathbb{K}}(H_0^{\mathbb{K}})$ as \mathbb{K} -algebras, we need to verify that $U \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$.

Set $\Phi_w := \tau(e, w)\Phi_{\kappa}$ and $\Psi_w := \tau(e, w)\Psi$ for $w \in S_N$, so that $\Phi_w(t, \gamma) = W_{\kappa}(t, w^{-1}\gamma)\Psi_w(t, \gamma)$. Since $C_{(e,w)}(t, \gamma)$ is independent of $t \in T$, we simply write it as $C_{(e,w)}(\gamma)$. Recall the W-invariant subset $\mathcal{S} \subset T$ (see (4.6)), which contains \mathcal{S}_+ . By Lemma 4.4 and Lemma 5.7, we have for some $\epsilon > 0$ the power series expansion

$$\Psi_w(t,\gamma) = \sum_{\alpha \in Q_+} C_{(e,w)}(\gamma) \big(\Gamma_\alpha(w^{-1}\gamma) \big) t^{-\alpha}$$

for $(t, \gamma) \in B_{\epsilon}^{-1} \times T \setminus S$, converging normally on compact of $B_{\epsilon}^{-1} \times T \setminus S$. We write $\Gamma_{\alpha}^{w}(\gamma) := C_{(e,w)}(\gamma) (\Gamma_{\alpha}(w^{-1}\gamma))$ in the remainder of the proof. It is a meromorphic function in $\gamma \in T$, analytic on $T \setminus S$, and the power series expansion of Ψ_w becomes

(5.19)
$$\Psi_w(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma^w_\alpha(\gamma) t^{-\alpha}.$$

Observe that

$$\Gamma_0^w(\gamma) \to C_{(e,w)}^{(0)}(T_{w_0}) = k^{-\ell(w)} T_{w_0} T_{w^{-1}},$$

in the limit $\gamma^{\alpha_i} \to 0 \ (1 \le i < N)$, in view of the previous lemma.

Write $U = V\Xi$ with V, Ξ the K-linear endomorphisms of $H_0^{\mathbb{K}}$ given by

$$\Xi \left(k^{-\ell(w)} T_{w_0} T_{w^{-1}} \right) (t, \gamma) = W_{\kappa}(t, w^{-1} \gamma) k^{-\ell(w)} T_{w_0} T_{w^{-1}},$$
$$V \left(k^{-\ell(w)} T_{w_0} T_{w^{-1}} \right) = \Psi_w,$$

for $w \in S_N$. Since $\Xi \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$ it suffices to show that $V \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$. Let Mbe the matrix of V with respect to the \mathbb{K} -basis $k^{-\ell(w)}T_{w_0}T_{w^{-1}}$ ($w \in S_N$) of $H_0^{\mathbb{K}}$. Now fix $\zeta \in T \setminus S$ such that $\zeta^{\alpha} \notin q^{\mathbb{Z}}$ for all $\alpha \in R$. The matrix $M(t,\gamma)$ may be specialized at $\gamma = \zeta$ and the limit of $M(t,\zeta)$ as $t^{-\alpha_i} \to 0$ ($1 \le i < N$) exists. We write $M^{(0)}(\zeta)$ for the limit and $V^{(0)}(\zeta)$ for the corresponding linear endomorphism of H_0 . We then have

$$V^{(0)}(\zeta) \left(k^{-\ell(w)} T_{w_0} T_{w^{-1}} \right) = \Gamma_0^w(\zeta) = K(w^{-1}\zeta) C_{(e,w)}(\zeta) T_{w_0}$$

with $K(\gamma)$ given by (5.16). Note that $K(w^{-1}\zeta) \neq 0$ since $\zeta^{\alpha} \notin q^{\mathbb{Z}}$ for all $\alpha \in R$. By the explicit expression for the cocycle value $C_{(e,w)}(\zeta) \in \operatorname{End}(H_0)$ (see the proof of the previous lemma) we have

$$C_{(e,w)}(\zeta)(T_{w_0}) = \sum_{v \le w} a_v^w(\zeta) T_{w_0} T_{v^{-1}},$$

with $a_w^w(\zeta) \neq 0$ and with \leq the Bruhat order on S_N . This implies that $V^{(0)}(\zeta)$ is a linear automorphism of H_0 , hence $\det(M^{(0)}(\zeta)) \neq 0$. Consequently, $\det(M) \in \mathbb{K}^{\times}$ and $V \in \operatorname{GL}_{\mathbb{K}}(H_0^{\mathbb{K}})$.

Proposition 5.13. (i) $U' \in \operatorname{End}(H_0)^{\mathbb{K}}$ is an $\operatorname{End}(H_0)$ -valued meromorphic solution of BqKZ if and only if U' = UF for some $F \in \operatorname{End}(H_0)^{\mathbb{F}}$.

(ii) U, viewed as \mathbb{K} -linear endomorphism of $H_0^{\mathbb{K}}$, restricts to an \mathbb{F} -linear isomorphism $U: H_0^{\mathbb{F}} \to \text{SOL}$.

(iii) $\{\tau(e, w)\Phi_{\kappa}\}_{w\in S_N}$ is an \mathbb{F} -basis of SOL.

Proof. (i) If U' is an End (H_0) -valued meromorphic solution of BqKZ then, since U is invertible, we have for all $\lambda, \mu \in \mathbb{Z}^N$,

$$U(q^{-\lambda}t,q^{\mu}\gamma)^{-1}U'(q^{-\lambda}t,q^{\mu}\gamma) = U(t,\gamma)^{-1}U'(t,\gamma).$$

Hence U' = UF with $F \in \text{End}(H_0)^{\mathbb{F}}$. The converse implication is clear.

(ii) By the previous lemma we have $U : H_0^{\mathbb{F}} \hookrightarrow \text{SOL}$. It is surjective, since for $g \in \text{SOL}$, $f := U^{-1}g \in H_0^{\mathbb{K}}$ satisfies $f(q^{-\lambda}t, q^{\mu}\gamma) = f(t, \gamma)$ for all $\lambda, \mu \in \mathbb{Z}^N$ (cf. the proof of (i)), hence $f \in H_0^{\mathbb{F}}$.

(iii) This is clear from (ii) and from the definition of U.

By Proposition 5.9 and by the proofs of Lemma 5.12 and Proposition 5.13 we obtain the following consistency statement for the quantum KZ equation (4.10) with specialized central character (see [7] and [9]). Recall the W-invariant subset $\mathcal{S} \subset T$ given by (4.6). Recall furthermore that $C_{(e,w)}(t,\gamma)$ for $w \in S_N$ only depends on γ , so we simply write it as $C_{(e,w)}(\gamma)$.

Corollary 5.14. Fix $\zeta \in T \setminus S$ such that $\zeta^{\alpha} \notin q^{\mathbb{Z}}$ for all $\alpha \in R$. For generic $\kappa \in \mathbb{C}^{\times}$, the H_0 -valued meromorphic functions $(\tau(e, w)\Phi_{\kappa})(t, \gamma)$ in $(t, \gamma) \in T \times T$ $(w \in S_N)$ can be specialized at $\gamma = \zeta$, giving rise to

- (i) a basis $\{C_{(e,w)}(\zeta)\Phi_{\kappa}(\cdot,w^{-1}\zeta)\}_{w\in S_N}$ of SOL $_{\zeta}$ over $\mathcal{E}(T)$;
- (ii) an invertible End(H_0)-valued meromorphic solution U_{ζ} of the quantum KZ equations (4.10), where $U_{\zeta} \in \text{End}(H_0)^{\mathcal{M}(T)}$ is explicitly defined by

 $U_{\zeta}(k^{-\ell(w)}T_{w_0}T_{w^{-1}}) := C_{(e,w)}(\zeta)\Phi_{\kappa}(\cdot, w^{-1}\zeta), \qquad w \in S_N.$

6. The correspondence with bispectral problems

For a finite dimensional affine Hecke algebra module M, Cherednik [7, Thm. 4.2] derived, for arbitrary root systems, a correspondence between solutions of the quantum affine KZ equations associated to M and solutions of a system of (possibly matrix-valued) q-difference equations. This correspondence was considered before in [6, Thm. 3.4] for a special class of modules M, in which case the corresponding q-difference operators contain the Macdonald q-difference operator (the latter observation is due to Cherednik [6, Thm. 4.4] for GL_N, and due to Kato [26, Thm. 4.6] and Cherednik [7] for general root systems). In the classical setting (q = 1) it goes back to Matsuo [35].

In the present GL_N setting, with $M = M(\zeta)$ the minimal principal series module with specialized central character $\zeta \in T$, these constructions give rise to an explicit map χ_+ (independent of ζ) from the solution space $\operatorname{SOL}_{\zeta}$ of the quantum KZ equation (4.10) to the solution space of the spectral problem for the commuting Ruijsenaars' [39] trigonometric q-difference operators with spectral parameter ζ^{-1} , see [6, Thm. 4.4] (for GL_N the Macdonald q-difference operators coincide with the Ruijsenaars' operators). We recall this result in detail in Subsection 6.4.

In this section we investigate the map χ_+ when applied to solutions of the bispectral extension of the quantum KZ equations; see Subsection 3.2. In order to do so, we need to replace in the correspondence as described in the previous paragraph, the role of $M(\zeta)$ by the infinite dimensional, formal principal series module of H as defined in Subsection 4.1. We then show that the same map χ_+ gives rise to an embedding of the solution space SOL of BqKZ into the solution space of a bispectral problem for the Ruijsenaars operators. The techniques employed in this section are analogous to the ones for the usual correspondence (see [9, Chpt. 1]). For the convenience of the reader we provide full details of the arguments involved.

We fix $\kappa \in \mathbb{C}^{\times}$ throughout Subsections 6.1-6.3.

6.1. The monodromy cocycle. Observe that $F \in \text{End}(H_0)^{\mathbb{K}}$ is a $\text{End}(H_0)$ -valued meromorphic solution of BqKZ if and only if

$$\tau(\mathbf{w})F = F, \qquad \mathbf{w} \in \mathbb{Z}^N \times \mathbb{Z}^N,$$

where the W-action τ on $\operatorname{End}(H_0)^{\mathbb{K}}$ is defined by

$$(\tau(\mathbf{w})F)(t,\gamma) := C_{\mathbf{w}}(t,\gamma)(\mathbf{w}F)(t,\gamma) = C_{\mathbf{w}}(t,\gamma)F(\mathbf{w}^{-1}(t,\gamma))$$

for $w \in \mathbb{W}$ and $F \in \text{End}(H_0)^{\mathbb{K}}$, viewed as identities between $\text{End}(H_0)$ -valued meromorphic functions in $(t, \gamma) \in T \times T$.

By Proposition 5.13(i), given an $\operatorname{End}(H_0)$ -valued meromorphic solution F of BqKZ, there exists a unique $G \in \operatorname{End}(H_0)^{\mathbb{F}}$ such that F = UG. Accordingly, G describes the deviation of F from the fundamental solution U of BqKZ, and therefore can be thought of as a connection matrix. We will consider the special cases when F are the $\operatorname{End}(H_0)$ -valued meromorphic solutions $\tau(w)U$ ($w \in W$) of BqKZ.

For $w \in W$ we set

$$\mathcal{T}_{\mathbf{w}} := U^{-1}(\tau(\mathbf{w})U) \in \operatorname{End}(H_0)^{\mathbb{F}}$$

that is, $\mathcal{T}_{\mathbf{w}}$ ($\mathbf{w} \in \mathbb{W}$) is the unique element of $\operatorname{End}(H_0)^{\mathbb{F}}$ such that

$$\tau(\mathbf{w})U = U\mathcal{T}_{\mathbf{w}}.$$

Note that $\operatorname{End}(H_0)^{\mathbb{F}}$ is a W-stable subalgebra of $\operatorname{End}(H_0)^{\mathbb{K}}$ with respect to the action $(wF)(t,\gamma) = F(w^{-1}(t,\gamma))$. The following lemma now shows that the \mathcal{T}_w $(w \in W)$ define a cocycle of W with values in the group of units of $\operatorname{End}(H_0)^{\mathbb{F}}$.

Lemma 6.1. (i) T_w = id for w ∈ Z^N × Z^N.
(ii) For w, w' ∈ W we have the cocycle relation
T_{ww'} = T_ww(T_{w'})

in $\operatorname{End}(H_0)^{\mathbb{F}}$.

Proof. (i) This follows immediately from the fact that U is an $End(H_0)$ -valued meromorphic solution of BqKZ.

(ii) Note that $\mathcal{T}_{\mathbf{w}} = U^{-1}C_{\mathbf{w}}\mathbf{w}(U)$ for $\mathbf{w} \in \mathbb{W}$. By the cocycle condition for $C_{\mathbf{w}} \in \mathrm{End}(H_0)^{\mathbb{K}}$, which reads in the present notations as $C_{\mathbf{w}\mathbf{w}'} = C_{\mathbf{w}}\mathbf{w}(C_{\mathbf{w}'})$ for $\mathbf{w}, \mathbf{w}' \in \mathbb{W}$, we have

$$\mathcal{T}_{\mathbf{w}\mathbf{w}'} = U^{-1}C_{\mathbf{w}\mathbf{w}'}\mathbf{w}\mathbf{w}'(U) = U^{-1}C_{\mathbf{w}}\mathbf{w}(C_{\mathbf{w}'}\mathbf{w}'(U))$$
$$= U^{-1}C_{\mathbf{w}}\mathbf{w}(U)\mathbf{w}(U^{-1}C_{\mathbf{w}'}\mathbf{w}'(U)) = \mathcal{T}_{\mathbf{w}}\mathbf{w}(\mathcal{T}_{\mathbf{w}'})$$

for all $w, w' \in W$.

Definition 6.2. In analogy with the terminology in [9, §1.3.3] for the quantum KZ equation, we call $\{\mathcal{T}_w\}_{w\in\mathbb{W}}$ the monodromy cocycle of the BqKZ.

Remark 6.3. Connection matrices and Riemann-Hilbert problems for ordinary linear q-difference equations have been extensively studied; see, e.g., [3] and [41]. For quantum KZ equations, connection matrices have been computed explicitly in, e.g., [20], [14, §12], and [27].

6.2. The correspondence. Consider the algebra $\mathbb{C}(T \times T) \# \mathbb{W}$, where \mathbb{W} acts as field automorphisms on $\mathbb{C}(T \times T)$ by the formula (3.1). Recall that $\mathbb{C}(T \times T) \# \mathbb{W}$ naturally acts on \mathbb{K} . We write Df for the action of $D \in \mathbb{C}(T \times T) \# \mathbb{W}$ on $f \in \mathbb{K}$.

We have a representation $\vartheta \colon \mathbb{C}(T \times T) \# \mathbb{W} \to \operatorname{End}(\operatorname{End}(H_0)^{\mathbb{K}})$ given by

$$\begin{split} \vartheta(f)F &= fF, \qquad f \in \mathbb{C}(T \times T), \\ \vartheta(\mathbf{w})F &= \mathbf{w}(F), \qquad \mathbf{w} \in \mathbb{W} \end{split}$$

for $F \in \operatorname{End}(H_0)^{\mathbb{K}}$. Let \mathbb{D} be the subalgebra $\mathbb{C}(T \times T) # (\mathbb{Z}^N \times \mathbb{Z}^N)$ of $\mathbb{C}(T \times T) # \mathbb{W}$. Under the natural action of $\mathbb{C}(T \times T) # \mathbb{W}$ on $\mathbb{C}(T \times T)$, the subalgebra \mathbb{D} identifies with the algebra of q-difference operators on $T \times T$ with rational coefficients.

Set $H_0^* := \operatorname{Hom}(H_0, \mathbb{C})$. We will regard a linear functional $\chi \in H_0^*$ also as an element of $\operatorname{Hom}_{\mathbb{K}}(H_0^{\mathbb{K}}, \mathbb{K})$ by \mathbb{K} -linear extension. For $F \in \operatorname{End}(H_0)^{\mathbb{K}}$, denote

$$\phi_{\chi,v}^F := \chi(Fv) \in \mathbb{K}, \qquad \chi \in H_0^*, v \in H_0$$

for its matrix coefficients. Note that for any $D \in \mathbb{C}(T \times T) \# \mathbb{W}, \chi \in H_0^*$ and $v \in H_0$,

$$(6.1) D\phi_{\chi,v}^F = \phi_{\chi,v}^{\vartheta(D)F}$$

for all $F \in \operatorname{End}(H_0)^{\mathbb{K}}$.

Lemma 6.4. For $w \in W$ we have

$$\vartheta(\mathbf{w})U = C_{\mathbf{w}}^{-1}U\mathcal{T}_{\mathbf{w}}.$$

In particular, $\vartheta(\mathbf{w})U = C_{\mathbf{w}}^{-1}U$ for $\mathbf{w} \in \mathbb{Z}^N \times \mathbb{Z}^N$.

Proof. For $w \in W$

$$\vartheta(\mathbf{w})U = \mathbf{w}(U) = C_{\mathbf{w}}^{-1}(\tau(\mathbf{w})U) = C_{\mathbf{w}}^{-1}U\mathcal{T}_{\mathbf{w}}$$

The second claim follows from the fact that $\mathcal{T}_{w} = id$ for $w \in \mathbb{Z}^{N} \times \mathbb{Z}^{N}$.

We now are going to look for a particular linear functional χ such that the matrix coefficients $\phi_{\chi,v}^U$ ($v \in H_0$) of U solves a bispectral problem with respect to two commuting families of Ruijsenaars' trigonometric q-difference operators (one family acting on the first torus component, the second on the second torus component). In view of (6.1) and the previous lemma, to obtain q-difference equations for $\phi_{\chi,v}$ we have to deal with the cocycle value C_w and the monodromy matrix \mathcal{T}_w in the equations $w\phi_{\chi,v}^U = \phi_{\chi,v}^{C_w^{-1}U\mathcal{T}_w}$. It is convenient to postpone the analysis of the monodromy cocycle by initially absorbing it into the action ϑ of $\mathbb{C}(T \times T) \# \mathbb{W}$ via the twisted algebra homomorphism

$$\vartheta_{\mathcal{T}} \colon \mathbb{C}(T \times T) \# \mathbb{W} \to \operatorname{End}(\operatorname{End}(H_0)^{\mathbb{K}}),$$

defined by

$$\vartheta_{\mathcal{T}}(f)F = fF, \qquad f \in \mathbb{C}(T \times T)$$

$$\vartheta_{\mathcal{T}}(\mathbf{w})F = \mathbf{w}(F)\mathcal{T}_{\mathbf{w}}^{-1}, \qquad \mathbf{w} \in \mathbb{W}$$

for $F \in \operatorname{End}(H_0)^{\mathbb{K}}$. Note that $\vartheta_{\mathcal{T}}$ is indeed an algebra homomorphism, thanks to the cocycle condition for \mathcal{T} . Moreover, $\vartheta_{\mathcal{T}}|_{\mathbb{D}} = \vartheta|_{\mathbb{D}}$.

For $D \in \mathbb{C}(T \times T) \# \mathbb{W}$ we will occasionally use the notations

(6.2)
$$D = \sum_{\mathbf{w} \in \mathbb{W}} d_{\mathbf{w}} \mathbf{w} = \sum_{\mathbf{v} \in \mathbb{S}_N} D_{\mathbf{v}} \mathbf{v},$$

where $d_{\mathbf{w}} \in \mathbb{C}(T \times T)$ ($\mathbf{w} \in \mathbb{W}$) and $D_{\mathbf{v}} = \sum_{\mathbf{u} \in \mathbb{Z}^N \times \mathbb{Z}^N} d_{\mathbf{u}\mathbf{v}}\mathbf{u} \in \mathbb{D}$ ($\mathbf{v} \in \mathbb{S}_N$). Reformulating (6.1) and Lemma 6.4 in terms of the twisted action $\vartheta_{\mathcal{T}}$ yields the following result.

Lemma 6.5. (i) For $w \in W$ we have

$$\vartheta_{\mathcal{T}}(\mathbf{w})U = C_{\mathbf{w}}^{-1}U.$$

(ii) For $D \in \mathbb{C}(T \times T) \# \mathbb{W}$ we have

$$\phi_{\chi,v}^{\vartheta_{\mathcal{T}}(D)U} = \sum_{\mathbf{v}\in\mathbb{S}_N} D_{\mathbf{v}}(\phi_{\chi,v}^{C_{\mathbf{v}}^{-1}U})$$

for all $\chi \in H_0^*$ and $v \in H_0$.

Proof. (i) This is clear from Lemma 6.4 and the definition of $\vartheta_{\mathcal{T}}$. (ii) By Lemma 6.4 and (6.1), we obtain

$$\phi_{\chi,v}^{\vartheta_{\mathcal{T}}(D)U} = \sum_{\mathbf{v}\in\mathbb{S}_N} \phi_{\chi,v}^{\vartheta(D_{\mathbf{v}})\vartheta_{\mathcal{T}}(\mathbf{v})U} = \sum_{\mathbf{v}\in\mathbb{S}_N} D_{\mathbf{v}}(\phi_{\chi,v}^{\vartheta_{\mathcal{T}}(\mathbf{v})U}).$$

The result now follows from (i).

We define the restriction map Res: $\mathbb{C}(T \times T) \# \mathbb{W} \to \mathbb{D}$ to be the $\mathbb{C}(T \times T)$ -linear map

$$\operatorname{Res}(D) := \sum_{\mathbf{v} \in \mathbb{S}_N} D_{\mathbf{v}}, \qquad D \in \mathbb{D}.$$

Lemma 6.4(ii) implies that if we have a linear functional $\chi_+ \in H_0^*$ such that $\chi_+(C_v^{-1}U) = \chi_+(U)$ for all $v \in \mathbb{S}_N$, then the corresponding matrix coefficients $\phi_{\chi_+,v}^U$ $(v \in H_0)$ satisfy

(6.3)
$$\operatorname{Res}(D)(\phi_{\chi_+,v}^U) = \phi_{\chi_+,v}^{\vartheta_{\mathcal{T}}(D)U}$$

for all $D \in \mathbb{C}(T \times T) \# \mathbb{W}$.

Lemma 6.6. Define $\chi_+ \in H_0^*$ by $\chi_+(T_w) = k^{\ell(w)}$ for all $w \in S_N$. Then

$$\chi_+(C_{\rm v}^{-1}F) = \chi_+(F)$$

for $F \in \text{End}(H_0)^{\mathbb{K}}$ and $\mathbf{v} \in \mathbb{S}_N$.

Proof. Since $C_{\iota}(T_w) = T_{w^{-1}}$ for $w \in S_N$ we have $\chi_+ \circ C_{\iota} = \chi_+$. By the cocycle condition for C_w ($w \in \mathbb{S}_N$) it remains to prove that $\chi_+ \circ C_{(s_i,e)} = \chi_+$ for $1 \leq i < N$. But this follows from the expression $C_{(s_i,e)}(t,\gamma) = c_k(t_i/t_{i+1})^{-1}(\eta(T_i)-k)+1$ (see Lemma 4.3) since

(6.4)
$$\chi_+((T_i - k)h) = 0$$

for $1 \leq i < N$ and $h \in H_0$.

If $D \in \mathbb{C}(T \times T) \# \mathbb{W}$ satisfies $\vartheta_{\mathcal{T}}(D)U = \lambda U$ for some $\lambda \in \mathbb{K}$, then it follows from (6.3) that the matrix coefficients $\phi_{\chi_+,v}^U$ ($v \in H_0$) are eigenfunctions of $\operatorname{Res}(D)$ with eigenvalue λ . We will now construct such a commuting family of D's. It leads to the interpretation of the $\phi_{\chi_+,v}$ ($v \in H_0$) as solutions of a bispectral problem.

The appropriate elements $D \in \mathbb{C}(T \times T) \# \mathbb{W}$ are obtained as images of elements from the center Z(H) of the affine Hecke algebra H under the faithful algebra homomorphism ρ from Theorem 2.4. Since we aim at a bispectral version, we will interpret ρ as algebra map $\rho : H \to \mathbb{C}(T \times T) \# \mathbb{W}$ in two different ways. We have, on the one hand, the algebra homomorphism

$$\rho_{k^{-1},q}^x: H(k^{-1}) \to \mathbb{C}(T \times T) \# \mathbb{W},$$

which is the map $\rho_{k^{-1},q}$ from Theorem 2.4, interpreted as algebra homomorphism from $H(k^{-1})$ to the subalgebra $\mathbb{C}(T \times \{1\}) \#(W \times \{e\})$ of $\mathbb{C}(T \times T) \# \mathbb{W}$. On the other hand, we have an algebra homomorphism

$$\rho^y_{k,q^{-1}}: H(k) \to \mathbb{C}(T \times T) \# \mathbb{W},$$

defined as the map $\rho_{k,q^{-1}}$ from Theorem 2.4, interpreted as algebra homomorphism from H(k) to the subalgebra $\mathbb{C}(\{1\} \times T) \# (\{e\} \times W)$ of $\mathbb{C}(T \times T) \# \mathbb{W}$. Note that they can be combined into an algebra homomorphism

$$\rho_{k^{-1},q}^x \times \rho_{k,q^{-1}}^y : H(k^{-1}) \otimes H(k) \to \mathbb{C}(T \times T) \# \mathbb{W}.$$

Definition 6.7. (i) For $h \in H(k^{-1})$, define

$$D_h^x := \rho_{k^{-1} a}^x(h) \in \mathbb{C}(T \times T) \# \mathbb{W}.$$

(ii) For $h \in H(k)$, define

$$D_h^y := \rho_{k, q^{-1}}^y(h) \in \mathbb{C}(T \times T) \# \mathbb{W}$$

Remark 6.8. Let °: $H(k^{-1}) \to H(k)$ be the algebra isomorphism defined by $\pi^{\circ} = \pi$ and $T_i^{\circ} = T_i^{-1}$ for $1 \le i < N$. Then

(6.5)
$$D_{h^{\circ}}^{y} = \iota D_{h}^{x}\iota, \qquad \forall h \in H(k^{-1}).$$

This follows by verifying the identity

$$\rho^{y}_{k,q^{-1}}(h^{\circ}) = \iota \rho^{x}_{k^{-1},q}(h)\iota$$

for the algebraic generators π and T_i $(1 \le i < N)$ of $H(k^{-1})$ using Theorem 2.4.

Recall the formal principal series representation, encoded by the algebra homomorphism $\eta: H(k) \to \operatorname{End}(H_0)^{\mathbb{K}}$ (see Subsection 4.1).

Proposition 6.9. (i) For $h \in H(k^{-1})$ we have

(6.6)
$$\vartheta_{\mathcal{T}}(D_h^x)U = \eta(h^{\dagger})U,$$

where $\dagger \colon H(k^{-1}) \to H(k)$ is the unique anti-algebra isomorphism satisfying

$$T_i^{\dagger} = T_i^{-1}, \qquad \pi^{\dagger} = \pi^{-1}$$

for $1 \le i < N$. (ii) For $h \in H(k)$ we have

(6.7)
$$\vartheta_{\mathcal{T}}(D_h^y)U = C_{\iota}\iota(\eta(h^{\ddagger}))C_{\iota}U,$$

where $\ddagger: H(k) \rightarrow H(k)$ is the unique anti-algebra involution satisfying

$$T_i^{\ddagger} = T_i, \qquad \pi^{\ddagger} = \pi^{-1}$$

for $1 \leq i < N$ (note that $\dagger = \ddagger \circ \circ$).

Proof. (i) We first show that it suffices to prove (6.6) for algebraic generators of $H(k^{-1})$. Indeed, if (6.6) is valid for $h, h' \in H(k^{-1})$, then we have

$$\begin{split} \vartheta_{\mathcal{T}}(D_{hh'}^x)U &= \vartheta_{\mathcal{T}}(D_h^t)\vartheta_{\mathcal{T}}(D_{h'}^x)U = \vartheta_{\mathcal{T}}(D_h^x)\eta(h'^{\dagger})U \\ &= \eta(h'^{\dagger})\vartheta_{\mathcal{T}}(D_h^x)U = \eta(h'^{\dagger})\eta(h^{\dagger})U \\ &= \eta((hh')^{\dagger})U, \end{split}$$

where the third equality follows since $[\vartheta_{\mathcal{T}}(D_h^x), \eta(h'^{\dagger})] = 0$ as endomorphisms of $\operatorname{End}_{\mathbb{K}}(H_0^{\mathbb{K}})$ (here $\eta(h'^{\dagger})$ should be viewed as element in $\operatorname{End}(\operatorname{End}(H_0)^{\mathbb{K}})$ by left multiplication). Indeed, since $\eta(h'^{\dagger})(t,\gamma)$ does not depend on the torus parameter $t \in T$, it commutes with $\vartheta_{\mathcal{T}}(D_h^x) \in \vartheta_{\mathcal{T}}(\mathbb{C}(T \times \{1\}) \#(W \times \{e\}))$ (which involves, besides the action of $W \times \{e\}$, only right multiplication by the monodromy cocycle).

So it remains to verify (6.6) for $h = \pi \in H(k^{-1})$ and for $h = T_i \in H(k^{-1})$ $(1 \le i < N)$. For $h = \pi \in H(k^{-1})$ we have

$$\vartheta_{\mathcal{T}}(D^x_{\pi})U = \vartheta_{\mathcal{T}}((\pi, e))U = C^{-1}_{(\pi, e)}U = \eta(\pi^{\dagger})U,$$

where the last equality follows from (4.5). For $h = T_i \in H(k^{-1})$ $(1 \le i < N)$, we have

$$\vartheta_{\mathcal{T}}(D_{T_{i}}^{x})U = (k^{-1} - c_{k}(x_{i+1}/x_{i}))U + c_{k}(x_{i+1}/x_{i})\vartheta_{\mathcal{T}}((s_{i},e))U$$

= $(k^{-1} - c_{k}(x_{i+1}/x_{i})U + c_{k}(x_{i+1}/x_{i})C_{(s_{i},e)}^{-1}U$
= $\eta(T_{i}^{\dagger})U$,

where we used that $c_{k^{-1}}(z^{-1}) = c_k(z)$ in the first equality, while the second equality follows from Lemma 6.5(i) and the third equality from Lemma 4.3.

(ii) Unfortunately it is not possible to derive (ii) directly from (i) and from (6.5). Instead, one has to repeat the steps of the proof of (i). It again amounts to verifying (6.7) for $h = \pi \in H(k)$ and for $h = T_i \in H(k)$ ($1 \le i < N$). We show the second case, the first case is left to the reader.

Let $1 \leq i < N$. Then we have for $T_i \in H(k)$,

(6.8)
$$\begin{aligned} \vartheta_{\mathcal{T}}(D_{T_i}^y)U &= (k - c_k(y_i/y_{i+1}))U + c_k(y_i/y_{i+1})\vartheta_{\mathcal{T}}((e,s_i))U \\ &= (k - c_k(y_i/y_{i+1}))U + c_k(y_i/y_{i+1})C_{(e,s_i)}^{-1}U. \end{aligned}$$

Since

$$C_{(e,s_i)} = C_{\iota} \iota(C_{(s_i,e)}) C_{\iota}$$

by the cocycle condition (recall Remark 3.5) and since $C_{\iota}^2 = \text{id}$ and $C_{(s_i,e)}(t,\gamma)^{-1} = C_{(s_i,e)}(s_it,\gamma)$, Lemma 4.3 implies that

$$C_{(e,s_i)}^{-1} = c_k (y_i / y_{i+1})^{-1} (C_\iota \iota(\eta(T_i)) C_\iota - k) + 1.$$

Substituting in (6.8) gives $\vartheta_{\mathcal{T}}(D_{T_i}^y)U = C_{\iota}\iota(\eta(T_i^{\ddagger}))C_{\iota}U$, as desired.

The following lemma plays an important role in the bispectral version of the correspondence. Recall that the center Z(H) of the affine Hecke algebra H is given by $\mathbb{C}_Y[T]^{S_N}$ (Bernstein, see [30]).

Lemma 6.10. For $p \in \mathbb{C}[T]^{S_N}$ we have

$$p(Y)^{\dagger} = p(Y^{-1}), \qquad p(Y)^{\ddagger} = p(Y^{-1}).$$

Proof. By (2.4) it immediately follows that $Y_i^{\dagger} = Y_i^{-1}$ for $1 \le i \le N$. This implies the first formula.

For the second formula, it suffices to show that

(6.9)
$$Y_i^{\ddagger} = T_{w_0} Y_{N-i+1}^{-1} T_{w_0}^{-1}$$

in H(k) for $1 \leq i \leq N$, since we then have, for $p \in \mathbb{C}[T]^{S_N}$,

$$p(Y)^{\ddagger} = T_{w_0} p(Y_N^{-1}, \dots, Y_1^{-1}) T_{w_0}^{-1} = T_{w_0} p(Y^{-1}) T_{w_0}^{-1} = p(Y^{-1})$$

where the last equality follows from the fact that $p(Y^{-1}) \in Z(H(k))$. To prove (6.9), note that $T_{w_0}T_i^{-1} = T_{N-i}^{-1}T_{w_0}$, and $Y_{i+1} = T_i^{-1}Y_iT_i^{-1}$ by (2.4), for $1 \le i < N$. Hence (6.9) holds for Y_{i+1} if it is true for Y_i . It thus remains to prove (6.9) for i = 1. We will use the following observation. Write $\sigma_i = s_i s_{i+1} \cdots s_{N-1}$ $(1 \le i < N)$ and $\tau_i = s_j \cdots s_{N-2}$ $(1 \le j < N - 1)$, which are reduced expressions in S_N . Then the longest Weyl group element $w_0 \in S_N$ can be written as

(6.10)
$$w_0 = \sigma_{N-1}\sigma_{N-2}\cdots\sigma_1$$
$$= \sigma_1(\tau_{N-2}\tau_{N-3}\cdots\tau_1),$$

and $\ell(w_0)$ is the sum of the lengths of the factors in the respective products in (6.10).

By (2.4), formula (6.9) for i = 1 will be valid if

(6.11)
$$T_{\sigma_1}\pi^{-1} = T_{w_0}\pi^{-1}T_{\sigma_1}T_{w_0}^{-1}$$

in H(k). By the first expression in (6.10) and the fact that $\pi^{-1}T_{i+1}\pi = T_i$ for $1 \leq i < N-1$, we have in H(k),

$$\pi^{-1}T_{\sigma_1}T_{w_0}^{-1} = \pi^{-1}T_{\sigma_2}^{-1}T_{\sigma_3}^{-1}\cdots T_{\sigma_{N-1}}^{-1} = T_{\tau_1}^{-1}T_{\tau_2}^{-1}\cdots T_{\tau_{N-2}}^{-1}\pi^{-1},$$

so that (6.11) will follow from

$$T_{\sigma_1} = T_{w_0} T_{\tau_1}^{-1} T_{\tau_2}^{-1} \cdots T_{\tau_{N-2}}^{-1}$$

in H(k). But this is a direct consequence of the second expression of w_0 in (6.10).

Corollary 6.11. For $p \in \mathbb{C}[T]^{S_N}$, we have $p(Y)^\circ = p(Y)$, where $\circ: H(k^{-1}) \to H(k)$ is the algebra isomorphism defined in Remark 6.8.

Proof. This follows from the previous lemma and the fact that $\dagger = \ddagger \circ \circ$.

Definition 6.12. (i) Define

$$L_p^x := \operatorname{Res}(D_{p(Y)}^x) \in \mathbb{D}, \qquad p \in \mathbb{C}[T]^{S_N},$$

where p(Y) is the corresponding element in $\mathbb{C}_Y[T]^{S_N} = Z(H(k^{-1}))$. (ii) Define

$$L_p^y := \operatorname{Res}(D_{p(Y)}^y) \in \mathbb{D}, \qquad p \in \mathbb{C}[T]^{S_N},$$

where p(Y) is the corresponding element in Z(H(k)).

By Corollary 6.11 and (6.5) we have

$$L_p^y = \iota L_p^x \iota, \qquad \forall \, p \in \mathbb{C}[T]^{S_N}$$

Furthermore, it is well-known (see [9] and [34]) that the $L_p^x \in \mathbb{C}(T \times \{1\}) # (\mathbb{Z}^N \times \{e\}) \subset \mathbb{D}$ are pairwise commuting and $S_N \times S_N$ -invariant,

$$\mathbf{w}L_p^x \mathbf{w}^{-1} = L_p^x, \qquad \forall \mathbf{w} \in S_N \times S_N.$$

Similarly, the $L_p^y = \iota L_p^x \iota \in \mathbb{C}(\{1\} \times T) \#(\{e\} \times \mathbb{Z}^N) \subset \mathbb{D}$ are pairwise commuting and $S_N \times S_N$ -invariant. Clearly also $[L_p^x, L_{p'}^y] = 0$ for all $p, p' \in \mathbb{C}[T]^{S_N}$ in \mathbb{D} .

For the elementary symmetric functions $e_i \in \mathbb{C}[T]^{S_N}$ $(1 \le i \le N)$ given by

$$e_i(t) = \sum_{\substack{I \subseteq \{1,\dots,N\}\\ \#I=i}} \prod_{j \in I} t_j,$$

the corresponding $L_{e_i}^x$, viewed as elements in

$$\mathbb{C}(T)\#_q\mathbb{Z}^N\simeq\mathbb{C}(T\times\{1\})\#(\mathbb{Z}^N\times\{e\})\subset\mathbb{D},$$

are explicitly given by

(6.12)
$$L_{e_i}^x = \sum_{\substack{I \subseteq \{1,\dots,N\} \\ \#I=i}} \left(\prod_{\substack{r \in I \\ s \notin I}} \frac{kx_r - k^{-1}x_s}{x_r - x_s} \right) \sum_{r \in I} \epsilon_r \in \mathbb{C}(T) \#_q \mathbb{Z}^N, \qquad 1 \le i \le N;$$

see, e.g., [9, §1.3.5] and [28]. Hence, the $L_{e_i}^x$ $(1 \le i \le N)$ are, under their natural interpretation as q-difference operators on $\mathcal{M}(T)$, Ruijsenaars' commuting, trigonometric q-difference operators from [39].

Definition 6.13. Consider the bispectral problem

(6.13)
$$(L_p^x f)(t,\gamma) = p(\gamma^{-1})f(t,\gamma), \qquad \forall p \in \mathbb{C}[T]^{S_N}, \\ (L_p^y f)(t,\gamma) = p(t)f(t,\gamma), \qquad \forall p \in \mathbb{C}[T]^{S_N}$$

for $f \in \mathbb{K}$, where the equations (6.13) are viewed as identities between meromorphic functions in $(t, \gamma) \in T \times T$. We write BiSP $\subset \mathbb{K}$ for the set of solutions $f \in \mathbb{K}$ of (6.13).

Remark 6.14. The bispectral problem for ordinary linear differential operators was introduced by Duistermaat and Grünbaum in [12]. Many different types of bispectral problems have since been considered. In particular, in [21] the bispectral problem for ordinary linear second-order q-difference operators is investigated. For N = 2, our bispectral problem belongs to this class.

The preceding remarks on the invariance properties of the L_p^x and the L_p^y $(p \in \mathbb{C}[T]^{S_N})$ directly give

Lemma 6.15. BiSP is an \mathbb{S}_N -invariant \mathbb{F} -subspace of \mathbb{K} with respect to the usual \mathbb{S}_N -action $(wf)(t, \gamma) = f(w^{-1}(t, \gamma))$ on $f \in \mathbb{K}$.

We can now prove the following bispectral version of the correspondence between solutions of the quantum KZ equations and the spectral problem of the L_p^x $(p \in \mathbb{C}[T]^{S_N})$.

Theorem 6.16. The linear functional $\chi_+ \in H_0^*$ (see Lemma 6.6) defines a \mathbb{S}_N -equivariant \mathbb{F} -linear map

$$\chi_+ \colon \text{SOL} \to \text{BiSP}.$$

Proof. The K-linear extended linear functional χ_+ defines an \mathbb{S}_N -equivariant, F-linear map $\chi_+ : H_0^{\mathbb{K}} \to \mathbb{K}$, since Lemma 6.6 implies that $\chi_+(\tau(\mathbf{w})f) = \mathbf{w}(\chi_+f)$ for $\mathbf{w} \in \mathbb{S}_N$ and $f \in H_0^{\mathbb{K}}$. Hence χ_+ restricts to an \mathbb{S}_N -equivariant, F-linear map $\chi_+ : \text{SOL} \to \mathbb{K}$.

It remains to show that $\chi_+(f) \in \text{BiSP}$ if $f \in \text{SOL}$. Let $f \in \text{SOL}$. By Proposition 5.13 and \mathbb{F} -linearity, it suffices only to consider f of the form f = Uv for $v \in H_0$. Then $\chi_+(f) = \chi_+(Uv) = \phi_{\chi_+,v}^U$. For $p \in \mathbb{C}[T]^{S_N}$ we have

$$\begin{pmatrix} L_p^x \phi_{\chi_+,v}^U \end{pmatrix}(t,\gamma) = \left(\operatorname{Res}(D_{p(Y)}^x)(\phi_{\chi_+,v}^U) \right)(t,\gamma) = \phi_{\chi_+,v}^{\vartheta \tau (D_{p(Y)}^x)U}(t,\gamma)$$
$$= \phi_{\chi_+,v}^{\eta (p(Y)^\dagger)U}(t,\gamma) = p(\gamma^{-1})\phi_{\chi_+,v}^U(t,\gamma)$$

as meromorphic functions in $(t, \gamma) \in T \times T$, where the last equality follows from Lemma 6.10, (4.3) and the fact $p \in \mathbb{C}[T]^{S_N}$. Similarly,

$$\begin{pmatrix} L_p^y \phi_{\chi_+,v}^U \end{pmatrix}(t,\gamma) = \left(\operatorname{Res}(D_{p(Y)}^y)(\phi_{\chi_+,v}^U) \right)(t,\gamma) = \phi_{\chi_+,v}^{\vartheta_{\tau}(D_{p(Y)}^y)U}(t,\gamma)$$
$$= \phi_{\chi_+,v}^{C_\iota(\eta(p(Y)^{\ddagger}))C_\iota U}(t,\gamma) = p(t)\phi_{\chi_+,v}^U(t,\gamma)$$

as meromorphic functions in $(t, \gamma) \in T \times T$, hence $f = \phi_{\chi_+, v}^U \in \text{BiSP}$.

6.3. Bispectral Harish-Chandra series.

Definition 6.17. We call $\Phi_{\kappa}^+ := \chi_+(\Phi_{\kappa}) \in \text{BiSP}$ the basic Harish-Chandra series solution of the bispectral problem.

Corollary 6.18. The solution $\Phi_{\kappa}^+ \in \text{BiSP}$ of the bispectral problem is selfdual, i.e., $\Phi^+(t, \infty) = \Phi^+(\infty^{-1}, t^{-1})$

$$\Phi_{\kappa}^{+}(t,\gamma) = \Phi_{\kappa}^{+}(\gamma^{-1},t^{-1})$$

as meromorphic functions in $(t, \gamma) \in T \times T$.

Proof. By Theorem 5.6, we have

$$\Phi_{\kappa}^+(t,\gamma) = \chi_+(C_{\iota}\Phi_{\kappa}(\gamma^{-1},t^{-1})).$$

But $\chi_+ C_\iota = \chi_+$, hence the result.

Remark 6.19. In [17], for special values of k, the function Φ_{κ}^+ is constructed as formal power series in terms of generalized characters of Verma modules over the quantum group $\mathcal{U}_q(\mathfrak{sl}_N)$ (see also [15], [16]). The quantum group approach also leads to the self-duality of Φ_{κ}^+ ; see [17, Thm. 5.6] (see [16]).

Note that $\Phi_{\kappa}^+ = W_{\kappa} \Psi^+$ with $\Psi^+ = \chi_+(\Psi)$. For $\alpha \in Q_+$ set

$$\Gamma^+_{\alpha}(\gamma) = \chi_+(\Gamma_{\alpha}(\gamma))$$

as meromorphic function in $\gamma \in T$. By Lemma 5.7 and Theorem 5.10, Γ^+_{α} is analytic at $T \setminus S_+$ and

$$\Gamma_0^+(\gamma) = k^{\binom{N}{2}} K(\gamma)$$

with K given by (5.16). Recall that the solution space BiSP of the bispectral problem is S_N -stable. In particular, we have solutions $\Phi_w^+ \in \text{BiSP}$ given by

(6.14)
$$\Phi_w^+(t,\gamma) := \Phi_\kappa^+(t,w^{-1}\gamma),$$

These are solutions of the bispectral problem which are asymptotically free in the asymptotic sector $\{t \in T \mid |t^{\alpha_i}| \gg 0 \forall 1 \le i < N\}$ in the following sense: by Lemma 5.7 we have $\Phi_w^+(t,\gamma) = W_\kappa(t,w^{-1}\gamma)\Psi_w^+(t,\gamma)$ with $\Psi_w^+(t,\gamma) := \Psi^+(t,w^{-1}\gamma)$ admitting, for $\epsilon > 0$ sufficiently small, the power series expansion

(6.15)
$$\Psi_w^+(t,\gamma) = \sum_{\alpha \in Q_+} \Gamma_\alpha^+(w^{-1}\gamma)t^{-1}$$

for $(t, \gamma) \in B_{\epsilon}^{-1} \times T \setminus w(\mathcal{S}_+)$, converging normally in compact of $B_{\epsilon}^{-1} \times T \setminus w(\mathcal{S}_+)$.

 α

Proposition 6.20. The set of asymptotic solutions $\{\Phi_w^+\}_{w \in S_N} \subset \text{BiSP}$ of the bispectral problem is \mathbb{F} -linearly independent.

Proof. Suppose that

$$\sum_{w \in S_N} a_w(t,\gamma) \Phi_w^+(t,\gamma) = 0$$

as meromorphic functions in $(t, \gamma) \in T \times T$ with coefficients $a_w \in \mathbb{F}$ $(w \in S_N)$. Replacing t by $q^{-m\delta}t$ $(m \in \mathbb{N})$ and using (5.5) we obtain

(6.16)
$$\sum_{w \in S_N} k^{-m\langle \delta, \delta \rangle} \gamma^{-mww_0(\delta)} a_w(t,\gamma) W_\kappa(t,w^{-1}\gamma) \Psi_w^+(q^{-m\delta}t,\gamma) = 0$$

as meromorphic functions in $(t, \gamma) \in T \times T$. Fix $u \in S_N$. We are going to derive from (6.16) that $a_u = 0$. For this we will use the fact that for $w \neq u$,

(6.17)
$$\lim_{m \to \infty} \zeta^{m(uw_0(\delta) - ww_0(\delta))} = \lim_{m \to \infty} (w_0 u^{-1} \zeta)^{m(\delta - w_0 u^{-1} ww_0(\delta))} = 0$$

if $\zeta \in uw_0(B_1)$.

Recall the W-invariant subset $S \subset T$ (see (4.6)), which contains S_+ . For generic $\zeta \in T$ (concretely, $\zeta \notin S$, and $a_w(t, \gamma)$ and $W_{\kappa}(t, w^{-1}\gamma)$ specializable at $\gamma = \zeta$ for all $w \in S_N$), it follows from Proposition 5.9 and (6.16) that, for all $m \in \mathbb{N}$,

(6.18)
$$\sum_{w \in S_N} \zeta^{m(uw_0(\delta) - ww_0(\delta))} a_w(t,\zeta) W_\kappa(t,w^{-1}\zeta) \Psi_w^+(q^{-m\delta}t,\zeta) = 0$$

as meromorphic function in $t \in T$. Using (6.17) and the power series expansion (6.15), the limit $m \to \infty$ of (6.18) yields, for generic $\zeta \in uw_0(B_1)$,

$$a_u(t,\zeta)W_\kappa(t,u^{-1}\zeta)\Gamma_0^+(u^{-1}\zeta) = 0$$

as meromorphic function in $t \in T$. This implies $a_u = 0$, as desired.

Corollary 6.21. The map χ_+ : SOL \rightarrow BiSP is injective.

Proof. Note that $\chi_+(\tau(e, w)\Phi_{\kappa}) = \Phi_w^+$ ($w \in S_N$). The statement follows now directly from Proposition 6.20 and Proposition 5.13.

6.4. Specialized central character and Harish-Chandra series. We write

$$\operatorname{SP}_{\zeta} = \{ f \in \mathcal{M}(T) \mid L_p^x f = p(\zeta^{-1}) f \quad \forall p \in \mathbb{C}[T]^{S_N} \}$$

for the spectral problem of the Ruijsenaars q-difference operators with fixed spectral parameter $\zeta \in T$. Note that $SP_{\zeta} \subset H_0^{\mathcal{M}(T)}$ is S_N -stable, with S_N -action on $H_0^{\mathcal{M}(T)}$ given by $(wf)(t) = f(w^{-1}t)$ for $f \in H_0^{\mathcal{M}(T)}$ and $w \in S_N$.

By [13, Prop. 5.2], the quantum KZ equations (4.10) are consistent for all values $\zeta \in T$ of the central character. The arguments from Subsection 6.2, applied to the quantum KZ equations (4.10) for fixed ζ and with the role of U taken over by an invertible matrix solution U_{ζ} of (4.10), result in the following special case of the Cherednik-Matsuo correspondence from [6, 7] (concretely, in the notations of [7], take the principal series module $V = M(\zeta)$ in [7, Thm. 4.2] and let τ be the projection from $M(\zeta)$, along the direct sum decomposition of $M(\zeta)$ in H_0 -isotypical components, onto the trivial component).

Proposition 6.22. Let $\zeta \in T$. Then χ_+ defines an $\mathcal{E}(T)$ -linear S_N -equivariant map $\chi_+ : \operatorname{SOL}_{\zeta} \to \operatorname{SP}_{\zeta}$.

For a further analysis of the map χ_+ : SOL $\zeta \to$ SP $_{\zeta}$, we refer to [7] and [9, §1.3.4].

Harish-Chandra type series solutions of the spectral problem of the Ruijsenaars q-difference operators L_p^x ($p \in \mathbb{C}[T]^{S_N}$) with fixed spectral parameter $\zeta \in T$ were studied in, e.g., [15] and [27] (see also [31] for arbitrary root systems). The results of the previous subsection allow us to reobtain these solutions by specialization of the basic Harish-Chandra series Φ_{κ}^+ . It leads to new results on the convergence and singularities of these solutions, which we state now explicitly.

By Subsection 5.4, for generic $\kappa \in \mathbb{C}^{\times}$ the basic Harish-Chandra series $\Phi_{\kappa}^{+}(t, \gamma)$ is specializable at $\gamma = \zeta$ when $\zeta \in T \setminus S_{+}$. Concretely, for $\zeta \in T \setminus S_{+}$ and generic $\kappa \in \mathbb{C}^{\times}$, we can write

$$\Phi_{\kappa}^{+}(t,\zeta) = W_{\kappa}(t,\zeta)\Psi^{+}(t,\zeta)$$

as meromorphic function in $t \in T$, where $\Psi^+ = \chi_+(\Psi)$ (see Subsection 6.3). Due to the results in Subsection 5.4 (see Proposition 5.8) we obtain the following result.

Corollary 6.23. For $\zeta \in T \setminus S_+$, the meromorphic function $\Psi^+(t,\zeta)$ in $t \in T$ is analytic at $t \in T \setminus S_+^{-1}$.

Let $\zeta \in T \setminus S$, where $S \subset T$ is the *W*-invariant set (4.6). For $\kappa \in \mathbb{C}^{\times}$ such that $W_{\kappa}(t, w^{-1}\gamma)$ may be specialized at $\gamma = \zeta$ for all $w \in S_N$, the asymptotic solutions $\Phi_w^+(t, \gamma)$ ($w \in S_N$) of the bispectral problem (see (6.14)) may thus be specialized at $\gamma = \zeta$, giving rise to solutions $\Phi_w^+(\cdot; \zeta) \in SP_{\zeta}$ ($w \in S_N$); see Corollary 5.14 and Proposition 6.22. Observe that for $\epsilon > 0$ sufficiently small,

$$\Phi_w^+(t,\zeta) = W_\kappa(t,w^{-1}\zeta) \sum_{\alpha \in Q_+} \Gamma_\alpha^+(w^{-1}\zeta)t^{-\alpha}$$

for $t \in B_{\epsilon}^{-1}$, with normal convergence of the power series on compact of B_{ϵ}^{-1} . Since $\zeta \notin S$ we furthermore have

$$\Gamma_0^+(w^{-1}\zeta) = k^{\binom{N}{2}} K(w^{-1}\zeta) \neq 0,$$

with K given by (5.16).

Definition 6.24. Let $\zeta \in T \setminus S$. The $\Phi_w^+(\cdot; \zeta) \in SP_{\zeta}$ ($w \in S_N$) are the Harish-Chandra series solutions of the spectral problem $L_p^x f = p(\zeta^{-1})f$ ($p \in \mathbb{C}[T]^{S_N}$).

Remark 6.25. In [15] (and [31]) the Harish-Chandra series are investigated as formal power series solutions to the spectral problem of the Ruijsenaars operators. The advantage of the present approach is the fact that it implies the convergence of the formal power series, basically as a consequence of a general statement about convergence of formal power series solutions of holonomic systems of q-difference equations (see the appendix). Chalykh's [5] Baker-Akhiezer functions arise as Harish-Chandra series solutions for special values of k; see [31, §4.4]. In [27], the Harish-Chandra series solutions of the Ruijsenaars operators are constructed as matrix coefficients of products of vertex operators. By this approach, one obtains an explicit integral representation of the Harish-Chandra series.

Remark 6.26. Observe that

(6.19)
$$\lim_{\substack{\lambda \in \Lambda:\\ \lambda \to \infty}} \frac{\Phi_{\kappa}^+(t, q^{\lambda} k^{-\delta})}{W_{\kappa}(t, q^{\lambda} k^{-\delta})} = \Gamma_0^+(t^{-1}) = k^{\binom{N}{2}} K(t^{-1}),$$

with $\lambda \to \infty$ meaning $\lambda_i - \lambda_{i+1} \to \infty$ for all $1 \leq i < N$. Thus, K (see (5.16)) is a normalized limit of the asymptotic solutions $\Phi_{\kappa}^+(\cdot, q^{\lambda}k^{-\delta}) \in \operatorname{SP}_{q^{\lambda}k^{-\delta}}$. The solution space $\operatorname{SP}_{q^{\lambda}k^{-\delta}}$ contains the symmetric Macdonald polynomial of degree $\lambda \in \Lambda$. It turns out though that $\Phi_{\kappa}^+(\cdot, q^{\lambda}k^{-\delta})$ is not a multiple of the Macdonald polynomial of degree $\lambda \in \Lambda$, but $\Phi_{\kappa}^+(\cdot, q^{w_0(\lambda)}k^{\delta})$ is (this will become apparent in the next section). On the other hand, the leading coefficient K (see (5.16)) also naturally appears as a normalized limit of the Macdonald polynomial when the degree $\lambda \in \Lambda$ of the polynomial tends to infinity; see [10, Lemma 4.3] (this limit was proven in [40] in the L^2 -sense).

7. Polynomial theory

We assume throughout this section that $k \in \mathbb{C}^{\times}$ satisfies the generic conditions

(7.1)
$$\begin{aligned} k^{2j} \not\in q^{\mathbb{Z}}, & \forall 1 \le j \le N, \\ k^{\langle \delta, \varpi_j - w(\varpi_j) \rangle} \not\in q^{\mathbb{Z}}, & \forall 1 \le j < N, \forall w \in S_N : w(\varpi_j) \ne \varpi_j. \end{aligned}$$

7.1. Polynomial solutions of the quantum KZ equation. We are going to use a special case of Proposition 4.9 to create S_N -invariant (with respect to the S_N action ς on SOL_{ζ}; see (4.11)) polynomial solutions of the quantum KZ equations.

Lemma 7.1. Let $\lambda \in \Lambda$. The possible poles of the $\mathbb{C}[T] \otimes \operatorname{End}(H_0)$ -valued rational function

$$\gamma \mapsto C_{(e,-\lambda)}(\cdot, q^{\lambda}\gamma) = C_{(e,\lambda)}(\cdot, \gamma)^{-1}$$

in $\gamma \in T$ are at $\gamma^{\alpha} \in k^2 q^{-\mathbb{N}}$ for some $\alpha \in R^+$. The possible poles of

$$\gamma \mapsto C_{(e,\lambda)}(\cdot,\gamma)$$

are at $\gamma^{\alpha} \in k^{-2}q^{-\mathbb{N}}$ for some $\alpha \in R^+$.

Proof. Since $R_i(z)$ has only a (simple) pole at $z = k^{-2}$, this follows from (5.13) and the cocycle property of C; see Lemma 5.7.

Since k satisfies $k^{2j} \notin q^{\mathbb{Z}}$ for $1 \leq j \leq N$ by (7.1), the spectrum of $\eta_{q^{\lambda}k^{-\delta}}(\mathbb{C}_Y[T])$ is simple and the $\xi_w(q^{\lambda}k^{-\delta})$ ($w \in S_N$) form a \mathbb{C} -basis of H_0 for all $\lambda \in \Lambda$. Furthermore, for such k we have that $\gamma \mapsto C_{(e,\lambda)}(\cdot, \gamma)^{\pm 1}$ is regular at $\gamma = k^{-\delta}$ for all $\lambda \in \Lambda$; see Lemma 7.1. The additional conditions on k in (7.1) will play a role in Subsection 7.3 and Subsection 7.4.

Proposition 4.9 now immediately implies the following result.

Corollary 7.2. Let $\lambda \in \Lambda$. Then $f \mapsto C_{(e,\lambda)}(\cdot, k^{-\delta})^{-1}f$ defines an S_N -equivariant isomorphism $\operatorname{SOL}_{k^{-\delta}} \to \operatorname{SOL}_{q^{\lambda}k^{-\delta}}$.

The special interest in the quantum KZ equations for the particular central characters $\gamma = q^{\lambda} k^{-\delta}$ ($\lambda \in \Lambda$) comes from the fact that it admits S_N -invariant polynomial solutions. The key step in deriving this result is the following lemma.

Lemma 7.3. The element $v_+ := \sum_{w \in S_N} k^{\ell(w)} T_w \in H_0$ is a constant S_N -invariant solution of the quantum KZ equation with central character $k^{-\delta}$. In other words,

$$C_{\lambda}^{k^{-\delta}}(t)v_{+} = v_{+}, \qquad \forall \lambda \in \mathbb{Z}^{N}.$$

Proof. Note that $R_i(z)v_+ = v_+$, so by Lemma 4.1 and Lemma 4.5(ii), for any $\zeta \in T$,

$$C_{\varpi_{i}}^{\zeta}(t)v_{+} = \eta_{\zeta}(\pi)^{i}v_{+} = \sum_{w \in S_{N}} k^{\ell(w)} \zeta^{w^{-1}w_{0}\varpi_{i}} T_{\sigma^{i}w} = \sum_{w \in S_{N}} k^{\ell(\sigma^{-i}w)} \zeta^{w^{-1}\varpi_{i}} T_{w}.$$

Then use $\ell(\sigma^{-i}w) - \ell(w) = \langle \delta, w^{-1}\varpi_i \rangle$ for $1 \leq i \leq N$ (for the proof of this formula, it suffices to prove it for i = 1. In that case, look at the positive roots that are mapped to negative roots by $\sigma^{-1}w$). It implies that

$$C_{\varpi_i}^{\zeta}(t)v_+ = \sum_{w \in S_N} k^{\ell(w)} (k^{\delta}\zeta)^{w^{-1}\varpi_i} T_w.$$

In particular, $C_{\varpi_i}^{k^{-\delta}}(t)v_+ = v_+$ for all *i*. Note, furthermore, that $R_i(z)v_+ = v_+$ implies that $\varsigma(s_i)v_+ = C_{s_i}^{k^{-\delta}}(t)v_+ = v_+$ for all $1 \le i < N$ (with ς given by (4.11)). Hence, $v_+ \in \text{SOL}_{k^{-\delta}}$ is S_N -invariant.

Proposition 7.4. For $\lambda \in \Lambda$, the nonzero S_N -invariant solution

$$Q_{\lambda} := C_{(e,\lambda)}(\cdot, k^{-\delta})^{-1}v_{+} \in \mathrm{SOL}_{q^{\lambda}k^{-\delta}}$$

of the quantum KZ equation is an H_0 -valued Laurent polynomial on T satisfying

(7.2)
$$Q_{\lambda}(t) = \sum_{\alpha \in Q_{+}} K_{\alpha}(\lambda) t^{\lambda - \alpha}$$

with $K_{\alpha}(\lambda) \in H_0$ (all but finitely many terms zero).

Proof. Note that Corollary 7.2 and Lemma 7.3 imply that $0 \neq Q_{\lambda} \in \text{SOL}_{q^{\lambda}k^{-\delta}}$ and that Q_{λ} is S_N -invariant. The triangularity property (7.2) follows from the cocycle property, (5.13), the explicit form of the $R_i(z)$ and the fact that

$$\eta(\pi)(t^{-1})^{-i}T_w = t^{w^{-1}\varpi_i}T_{\sigma^{-i}w}, \qquad w \in S_N,$$

which is a direct consequence of Lemma 4.1.

7.2. Duality.

Lemma 7.5. For $\lambda \in \Lambda$, we have $Q_{\lambda}(k^{\delta}) = v_+$.

Proof. Since $v_+ \in \text{SOL}_{k^{-\delta}}$ and $C_{\iota}v_+ = v_+$, we obtain

$$Q_{\lambda}(k^{\delta}) = C_{(e,-\lambda)}(k^{\delta}, q^{\lambda}k^{-\delta})v_{+}$$

= $C_{\iota}C_{(-\lambda,e)}(q^{-\lambda}k^{\delta}, k^{-\delta})C_{\iota}v_{+} = v_{+},$

for $\lambda \in \Lambda$.

The polynomial solutions Q_{λ} of the quantum KZ equations are self-dual in the following sense.

Proposition 7.6. For $\lambda, \mu \in \Lambda$, we have

$$Q_{\lambda}(q^{-\mu}k^{\delta}) = C_{\iota}Q_{\mu}(q^{-\lambda}k^{\delta}).$$

Proof. For $\lambda, \mu \in \Lambda$ we have, using $v_+ \in \text{SOL}_{k^{-\delta}}$ and the previous lemma,

(7.3)

$$Q_{\lambda}(q^{-\mu}k^{\circ}) = C_{(e,-\lambda)}(q^{-\mu}k^{\circ},q^{\lambda}k^{-\circ})v_{+}$$

$$= C_{(e,-\lambda)}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta})C_{(-\mu,e)}(q^{-\mu}k^{\delta},k^{-\delta})v_{+}$$

$$= C_{(-\mu,-\lambda)}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta})v_{+}.$$

Since $C_{(-\mu,-\lambda)}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta}) = C_{\iota}C_{(-\lambda,-\mu)}(q^{-\lambda}k^{\delta},q^{\mu}k^{-\delta})C_{\iota}$ and $C_{\iota}v_{+} = v_{+}$, we conclude from (7.3) that $Q_{\lambda}(q^{-\mu}k^{\delta}) = C_{\iota}Q_{\mu}(q^{-\lambda}k^{\delta}).$

7.3. Relation to the basic asymptotically free solution. In this subsection, we relate the polynomial solutions Q_{λ} ($\lambda \in \Lambda$) of the quantum KZ equations to the basic asymptotic solution Φ_{κ} . Some care is needed though: it is not possible to specialize all the asymptotic solutions $C_{(e,w)}(t,\gamma)\Phi_{\kappa}(t,w^{-1}\gamma)$ ($w \in S_N$) to $\gamma = q^{\lambda}k^{-\delta}$ ($\lambda \in \Lambda$) since $q^{\lambda}k^{-\delta} \in S$; see Corollary 5.14. We shall see that $C_{(e,w_0)}(t,\gamma)\Phi_{\kappa}(t,w_0\gamma)$ can be specialized at $\gamma = q^{\lambda}k^{-\delta}$, which is sufficient for our purposes.

Lemma 7.7. Let $\lambda \in \Lambda$. There exists a unique $\Xi_{\lambda} \in \text{SOL}_{q^{\lambda}k^{-\delta}}$ such that, for $\epsilon > 0$ sufficiently small, we have an H_0 -valued power series expansion

$$\Xi_{\lambda}(t) = \sum_{\alpha \in Q^+} \widetilde{\Gamma}_{\alpha}(\lambda) t^{\lambda - \alpha}$$

converging normally on compact of B_{ϵ}^{-1} and with leading coefficient

$$\widetilde{\Gamma}_0(\lambda) = \eta_{q^{\lambda}k^{-\delta}}(T_{w_0})\xi_{w_0}(q^{\lambda}k^{-\delta}).$$

Proof. Consider the gauged quantum KZ equations for $1 \le i \le N$,

(7.4)
$$\widetilde{A}_i(t)\widetilde{\Xi}(q^{-\varpi_i}t) = \widetilde{\Xi}(t), \qquad \widetilde{\Xi} \in H_0^{\mathcal{M}(T)}$$

with q-connection matrices $\widetilde{A}_i(t) = q^{-\langle \lambda, \varpi_i \rangle} C_{(\varpi_i, e)}(t, q^{\lambda} k^{-\delta})$. Note that $\widetilde{A}_N(t) =$ id; see Lemma 5.2. Observe that $\widetilde{\Xi}$ is a solution of the holonomic system (7.4) of q-difference equations if and only if $x^{\lambda} \widetilde{\Xi} \in \text{SOL}_{q^{\lambda} k^{-\delta}}$. By Corollary 4.6, we have $\widetilde{A}_i \in Q_0(\mathcal{A}) \otimes \text{End}(H_0)$ and

(7.5)
$$\widetilde{A}_{i}^{(0)} = q^{-\langle\lambda,\varpi_{i}\rangle} k^{\langle\delta,\varpi_{i}\rangle} \eta_{q^{\lambda}k^{-\delta}} \left(T_{w_{0}} Y^{w_{0}(\varpi_{i})} T_{w_{0}}^{-1} \right).$$

The $\widetilde{A}_i^{(0)}$ $(1 \le i < N)$ are semisimple endomorphisms of H_0 . A basis of simultaneous eigenvectors of H_0 is given by $\eta_{q^{\lambda}k^{-\delta}}(T_{w_0})\xi_w(q^{\lambda}k^{-\delta})$ $(w \in S_N)$. In fact,

$$\widetilde{A}_{i}^{(0)}\left(\eta_{q^{\lambda}k^{-\delta}}(T_{w_{0}})\xi_{w}(q^{\lambda}k^{-\delta})\right) = \gamma_{w,i}\eta_{q^{\lambda}k^{-\delta}}(T_{w_{0}})\xi_{w}(q^{\lambda}k^{-\delta})$$

for all $1 \leq i < N$ and $w \in S_N$ with

$$\gamma_{w,i} = q^{\langle \lambda, w^{-1}w_0(\varpi_i) - \varpi_i \rangle} k^{\langle \delta, \varpi_i - w^{-1}w_0(\varpi_i) \rangle}$$

see (5.10). Note, in particular, that $\gamma_{w,i} \notin q^{-\mathbb{N}}$ for all $w \in S_N$ and all $1 \leq i < N$ by the generic conditions (7.1) on k, and that $\gamma_{w_0,i} = 1$ for all $1 \leq i < N$. Hence, Theorem 8.6 in the appendix, applied to (7.4) by taking M = N - 1, $q_i = q$ and variables $z_i = x^{-\alpha_i}$ $(1 \leq i < N)$ shows that there exists a unique $\widetilde{\Xi} \in \mathcal{M}(T) \otimes H_0$ satisfying (7.4) and admitting an H_0 -valued power series expansion

$$\widetilde{\Xi}(t) = \sum_{\alpha \in Q^+} \widetilde{\Gamma}_{\alpha}(\lambda) t^{-\alpha}$$

converging normally on compact of B_{ϵ}^{-1} for some $\epsilon > 0$ small enough, and having as leading coefficient $\widetilde{\Gamma}_0(\lambda) = \eta_{q^{\lambda}k^{-\delta}}(T_{w_0})\xi_{w_0}(q^{\lambda}k^{-\delta})$. This directly implies the lemma.

Recall that the cocycle values $C_{(e,w)}(t,\gamma)$ ($w \in S_N$) are independent of $t \in T$. We suppress t from the notation and simply write $C_{(e,w)}(\gamma)$. Recall that $C_{(e,w)}(\gamma)$ for $w \in S_N$ is an End(H_0)-valued regular function in $\gamma \in T$.

Theorem 7.8. Fix $\lambda \in \Lambda$. For $\kappa \notin q^{\mathbb{Z}}$, the basic asymptotic solution $\Phi_{\kappa}(t,\gamma)$ of BqKZ can be specialized at $\gamma = q^{w_0(\lambda)}k^{\delta}$, giving rise to a H_0 -valued meromorphic function $\Phi_{\kappa}(t, q^{w_0(\lambda)}k^{\delta})$ in $t \in T$. Then

(7.6)
$$Q_{\lambda}(t) = r_{\kappa} C_{(e,w_0)}(q^{\lambda} k^{-\delta}) \Phi_{\kappa}(t, q^{w_0(\lambda)} k^{\delta})$$

with

(7.7)
$$r_{\kappa} = \theta(\kappa)^{N} k^{-\binom{N}{2}} \prod_{1 \le i < j \le N} \frac{\left(k^{2(j-i+1)}; q\right)_{\infty}}{\left(k^{2(j-i)}; q\right)_{\infty}}.$$

Proof. We first show that both Q_{λ} and the right-hand side of (7.6) are nonzero scalar multiples of Ξ_{λ} .

We start with the right-hand side of (7.6). Since Φ is \mathbb{S}_N -stable, we have

$$\Phi_{w_0} := \tau(e, w_0) \Phi_{\kappa} \in \text{SOL}.$$

Concretely, it is given by

$$\Phi_{w_0}(t,\gamma) = C_{(e,w_0)}(\gamma)\Phi_{\kappa}(t,w_0(\gamma)) = W_{\kappa}(t,w_0(\gamma))C_{(e,w_0)}(\gamma)\Psi(t,w_0(\gamma)).$$

Since $w_0(q^{\lambda}k^{-\delta}) = q^{w_0(\lambda)}k^{\delta} \notin S_+$ by (7.1), we may, in view of Proposition 5.9, specialize $\Phi_{w_0}(t,\gamma)$ at $\gamma = q^{\lambda}k^{-\delta}$, obtaining $\Phi_{w_0}(\cdot,q^{\lambda}k^{-\delta}) \in \text{SOL}_{q^{\lambda}k^{-\delta}}$. By (5.3) we have

(7.8)
$$W_{\kappa}(t, w_0(q^{\lambda}k^{-\delta})) = k^{\langle \delta, \lambda \rangle} \theta(\kappa)^{-N} t^{\lambda},$$

hence by Proposition 5.9,

$$\Phi_{w_0}(t,q^{\lambda}k^{-\delta}) = k^{\langle \delta,\lambda\rangle} \theta(\kappa)^{-N} \sum_{\alpha \in Q_+} \Gamma_{\alpha}^{w_0} t^{\lambda-\alpha}$$

with $\Gamma_{\alpha}^{w_0} = C_{(e,w_0)}(q^{\lambda}k^{-\delta})\Gamma_{\alpha}(q^{w_0(\lambda)}k^{\delta})$. From the definitions of $C_{(e,w_0)}$, d_w , η and ξ_{w_0} (see Subsections 2.3, 3.1 and 4.1) we have

(7.9)

$$C_{(e,w_0)}(\gamma)T_{w_0} = d_{w_0}(\gamma^{-1})^{-1}\eta(T_{w_0})\xi_{w_0}(\gamma)$$

$$= \left(\prod_{\alpha \in R^+} \frac{1}{k - k^{-1}\gamma^{\alpha}}\right)\eta(T_{w_0})\xi_{w_0}(\gamma)$$

as H_0 -valued regular functions in $\gamma \in T$. By Theorem 5.10, the leading coefficient $\Gamma_0^{w_0}$ thus simplifies to

$$\Gamma_{0}^{w_{0}} = K(q^{w_{0}(\lambda)}k^{\delta})C_{(e,w_{0})}(q^{\lambda}k^{-\delta})T_{w_{0}}
= K(q^{w_{0}(\lambda)}k^{\delta})d_{w_{0}}(q^{-\lambda}k^{\delta})^{-1}\eta_{q^{\lambda}k^{-\delta}}(T_{w_{0}})\xi_{w_{0}}(q^{\lambda}k^{-\delta}),$$

where K is given by (5.16). Combined with the previous lemma, we conclude that

(7.10)
$$\Phi_{w_0}(t, q^{\lambda} k^{-\delta}) = k^{\langle \delta, \lambda \rangle} \theta(\kappa)^{-N} K(q^{w_0(\lambda)} k^{\delta}) d_{w_0}(q^{-\lambda} k^{\delta})^{-1} \Xi_{\lambda}(t).$$

In view of (7.1), $\Phi_{w_0}(t, q^{\lambda} k^{-\delta})$ thus is a nonzero constant multiple of $\Xi_{\lambda}(t)$.

Next, we consider $0 \neq Q_{\lambda} \in \text{SOL}_{q^{\lambda}k^{-\delta}}$. By Lemma 7.7 and (7.2), it suffices to note that $K_0(\lambda)$ is a constant multiple of $\eta_{q^{\lambda}k^{-\delta}}(T_{w_0})\xi_{w_0}(q^{\lambda}k^{-\delta})$, which follows directly from the fact that $K_0(\lambda) \in H_0$ satisfies

$$\widetilde{A}_i^{(0)} K_0(\lambda) = K_0(\lambda), \qquad \forall 1 \le i \le N,$$

where \widetilde{A}_i is given by (7.5); see the proof Lemma of 7.7. Thus, $Q_{\lambda}(t)$ is a nonzero constant multiple of $\Xi_{\lambda}(t)$, and we conclude that

$$Q_{\lambda}(t) = r_{\kappa}(\lambda)\Phi_{w_0}(t, q^{\lambda}k^{-\delta}),$$

for some $r_{\kappa}(\lambda) \in \mathbb{C}^{\times}$. We first show that $r_{\kappa}(\lambda)$ is independent of $\lambda \in \Lambda$.

For $w \in S_N$, we write $C_{(w,e)}(t)$ for the γ -independent value $C_{(w,e)}(t,\gamma)$ of the cocycle. Let $\lambda, \mu \in \Lambda$. By the S_N -invariance of Q_{λ} , we then have, on the one hand,

$$Q_{\lambda}(q^{-\mu}k^{\delta}) = C_{(w_{0},e)}(q^{-\mu}k^{\delta})Q_{\lambda}(q^{-w_{0}(\mu)}k^{-\delta})$$

= $r_{\kappa}(\lambda)C_{(w_{0},e)}(q^{-\mu}k^{\delta})C_{(e,w_{0})}(q^{\lambda}k^{-\delta})\Phi_{\kappa}(q^{-w_{0}(\mu)}k^{-\delta},q^{w_{0}(\lambda)}k^{\delta})$
= $r_{\kappa}(\lambda)C_{(w_{0},w_{0})}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta})\Phi_{\kappa}(q^{-w_{0}(\mu)}k^{-\delta},q^{w_{0}(\lambda)}k^{\delta}).$

On the other hand, using the self-duality of Q_{λ} (see Proposition 7.6) and of Φ_{κ} (see Theorem 5.6),

$$\begin{aligned} Q_{\lambda}(q^{-\mu}k^{\delta}) &= C_{\iota}Q_{\mu}(q^{-\lambda}k^{\delta}) = C_{\iota}C_{(w_{0},e)}(q^{-\lambda}k^{\delta})Q_{\mu}(q^{-w_{0}(\lambda)}k^{-\delta}) \\ &= r_{\kappa}(\mu)C_{\iota}C_{(w_{0},e)}(q^{-\lambda}k^{\delta})C_{(e,w_{0})}(q^{\mu}k^{-\delta})\Phi_{\kappa}(q^{-w_{0}(\lambda)}k^{-\delta},q^{w_{0}(\mu)}k^{\delta}) \\ &= r_{\kappa}(\mu)C_{\iota}C_{(w_{0},w_{0})}(q^{-\lambda}k^{\delta},q^{\mu}k^{-\delta})\Phi_{\kappa}(q^{-w_{0}(\lambda)}k^{-\delta},q^{w_{0}(\mu)}k^{\delta}) \\ &= r_{\kappa}(\mu)C_{(w_{0},w_{0})}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta})C_{\iota}\Phi_{\kappa}(q^{-w_{0}(\lambda)}k^{-\delta},q^{w_{0}(\mu)}k^{\delta}) \\ &= r_{\kappa}(\mu)C_{(w_{0},w_{0})}(q^{-\mu}k^{\delta},q^{\lambda}k^{-\delta})\Phi_{\kappa}(q^{-w_{0}(\mu)}k^{-\delta},q^{w_{0}(\lambda)}k^{\delta}). \end{aligned}$$

We conclude that $r_{\kappa}(\lambda) = r_{\kappa}(\mu)$ if $Q_{\lambda}(q^{-\mu}k^{\delta}) \neq 0$. In particular, since $Q_{\lambda}(k^{\delta}) = v_{+} \neq 0$, we have $r_{\kappa}(\lambda) = r_{\kappa}(0)$ for all $\lambda \in \Lambda$.

It remains to compute $r_{\kappa} := r_{\kappa}(0)$. Using the fact that $C_{(e,s_i)}(\gamma) = C_{\iota}R_i(\gamma^{-\alpha_i})C_{\iota}$, with $R_i(z) = c_k(z)^{-1}(\eta(T_i^{-1})-k^{-1})+1$ for $1 \leq i < N$, as well as that $C_{\iota}(T_{w^{-1}}^{-1}T_{w_0}) =$ $T_{w_0w^{-1}}$ for all $w \in S_N$, we get $C_{(e,w_0)}(\gamma)T_{w_0} = \sum_{w \leq w_0} e_w(\gamma)T_{w_0w^{-1}}$ as H_0 -valued regular function in $\gamma \in T$ with $e_w \in \mathbb{C}[T]$ and with

$$e_{w_0}(\gamma) = \prod_{\beta \in R^+} c_k(\gamma^{-\beta})^{-1}.$$

Taking the T_e -coefficient in the expansion of the formula

$$v_{+} = Q_{0} = r_{\kappa} \Phi_{w_{0}}(\cdot, k^{-\delta}) = r_{\kappa} \theta(\kappa)^{-N} K(k^{\delta}) C_{(e,w_{0})}(k^{-\delta}) T_{w_{0}}$$

with respect to the \mathbb{C} -basis $\{T_w\}_{w \in S_N}$ of H_0 , we conclude that

$$r_{\kappa} = \theta(\kappa)^{N} K(k^{\delta})^{-1} \prod_{\beta \in R^{+}} c_{k}(k^{\langle \delta, \beta \rangle})$$

Substituting the explicit expressions (2.5) and (5.16) of c_k and K, respectively, we get the desired formula (7.7) for r_{κ} .

The following formula is an analog for the Q_{λ} ($\lambda \in \Lambda$) of the evaluation formula for the self-dual symmetric Macdonald polynomials (see Subsection 7.4).

Corollary 7.9. Let $\lambda \in \Lambda$ and write $Q_{\lambda}(t) = \sum_{\alpha \in Q_+} K_{\alpha}(\lambda)t^{\lambda-\alpha}$ with $K_{\alpha}(\lambda) \in H_0$ (see Proposition 7.4). The leading coefficient $K_0(\lambda)$ is given by

$$K_{0}(\lambda) = k^{\langle \delta, \lambda \rangle} \left(\prod_{1 \le i < j \le N} \prod_{m=0}^{\lambda_{i} - \lambda_{j} - 1} \frac{1 - q^{-m} k^{2(j-i)}}{1 - q^{-m} k^{2(j-i+1)}} \right) k^{-\binom{N}{2}} P(k^{2}) C_{(e,w_{0})}(q^{\lambda} k^{-\delta}) T_{w_{0}}(q^{\lambda} k^{-\delta}) = 0$$

with

(7.11)
$$P(k^2) = \prod_{1 \le i < j \le N} \frac{1 - k^{2(j-i+1)}}{1 - k^{2(j-i)}}.$$

Proof. By (7.6) and Theorem 5.10, we have for $\lambda \in \Lambda$,

$$K_0(\lambda) = r_{\kappa} k^{\langle \delta, \lambda \rangle} \theta(\kappa)^{-N} K(q^{w_0(\lambda)} k^{\delta}) C_{(e,w_0)}(q^{\lambda} k^{-\delta}) T_{w_0},$$

with K given by (5.16) and r_{κ} given by (7.7). Substituting the explicit expressions for K and r_{κ} we get the desired expression.

The following consequence should be compared with the general expansion formula of $v_+ = \sum_{w \in S_N} k^{\ell(w)} T_w \in H_0$ in terms of the $\xi_w(\gamma)$ ($w \in S_N$); see [37, Lemma 2.27 (2)].

Corollary 7.10. The element $v_+ = \sum_{w \in S_N} k^{\ell(w)} T_w \in H_0$ can be written as

(7.12)
$$v_{+} = k^{-\binom{N}{2}} P(k^{2}) C_{(e,w_{0})}(k^{-\delta}) T_{w_{0}}$$
$$= \left(\prod_{1 \le i < j \le N} \frac{1}{1 - k^{2(i-j)}}\right) \eta_{k^{-\delta}}(T_{w_{0}}) \xi_{w_{0}}(k^{-\delta})$$

Proof. We have $v_+ = Q_0 = K_0(0)$, hence the previous corollary gives the first equality of (7.12). The second equality then follows from (7.9).

Applying the map χ_+ to the first line of (7.12) gives

$$\sum_{w \in S_N} k^{2\ell(w)} = P(k^2)$$

with $P(k^2)$ given by (7.11), which is a well-known product formula for the Poincaré series of S_N ; see [33, Cor. (2.5)].

7.4. Relation to symmetric self-dual Macdonald polynomials. In this subsection we collect various consequences of the previous subsections for the symmetric Laurent polynomials $\chi_+(Q_\lambda) \in \mathbb{C}[T]^{S_N}$ ($\lambda \in \Lambda$). We keep the generic conditions (7.1) on $k \in \mathbb{C}^{\times}$. We denote

$$E_{\lambda} := P(k^2)^{-1} \chi_+(Q_{\lambda}) \in \mathbb{C}[T]^{S_N}, \qquad \lambda \in \Lambda.$$

By Proposition, 7.4 we have

$$E_{\lambda}(t) = \sum_{\alpha \in Q_{+}} K_{\alpha}^{+}(\lambda) t^{\lambda - \alpha}$$

with $K^+_{\alpha}(\lambda) = P(k^2)^{-1}\chi_+(K_{\alpha}(\lambda)) \in \mathbb{C}$ all but finitely many zero, and with leading coefficient $K^+_0(\lambda) \neq 0$ by Corollary 7.9 and (7.1).

Theorem 7.11. The $E_{\lambda} \in \mathbb{C}[T]^{S_N}$ ($\lambda \in \Lambda$) are the symmetric self-dual Macdonald polynomials. In other words, the E_{λ} are the unique symmetric regular functions on T satisfying

(7.13)
$$L_p^x(E_\lambda) = p(q^{-\lambda}k^{\delta})E_\lambda \quad \forall p \in \mathbb{C}[T]^{S_N}$$

and $E_{\lambda}(k^{\delta}) = 1$ for all $\lambda \in \Lambda$.

Proof. By Proposition 6.22, $E_{\lambda} \in \mathbb{C}[T]^{S_N}$ satisfies (7.13). Since the S_N -orbits $S_N(q^{-\lambda}k^{\delta})$ ($\lambda \in \Lambda$) in T are pairwise different by (7.1), the eigenvalue equations (7.13) uniquely characterize $E_{\lambda} \in \mathbb{C}[T]^{S_N}$ up to a nonzero constant multiple. Now

$$E_{\lambda}(k^{\delta}) = P(k^2)^{-1}\chi_+(Q_{\lambda}(k^{\delta})) =$$

by Lemma 7.5, which fixes the normalization of the solution $E_{\lambda} \in \mathbb{C}[T]^{S_N}$ of (7.13) uniquely.

The duality property of Q_{λ} (see Proposition 7.6) immediately gives the well-known duality property of the Macdonald polynomials.

Corollary 7.12. The Macdonald polynomials E_{λ} ($\lambda \in \Lambda$) are self-dual, in the sense that

$$E_{\lambda}(q^{-\mu}k^{\delta}) = E_{\mu}(q^{-\lambda}k^{\delta})$$

for all $\lambda, \mu \in \Lambda$.

Remark 7.13. The self-duality of (the suitably normalized) Macdonald polynomials was initially proved by Koornwinder using Pieri formulas in an unpublished manuscript (the argument is reproduced in [32, VI (6.6)]). Cherednik ([9, Thm. 1.4.6] and [8, Thm. 3.2]) reproduced the self-duality of the Macdonald polynomials using the anti-involution * (see Theorem 2.8) on the double affine Hecke algebra.

We also immediately reobtain the well-known evaluation formula for the symmetric Macdonald polynomials; see [32, VI (6.11)] (the parameters (n, q, t) in [32, Chpt. VI] correspond to (N, q^{-1}, k^2) in our notations).

Corollary 7.14. For $\lambda \in \Lambda$ let $P_{\lambda} := K_0^+(\lambda)^{-1}E_{\lambda} \in \mathbb{C}[T]^{S_N}$ be the monic symmetric Macdonald polynomial of degree λ . Then

(7.14)
$$P_{\lambda}(k^{\delta}) = k^{-\langle \delta, \lambda \rangle} \prod_{1 \le i < j \le N} \prod_{m=0}^{\lambda_i - \lambda_j - 1} \frac{1 - q^{-m} k^{2(j-i+1)}}{1 - q^{-m} k^{2(j-i)}}.$$

Proof. By the previous theorem we have $P_{\lambda}(k^{\delta}) = K_0^+(\lambda)^{-1}$. Corollary 7.9 gives

$$K_0^+(\lambda) = k^{\langle \delta, \lambda \rangle} \prod_{1 \le i < j \le N} \prod_{m=0}^{\lambda_i - \lambda_j - 1} \frac{1 - q^{-m} k^{2(j-i)}}{1 - q^{-m} k^{2(j-i+1)}},$$

which implies the desired result.

8. Appendix on holonomic systems of q-difference equations

In the appendix we detail the construction of power series solutions of holonomic systems of q-difference equations. Special cases have been investigated in, e.g., [2], [20] and [14, §12]. Many arguments go back to classical works [1], [3], [4], [48] on ordinary linear q-difference equations.

We begin with the construction of formal asymptotic solutions to holonomic systems of q-difference equations. Let $\mathbb{C}[[z]] = \mathbb{C}[[z_1, \ldots, z_M]]$ denote the ring of formal power series in M indeterminates z_1, \ldots, z_M over the complex numbers. Let V be a finite-dimensional complex vector space and let

$$A_i \in \mathbb{C}[[z]] \otimes \operatorname{End}(V)$$

for i = 1, ..., M. Since $\mathbb{C}[[z]] \otimes \operatorname{End}(V)$ is isomorphic to $\operatorname{End}_{\mathbb{C}[[z]]}(\mathbb{C}[[z]] \otimes V)$ as $\mathbb{C}[[z]]$ -module, we can view the A_i as $\mathbb{C}[[z]]$ -linear endomorphisms of $\mathbb{C}[[z]] \otimes V$. Fix $0 < q_i < 1$ for $1 \le i \le M$. Define the q_i -dilation operators

$$\mathcal{T}_i \colon \mathbb{C}[[z]] \to \mathbb{C}[[z]]$$

for $i = 1, \ldots, M$ as the complex linear maps

$$\mathcal{T}_i(\sum_{\mathbf{m}} d_{\mathbf{m}} z^{\mathbf{m}}) := \sum_{\mathbf{m}} q_i^{m_i} d_{\mathbf{m}} z^{\mathbf{m}} \qquad (d_{\mathbf{m}} \in \mathbb{C}),$$

where we use multi-index notation $z^{\mathbf{m}} = z_1^{m_1} \cdots z_M^{m_M}$ for $\mathbf{m} = (m_1, \ldots, m_M)$ with $m_j \in \mathbb{Z}_{\geq 0}$. We also view \mathcal{T}_i as operators on $\mathbb{C}[[z]] \otimes V$ and on $\mathbb{C}[[z]] \otimes \text{End}(V)$. Consider the system of first-order linear q-difference equations

(8.1)
$$A_i \mathcal{T}_i f = f, \qquad (i = 1, \dots, M)$$

for $f \in \mathbb{C}[[z]] \otimes V$.

For $f \in \mathbb{C}[[z]] \otimes V$ and $A \in \mathbb{C}[[z]] \otimes \text{End}(V)$, we introduce the notations

$$f^{(m)} := f|_{z_{m+1}=...=z_M=0} \in \mathbb{C}[[z_1,...,z_m]] \otimes V$$
$$A^{(m)} := A|_{z_{m+1}=...=z_M=0} \in \mathbb{C}[[z_1,...,z_m]] \otimes \text{End}(V)$$

for $0 \le m \le M$, with the convention that $f^{(M)} = f$ and $A^{(M)} = A$. We make the following assumptions on the system of q-difference equations (8.1):

(a) The system (8.1) is holonomic, that is

(8.2)
$$A_i \mathcal{T}_i(A_j) = A_j \mathcal{T}_j(A_i)$$

for all $1 \leq i, j \leq M$. Note that the holonomy implies that the leading coefficients $A_i^{(0)} \in \text{End}(V)$ of A_i mutually commute, i.e.,

$$[A_i^{(0)}, A_j^{(0)}] = 0$$

for all $1 \leq i, j \leq M$.

(b) The complex linear endomorphisms $A_1^{(0)}, \ldots, A_M^{(0)}$ of V are semisimple. Combined with (a) we thus have

$$V = \bigoplus_{\gamma \in S} V[\gamma]$$

with $V[\gamma] := \{ v \in V \mid A_i^{(0)}v = \gamma_i v \ \forall i \} \ (\gamma \in \mathbb{C}^M) \text{ and } S := \{ \gamma \in \mathbb{C}^M \mid V[\gamma] \neq \{0\} \}.$ (c) $(1^M) := (1, \dots, 1) \in \mathbb{C}^M$ belongs to S. (d) $\gamma_k \notin q_k^{-\mathbb{N}}$ for all $\gamma \in S$ and $1 \le k \le M$.

Proposition 8.1. Fix $v \in V[(1^M)]$. Consider the system (8.1) of q-difference equations and suppose that (a)-(d) are satisfied. Then there exists a unique solution $\Phi_v \in \mathbb{C}[[z]] \otimes V$ of (8.1) such that

$$\Phi_{v}^{(0)} = v.$$

Proof. The proposition is a consequence of the following lemma.

Lemma 8.2. Let $0 \le m \le M$. Suppose one has a solution

$$f_m \in \mathbb{C}[[z_1, \dots, z_m]] \otimes V$$

of the system of equations

(8.3)
$$\begin{aligned} A_r^{(m)} \mathcal{T}_r f_m &= f_m, \qquad 1 \le r \le m, \\ A_s^{(m)} f_m &= f_m, \qquad m < s \le M. \end{aligned}$$

Then there exists a unique

$$f_{m+1} = \sum_{n \ge 0} f_{m;n} z_{m+1}^n \in \mathbb{C}[[z_1, \dots, z_{m+1}]] \otimes V$$

with $f_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $f_{m;0} = f_m$ satisfying (8.3) with the role of m replaced by m + 1:

(8.4)
$$A_r^{(m+1)} \mathcal{T}_r f_{m+1} = f_{m+1}, \qquad 1 \le r \le m+1, \\ A_s^{(m+1)} f_{m+1} = f_{m+1}, \qquad m+1 < s \le M.$$

The proposition follows directly from the lemma as follows. Note that $f_0 := v \in$ $V[(1^M)]$ is a solution of (8.3) for m = 0. The formal V-valued series $f_M \in \mathbb{C}[[z]] \otimes V$, obtained by repeated application of the lemma starting from $f_0 = v$, gives a formal V-valued series solution of (8.1) satisfying $f_M^{(0)} = v$. For uniqueness, assume that $f \in \mathbb{C}[[z]] \otimes V$ is another formal V-valued series satisfying $f^{(0)} = v$ and solving (8.1). We have $f^{(0)} = v = f_0$ and $f^{(m)}$ solves (8.3) for all $0 \le m < M$. Hence, by the uniqueness part of the lemma, $f = f^{(M)} = f_M$.

We now proceed to prove the lemma. We assume that we have a formal power series solution $f_m \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ of (8.3) for some $0 \leq m < M$. We write

(8.5)
$$A_r^{(m+1)} = \sum_{n \ge 0} A_{r;n}^{(m)} z_{m+1}^n,$$

where $A_{r;n}^{(m)} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes \operatorname{End}(V)$ and $A_{r;0}^{(m)} = A_r^{(m)}$. By a direct computation one verifies that

$$f_{m+1} = \sum_{n \ge 0} f_{m;n} z_{m+1}^n \in \mathbb{C}[[z_1, \dots, z_{m+1}]] \otimes V$$

with $f_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $f_{m;0} = f_m$ satisfies the q-difference equation

$$A_{m+1}^{(m+1)}\mathcal{T}_{m+1}f_{m+1} = f_{m+1}$$

if and only if

(8.6)
$$(1 - q_{m+1}^n A_{m+1}^{(m)}) f_{m;n} = \sum_{l=1}^n q_{m+1}^{n-l} A_{m+1;l}^{(m)} f_{m;n-l}$$

for all $n \in \mathbb{Z}_{\geq 0}$. The recurrence relations (8.6) admit a unique solution $(f_{m;n})_{n \in \mathbb{Z}_{\geq 0}}$ with $f_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and with initial condition $f_{m;0} = f_m$. Indeed, note that (8.6) is valid for n = 0 since $f_{m;0} = f_m$ satisfies (8.3). For $n \geq 1$, we have

$$\det\left(1-q_{m+1}^{n}A_{m+1}^{(m)}\right)\in\mathbb{C}[[z_{1},\ldots,z_{m}]]^{\times}$$

since

$$\det\left(1-q_{m+1}^n A_{m+1}^{(m)}\right)|_{z_1=\ldots=z_m=0} = \det\left(1-q_{m+1}^n A_{m+1}^{(0)}\right) = \prod_{\gamma \in S} (1-q_{m+1}^n \gamma_{m+1}^{\dim(V[\gamma])}) \neq 0$$

by assumption (d). Cramer's rule then implies that (8.6) admits a unique solution $(f_{m;n})_{n \in \mathbb{Z}_{\geq 0}}$ with $f_{m;0} = f_m$.

We conclude that there exists a unique

$$f_{m+1} = \sum_{n \ge 0} f_{m;n} z_{m+1}^n$$

with $f_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $f_{m;0} = f_m$ satisfying the q-difference equation (8.7)

It remains to show that f_{m+1} also satisfies (8.4) for $r = 1, \ldots, m$ and for $s = m+2, \ldots, M$.

Fix $1 \leq r \leq m$ and write $g_r := A_r^{(m+1)} \mathcal{T}_r f_{m+1}$. Its expansion in powers of z_{m+1} is written as

$$g_r = \sum_{n \ge 0} g_{r;n} z_{m+1}^n$$

with $g_{r;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $g_{r;0} = A_r^{(m)} \mathcal{T}_r f_m = f_m$, where the last equality follows from the fact that f_m is assumed to satisfy (8.3). Furthermore, using the holonomy (8.2) and the q-difference equation (8.7) in z_{m+1} satisfied by f_{m+1} , we have

$$A_{m+1}^{(m+1)} \mathcal{T}_{m+1} g_r = A_{m+1}^{(m+1)} \mathcal{T}_{m+1} (A_r^{(m+1)}) \mathcal{T}_r \mathcal{T}_{m+1} f_{m+1}$$

= $A_r^{(m+1)} \mathcal{T}_r (A_{m+1}^{(m+1)}) \mathcal{T}_r \mathcal{T}_{m+1} f_{m+1}$
= $A_r^{(m+1)} \mathcal{T}_r (A_{m+1}^{(m+1)} \mathcal{T}_{m+1} f_{m+1})$
= $A_r^{(m+1)} \mathcal{T}_r f_{m+1} = g_r.$

We conclude that g_r satisfies the characterizing properties of f_{m+1} . Hence $g_r = f_{m+1}$, i.e.,

$$A_r^{(m+1)}\mathcal{T}_r f_{m+1} = f_{m+1}.$$

Fix $m+1 < s \leq M$ and write $g_s := A_s^{(m+1)} f_{m+1}$. By a similar argument as used in the previous paragraph, we now show that $g_s = f_{m+1}$. We write

$$g_s = \sum_{n \ge 0} g_{s;n} z_{m+1}^n$$

with $g_{s;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $g_{s;0} = A_s^{(m)} f_m = f_m$, where the last equality follows by the assumption that f_m satisfies (8.3). Using the holonomy (8.2), the *q*-difference equation (8.7), and the obvious fact that $\mathcal{T}_s(A_{m+1}^{(m+1)}) = A_{m+1}^{(m+1)}$ since s > m+1, we have

$$A_{m+1}^{(m+1)} \mathcal{T}_{m+1} g_s = A_{m+1}^{(m+1)} \mathcal{T}_{m+1} (A_s^{(m+1)}) \mathcal{T}_{m+1} f_{m+1}$$

= $A_s^{(m+1)} \mathcal{T}_s (A_{m+1}^{(m+1)}) \mathcal{T}_{m+1} f_{m+1}$
= $A_s^{(m+1)} A_{m+1}^{(m+1)} \mathcal{T}_{m+1} f_{m+1}$
= $A_s^{(m+1)} f_{m+1} = g_s.$

We conclude that g_s satisfies the characterizing properties of f_{m+1} . Hence $g_s = f_{m+1}$, i.e.

$$A_s^{(m+1)}f_{m+1} = f_{m+1}$$

This completes the proof of Lemma 8.2, and hence the proof of Proposition 8.1. \Box

We investigate the analytical properties of the solution Φ_v when the *q*-connection matrices A_i $(1 \leq i \leq M)$ satisfy, besides the conditions (a)-(d), the following analyticity condition:

(e) For some $\epsilon > 0$ the formal $\operatorname{End}(V)$ -valued series $A_i \in \mathbb{C}[[z]] \otimes \operatorname{End}(V)$ $(1 \leq i \leq M)$ converges normally on compact of the open polydisc $D_{\epsilon}^M := \{z \in \mathbb{C}^M \mid |z_i| < \epsilon \forall i\}.$

In other words, if we expand A_i along a basis of $\operatorname{End}(V)$, condition (e) requires its coefficients in $\mathbb{C}[[z]]$ to converge normally on compact of D_{ϵ}^M .

Proposition 8.3. Suppose that the q-connection matrices $A_i \in \mathbb{C}[[z]] \otimes \text{End}(V)$ $(1 \leq i \leq M)$ satisfy (a)-(e). Let $v \in V[(1^M)]$. There exists an $\epsilon > 0$ such that the formal V-valued series $\Phi_v \in \mathbb{C}[[z]] \otimes V$ converges normally on compact of D_{ϵ}^M .

Proof. For ease of notation, we will write Φ instead of Φ_v . By induction on $m = 0, \ldots, M$ we prove that there exists $\epsilon > 0$ such that $\Phi^{(m)} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ converges normally on compact of D_{ϵ}^m .

For m = 0, there is nothing to prove. Fix $0 \le m < M$ and suppose $\Phi^{(m)}$ converges normally on compact of D^m_{δ} for some $\delta > 0$. Write

$$\Phi^{(m+1)} = \sum_{n \ge 0} \Phi_{m;n} z_{m+1}^n$$

with $\Phi_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ and $\Phi_{m;0} = \Phi^{(m)}$. Recall from the proof of Lemma 8.2 that the formal V-valued power series $\Phi_{m;n}$ $(n \ge 1)$ are unique characterized by the recurrence relations

(8.8)
$$\Phi_{m;n} = \sum_{l=1}^{n} q_{m+1}^{n-l} \left(1 - q_{m+1}^{n} A_{m+1}^{(m)} \right)^{-1} A_{m+1;l}^{(m)} \Phi_{m;n-l}$$

for all $n \geq 1$. We use this recurrence formula to find bounds for $\Phi_{m;n}$ in a neighborhood of $0 \in \mathbb{C}^m$.

Turn the finite-dimensional complex vector space V into an inner product space, with corresponding norm denoted by $\|\cdot\|$. We also write $\|\cdot\|$ for the operator norm of the associated finite-dimensional normed space $\operatorname{End}(V)$. We continue the proof of the proposition with two technical sublemmas. First we find a proper uniform bound for $A_{m+1:l}^{(m)}$ for all l (see (8.8)).

Lemma 8.4. There exists $\epsilon > 0$ and M > 0 such that $||A_{m+1;l}^{(m)}|| \leq M\epsilon^{-l}$ on $\overline{D}_{\epsilon}^{m}$ for all $l \geq 0$.

Proof. By (e) there exists an $\epsilon > 0$ such that $A_{m+1}^{(m+1)} \in \mathbb{C}[[z_1, \ldots, z_{m+1}]] \otimes \operatorname{End}(V)$ converges normally on compacta of the polydisc $D_{2\epsilon}^{m+1}$. Consequently, for $\epsilon < \epsilon' < 2\epsilon$ we have that $||A_{m+1}^{(m+1)}||$ is uniformly bounded on the polydisc $D_{\epsilon'}^{m+1}$, say by M > 0. In particular, we get

$$\|(\partial_{z_{m+1}}^l A_{m+1}^{(m+1)})(z_1, \dots, z_{m+1})|_{z_{m+1}=0}\| \le M\epsilon^{-l}l!$$

for all $(z_1, \ldots, z_m) \in \overline{D}_{\epsilon}^m$ and for all $l \ge 0$ (see, e.g., [23] Theorem 2.2.7). This proves the lemma in view of the definition (8.5) of $A_{m+1;l}^{(m)}$.

Lemma 8.5. There exists an $\epsilon > 0$ such that $\Phi_{m;n} \in \mathbb{C}[[z_1, \ldots, z_m]] \otimes V$ converges normally on compact of D_{ϵ}^m for all $n \geq 0$. Furthermore, there exists a constant C > 0 (independent of n) such that

$$\|\Phi_{m;n}\| \le \frac{C}{1+C} \left(\frac{1+C}{q_{m+1}\epsilon}\right)^n \|\Phi^{(m)}\|$$

on D_{ϵ}^m for all $n \geq 1$.

Proof. In the proof of this lemma, we write q instead of q_{m+1} . By assumption, $\Phi_{m;0} = \Phi^{(m)}$ converges normally on compact of D_{ϵ}^m if $0 < \epsilon < \delta$. We now use the recurrence relation (8.8) to obtain the desired results for $\Phi_{m;n}$ with $n \ge 1$.

By the proof of Lemma 8.2 and since 0 < q < 1, there exists some $\epsilon > 0$ (independent of $n \ge 1$) such that $\det(1 - q^n A_{m+1}^{(m)})^{-1}$ is analytic on D_{ϵ}^m for all $n \ge 1$ and such that $|\det(1 - q^n A_{m+1}^{(m)})^{-1}|$ is bounded on the closure $\overline{D}_{\epsilon}^m$ of D_{ϵ}^m , with bound independent of $n \ge 1$. For such ϵ , it follows from (8.8) that $\Phi_{m;n}$ converges normally on compacta of D_{ϵ}^m for all $n \ge 1$. Furthermore, by (e), 0 < q < 1, and Cramer's rule, it implies that for $\epsilon > 0$ small enough,

$$\|(1 - q^n A_{m+1}^{(m)})^{-1}\| \le C'$$

on $\overline{D}_{\epsilon}^{m}$ for all $n \geq 1$, with C' > 0 also independent of n. By (8.8), 0 < q < 1 and the previous lemma, we thus obtain for $\epsilon > 0$ small enough,

(8.9)
$$\|\Phi_{m;n}\| \le C' \sum_{l=1}^{n} q^{n-l} \|A_{m+1;l}^{(m)}\| \|\Phi_{m;n-l}\| \le C \sum_{l=1}^{n} \left(\frac{1}{q\epsilon}\right)^{l} \|\Phi_{m;n-l}\|$$

on $\overline{D}_{\epsilon}^m$ for all $n \ge 1$ with the constant C = C'M > 0 (independent of n).

Now, we have the following claim (cf. [14] §10.6): the recurrence relation

$$g_n = C \sum_{l=1}^n \left(\frac{1}{q\epsilon}\right)^l g_{n-l}, \quad (n>0)$$

with $g_0 \in \mathbb{R}$ fixed is uniquely solved by

$$g_n = \frac{C}{C+1} \left(\frac{1+C}{q\epsilon}\right)^n g_0$$

for $n \ge 1$. Being obvious for n = 1, the claim follows using induction for n > 1 by

$$g_n = C\left(\frac{1}{q\epsilon}\right)^n g_0 + C\sum_{l=1}^{n-1} \left(\frac{1}{q\epsilon}\right)^l g_{n-l}$$

$$= C\left(\frac{1}{q\epsilon}\right)^n g_0 + \frac{C^2}{C+1} \sum_{l=1}^{n-1} \left(\frac{1}{q\epsilon}\right)^l \left(\frac{(1+C)}{q\epsilon}\right)^{n-l} g_0$$

$$= C\left(\frac{1}{q\epsilon}\right)^n g_0 \left(1 + C\sum_{l=0}^{n-2} (1+C)^l\right)$$

$$= C\left(\frac{1}{q\epsilon}\right)^n g_0 \left(1 + C\left(\frac{(1+C)^{n-1}-1}{1+C-1}\right)\right)$$

$$= \frac{C}{C+1} \left(\frac{1+C}{q\epsilon}\right)^n g_0.$$

Combined with (8.9), the lemma now follows immediately.

To conclude the proof of the proposition, note that the previous lemma shows that

$$\Phi^{(m+1)} = \sum_{n \ge 0} \Phi_{m;n} z_{m+1}^n \in \mathbb{C}[[z_1, \dots, z_{m+1}]] \otimes \operatorname{End}(V)$$

converges normally on compact of $D_{\epsilon'}^{m+1}$ if we take $\epsilon' > 0$ sufficiently small. This concludes the proof of the induction step.

We interpret the q-dilation operators \mathcal{T}_i as automorphisms of $\mathcal{M}(\mathbb{C}^M)$ by

 $(\mathcal{T}_i f)(z) = f(z_1, \ldots, z_{i-1}, q_i z_i, z_{i+1}, \ldots, z_M).$

Theorem 8.6. Suppose $A_i \in \mathcal{M}(\mathbb{C}^M) \otimes \operatorname{End}(V)$ $(1 \leq i \leq M)$ satisfy the holonomy conditions (8.2) as meromorphic $\operatorname{End}(V)$ -valued functions on \mathbb{C}^M . Suppose that the A_i are analytic at $0 \in \mathbb{C}^M$ and that their power series expansions at $0 \in \mathbb{C}^M$ satisfy the conditions (b)-(d).

Let $v \in V[(1^M)]$. There exists a unique $\Phi_v \in \mathcal{M}(\mathbb{C}^M) \otimes V$ solving the holonomic system (8.1) of q-difference equations and coinciding, in a small neighborhood of $0 \in \mathbb{C}^M$, with the converging V-valued power series solution Φ_v from Proposition 8.3.

Proof. Since the A_i are assumed to be analytic at $0 \in \mathbb{C}^M$, their power series expansions at $0 \in \mathbb{C}^M$ are converging normally on compact of some open polydisc D^M_{ϵ} ($\epsilon > 0$). Hence, condition (e) is automatically satisfied.

Let $\Phi_v \in \mathbb{C}[[z]] \otimes V$ be the power series solution from Proposition 8.3 and let $\epsilon > 0$ such that Φ_v converges normally on compacta of D_{ϵ}^M . Let $z' \in \mathbb{C}^M$ and $U \subset \mathbb{C}^M$ some open locally compact neighborhood of z'. Since $0 < q_i < 1$ $(1 \le i \le M)$, there exists a $\lambda \in \mathbb{Z}_{\geq 0}^M$ such that $q^{\lambda}U \subset D_{\epsilon}^M$, where $q^{\lambda}z = (q_1^{\lambda_1}z_1, \ldots, q_M^{\lambda_M}z_M)$. Define Φ_v as V-valued meromorphic function on $z \in U$ by

(8.10)
$$\Phi_v(z) = A_\lambda(z)\Phi_v(q^\lambda z),$$

where $A_{\lambda} \in \mathcal{M}(\mathbb{C}^M) \otimes \operatorname{End}(V)$ is defined inductively by

$$A_{\lambda+\mu}(z) = A_{\lambda}(z)A_{\mu}(q^{\lambda}z), \qquad \forall \lambda, \mu \in \mathbb{Z}_{\geq 0}^{M},$$

and $A_{\epsilon_i} = A_i \ (1 \le i \le M)$, where the $\epsilon_i \ (1 \le i \le M)$ are the standard generators of the additive monoid $\mathbb{Z}_{\ge 0}^M$. Of course, the definition of $A_{\lambda}(z)$ makes sense by the holonomy conditions for the A_i . Furthermore, (8.10) together with the holonomy conditions for the A_i show that the power series solution Φ_v of (8.2) has a unique extension to a meromorphic V-valued solution on \mathbb{C}^M of (8.2).

References

- [1] C.R. Adams, On the linear ordinary q-difference equation, Ann. Math. 30 (1929), 195–205.
- K. Aomoto, A note on holonomic q-difference systems, in: "Algebraic analysis", Vol. I, 25–28, Academic Press, Boston, MA, 1988.
- [3] G.D. Birkhoff, The generalized Riemann problem for linear differential equations and the allied problems for linear difference and q-difference equations, Proc. Amer. Acad. 49 (1913), 521–568.
- [4] R.D. Carmichael, The general theory of linear q-difference equations, Amer. J. Math. 34 (1912), 147–168.
- [5] O. Chalykh, Macdonald polynomials and algebraic integrability, Adv. Math. 166 (2002), no. 2, 193–259.
- [6] I. Cherednik, Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald's operators, Int. Math. Res. Not. 1992, No. 9, 171–180.
- [7] I. Cherednik, Induced representations of double affine Hecke algebras and applications, Math. Res. Lett. 1 (1994), 319–337.
- [8] I. Cherednik, Macdonald's evaluation conjectures and difference Fourier transform, Invent. Math. 122 (1995), 119–145.
- [9] I. Cherednik, *Double affine Hecke algebras*, Cambridge University Press, London Math. Soc. Lecture Note Series 319, 2005.
- [10] I. Cherednik, Whittaker limits of difference spherical functions, Int. Math. Res. Not. 2009, No. 20, 3793–842.
- [11] P. Di Francesco, P. Zinn-Justin, Quantum Knizhnik-Zamolodchikov equation, generalized Razumov-Stroganov sum rules and extended Joseph polynomials, J. Phys. A 38 (2005), L815– L822.
- [12] J.J. Duistermaat, F.A. Grünbaum, Differential equations in the spectral parameter, Comm. Math. Phys. 103 (1986), no. 2, 177–240.
- P.I. Etingof, Difference equations with elliptic coefficients and quantum affine algebras (1993), arXiv:hep-th/9312057.
- [14] P.I. Etingof, I.B. Frenkel, A.A. Kirillov Jr., Lectures on Representation Theory and Knizhnik-Zamolodchikov Equations, Math. Surveys and Monographs, Vol. 58, Amer. Math. Soc., 1998.
- [15] P. Etingof, A.A. Kirillov Jr., Macdonald's polynomials and representations of quantum groups, Math. Res. Lett. 1 (1994), no. 3, 279–296.
- [16] P. Etingof, A.A. Kirillov Jr., Representation-theoretic proof of the inner product and symmetry identities for Macdonald's polynomials, Compositio Math. 102 (1996), no. 2, 179–202.
- [17] P. Etingof, K. Styrkas, Algebraic integrability of Macdonald polynomials and representations of quantum groups, Compositio Math. 114 (1998), 125–152.
- [18] P. Etingof, A. Varchenko, Dynamical Weyl groups and applications, Adv. Math. 167 (2002), no. 1, 74–127.
- [19] G. Felder, Ya. Markov, V. Tarasov, A. Varchenko, Differential equations compatible with KZ equations, Math. Phys. Anal. Geom. 3 (2000), 139–177.
- [20] I.B. Frenkel, N.Yu. Reshetikhin, Quantum affine algebras and holonomic difference equations, Comm. Math. Phys. 146 (1992), no. 1, 1–60.
- [21] F.A. Grünbaum, L. Haine, Some functions that generalize the Askey-Wilson polynomials, Comm. Math. Phys. 184 (1997), no. 1, 173–202.
- [22] J. Harnad, Quantum Isomonodromic deformations and the Knizhnik-Zamolodchikov equations. In: "Symmetries and integrability of difference equations" (Estérel, PQ, 1994), 155– 161, CRM Proc. Lecture Notes, 9, Amser. Math. Soc., Providence, RI, 1996.
- [23] L. Hörmander, An Introduction to Complex Analysis in Several Variables. Second revised edition. North-Holland Mathematical Library, Vol. 7. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [24] M. Jimbo, T. Miwa, Algebraic analysis of solvable lattice models. CBMS Regional Cinference Series in Mathematics 85. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1995.

- [25] M. Kasatani, Y. Takeyama, The quantum Knizhnik-Zamolodchikov equation and nonsymmetric Macdonald polynomials, Funkcialaj Ekvacioj 50 (2007), 491–509.
- [26] S. Kato, R-matric arising from affine Hecke algebras and its applications to Macdonald's difference operators, Comm. Math. Phys. 165 (1994), no. 3, 533-553.
- [27] A. Kazarnovski-Krol, Matrix elements of vertes operators of the deformed WA_n-algebra and the Harish-Chandra solutions to Macdonald's difference equations, Selecta Math. (N.S.) 5 (1999), no. 2, 257–301.
- [28] A.N. Kirillov, M. Noumi, Affine Hecke algebras and raising operators for Macdonald polynomials, Duke Math. J. 93 (1998), no. 1, 1–39.
- [29] V.P. Leksin, Isomonodromic and bispectral properties of the generalized Knizhnik-Zamolodchikov equations, J. Math. Sci. (N.Y.) 129 (2005), no. 1, 3635–3641.
- [30] G. Lusztig, Affine Hecke Algebras and their Graded Version, J. Amer. Math. Soc. 2 (1989), no. 3, 599–635.
- [31] G. Letzter, J.V. Stokman, Macdonald difference operators and Harish-Chandra series, Proc. Lond. Math. Soc. 97 (2008), no. 1, 60–96.
- [32] I.G. Macdonald, Symmetric functions and Hall polynomials. Second edition, Oxford Math. Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995.
- [33] I.G. Macdonald, The Poincaré series of a Coxeter group, Math. Ann. 199 (1972), 161–174.
- [34] I.G. Macdonald, Affine Hecke algebras and orthogonal polynomials, Cambridge Tracts in Mathematics 157, Cambridge University Press, Cambridge, 2003.
- [35] A. Matsuo, Integrable connections related to zonal spherical functions, Invent. Math. 110 (1992), 96–121.
- [36] K. Mimachi, A solution to quantum Knizhnik-Zamolodchikov equations and its applications to eigenvalue problems of the Macdonald type, Duke Math. J. 85 (1996), no. 3, 635–658.
- [37] E.M. Opdam, A generating function for the trace of the Iwahori-Hecke algebra, in: Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), 301–323, Progr. Math., 210, Birkhäuser Boston, Boston, MA, 2003.
- [38] N. Reshetikhin, The Knizhnik-Zamolodchikov system as a deformation of the isomonodromy problem, Lett. Math. Phys. 26 (1992), 167–177.
- [39] S.N.M. Ruijsenaars, Complete integrability of relativistic Calogero-Moser systems and elliptic function identities, Comm. Math. Phys. 110 (1987), 191–213.
- [40] S.N.M. Ruijsenaars, Factorized weight functions vs. factorized scattering, Comm. Math. Phys. 228 (2002), no. 3, 467–494.
- [41] J. Sauloy, Systemes aux q-différences singuliers réguliers: classification, matrice de connexion et monodromie, Ann. Inst. Fourier (Grenoble) 50 (2000), no. 4, 1021–1071.
- [42] F. Smirnov, General formula for soliton form factors in sine-Gordon model, J. Phys. A 19 (1986), no. 10, L575–L578.
- [43] J.-P. Serre, Local Fields, Graduate Texts in Mathematics 67, Springer-Verlag, New York-Berlin, 1979.
- [44] V. Tarasov, Duality for Knizhnik-Zamolodchikov and dynamical equations, and hypergeometric integrals, in "Infinite dimensional algebras and quantum integrable systems", 235–263, Prog. Math., 237, Birkhäuser, Basel, 2005.
- [45] V. Tarasov, A. Varchenko, Difference equations compatible with trigonometric KZ differential equations, Int. Math. Res. Not. 2000, no. 15, 801–829.
- [46] V. Tarasov, A. Varchenko, Duality for Knizhnik-Zamolodchikov and dynamical equations. The 2000 Twente conference on Lie groups (Enschede) Acta Appl. Math. 73 (2002), no. 1-2, 141–154.
- [47] V. Toledano Laredo, A Kohno-Drinfeld theorem for quantum Weyl groups, Duke Math. J. 112 (2002), no. 3, 421–451.
- [48] W.J. Trjitzinsky, Analytic theory of linear q-difference equations, Acta Math. 61 (1933), 1–38.

KdV Institute for Mathematics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands.

E-mail address: m.vanmeer@uva.nl, j.v.stokman@uva.nl