

Howes et al. reply: Matthaeus, Servidio, and Dmitruk (hereafter MSD) [1] comment on our Letter [2], arguing that our resolution is insufficient to claim (i) that our gyrokinetic (GK) simulations capture simultaneously the MHD and kinetic sub-ion-gyroscale plasma turbulent cascades. Our spatial resolution ($64 \times 64 \times 128$) is insufficient to determine converged values of spectral indices—a claim we do not make—but is sufficient to capture the nonlinear energy transfer from large to small scales, see a qualitative transition of electric and magnetic energy spectra at the ion gyroscale, and yield results consistent with theory above and below this scale (see Figs. 2 and 3 of [2]). MSD speculate that the low-wave-number spectrum in our simulations is imposed by the driving. The driving occurs at the lowest mode numbers $(k_x, k_y)L_\perp/(2\pi) = (1, 0)$, $(0, 1)$, and $(-1, 0)$; higher-wave-number amplitudes are determined by nonlinear interactions.

Our Letter presents the first kinetic simulation showing that the spectra of anisotropic turbulence in a weakly collisional plasma are qualitatively consistent with measurements in the solar wind (SW) [3], a result not *a priori* obvious ([4] addresses a different problem). This first step in tackling SW turbulence at kinetic scales suggests that GK modeling is a promising line of inquiry.

MSD allege that we claim (ii) that dissipation is required to achieve our result. This apparently arises from a misunderstanding of the nature of kinetic Alfvén waves (KAW), not a dissipative effect but dispersive waves weakly Landau-damped at $k_\perp \rho_i > 1$. We emphasize that damping is small for the parameters used in our simulation (see Fig. 1 of [2]), so the spectrum attributed to KAW turbulence is near-dissipationless (agreeing with suggestions in Refs. [3, 5], which MSD incorrectly claim to contradict our results).

Although GK cannot describe all features of SW turbulence (as we acknowledge [6]), it is a rigorous approximation for small-scale anisotropic low-frequency fluctuations (see [7] and references therein). The anisotropy assumption is based on a wealth of theoretical, numerical, and observational evidence (see [2]).

While questioning the validity of GK, MSD appear to suggest that Hall MHD (HMHD) is a preferable model for SW turbulence. Much has been learned from this temptingly simple model [8]. The fully nonlinear anisotropic limit ($k_\parallel \ll k_\perp$) of HMHD can be derived from GK in the limit $\beta_i \ll 1$ and $\beta_e \sim 1$ (see Appendix E of Ref. [9]). The transition from the MHD limit to KAW occurs at $l_t \sim d_i/\sqrt{1+2/\beta_e}$, where $d_i = \rho_i/\sqrt{\beta_i}$ is the ion inertial scale. For $l_t \gg \rho_i$ (enabling a fluid approximation), we must have $\beta_i \ll 1$ and $T_i \ll T_e$ —it is well known from kinetic theory [10] that this *cold ion* limit is the only one in which HMHD is strictly valid, a limit not universally

applicable to the SW. Thus, neither the transition to the dispersive waves nor the effects of collisionless damping (not captured by any fluid-like limit) are correctly described by HMHD in the SW parameter regime.

While it is true that the transition to dispersive waves in HMHD causes flatter electric and steeper magnetic energy spectra, it is quite unclear whether the simulations MSD present are directly applicable to the SW (much more unclear, we believe, than in the case of GK). Without a kinetic simulation to assess the importance of finite-ion-gyroscale effects and collisionless damping, a numerical HMHD simulation may be inconsistent with the underlying kinetic plasma physics, little more than an *ad hoc* model. It is possible to get what appears to be the right answer for the wrong reason, as demonstrated by the 1-D HMHD result shown by MSD: surely they are not claiming that SW turbulence is accurately modeled as a 1-D cascade in k_\parallel ! That qualitatively correct spectra seem to emerge in such studies does not invalidate the much more general GK approach or remove the need for kinetic simulations.

Obviously, the success of GK in qualitatively reproducing the observed spectra does not in itself prove its applicability in space. We make no such claim and accept that further analytical and numerical work is needed. In presenting the first numerical results of our long-term project on kinetic turbulence, we phrased our conclusions carefully. The hyperbolic claims attributed to us by MSD—*e.g.*, that we prove the applicability of GK to the SW, or show that the spectra reported in Ref. [3] require dissipation—are an incorrect representation of our work.

G. G. Howes, S. C. Cowley, W. Dorland, G. W. Hammett, E. Quataert, A. A. Schekochihin and T. Tatsuno

-
- [1] W. H. Matthaeus, S. Servidio, and P. Dmitruk, preceding Comment, Phys. Rev. Lett. **101**, LRK1046 (2008).
 - [2] G. G. Howes *et al.*, Phys. Rev. Lett. **100**, 065004 (2008)
 - [3] S. D. Bale *et al.*, Phys. Rev. Lett. **94**, 215002 (2005)
 - [4] K. Bowers and H. Li, Phys. Rev. Lett. **98**, 035002 (2007)
 - [5] O. Alexandrova *et al.*, Planet. Space Sci. **55**, 2224 (2007)
 - [6] G. G. Howes *et al.*, J. Geophys. Res. **113**, 5103 (2008)
 - [7] G. G. Howes *et al.*, Astrophys. J. **651**, 590 (2006)
 - [8] J. Birn *et al.*, J. Geophys. Res. **106**, 3715 (2001); S. Galtier *et al.*, Phys. Plasmas **10** 3065 (2003).
 - [9] A. A. Schekochihin *et al.*, Astrophys. J. Supp., submitted (2007) [e-print arXiv:0704.0044].
 - [10] A. Ito *et al.*, Phys. Plasmas **11**, 5643 (2004); A. Hirose *et al.*, Phys. Lett. A **330**, 474 (2004)