

A NOTE ON LOW ENERGY SCATTERING FOR HOMOGENEOUS LONG RANGE POTENTIALS

RUPERT L. FRANK

The very interesting, recent paper [DeSk2] is concerned with scattering at zero energy for long-range potentials. It is shown that if $V = -\alpha|x|^{-\mu} + W$ for some $\alpha > 0$, $0 < \mu < 2$ and a sufficiently fast decaying, radial function W , then the scattering matrix for the pair $(-\Delta + V, -\Delta)$ at zero energy is given by

$$\exp\left(-i\frac{\pi\mu}{2-\mu}\sqrt{-\Delta_{\mathbb{S}^{d-1}} + \left(\frac{d-2}{2}\right)^2}\right) + K$$

for some compact operator K (depending on V). Moreover, it is conjectured that if $V = -\alpha|x|^{-\mu}$ (that is, if $W \equiv 0$), then $K \equiv 0$. The purpose of this note is to prove this conjecture.

Theorem 1. *Let $d \geq 2$, $0 < \mu < 2$ and $\alpha > 0$. For $\lambda > 0$ let $S(\lambda)$ be the scattering matrix for the pair $(-\Delta - \alpha|x|^{-\mu}, -\Delta)$ at energy λ in the sense of [DeSk1]. Then*

$$s - \lim_{\lambda \rightarrow 0^+} S(\lambda) = \exp\left(-i\frac{\pi\mu}{2-\mu}\sqrt{-\Delta_{\mathbb{S}^{d-1}} + \left(\frac{d-2}{2}\right)^2}\right).$$

It was shown in [DeSk1] that $S(0) := s - \lim_{\lambda \rightarrow 0^+} S(\lambda)$ exists and is given through the phase shifts of the solutions of the energy zero equations corresponding to fixed angular momentum. To be more precise, for any $l \in \mathbb{N}_0$ there exists a function f satisfying

$$\begin{aligned} -f'' + \left(\frac{(l-1+d/2)^2 - 1/4}{r^2} - \frac{\alpha}{r^\mu}\right) f &= 0 \quad \text{in } \mathbb{R}_+, \\ \lim_{r \rightarrow 0} r^{-l - \frac{d-1}{2}} f(r) &= 1, \\ \lim_{r \rightarrow \infty} \left(r^{-\mu/4} f(r) - C_l \sin\left(\frac{2\sqrt{\alpha}}{2-\mu} r^{\frac{2-\mu}{2}} + D_l\right)\right) &= 0 \end{aligned}$$

for some $C_l > 0$, and in terms of this function the action of $S(0)$ on a spherical harmonic Y of order l is given by

$$S(0)Y = \exp\left(i2\left(D_l + \frac{\pi}{4}(d-3+2l)\right)\right) Y.$$

(Note that the 2π -ambiguity in the definition of D_l does not affect the formula for $S(0)$.) In view of these facts and recalling that the Laplacian acts on spherical harmonics of order l as multiplication by $l(l+d-2)$, Theorem 1 will follow if we can prove

that $D_l = -\frac{\pi(d-2+2l)}{2(2-\mu)} + \frac{\pi}{4}$ modulo 2π . This equality is the assertion of the following lemma applied to $\nu = l + \frac{d-2}{2}$.

Lemma 2. *Let $\nu \geq 0$, $0 < \mu < 2$ and $\alpha > 0$. The function*

$$f(r) := \Gamma\left(\frac{2\nu}{2-\mu} + 1\right) \left(\frac{2-\mu}{\sqrt{\alpha}}\right)^{\frac{2\nu}{2-\mu}} r^{1/2} J_{\frac{2\nu}{2-\mu}}\left(\frac{2\sqrt{\alpha}}{2-\mu} r^{\frac{2-\mu}{2}}\right) \quad (1)$$

satisfies

$$-f'' + \left(\frac{\nu^2 - 1/4}{r^2} - \frac{\alpha}{r^\mu}\right) f = 0 \quad \text{in } \mathbb{R}_+, \quad (2)$$

$$\lim_{r \rightarrow 0} r^{-\nu-1/2} f(r) = 1, \quad (3)$$

$$\lim_{r \rightarrow \infty} \left(r^{-\mu/4} f(r) - C \sin\left(\frac{2\sqrt{\alpha}}{2-\mu} r^{\frac{2-\mu}{2}} - \frac{\pi\nu}{2-\mu} + \frac{\pi}{4}\right) \right) = 0 \quad (4)$$

with $C := \pi^{-1/2} \Gamma\left(\frac{2\nu}{2-\mu} + 1\right) ((2-\nu)/\sqrt{\alpha})^{\frac{2\nu}{2-\mu} + \frac{1}{2}} > 0$.

Proof. We will show that the function f given in (1) is the unique solution of the initial value problem (2) – (3). Then the asymptotics (4) follow from the asymptotics

$$J_{\tilde{\nu}}(s) = \sqrt{\frac{2}{\pi s}} \left(\sin\left(s - \frac{\pi\tilde{\nu}}{2} + \frac{\pi}{4}\right) + o(1) \right), \quad s \rightarrow \infty,$$

of Bessel functions [AbSt, (9.2.1)].

Let f denote any solution of the initial value problem (2) – (3) and define g by

$$f(r) =: r^{1/2} g(br^{\frac{2-\mu}{2}}), \quad b := \frac{2\sqrt{\alpha}}{2-\mu}.$$

This definition is motivated by the asymptotics (4): If we want to arrive at a function which behaves asymptotically like an inverse square root times an oscillating function – which is the asymptotic behavior of any Bessel function –, then (4) suggests to look at $r^{-1/2} f(r)$ as a function of $r^{\frac{2-\mu}{2}}$. This is essentially what we call g , and we are about to show that g is indeed a Bessel function.

A short computation shows that equation (2) in terms of g becomes

$$g'' + s^{-1} g' - \left(\frac{2\nu}{2-\mu}\right)^2 s^{-2} g + g = 0,$$

which is Bessel's equation with parameter $\tilde{\nu} := 2\nu/(2-\mu)$. Hence g is a linear combination of $J_{\tilde{\nu}}$ and $Y_{\tilde{\nu}}$. Boundary condition (3) in terms of g becomes

$$\lim_{s \rightarrow 0} s^{-\tilde{\nu}} g(s) = b^{-\tilde{\nu}}.$$

Since $J_{\tilde{\nu}}(s) \sim (s/2)^{\tilde{\nu}}/\Gamma(\tilde{\nu}+1)$ for all $\tilde{\nu} \geq 0$ and $Y_{\tilde{\nu}}(s) \sim -(1/\pi)\Gamma(\tilde{\nu})(s/2)^{-\tilde{\nu}}$ for $\tilde{\nu} > 0$, resp. $Y_0(s) \sim (2/\pi) \ln s$ as $s \rightarrow 0$ [AbSt, (9.1.7–9)], we conclude that

$$g(s) = \Gamma(\tilde{\nu}+1)(2/b)^{\tilde{\nu}} J_{\tilde{\nu}}(s).$$

This proves the lemma. □

Acknowledgements. The author wishes to thank R. Seiringer for useful discussions. Support through DFG grant FR 2664/1-1 and U.S. NSF grant PHY 06 52854 is gratefully acknowledged.

REFERENCES

- [AbSt] M. Abramowitz, I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Reprint of the 1972 edition. Dover Publications, New York, 1992.
- [DeSk1] J. Dereziński, E. Skibsted, *Quantum scattering at low energies*. Preprint (2008).
- [DeSk2] J. Dereziński, E. Skibsted, *Scattering at zero energy for attractive homogeneous potentials*. Preprint (2008).

RUPERT L. FRANK, DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, WASHINGTON ROAD, PRINCETON, NJ 08544, USA

E-mail address: `rlfrank@math.princeton.edu`