Artur Khojabaghyan^a * and Vahan V. Mkrtchyan^{a b} †

^aDepartment of Informatics and Applied Mathematics, Yerevan State University, Yerevan, 0025, Armenia

^bInstitute for Informatics and Automation Problems, National Academy of Sciences of Republic of Armenia, 0014, Armenia

For a graph G let L(G) and l(G) denote the size of the largest and smallest maximum matching of a graph obtained from G by removing a maximum matching of G. We show that $L(G) \leq 2l(G)$, and $L(G) \leq \frac{3}{2}l(G)$ provided that G contains a perfect matching. We also characterize the class of graphs for which L(G) = 2l(G). Our characterization implies the existence of a polynomial algorithm for testing the property L(G) = 2l(G). Finally we show that it is NP-complete to test whether a graph G containing a perfect matching satisfies $L(G) = \frac{3}{2}l(G)$.

1. Introduction

In the paper graphs are assumed to be finite, undirected, without loops or multiple edges. Let V(G) and E(G) denote the sets of vertices and edges of a graph G, respectively. If $v \in V(G)$ and $e \in E(G)$, then e is said to cover v if e is incident to v. For $V' \subseteq V(G)$ and $E' \subseteq E(G)$ let $G \setminus V'$ and $G \setminus E'$ denote the graphs obtained from G by removing V'and E', respectively. Moreover, let V(E') denote the set of vertices of G that are covered by an edge from E'. A subgraph H of G is said to be spanning for G, if V(E(H)) = V(G).

The length of a path (cycle) is the number of its edges. A k-path (k-cycle) is a path (cycle) of length k. A 3-cycle is called a triangle.

A set $V' \subseteq V(G)$ $(E' \subseteq E(G))$ is said to be independent, if V'(E') contains no adjacent vertices (edges). An independent set of edges is called matching. A matching of G is called perfect, if it covers all vertices of G. Let $\nu(G)$ denote the cardinality of a largest matching of G. A matching of G is maximum, if it contains $\nu(G)$ edges.

For a positive integer k and a matching M of G, a (2k - 1)-path P is called Maugmenting, if the 2^{nd} , 4^{th} , 6^{th} ,..., $(2k-2)^{th}$ edges of P belong to M, while the endvertices of P are not covered by an edge of M. The following theorem of Berge gives a sufficient and necessary condition for a matching to be maximum:

Theorem 1 (Berge [2]) A matching M of G is maximum, if G contains no M-augmenting path.

^{*}email: arturkhojabaghyan@gmail.com

[†]email: vahanmkrtchyan2002@{ysu.am, ipia.sci.am, yahoo.com}

For two matchings M and M' of G consider the subgraph H of G, where $V(H) = V(M \triangle M')$ and $E(H) = M \triangle M'$. The connected components of H are called $M \triangle M'$ -alternating components. Note that $M \triangle M'$ alternating components are always paths or cycles of even length. For a graph G define:

 $L(G) \equiv \max\{\nu(G \setminus F) : F \text{ is a maximum matching of } G\},\$ $l(G) \equiv \min\{\nu(G \setminus F) : F \text{ is a maximum matching of } G\}.$

It is known that L(G) and l(G) are NP-hard calculable even for connected bipartite graphs G with maximum degree three [4], though there are polynomial algorithms which construct a maximum matching F of a tree G such that $\nu(G \setminus F) = L(G)$ and $\nu(G \setminus F) = l(G)$ (to be presented in [5]).

In the same paper [5] it is shown that $L(G) \leq 2l(G)$. In the present paper we reprove this equality, and also show that $L(G) \leq \frac{3}{2}l(G)$ provided that G contains a perfect matching.

A naturally arising question is the characterization of graphs G with L(G) = 2l(G) and the graphs G with a perfect matching that satisfy $L(G) = \frac{3}{2}l(G)$. In this paper we solve these problems by giving a characterization of graphs G with L(G) = 2l(G) that implies the existence of a polynomial algorithm for testing this property, and by showing that it is NP-complete to test whether a bridgeless cubic graph G satisfies $L(G) = \frac{3}{2}l(G)$. Recall that by Petersen theorem any bridgeless cubic graph contains a perfect matching (see, for example, theorem 3.4.1 of [6]).

Terms and concepts that we do not define can be found in [1, 2, 6, 8].

2. Some auxiliarly results

We will need the following:

Theorem 2 Let G be a graph. Then:

(a) for any two maximum matchings F, F' of G, we have $\nu(G \setminus F') \leq 2\nu(G \setminus F)$;

- (b) $L(G) \leq 2l(G);$
- (c) If L(G) = 2l(G), F_L , F_l are two maximum matchings of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, and H_L is **any** maximum matching of the graph $G \setminus F_L$, then:
 - (c1) $F_l \setminus F_L \subset H_L;$
 - (c2) $H_L \setminus F_l$ is a maximum matching of $G \setminus F_l$;
 - (c3) $F_L \setminus F_l$ is a maximum matching of $G \setminus F_l$;
- (d) if G contains a perfect matching, then $L(G) \leq \frac{3}{2}l(G)$.

Proof. (a)Let H' be any maximum matching in the graph $G \setminus F'$. Then:

$$\nu(G \setminus F') = |H'| = |H' \cap F| + |H' \setminus F| \le |F \setminus F'| + \nu(G \setminus F) = |F' \setminus F| + \nu(G \setminus F) \le 2\nu(G \setminus F).$$

(b) follows from (a).

(c) Consider the proof of (a) and take $F' = F_L$, $H' = H_L$ and $F = F_l$. Since L(G) = 2l(G), we must have equalities throughout, thus properties (c1)-(c3) should be true.

(d) Let F_L , F_l be two perfect matchings of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, and assume H_L to be a maximum matching of the graph $G \setminus F_L$. Define:

$$X = \{e = (u, v) \in F_L : u \text{ and } v \text{ are incident to an edge from } H_L \cap F_l\},\$$
$$x = |X|, k = |H_L \cap F_l|;$$

Clearly, $(H_L \setminus F_l) \cup X$ is a matching of the graph $G \setminus F_l$, therefore, taking into account that $(H_L \setminus F_l) \cap X = \emptyset$, we deduce

$$l(G) = \nu(G \setminus F_l) \ge |H_L \setminus F_l| + |X| = |H_L| - |H_L \cap F_l| + |X| = L(G) - k + x.$$

Since F_L is a perfect matching, it covers the set $V(H_L \cap F_l) \setminus V(X)$, which contains

$$|V(H_L \cap F_l) \setminus V(X)| = 2|(H_L \cap F_l)| - 2|X| = 2k - 2x$$

vertices. Define the set E_{F_L} as follows:

 $E_{F_L} = \{ e \in F_L : e \text{ covers a vertex from } V(H_L \cap F_l) \setminus V(X) \}.$

Clearly, E_{F_L} is a matching of $G \setminus F_l$, too, and therefore

$$l(G) = \nu(G \setminus F_l) \ge |E_{F_L}| = 2k - 2x.$$

Let us show that

$$\max\{L(G) - k + x, 2k - 2x\} \ge \frac{2L(G)}{3}.$$

Note that

if $x \ge k - \frac{L(G)}{3}$ then $L(G) - k + x \ge L(G) - k + k - \frac{L(G)}{3} = \frac{2L(G)}{3}$; if $x \le k - \frac{L(G)}{3}$ then $2k - 2x \ge \frac{2L(G)}{3}$, thus in both cases we have $l(G) \ge \frac{2L(G)}{3}$ or

$$\frac{L(G)}{l(G)} \le \frac{3}{2}.$$

The proof of the theorem 2 is completed. \Box

Lemma 1 (Lemma 2.20, 2.41 of [5]) Let G be a graph, and assume that u and v are vertices of degree one sharing a neighbour $w \in V(G)$. Then:

$$L(G) = L(G \setminus \{u, v, w\}) + 1, l(G) = l(G \setminus \{u, v, w\}) + 1.$$

Proof. The proofs of these two equalities are similar, thus we will stop only on the proof of the first one. Our proof is based on the ideas of [5].

First of all, observe that $\nu(G) = \nu(G \setminus \{u, v, w\}) + 1$. Let us show that $L(G) \geq L(G \setminus \{u, v, w\}) + 1$. Take any maximum matching F of $G \setminus \{u, v, w\}$ with $\nu((G \setminus \{u, v, w\}) \setminus F) = L(G \setminus \{u, v, w\})$, and let H be any maximum matching of $(G \setminus \{u, v, w\}) \setminus F$. Define:

$$F' = F \cup \{(u, w)\}, H' = H \cup \{(v, w)\}.$$

Observe that F' is a maximum matching of G, and H' is a matching of $G \setminus F'$. Thus,

$$L(G) \ge \nu(G \setminus F') \ge |H'| = 1 + |H| = 1 + \nu((G \setminus \{u, v, w\}) \setminus F) = 1 + L(G \setminus \{u, v, w\}).$$

To complete the proof of the first equality, it suffices to show that $L(G) \leq L(G \setminus \{u, v, w\}) + 1$. 1. First of all, let us show that there is a maximum matching F' of G with $\nu(G \setminus F') = L(G)$, such that F' contains one of the edges (u, w) and (v, w).

Take any maximum matching F' of G with $\nu(G \setminus F') = L(G)$, and assume that $F' \cap \{(u, w), (v, w)\} = \emptyset$. Morever, let H' be a maximum matching of $G \setminus F'$. Since F' is a maximum matching of G, F' must contain an edge (w, z), where $z \neq u, v$. Define:

$$F'' = \begin{cases} (F' \setminus \{(w, z)\}) \cup \{(w, u)\}, & \text{if } (w, u) \notin H; \\ (F' \setminus \{(w, z)\}) \cup \{(w, v)\}, & \text{if } (w, u) \in H. \end{cases}$$

Observe that F'' is a maximum matching of G, and H' is a matching of $G \setminus F''$. Thus,

$$\nu(G \setminus F'') \ge |H'| = \nu(G \setminus F') = L(G).$$

The last inequality implies that $\nu(G \setminus F'') = L(G)$. Moreover, $F'' \cap \{(u, w), (v, w)\} \neq \emptyset$.

Thus, initially we can assume that F' is a maximum matching of G with $\nu(G \setminus F') = L(G)$, such that F' contains one of the edges (u, w) and (v, w). Without loss of generality, we can also assume that this edge is (u, w). Now, we claim that there is a maximum matching H' of $G \setminus F'$ that contains the edge (v, w).

Take any maximum matching H' of $G \setminus F'$ and suppose that $(v, w) \notin H'$. Since H' is a maximum matching of $G \setminus F'$, there must exist an edge $(w, z) \in H'$, where $z \neq u, v$. Define:

$$H'' = (H' \setminus \{(w, z)\}) \cup \{(w, v)\}.$$

Observe that H'' is a maximum matching of $G \setminus F'$, since $|H''| = |H'| = \nu(G \setminus F') = L(G)$. Moreover, it contains the edge (w, v). Thus, initially we can assume that H' is a maximum matching of $G \setminus F'$ that contains the edge (v, w).

We are ready to show that $L(G) \leq L(G \setminus \{u, v, w\}) + 1$. Since $\nu(G) = \nu(G \setminus \{u, v, w\}) + 1$, we have that $F' \setminus \{(u, w)\}$ is a maximum matching of $G \setminus \{u, v, w\}$. Taking into account that $H' \setminus \{(v, w)\}$ is a matching of $(G \setminus \{u, v, w\}) \setminus (F' \setminus \{(u, w)\})$, we deduce:

$$L(G) = \nu(G \setminus F') = |H'| = 1 + |H' \setminus \{(v, w)\}| \le 1 + \nu((G \setminus \{u, v, w\}) \setminus (F' \setminus \{(u, w)\})) \le 1 + L(G \setminus \{u, v, w\}).$$

Corollary 1 Let G be a graph with L(G) = 2l(G). Then there are no vertices u, v of degree one, that are adjacent to the same vertex w.

Proof. Suppose not. Then lemma 1 and (b) of theorem 2 imply

$$L(G) = 1 + L(G - \{u, v, w\}) \le 1 + 2l(G - \{u, v, w\}) = 1 + 2(l(G) - 1) < 2l(G)$$

a contradiction. \Box

3. Characterization of graphs G satisfying L(G) = 2l(G)

Let T be the set of all triangles of G that contain at least two vertices of degree two. Note that any vertex of degree two lies in at most one triangle from T. From each triangle $t \in T$ choose a vertex v_t of degree two, and define $V_1(G)$ as follows:

 $V_1(G) = \{v : d_G(v) = 1\} \cup \{v_t : t \in T\}$

Theorem 3 Let G be a connected graph with $|V(G)| \ge 3$. Then L(G) = 2l(G) if and only if

(1) $G \setminus V_1(G)$ is a bipartite graph with a bipartition (X, Y);

(2) $|V_1(G)| = |Y|$ and any $y \in Y$ has exactly one neighbour in $V_1(G)$;

(3) the graph $G \setminus V_1(G)$ contains |X| vertex disjoint 2-paths.

Proof. Sufficiency. Let G be a connected graph with $|V(G)| \ge 3$ satisfying the conditions (1)-(3). Let us show that L(G) = 2l(G).

For each vertex v with d(v) = 1 take the edge incident to it and define F_1 as the union of all these edges. For each vertex $v_t \in V_1(G)$ take the edge that connects v_t to a vertex of degree two, and define F_2 as the union of all those edges. Set:

$$F = F_1 \cup F_2.$$

Note that F is a matching with $|F| = |V_1(G)| = |Y|$. Moreover, since G is bipartite and $|V_1(G)| = |Y|$, the definitions of F_1 and F_2 imply that there is no F-augmenting path in G. Thus, by Berge theorem, F is a maximum matching of G, and

$$\nu(G) = |F| = |V_1(G)| = |Y|.$$

Observe that the graph $G \setminus F$ is a bipartite graph with $\nu(G \setminus F) \leq |X|$, thus

$$l(G) \le \nu(G \setminus F) \le |X|.$$

Now, consider the |X| vertex disjoint 2-paths of the graph $G \setminus V_1(G)$ guaranteed by (3). (2) implies that these 2-paths together with the $|F| = |V_1(G)| = |Y|$ edges of F form |X| vertex disjoint 4-paths of the graph G.

Consider matchings M_1 and M_2 of G obtained from these 4-paths by adding the first and the third, the second and the fourth edges of these 4-paths to M_1 and M_2 , respectively. Define:

$$F' = (F \setminus M_2) \cup (M_1 \setminus F).$$

Note that F' is a matching of G and |F'| = |F|, thus F' is a maximum matching of G. Since $F' \cap M_2 = \emptyset$, we have

$$L(G) \ge \nu(G \setminus F') \ge |M_2| = 2|X| \ge 2l(G).$$

(b) of theorem 2 implies that L(G) = 2l(G).

Necessity. Now, assume that G is a connected graph with $|V(G)| \ge 3$ and L(G) = 2l(G). By proving a series of claims, we show that $G \setminus V_1(G)$ satisfies the conditions (1)-(3) of the theorem. **Claim 1** For any maximum matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, $F_L \cup F_l$ induces a spanning subgraph, that is $V(F_L) \cup V(F_l) = V(G)$.

Proof. Suppose that there is a vertex $v \in V(G)$ that is covered neither by F_L nor by F_l . Since F_L and F_l are maximum matchings of G, for each edge e = (u, v) the vertex u is incident to an edge from F_L and to an edge from F_l .

Case 1: there is an edge e = (u, v) such that u is incident to an edge from $F_L \cap F_l$.

Note that $\{e\} \cup (F_L \setminus F_l)$ is a matching of $G \setminus F_l$ which contradicts (c3) of the theorem 2. Case 2: for each edge e = (u, v) u is incident to an edge $f_L \in F_L \setminus F_l$ and to an edge $f_l \in F_l \setminus F_L$.

Let H_L be any maximum matching of $G \setminus F_L$. Due to (c1) of theorem 2 $f_l \in H_L$. Define:

 $H'_L = (H_L \setminus \{f_l\}) \cup \{e\}.$

Note that H'_L is a maximum matching of $G \setminus F_L$ such that $F_l \setminus F_L$ is not a subset of H'_L contradicting (c1) of theorem 2. \Box

Claim 2 For any maximum matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, the alternating components $F_L \triangle F_l$ are 2-paths.

Proof. It suffices to show that there is no edge $f_L \in F_L$ that is adjacent to two edges from F_l . Suppose that some edge $f_L \in F_L$ is adjacent to edges f'_l and f''_l from F_l . Let H_L be any maximum matching of $G \setminus F_L$. Due to (c1) of theorem 2 $f'_l, f''_l \in H_L$. This implies that $\{f_L\} \cup (H_L \setminus F_l)$ is a matching of $G \setminus F_l$ which contradicts (c2) of theorem 2. \Box

Claim 3 For any maximum matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$

- (a) if $u \in V(F_l) \setminus V(F_L)$ then d(u) = 1 or d(u) = 2. Moreover, in the latter case, if v and w denote the two neighbours of u, where $(u, w) \in F_l$, then d(w) = 2 and $(v, w) \in F_L$.
- (b) if $u \in V(F_L) \setminus V(F_l)$ then $d(u) \ge 2$.

Proof. (a) Assume that u is covered by an edge $e_l \in F_l$ and $u \notin V(F_L)$. Suppose that $d(u) \geq 2$, and there is an edge e = (u, v) such that $e \notin F_l$. Taking into account the claim 1, we need only to consider the following four cases:

Case 1: $v \in V(F_l) \setminus V(F_L)$.

This is impossible, since F_L is a maximum matching.

Case 2: v is covered by an edge $f \in F_L \cap F_l$;

Let H_L be any maximum matching of $G \setminus F_L$. Due to (c1) of theorem $2 e_l \in H_L$, thus $e \notin H_L$.

Define:

$$F'_L = (F_L \setminus \{f\}) \cup \{e\}.$$

Note that F'_L is a maximum matching, and H_L is a matching of $G \setminus F'_L$. Moreover,

$$\nu(G \setminus F'_L) \ge |H_L| = \nu(G \setminus F_L) = L(G),$$

thus H_L is a maximum matching of $G \setminus F'_L$ and $\nu(G \setminus F'_L) = L(G)$. This is a contradiction because $F'_L \triangle F_l$ contains a component which is not a 2-path contradicting claim 2.

Case 3: v is incident to an edge $f_L \in F_L$, $f_l \in F_l$ and $f_L \neq f_l$.

Let H_L be any maximum matching of $G \setminus F_L$. Due to (c1) of theorem 2, $e_l, f_l \in H_L$. Define:

$$F'_L = (F_L \setminus \{f_L\}) \cup \{e\}.$$

Note that F'_L is a maximum matching, and H_L is a matching of $G \setminus F'_L$. Moreover,

$$\nu(G \setminus F'_L) \ge |H_L| = \nu(G \setminus F_L) = L(G),$$

thus H_L is a maximum matching of $G \setminus F'_L$ and $\nu(G \setminus F'_L) = L(G)$. This is a contradiction because $F'_L \triangle F_l$ contains a component which is not a 2-path contradicting claim 2.

Case 4: v is covered by an edge $e_L \in F_L$ and $v \notin V(F_l)$.

Note that if e_L is not adjacent to e_l then claim 2 implies that the edges e, e_L and the edge $\tilde{e} \in F_l \setminus F_L$ that is adjacent to e_L would form an augmenting 3-path with respect to F_L , which would contradict the maximality of F_L .

Thus it remains to consider the case when e_L is adjacent to e_l and d(u) = 2. Let w be the vertex adjacent to both e_l and e_L . Let us show that d(w) = 2. Let H_L be any maximum matching of $G \setminus F_L$. Due to (c1) of theorem 2, $e_l \in H_L$. Define:

$$F'_L = (F_L \setminus \{e_L\}) \cup \{e\}.$$

Note that F'_L is a maximum matching, and H_L is a matching of $G \setminus F'_L$. Moreover,

$$\nu(G \setminus F'_L) \ge |H_L| = \nu(G \setminus F_L) = L(G),$$

thus H_L is a maximum matching of $G \setminus F'_L$ and $\nu(G \setminus F'_L) = L(G)$. If $d(w) \ge 3$ there is a vertex $w' \ne u, v$ such that $(w, w') \in E(G)$ and w' satisfies one of the conditions of cases 1,2 and 3 with respect to F'_L and F_l . A contradiction. Thus d(w) = 2.

Clearly, $(v, w) = e_L \in F_L$.

(b) This follows from (a) of claim 3 and corollary 1. \Box

Claim 4 Let F_L , F_l be any maximum matchings of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$. Then for any maximum matching H_L of the graph $G \setminus F_L$ there is no edge of $F_L \cap F_l$ which is adjacent to two edges from H_L .

Proof. Due to (c3) of theorem 2 any edge from H_L that is incident to a vertex covered by an edge of $F_L \cap F_l$ is also incident to a vertex from $V(F_L) \setminus V(F_l)$. If there were an edge $e \in F_L \cap F_l$ which is adjacent to two edges $h_L, h'_L \in H_L$, then the edges h_L, e and h'_L would form an augmenting 3-path with respect to F_l , which would contradict the maximality of F_l . \Box

Claim 5 (1) for any maximum matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, we have $(V(F_L) \setminus V(F_l)) \cap V_1(G) = \emptyset$; (2) there is a maximum matching F_l of G with $\nu(G \setminus F_l) = l(G)$ and a maximum matching F_L of the graph G with $\nu(G \setminus F_L) = L(G)$, such that $V_1(G) \subseteq V(F_L \cap F_l) \cup (V(F_l) \setminus V(F_L))$.

Proof. (1) On the opposite assumption, consider a vertex $x \in (V(F_L) \setminus V(F_l)) \cap V_1(G)$. Since $x \in V_1(G)$ then $d(x) \leq 2$. On the other hand, (b) of claim 3 implies that $d(x) \geq 2$, thus d(x) = 2. Then there are vertices y, z such that $(x, z) \in F_L$, $(z, y) \in F_l$. Note that due to (a) of claim 3, we have $d(y) \leq 2$. Let us show that d(y) = 1. Suppose that d(y) = 2. Then due to (a) of claim 3, we have that d(z) = 2, thus G is the triangle, which is a contradiction, since G does not satisfy L(G) = 2l(G).

Thus d(y) = 1. Since $x \in V_1(G)$, we imply that there is a vertex w with d(w) = 2 such that w, x, z form a triangle. Note that w is covered neither by F_L nor by F_l , which contradicts claim 1.

(2) Let e_t be an edge of a triangle $t \in T$ connecting the vertex $v_t \in V_1(G)$ to a vertex of degree two. Let us show that there is a maximum matching F_l of G with $\nu(G \setminus F_l) = l(G)$ such that $e_t \in F_l$ for each $t \in T$.

Choose a maximum matching F_l of G with $\nu(G \setminus F_l) = l(G)$ that contains as many edges e_t as possible. Let us show that F_l contains all edges e_t . Suppose that there is $t_0 \in T$ such that $e_{t_0} \notin F_l$. Define:

$$F_l' = (F_l \setminus \{e\}) \cup \{e_{t_0}\},$$

where e is the edge of F_l that is adjacent to e_{t_0} . Note that

$$\nu(G \setminus F_l) \le \nu(G \setminus F_l) = l(G),$$

thus F'_l is a maximum matching of G with $\nu(G \setminus F_l) = l(G)$. Note that F'_l contains more edges e_t than does F_l which contradicts the choice of F_l .

Thus, there is a maximum matching F_l of G with $\nu(G \setminus F_l) = l(G)$ such that $e_t \in F_l$ for all $t \in T$. Now, for this maximum matching F_l of G choose a maximum matching F_L of the graph G with $\nu(G \setminus F_L) = L(G)$, such that $V(F_L \cap F_l) \cup (V(F_l) \setminus V(F_L))$ covers maximum number of vertices from $V_1(G)$. Let us show that $V_1(G) \subseteq V(F_L \cap F_l) \cup (V(F_l) \setminus V(F_L))$.

Suppose that there is a vertex $x \in V_1(G)$ such that $x \notin V(F_L \cap F_l) \cup (V(F_l) \setminus V(F_L))$. Note that due to claim 1 and (b) of claim 3, any vertex of degree one is either incident to an edge from $F_L \cap F_l$ or to an edge $V(F_l) \setminus V(F_L)$. Thus due to definition of $V_1(G)$, d(x) = 2 and if y and z denote the two neighbors of x, then d(y) = 2 and $(y, z) \in E(G)$.

Since $x \notin V(F_L \cap F_l)$, we have that $(x, y) \notin F_L$, and since $x \notin (V(F_l) \setminus V(F_L))$, we have that $(y, z) \notin F_L$, thus $(x, z) \in F_L$, as F_L is a maximum matching. Let H_L be any maximum matching of $G \setminus F_L$. As L(G) = 2l(G), we have $(x, y) \in H_L$ ((c1) of theorem 2). Define:

$$F'_L = (F_L \setminus \{(x, z)\}) \cup \{(y, z)\}.$$

Note that F'_L is a maximum matching of G, H_L is a matching of $G \setminus F_L$, thus

$$\nu(G \setminus F'_L) \ge |H_L| = \nu(G \setminus F_L) = L(G).$$

Therefore F'_L is a maximum matching of G with $\nu(G \setminus F'_L) = L(G)$. Now, observe that $V(F'_L \cap F_l) \cup (V(F_l) \setminus V(F'_L))$ covers more vertices than does $V(F_L \cap F_l) \cup (V(F_l) \setminus V(F_L))$ which contradicts the choice of F_L . The proof of the claim 5 is completed. \Box

Claim 6 For any maximum matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$, we have

- (1) $V(F_L) \setminus V(F_l)$ is an independent set;
- (2) no edge of G connects two vertices that are covered by both $F_L \setminus F_l$ and $F_l \setminus F_L$;
- (3) no edge of G is adjacent to two different edges from $F_L \cap F_l$;
- (4) no edge of G connects a vertex covered by $F_L \cap F_l$ to a vertex covered by both $F_L \setminus F_l$ and $F_l \setminus F_L$;
- (5) if $(u, v) \in F_L \cap F_l$ then either $u \in V_1(G)$ or $v \in V_1(G)$.

Proof. (1)There is no edge of G connecting two vertices from $V(F_L) \setminus V(F_l)$ since F_l is a maximum matching.

- (2) follows from (c1) and (c2) of theorem 2.
- (3) follows from (c3) of theorem 2.

(4) Suppose that there is an edge $e = (y_1, y_2)$, such that y_1 is covered by an edge $(z, y_1) \in F_L \cap F_l$ and y_2 is covered by both $F_L \setminus F_l$ and $F_l \setminus F_L$. Consider a maximum matching H_L of the graph $G \setminus F_L$. Note that y_1 must be incident to an edge from H_L , as otherwise we could replace the edge of H_L that is adjacent to e and belongs also to $F_l \setminus F_L$ ((c1) of theorem 2) by the edge e to obtain a new maximum matching H'_L of the graph $G \setminus F_L$ which would not satisfy (c1) of theorem 2.

So let y_1 be incident to an edge $h_L \in H_L$, which connects y_1 with a vertex $x \in V(F_L) \setminus V(F_l)$. Note that due to claim 4, z is not incident to an edge from H_L . Now, let x_1 be a vertex such that $(x, x_1) \in F_L \setminus F_l$ (such a vertex exists since $x \in V(F_L) \setminus V(F_l)$). As F_l is a maximum matching, x_1 is incident to an edge $(x_1, x_2) \in F_l \setminus F_L$. By (c1) of theorem 2, $(x_1, x_2) \in H_L$. Moreover, by claim 2, x_2 is not adjacent to an edge from F_L . Thus the edges $(z, y_1), (y_1, x), (x, x_1)$ and (x_1, x_2) form an $F_L - H_L$ alternating 4-path P. Define:

$$F'_L = (F_L \setminus E(P)) \cup (H_L \cap E(P)),$$

$$H'_L = (H_L \setminus E(P)) \cup (F_L \cap E(P)).$$

Note that F'_L is a maximum matching of G, H'_L is a matching of $G \setminus F'_L$ of cardinality $|H_L|$, and

$$\nu(G \setminus F'_L) \ge |H'_L| = |H_L| = \nu(G \setminus F_L) = L(G),$$

thus H'_L is a maximum matching of $G \setminus F'_L$ and $\nu(G \setminus F'_L) = L(G)$. This is a contradiction since the edge *e* connects two vertices which are covered by $F'_L \setminus F_l$ and $F_l \setminus F'_L$ ((2) of claim 6).

(5)Suppose that $e = (u, v) \in F_L \cap F_l$. Since G is connected and $|V| \ge 3$, we, without loss of generality, may assume that $d(v) \ge 2$, and there is $w \in V(G), w \ne u$ such that $(w, v) \in E(G)$. Consider a maximum matching H_L of the graph $G \setminus F_L$. Note that, without loss of generality, we can assume that v is incident to an edge from H_L , as otherwise we could replace the edge of H_L that is incident to w (H_L is a maximum matching of $G \setminus F_L$) by the edge (w, v) to obtain a new maximum matching H'_L of the graph $G \setminus F_L$ such that v is incident to an edge from H'_L .

So we can assume that there is an edge $(v,q) \in H_L$, $q \neq u$. Note that due to claim 4, u is not incident to an edge from H_L . (c3) of theorem 2 implies that q is incident to an edge from $(q,q_1) \in F_L \setminus F_l$. As F_l is a maximum matching, q_1 is incident to an edge $(q_1,q_2) \in F_l \setminus F_L$. By (c1) of theorem 2, $(q_1,q_2) \in H_L$. Moreover, by claim 2, q_2 is not adjacent to an edge from F_L . Thus the edges $(u,v), (v,q), (q,q_1)$ and (q_1,q_2) form an $F_L - H_L$ alternating 4-path P. Define:

$$F'_L = (F_L \setminus E(P)) \cup (H_L \cap E(P)),$$

$$H'_L = (H_L \setminus E(P)) \cup (F_L \cap E(P)).$$

Note that F'_L is a maximum matching of G, H'_L is a matching of $G \setminus F'_L$ of cardinality $|H_L|$, and

$$\nu(G \setminus F'_L) \ge |H'_L| = |H_L| = \nu(G \setminus F_L) = L(G),$$

thus H'_L is a maximum matching of $G \setminus F'_L$ and $\nu(G \setminus F'_L) = L(G)$. Since $u \in V(F_l) \setminus V(F'_L)$ (a) of claim 3 implies that either d(u) = 1 and therefore $u \in V_1(G)$, or d(u) = d(v) = 2and therefore either $u \in V_1(G)$ or $v \in V_1(G)$. Proof of the claim 6 is completed. \Box

We are ready to complete the proof of the theorem. Take any maximum matchings F_L , F_l of the graph G guaranteed by the (2) of claim 5 and consider the following partition of $V(G \setminus V_1(G)) = V(G) \setminus V_1(G)$:

$$X = X(F_L, F_l) = V(F_L) \setminus V(F_l), Y = Y(F_L, F_l) = V(G) \setminus (V_1(G) \cup X).$$

Claim 6 implies that X and Y are independent sets of vertices of $G \setminus V_1(G)$, thus $G \setminus V_1(G)$ is a bipartite graph with a bipartition (X, Y).

The choice of maximum matchings F_L , F_l , (a) of claim 3, (5) of claim 6 and the definition of the set Y imply (2) of the theorem 3.

Let us show that it satisfies (3), too.

Consider the alternating 2-paths of

 $(H_L \setminus F_l) \triangle (F_L \setminus F_l).$

(c2), (c3) of theorem 2 and the definition of the set X imply that there are |X| such 2-paths. Moreover, these 2-paths are in fact 2-paths of the graph $G \setminus V_1(G)$. Thus G satisfies (3) of the theorem. The proof of the theorem 3 is completed. \Box

Corollary 2 The property of a graph L(G) = 2l(G) can be tested in polynomial time.

Proof. First of all note that the property L(G) = 2l(G) is additive, that is, a graph satisfies this property if and only if all its connected components do. Thus we can concentrate only on connected graphs.

All connected graphs with $|V(G)| \leq 2$ satisfy the equality L(G) = 2l(G), thus we can assume that $|V(G)| \geq 3$.

Next, we construct a set $V_1(G)$, which can be done in linear time. Now, we need to check whether the graph $G \setminus V_1(G)$ satisfies the conditions (1)-(3) of the theorem 3.

It is well-known that the properties (1) and (2) can be checked in polynomial time, so we will consider only the testing of (3).

From a graph $G \setminus V_1(G)$ with a bipartition (X, Y) we construct a network \vec{G} with new vertices s and t. The arcs of \vec{G} are defined as follows:

- connect s to every vertex of X with an arc of capacity 2;
- connect every vertex of Y to t by an arc of capacity 1;
- for every edge $(x, y) \in E(G)$, $x \in X$, $y \in Y$ add an arc connecting the vertex x to the vertex y which has capacity 1.

Note that

- the value of the maximum s t flow in \vec{G} is no more than 2|X| (the capacity of the cut (S, \bar{S}) , where $S = \{s\}, \bar{S} = V(\vec{G}) \setminus S$, is 2|X|);
- the value of the maximum s t flow in \overline{G} is 2|X| if and only if the graph $G \setminus V_1(G)$ contains |X| vertex disjoint 2-paths,

thus (3) also can be tested in polynomial time. \Box

Remark 1 Recently Monnot and Toulouse in [7] proved that 2-path partition problem remains NP-complete even for bipartite graphs of maximum degree three. Fortunately, in theorem 3 we are dealing with a special case of this problem which enables us to present a polynomial algorithm in corollary 2.

4. NP-completeness of testing $L(G) = \frac{3}{2}l(G)$ in the class of bridgeless cubic graphs

The reader may think that a result analogous to corollary 2 can be proved for the property $L(G) = \frac{3}{2}l(G)$ in the class of graphs containing a perfect matching. Unfortunately this fails already in the class of bridgeless cubic graphs, which by the well-known theorem of Petersen are known to possess a perfect matching (see theorem 3.4.1 of [6]).

Theorem 4 It is NP-complete to test the property $L(G) = \frac{3}{2}l(G)$ in the class of bridgeless cubic graphs.

Proof. Clearly, the problem of testing the property $L(G) = \frac{3}{2}l(G)$ for graphs containing a perfect matching is in NP, since if we are given perfect matchings F_L , F_l of the graph G with $\nu(G \setminus F_L) = L(G)$, $\nu(G \setminus F_l) = l(G)$ then we can calculate L(G) and l(G) in polynomial time.

We will use the well-known 3-edge-coloring problem ([3]) to establish the NP-completeness of our problem.

Let G be a bridgeless cubic graph. Consider a bridgeless cubic graph G_{\triangle} obtained from G by replacing every vertex of G by a triangle. We claim that G is 3-edge-colorable if and only if $L(G_{\triangle}) = \frac{3}{2}l(G_{\triangle})$.

Suppose that G is 3-edge-colorable. Then G_{\triangle} is also 3-edge-colorable, which means that G_{\triangle} contains two edge disjoint perfect matchings F and F'. This implies that

$$L(G_{\triangle}) \ge \nu(G_{\triangle} \setminus F) \ge |F'| = \frac{|V(G_{\triangle})|}{2}$$

On the other hand, the set E(G) forms a perfect matching of G_{Δ} , and

$$l(G_{\triangle}) \le \nu(G_{\triangle} \setminus E(G)) = \frac{|V(G_{\triangle})|}{3},$$

since every component of $G_{\triangle} \setminus E(G)$ is a triangle. Thus:

$$\frac{L(G_{\triangle})}{l(G_{\triangle})} \ge \frac{3}{2}$$

(d) of theorem 2 implies that $\frac{L(G_{\triangle})}{l(G_{\triangle})} = \frac{3}{2}$.

Now assume that $\frac{L(G_{\triangle})}{l(G_{\triangle})} = \frac{3}{2}$. Note that for every perfect matching F of the graph G_{\triangle} the graph $G_{\triangle} \setminus F$ is a 2-factor, therefore

$$L(G_{\Delta}) = \frac{|V(G_{\Delta})| - w(G_{\Delta})}{2},$$

$$l(G_{\Delta}) = \frac{|V(G_{\Delta})| - W(G_{\Delta})}{2}$$

where $w(G_{\triangle})$ and $W(G_{\triangle})$ denote the minimum and maximum number of odd cycles in a 2-factor of G_{\triangle} , respectively. Since $\frac{L(G_{\triangle})}{l(G_{\triangle})} = \frac{3}{2}$ we have

$$W(G_{\triangle}) = \frac{|V(G_{\triangle})| + 2w(G_{\triangle})}{3}.$$

Taking into account that $W(G_{\triangle}) \leq \frac{|V(G_{\triangle})|}{3}$, we have:

$$W(G_{\Delta}) = \frac{|V(G_{\Delta})|}{3},$$

$$w(G_{\Delta}) = 0.$$

Note that $w(G_{\triangle}) = 0$ means that G_{\triangle} is 3-edge-colorable, which in its turn implies that G is 3-edge-colorable. The proof of the theorem is completed. \Box

REFERENCES

- R. Diestel, Graph theory, Springer-Verlag Heidelberg, New York, 1997, 2000, 2005, 2010.
- 2. F. Harary, Graph Theory, Addison-Wesley, Reading, MA, 1969.

- 3. I. Holyer, The NP-completeness of edge coloring, SIAM J. Comput. 10, N4, 718-720, 1981 (available at: http://cs.bris.ac.uk/ian/graphs).
- 4. R.R. Kamalian, V. V. Mkrtchyan, On complexity of special maximum matchings constructing, Discrete Mathematics 308, (2008), pp. 1792-1800
- 5. R.R. Kamalian, V. V. Mkrtchyan, Two polynomial algorithms for special maximum matching constructing in trees, under construction (http://arxiv.org/abs/0707.2295).
- 6. L. Lovász, M.D. Plummer, Matching theory, Ann. Discrete Math. 29 (1986).
- 7. J. Monnot, S. Toulouse, The path partition problem and related problems in bipartite graphs, Operation Research Letters, 35, (2007), pp. 677-684
- 8. D. B. West, Introduction to Graph Theory, Prentice-Hall, Englewood Cliffs, 1996.