

Induced superstring cosmologies and moduli stabilization*

Tristan Catelin-Jullien¹, Costas Kounnas¹
Hervé Partouche² and Nicolaos Toumbas³

¹ Laboratoire de Physique Théorique, Ecole Normale Supérieure,[†]
24 rue Lhomond, F–75231 Paris cedex 05, France
catelin@lpt.ens.fr, Costas.Kounnas@lpt.ens.fr

² Centre de Physique Théorique, Ecole Polytechnique,[◇]
F–91128 Palaiseau, France
Herve.Partouche@cpt.polytechnique.fr

³ Department of Physics, University of Cyprus,
Nicosia 1678, Cyprus
nick@ucy.ac.cy

Abstract

We extend the analysis of the recently obtained stringy cosmological solutions induced by thermal and quantum effects, once space-time supersymmetry is spontaneously broken by geometrical fluxes. Cases in which more than one modulus participating in the supersymmetry breaking mechanism are investigated. The free energy is obtained at the full string level. In the intermediate cosmological region where the temperature and the supersymmetry breaking scale are sufficiently smaller than the Hagedorn temperature, the quantum and thermal corrections are under control and calculable. The reason is that the contributions to the effective potential of the moduli that are not participating in the supersymmetry breaking are exponentially suppressed. The backreaction on the initially flat background results in many cases into cosmological evolutions, where the dynamics of all complex structure moduli is frozen. The solutions describe effectively a radiation dominated era, where thermal effects are never negligible, even if the temperature tends to zero at late times. We analyze several types of supersymmetry breaking patterns and examine the stability of the corresponding radiation era.

* Research partially supported by the EU (under the contracts MRTN-CT-2004-005104, MRTN-CT-2004-512194, MRTN-CT-2004-503369, MEXT-CT-2003-509661), ANR (CNRS-USAR) contract 05-BLAN-0079-02, CNRS PICS 2530, 3059 and 3747, INTAS grant 03-51-6346 and INTER-REG IIIA Crete/Cyprus.

[†] Unité mixte du CNRS et de l'Ecole Normale Supérieure associée à l'Université Pierre et Marie Curie (Paris 6), UMR 8549.

[◇] Unité mixte du CNRS et de l'Ecole Polytechnique, UMR 7644.

1 Introduction

Perhaps the most natural setting for testing string theory is the cosmology of our Universe. By now, there is a plethora of cosmological data favoring the phenomenological model of hot Big Bang cosmology, where the cosmological evolution starts with a highly singular event, the Big Bang, followed by an initial period of rapid inflation, a very high temperature phase, a proportionally large amount of dark matter and dark energy [1]. Many features of this phenomenological model are not well understood, and the hope is that their explanation will arise from a fundamental theory of quantum gravity such as string theory.

In an effort to build a concrete theoretical framework for studying cosmology, a class of string theory vacua, where the backreaction of both thermal and quantum effects can be systematically taken into account, was recently examined in [2–5]. In particular, starting with four dimensional heterotic string models, with initial $N = 4$ or $N = 2$ space-time supersymmetry [5], and implementing the thermal and the quantum corrections due to spontaneous breaking of supersymmetry, cosmological solutions are found, at least when the temperature T and the supersymmetry breaking scale M are sufficiently below the Hagedorn temperature T_H . In these examples, all moduli that are not involved in the breaking of supersymmetry give exponentially suppressed contributions to the (thermal) effective potential. This is essentially the underlying reason for the no-scale structure [4, 5] characterizing these models. Due to this remarkable property, the thermal and quantum corrections are under control and calculable. We would like to stress here that the absence of both infrared and ultraviolet ambiguities leads to a well defined energy density and pressure [5].

To be more precise, we are forced to separate the cosmological evolution in four distinct phases, according to the value of the temperature, namely:

(i) The very early “Big Bang” phase, where the underlying string theory degrees of freedom are strongly coupled. Perhaps string dualities can be applied to understand this phase and resolve the classical Big Bang singularity. See e.g. [6–11, 19] and references therein.

(ii) A high temperature stringy phase, $T \lesssim T_H$, where string oscillators and winding states must be properly taken into account [13–17]. Often, these lead to a non-geometrical structure of the Universe, e.g. the T-fold cosmologies of [2], or even to a change of the topology and dimensionality of space [7, 11].

(iii) The third phase has features similar to that of a standard, radiation dominated Friedmann cosmology. Here, the Universe has cooled down to temperatures far below Hagedorn. The effects of string massive states are exponentially suppressed $\mathcal{O}(\exp(-M_s/T))$. In this phase, the ratio of the temperature T and supersymmetry breaking scale M is fixed, both evolving inversely proportional to the scale factor of the Universe [4, 5]. In cases with $N = 1$ initial supersymmetry, the behavior can be that of a cosmological constant dominated inflationary universe [3, 4].

(iv) At lower temperatures, new phenomena such as the electroweak phase transition, QCD confinement and structure formation are taking place. We expect also that in this phase, some dynamics becoming relevant at these lower temperatures will stabilize the no-scale modulus associated to the supersymmetry breaking scale [18].

Some interesting ideas concerning the first two stringy phases have been presented recently in [19], where it was argued that the introduction of certain chemical potentials in the standard canonical ensemble of superstrings removes the Hagedorn instabilities. These ensembles are characterized by a “Temperature duality,” $\mathcal{Z}(T/T_H) = \mathcal{Z}(T_H/T)$. (See also [12] for some related work.) Equally interesting are the models of [11], which possess a new kind of *massive boson-fermion degeneracy symmetry*. Type II, heterotic and orientifold models have been presented. Some proposals are put forward, in the framework of these theories, concerning the early structure of the Universe.

Here however, we would like to examine more thoroughly the generalization of supersymmetry breaking in the cases where more moduli are involved, not only in heterotic string but also in type II closed string theories, working in the intermediate region (iii). It is interesting that this intermediate phase can be studied with high precision at the full string level [5], thanks to the fact that just below the Hagedorn temperature, the theory possesses a no-scale structure. The free energy is set by a single, overall scale, which can be chosen to be either the temperature or the supersymmetry breaking scale, and the rest of the dependence is given in terms of functions of dimensionless, complex structure like ratios. The backreaction on the initially flat background induces the cosmological evolution [4, 5].

In the string models studied in [5], only a single modulus was participating in the supersymmetry breaking mechanism. In this work we extend the analysis for cases where more geometrical moduli participate in the spontaneous breaking of supersymmetry. As we already

remarked, all such moduli give non-exponentially suppressed contributions to the effective potential. We examine in more detail the case where the supersymmetry breaking arises via geometrical fluxes [20] induced by stringy Scherk-Schwarz [21] boundary conditions along two internal spatial cycles with radii denoted by R_4 and R_5 . The supersymmetry breaking scale M is proportional to the inverse of the volume modulus $\sqrt{R_4 R_5}$. The results can be easily extended to cases involving more internal cycles. The heterotic or type II geometrical fluxes are introduced by utilizing the helicity and/or other internal R-symmetry charges. In the type II models, left-moving and right-moving R-symmetry charges can be coupled symmetrically or asymmetrically to the two cycles.

As we will see, the low energy dynamics of some models admits a solution describing a radiation dominated era. During the evolution, the supersymmetry breaking scale M and the temperature T evolve in time the same way as the inverse of the scale factor of the Universe. It is then important to analyze the stability for the dimensionless modulus describing the shape of the internal space, $\mathcal{U} \equiv R_5/R_4$. We find a rich structure of phenomena, depending on whether the corresponding effective potential admits a minimum, a maximum or a run-away behavior. Complementary results can be found in [22]. There, it is shown that these solutions are attractors for the dynamics. That is, there are basins of initial conditions whose associated cosmological evolutions converge towards the radiation dominated era with stabilized complex structures.

The paper is organized as follows. In Section 2, we describe the basic principles of the construction, clarifying the domain of validity of our analysis. We examine different classes heterotic and type II string vacua and implement various ways that lead to a spontaneous breaking of supersymmetry. We calculate the free energy at the full string level, and obtain the generic structure of the thermal effective potential in the intermediate region where $T \ll T_H$. We use these general results to find their counterparts at zero temperature. We also derive in the Einstein frame the gravitational equations and the equations of motion for the main moduli participating in the supersymmetry breaking mechanism, including the dilaton field. We present solutions at zero temperature, where the time evolution of the scale factor and the stability of the complex structure \mathcal{U} are analyzed.

In Section 3, we present explicit thermal models leading to radiation dominated cosmological evolutions. The compatibility between the gravitational equations and the equations of mo-

tion of the relevant moduli leads to the equation of state $\rho = (3+n)P$, where n is the number of internal dimensions involved in the supersymmetry breaking mechanism. In addition, we analyze the stability for the modulus \mathcal{U} . We compare to the zero temperature situation, mainly to show the relevance of the thermal corrections to the cosmological evolution. We show that during the radiation era, even when the temperature is very small, thermal effects are never negligible. The qualitative behaviors of thermal and non-thermal evolutions are drastically different. This is due to the non-linear character of the gravitational and moduli equations.

The final Section is devoted to conclusions and perspectives.

2 General setup

We consider initially supersymmetric flat backgrounds within the context of four dimensional superstrings constructed via orbifolds [23] and/or via the free fermionic construction [24]. By turning on certain non-trivial geometrical fluxes, we can spontaneously break space-time supersymmetry [21]. The procedure that we follow involves coupling some of the internal lattice quantum numbers to the space-time Fermion number F and/or to any of the discrete R-symmetry charges Q_R . This is a generalization of the Scherk-Schwarz mechanism to superstrings [21]. In addition, the system is put at finite temperature [13–15]. Our aim is to study cosmological evolutions induced by the thermal and quantum corrections, as in [5], and in particular investigate how some of the geometrical moduli, participating in the breaking of supersymmetry, can be stabilized around a local minimum.

2.1 Heterotic models

We start with heterotic string compactifications on six-manifolds of the form $S^1(R_4) \times S^1(R_5) \times \mathcal{M}_4$. Here, the choice of the four-manifold \mathcal{M}_4 determines the initial amount of supersymmetry: $N_4 = 4$ for the case of T^4 , and $N_4 = 2$ for the cases of T^4/\mathbb{Z}_2 orbifold and K_3 compactifications. Two internal cycles, associated to the X^4 and X^5 directions, have been singled out, since these are to be utilized to break supersymmetry spontaneously.

We illustrate the derivation of the pressure on the simplest example. The initially super-

symmetric string partition function is given by

$$Z = \frac{V_3}{(2\pi)^3} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{5/2}} \Gamma_{(1,1)}(R_0) \frac{1}{2} \sum_{a,b} (-)^{a+b+ab} \theta \begin{bmatrix} a \\ b \end{bmatrix}^4 \frac{\Gamma_{(6,22)}}{\eta^{12} \bar{\eta}^{24}}, \quad (2.1)$$

where the Euclidean time direction is compactified on a circle of radius R_0 . V_3 is the volume of the three large spatial directions. The $\Gamma_{(6,22)}$ lattice is associated to the zero mode contribution of the internal six-manifold, along with the $E_8 \times E_8$ or $SO(32)$ right-moving lattice. For instance, the $E_8 \times E_8$ case on T^6 gives the block

$$\Gamma_{(6,22)} = \Gamma_{(6,6)} \frac{1}{2} \sum_{\gamma,\delta} \bar{\theta}[\gamma]_8^8 \frac{1}{2} \sum_{\gamma',\delta'} \bar{\theta}[\gamma']_8^8, \quad (2.2)$$

where γ , δ and γ' , δ' are integers defined modulo 2.

We wish to implement a non-zero temperature in the model. This is done by coupling the momentum and winding quantum numbers associated to the Euclidean time circle to the space-time fermion number F [13–15] [5]. The contributions of the right-moving world-sheet degrees of freedom to F are always even. Thus, at the level of the one-loop string partition function, the operation amounts to replacing the $\Gamma_{(1,1)}(R_0)$ lattice with

$$\Gamma_{(1,1)}(R_0) \rightarrow \sum_{h_0, \tilde{g}_0} \Gamma_{(1,1)}^{[h_0]}(R_0) (-)^{a\tilde{g}_0 + bh_0 + \tilde{g}_0 h_0}, \quad (2.3)$$

where $\Gamma_{(1,1)}^{[h]}_{[\tilde{g}]}$ is a \mathbb{Z}_2 -shifted lattice [25] given by [13–15] [5]

$$\Gamma_{1,1}^{[h]}_{[\tilde{g}]}(R) = \frac{R}{\sqrt{\tau_2}} \sum_{\tilde{m}, n} e^{-\frac{\pi R^2}{\tau_2} |(2\tilde{m} + \tilde{g}) + (2n + h)\tau|^2}. \quad (2.4)$$

The spontaneous breaking of space-time supersymmetry is done by coupling the two $\Gamma_{(1,1)}$ lattices associated with the internal circles of radii R_4 and R_5 to generic R-symmetry charges [5, 13–15, 21]. In the case of models with $N_4 = 4$ initial supersymmetry, all such charges associated to the left-moving world-sheet degrees of freedom are equivalent by symmetry. Different choices exist involving right-moving gauge R-charges [5]. For example, consider the $E_8 \times E_8$ models and decompose the E_8 representations in terms of $SO(16)$ ones. One can choose R-charges which are odd for the $SO(16)$ spinorial representations and even for the others¹.

¹In the $N_2 = 2$ orbifold models, one can choose R-charges associated to the twisted T^4 planes to which the left-moving world-sheet degrees of freedom contribute as well [5].

We will present a class of cases, where starting with $N_4 = 4$, $E_8 \times E_8$ vacua, space-time supersymmetry is broken if we couple the X^i lattices, $i = 4, 5$, to $F + \bar{Q}_i$, where the right-moving charges \bar{Q}_i are odd for the $SO(16)$ spinorial representations associated with one or both the E_8 factors. So we replace the X^4 and X^5 lattices as follows:

$$\Gamma_{(1,1)}(R_i) \rightarrow \sum_{h_i, \tilde{g}_i} \Gamma_{(1,1)}[\frac{h_i}{\tilde{g}_i}](R_i) (-)^{(a+\bar{Q}_i)\tilde{g}_i+(b+\bar{L}_i)h_i+\epsilon_i\tilde{g}_i h_i} . \quad (2.5)$$

\bar{Q}_i , $i = 4, 5$, can be identified to be either γ , γ' or $\gamma + \gamma'$. [γ and γ' are odd for the corresponding $SO(16)$ spinorial representations.] \bar{L}_i is equal to δ , δ' or $\delta + \delta'$ respectively, as dictated by modular invariance, and $\epsilon_i = 0, 1$ depending on the modular transformation $\tau \rightarrow \tau + 1$. Under this,

$$a + \bar{Q}_i \rightarrow a + \bar{Q}_i , \quad b + \bar{L}_i \rightarrow a + b + \bar{Q}_i + \bar{L}_i + \epsilon_i . \quad (2.6)$$

For instance, for $(\bar{Q}_i, \bar{L}_i) \equiv (\gamma + \gamma', \delta + \delta')$ one has $\epsilon_i = 1$, while for $(\bar{Q}_i, \bar{L}_i) \equiv (\gamma, \delta)$ or (γ', δ') , ϵ_i vanishes. With these modifications taken into account, the one-loop string partition function is given by

$$\begin{aligned} Z &= \frac{V_3}{(2\pi)^3} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{5/2}} \frac{\Gamma_{(4,4)}}{\eta^{12}\bar{\eta}^{24}} \frac{1}{2} \sum_{a,b} (-)^{a+b+ab} \theta \left[\frac{a}{b} \right]^4 \\ &\times \frac{1}{4} \sum_{\gamma, \delta} \sum_{\gamma', \delta'} \bar{\theta}[\frac{\gamma}{\delta}]^8 \bar{\theta}[\frac{\gamma'}{\delta'}]^8 \prod_{i=0,4,5} \sum_{h_i, \tilde{g}_i} \Gamma_{(1,1)}[\frac{h_i}{\tilde{g}_i}] (-)^{(a+\bar{Q}_i)\tilde{g}_i+(b+\bar{L}_i)h_i+\epsilon_i\tilde{g}_i h_i} . \end{aligned} \quad (2.7)$$

In this equation, we have $\bar{Q}_0 = \bar{L}_0 = 0$. Redefining $a = \hat{a} + \sum_i h_i$ and $b = \hat{b} + \sum_i \tilde{g}_i$, and using the Jacobi identity [5], one obtains

$$\begin{aligned} Z &= -\frac{V_3}{(2\pi)^3} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{5/2}} \frac{\Gamma_{(4,4)}}{\eta^{12}\bar{\eta}^{24}} \sum_{h_i, \tilde{g}_i} \theta \left[\frac{1+\sum_i h_i}{1+\sum_i \tilde{g}_i} \right]^4 (-)^{\sum_i h_i + \sum_i \tilde{g}_i + (\sum_i h_i)(\sum_i \tilde{g}_i)} \\ &\times \frac{1}{4} \sum_{\gamma, \delta} \sum_{\gamma', \delta'} \bar{\theta}[\frac{\gamma}{\delta}]^8 \bar{\theta}[\frac{\gamma'}{\delta'}]^8 \prod_i \Gamma_{(1,1)}[\frac{h_i}{\tilde{g}_i}] (-)^{\epsilon_i h_i \tilde{g}_i + \bar{Q}_i \tilde{g}_i + \bar{L}_i h_i} . \end{aligned} \quad (2.8)$$

In the large radii regime $R_i \gg R_H$, where R_H is the Hagedorn radius, the system is free of tachyons. The odd winding sectors, $h_i = 1$, are exponentially suppressed. In this regime only the sectors $h_i = 0$, $i = 0, 4, 5$ and $\tilde{g}_0 + \tilde{g}_4 + \tilde{g}_5 = 1$ modulo 2 contribute significantly (the latter condition due to the fact that $\theta[\frac{1}{1}]$ vanishes identically). Furthermore, if the internal lattice $\Gamma_{(4,4)}$ moduli are kept to be of order unity, we can express the leading contributions

as the following integral [5]:

$$\begin{aligned}
Z &= \frac{V_3}{(2\pi)^3} R_0 R_4 R_5 \sum_{\tilde{g}_i} \frac{1 - (-)^{\sum_i \tilde{g}_i}}{2} \sum_s (-)^{\tilde{Q}_4(s)\tilde{g}_4 + \tilde{Q}_5(s)\tilde{g}_5} \\
&\times \int_0^\infty \frac{d\tau_2}{\tau_2^4} \sum_{\tilde{m}_i} e^{-\frac{\pi}{\tau_2} [(2\tilde{m}_0 + \tilde{g}_0)^2 R_0^2 + (2\tilde{m}_4 + \tilde{g}_4)^2 R_4^2 + (2\tilde{m}_5 + \tilde{g}_5)^2 R_5^2]}. \tag{2.9}
\end{aligned}$$

In the first line the sum over s runs over the $2^3 \times 504$ massless boson/fermion pairs of the initially supersymmetric model. The contributions of massive states are exponentially suppressed, of order $e^{-\pi R_i}$. The integral gives the pressure in the string frame:

$$\begin{aligned}
P_{\text{string}} &= \frac{Z}{V_4} = \frac{R_4 R_5}{(2\pi)^4} \frac{2}{\pi^3} \sum_{\tilde{g}_i} \frac{1 - (-)^{\sum_i \tilde{g}_i}}{2} \sum_s (-)^{\tilde{g}_4 \tilde{Q}_4(s) + \tilde{g}_5 \tilde{Q}_5(s)} \\
&\times \sum_{\tilde{m}_i} \frac{1}{\left[(2\tilde{m}_0 + \tilde{g}_0)^2 R_0^2 + (2\tilde{m}_4 + \tilde{g}_4)^2 R_4^2 + (2\tilde{m}_5 + \tilde{g}_5)^2 R_5^2 \right]^3}. \tag{2.10}
\end{aligned}$$

We parameterize the various moduli as follows:

$$\begin{aligned}
T &:= \frac{1}{2\pi R_0 \sqrt{\text{Re}S}}, & M &:= \frac{1}{2\pi \sqrt{T} \text{Re}S}, & \text{Re}T_1 &:= R_4 R_5 \equiv \mathcal{T}, \\
\text{Re}U_1 &:= \frac{R_5}{R_4} \equiv \mathcal{U}, & u &:= \frac{R_0}{\sqrt{T}}, \tag{2.11}
\end{aligned}$$

where S is the 4d dilaton-axion modulus, $\text{Re}S = e^{-2\phi_D}$. The two supersymmetry breaking scales, in the Einstein frame, are the temperature T and the gravitino mass scale M . The pressure in this frame is related to the string frame pressure by:

$$P = \frac{1}{(\text{Re}S)^2} P_{\text{string}} = T^4 p(u, \mathcal{U}), \tag{2.12}$$

with

$$p(u, \mathcal{U}) = n_{100} p_{100}(u, \mathcal{U}) + n_{010} p_{010}(u, \mathcal{U}) + n_{001} p_{001}(u, \mathcal{U}) + n_{111} p_{111}(u, \mathcal{U}). \tag{2.13}$$

The coefficients $n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$ are given in terms of the supersymmetry breaking R-charges,

$$n_{100} = 2^3 \times 504, \quad n_{010} = \sum_s (-)^{\tilde{Q}_4(s)}, \quad n_{001} = \sum_s (-)^{\tilde{Q}_5(s)}, \quad n_{111} = \sum_s (-)^{\tilde{Q}_4(s) + \tilde{Q}_5(s)}, \tag{2.14}$$

while the dependence on the complex structure moduli u and \mathcal{U} involves the shifted Eisenstein functions:

$$p_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}(u, \mathcal{U}) = \frac{2}{\pi^3} \sum_{\tilde{m}_0, \tilde{m}_4, \tilde{m}_5} \frac{u^4}{\left[(2\tilde{m}_0 + \tilde{g}_0)^2 u^2 + (2\tilde{m}_4 + \tilde{g}_4)^2 \mathcal{U}^{-1} + (2\tilde{m}_5 + \tilde{g}_5)^2 \mathcal{U} \right]^3}. \tag{2.15}$$

2.2 Type II models

We construct Type II models with similar thermal and supersymmetry breaking properties. In these examples, the internal manifold involves either a T^4 factor for $N_4 = 8$ initial supersymmetry or a T^4/\mathbb{Z}_2 factor for $N_4 = 4$. Orientifolds of these lead to models with $N_4 = 4$ and $N_4 = 2$ initial supersymmetry respectively, and include open string matter sectors. At weak coupling in four dimensions, these are dual to heterotic models [26], some of which we considered in the previous section. Models with $N_4 = 2$ initial supersymmetry can also be constructed if we start with a $T^6/(\mathbb{Z}_2 \times \mathbb{Z}'_2)$ orbifold [27]. We illustrate the derivation of the pressure in the intermediate cosmological region, with $T \ll T_H$, for the type II $N_4 = 4$ models, but the results can be generalized to the other cases.

The $N_4 = 4$ partition function is

$$Z = \frac{V_3}{(2\pi)^3} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{5/2}} \frac{1}{(\eta\bar{\eta})^8} \prod_{i=0,4,5} \Gamma_{(1,1)}(R_i) \frac{1}{2} \sum_{H,G} Z_{(4,4)}[{}^H_G] \\ \times \frac{1}{2} \sum_{a,b} \theta_{[b]}^{[a]^2} \theta_{[b+G]}^{[a+H]} \theta_{[b-G]}^{[a-H]} (-)^{a+b+ab} \frac{1}{2} \sum_{\bar{a},\bar{b}} \bar{\theta}_{[\bar{b}]}^{[\bar{a}]^2} \bar{\theta}_{[\bar{b}+G]}^{[\bar{a}+H]} \bar{\theta}_{[\bar{b}-G]}^{[\bar{a}-H]} (-)^{\bar{a}+\bar{b}+\bar{a}\bar{b}}. \quad (2.16)$$

The T^4/\mathbb{Z}_2 part is given by [25]:

$$Z_{(4,4)}[{}^H_G] = \frac{\Gamma_{(4,4)}}{(\eta\bar{\eta})^4}, \quad \text{when } (H, G) = (0, 0) \\ Z_{(4,4)}[{}^H_G] = \frac{2^4 \eta^2 \bar{\eta}^2}{\theta_{[1-G]}^{[1-H]^2} \bar{\theta}_{[1-G]}^{[1-H]^2}}, \quad \text{when } (H, G) \neq (0, 0). \quad (2.17)$$

The characters H, G are integers defined modulo 2.

As usual, the finite temperature is implemented by inserting the thermal co-cycle and replacing the Euclidean time lattice as follows [13–15]:

$$\Gamma_{(1,1)}(R_0) \rightarrow \sum_{h_0, \tilde{g}_0} \Gamma_{(1,1)}[{}^{h_0}_{\tilde{g}_0}](R_0) (-)^{(a+\bar{a})\tilde{g}_0+(b+\bar{b})h_0}. \quad (2.18)$$

In contrast to the heterotic case, the contributions to the space-time fermion number F from both the left-moving and right-moving sectors can be odd or even. In the sequel, we denote by F_L the contribution of the world-sheet left-movers to the space-time fermion number and similarly for F_R .

There are several ways to break the initial $N_4 = 4$ supersymmetry spontaneously, either by symmetric or asymmetric geometrical fluxes [20, 21, 23]. The two left-moving space-time

supersymmetries can be broken if we couple either or both the X^4 and X^5 lattice charges to F_L , or to left-moving R-charges associated with the twisted planes: $F_L + Q_i$. Also, the two right-moving space-time supersymmetries are broken by coupling the lattice charges to F_R or to $F_R + \bar{Q}_i$. Each lattice is replaced as follows [19]:

$$\Gamma_{(1,1)}(R_i) \rightarrow \sum_{h_i, \tilde{g}_i} \Gamma_{(1,1)}^{[h_i]_{[\tilde{g}_i]}}(R_i) (-)^{[(a+Q_i)\tilde{g}_i+(b+L_i)h_i+\tilde{g}_i h_i]\epsilon_i} (-)^{[(\bar{a}+\bar{Q}_i)\tilde{g}_i+(\bar{b}+\bar{L}_i)h_i+\tilde{g}_i h_i]\bar{\epsilon}_i}, \quad (2.19)$$

where Q_i, \bar{Q}_i can be set to zero or identified with the twist charge H . Correspondingly L_i, \bar{L}_i can be set to zero or identified with the character G . Also, we have introduced the parameters $\epsilon_i, \bar{\epsilon}_i$, taking the values 0 or 1, to indicate whether we couple the circle i to the left- or right-movers.

In particular, we will examine 3 distinct cases where $N_4 = 4$ is spontaneously broken to $N_4 = 0$ (and then thermalized):

-*Case 1* : Two asymmetric breakings, e.g. $(\epsilon_4, \bar{\epsilon}_4) = (1, 0)$, $(\epsilon_5, \bar{\epsilon}_5) = (0, 1)$.

-*Case 2* : One symmetric and one asymmetric breaking, e.g. $(\epsilon_4, \bar{\epsilon}_4) = (1, 1)$, $(\epsilon_5, \bar{\epsilon}_5) = (0, 1)$.

-*Case 3* : Two symmetric breakings, $(\epsilon_4, \bar{\epsilon}_4) = (1, 1)$, $(\epsilon_5, \bar{\epsilon}_5) = (1, 1)$.

In addition, we will consider a case where the $N_4 = 4$ supersymmetry is partially broken to $N_4 = 2$. The remaining supersymmetries are then broken by thermal effects:

-*Case 1'* : Two left-moving asymmetric breakings, e.g. $(\epsilon_4, \bar{\epsilon}_4) = (1, 0)$, $(\epsilon_5, \bar{\epsilon}_5) = (1, 0)$.

The partition function can be written as follows:

$$\begin{aligned} Z &= \frac{V_3}{(2\pi)^3} R_0 R_4 R_5 \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^4} \frac{1}{(\eta\bar{\eta})^8} \frac{1}{2} \sum_{H,G} Z_{(4,4)}^{[H]_G} \\ &\times \frac{1}{2} \sum_{a,b} \theta_{[b]}^{[a]^2} \theta_{[b+G]}^{[a+H]} \theta_{[b-G]}^{[a-H]} (-)^{a+b+ab} \frac{1}{2} \sum_{\bar{a},\bar{b}} \bar{\theta}_{[\bar{b}]}^{[\bar{a}]^2} \bar{\theta}_{[\bar{b}+G]}^{[\bar{a}+H]} \bar{\theta}_{[\bar{b}-G]}^{[\bar{a}-H]} (-)^{\bar{a}+\bar{b}+\bar{a}\bar{b}} \\ &\times \prod_{i=0,4,5} \left\{ \sum_{h_i, \tilde{g}_i} \sum_{\tilde{m}_i, n_i} e^{-\frac{\pi R_i^2}{\tau_2} |(2\tilde{m}_i + \tilde{g}_i) + (2n_i + h_i)\tau|^2} (-)^{[(a+Q_i)\tilde{g}_i+(b+L_i)h_i+\tilde{g}_i h_i]\epsilon_i + [(\bar{a}+\bar{Q}_i)\tilde{g}_i+(\bar{b}+\bar{L}_i)h_i+\tilde{g}_i h_i]\bar{\epsilon}_i} \right\}. \end{aligned} \quad (2.20)$$

Here $(\epsilon_0, \bar{\epsilon}_0) = (1, 1)$ and $(Q_0, \bar{Q}_0) = (L_0, \bar{L}_0) = (0, 0)$.

As in the heterotic case, we are interested in the regime where the radii R_i , $i = 0, 4, 5$, are much bigger than the Hagedorn radius, $R_i \gg R_H$. In this intermediate cosmological

regime, the system is free of any tachyonic instabilities. The odd winding sectors, $h_i = 1$, are exponentially suppressed, and the only significant contributions to the partition function occur for $h_i = 0$, $\tilde{g}_0 + \epsilon_4 \tilde{g}_4 + \epsilon_5 \tilde{g}_5 = 1$ modulo 2 and $\tilde{g}_0 + \bar{\epsilon}_4 \tilde{g}_4 + \bar{\epsilon}_5 \tilde{g}_5 = 1$ modulo 2.

The pressure receives contributions from the untwisted sector, $H = 0$, and from the twisted sector $H = 1$. In the untwisted sector, the result is given by

$$\begin{aligned} \frac{Z_{\text{untwisted}}}{V_4} &= \frac{R_4 R_5}{(2\pi)^4} \sum_{\tilde{g}_i} \frac{1 - (-)^{\sum_i \epsilon_i \tilde{g}_i}}{2} \frac{1 - (-)^{\sum_i \bar{\epsilon}_i \tilde{g}_i}}{2} \\ &\times \frac{2}{\pi^3} \sum_{\tilde{m}_i} \frac{n_0^{\text{untwisted}}}{\left[(2\tilde{m}_0 + \tilde{g}_0)^2 R_0^2 + (2\tilde{m}_4 + \tilde{g}_4)^2 R_4^2 + (2\tilde{m}_5 + \tilde{g}_5)^2 R_5^2 \right]^3}. \end{aligned} \quad (2.21)$$

$n_0^{\text{untwisted}}$ is the number of massless boson/fermion pairs in the untwisted sector of the initially supersymmetric $N_4 = 4$ model. We have $n_0^{\text{untwisted}} = n_0/2$, where $n_0 = 2^7$ counts the massless pairs of the $N_4 = 8$ model; the factor of 1/2 is due to the orbifolding.

In the twisted sector we have

$$\begin{aligned} \frac{Z_{\text{twisted}}}{V_4} &= \frac{R_4 R_5}{(2\pi)^4} \sum_{\tilde{g}_i} \frac{1 - (-)^{\sum_i \epsilon_i \tilde{g}_i}}{2} \frac{1 - (-)^{\sum_i \bar{\epsilon}_i \tilde{g}_i}}{2} (-)^{(\epsilon_4 Q_4 + \bar{\epsilon}_4 \bar{Q}_4) \tilde{g}_4 + (\epsilon_5 Q_5 + \bar{\epsilon}_5 \bar{Q}_5) \tilde{g}_5} \\ &\times \frac{2}{\pi^3} \sum_{\tilde{m}_i} \frac{n_0^{\text{twisted}}}{\left[(2\tilde{m}_0 + \tilde{g}_0)^2 R_0^2 + (2\tilde{m}_4 + \tilde{g}_4)^2 R_4^2 + (2\tilde{m}_5 + \tilde{g}_5)^2 R_5^2 \right]^3}. \end{aligned} \quad (2.22)$$

$n_0^{\text{twisted}} = 2^8/2$ is the number of massless boson/fermion pairs in the twisted sector of the initially supersymmetric $N_4 = 4$ model. The $Q_4, \bar{Q}_4, Q_5, \bar{Q}_5$ appearing in Eq. (2.22) can be either zero or identified to the twist charge $H = 1$.

Using the definitions of the moduli introduced in Eq. (2.11), the pressure P is taking the same form as in Eq. (2.12) with

$$p(u, \mathcal{U}) = \sum_{\substack{\tilde{g}_0 + \epsilon_4 \tilde{g}_4 + \epsilon_5 \tilde{g}_5 = 1 \text{ mod } 2 \\ \tilde{g}_0 + \bar{\epsilon}_4 \tilde{g}_4 + \bar{\epsilon}_5 \tilde{g}_5 = 1 \text{ mod } 2}} n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5} p_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}(u, \mathcal{U}), \quad (2.23)$$

where the functions $p_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$ are given in Eq. (2.15), and the coefficients $n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$ are similarly defined, in terms of supersymmetry breaking R-charges, as in Eq. (2.14),

$$n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5} = n_0 = \frac{2^8}{2}, \quad \text{for } N_4 = 8, \quad (2.24)$$

$$n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5} = \frac{n_0}{2} \left(1 + 2(-)^{(\epsilon_4 Q_4 + \bar{\epsilon}_4 \bar{Q}_4) \tilde{g}_4 + (\epsilon_5 Q_5 + \bar{\epsilon}_5 \bar{Q}_5) \tilde{g}_5} \right), \quad \text{for } N_4 = 4. \quad (2.25)$$

In the $N_4 = 4$ cases, the coefficients $n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$ can take negative values as well. The results can be generalized to $N_4 = 2$ models.

2.3 The zero temperature limit

Setting $T = 0$, or $R_0 \rightarrow \infty$, in Eqs (2.10), (2.21) and (2.22), we can obtain the one-loop effective potential at zero temperature. It arises from quantum effects due to the spontaneous breaking of supersymmetry. For the heterotic models, the effective potential takes the form

$$\mathcal{V} = M^4 v(\mathcal{U}), \quad (2.26)$$

where M is defined in (2.11), and

$$v(\mathcal{U}) = n_{10} v_{10}(\mathcal{U}) + n_{01} v_{01}(\mathcal{U}). \quad (2.27)$$

The coefficients $n_{\tilde{g}_4 \tilde{g}_5}$ are determined in terms of the R-charges,

$$n_{\tilde{g}_4 \tilde{g}_5} = \sum_s (-)^{\tilde{g}_4 \bar{Q}_4(s) + \tilde{g}_5 \bar{Q}_5(s)}, \quad (2.28)$$

and

$$v_{\tilde{g}_4 \tilde{g}_5}(\mathcal{U}) = -\frac{2}{\pi^3} \sum_{\tilde{m}_4, \tilde{m}_5} \frac{1}{\left[(2\tilde{m}_4 + \tilde{g}_4)^2 \mathcal{U}^{-1} + (2\tilde{m}_5 + \tilde{g}_5)^2 \mathcal{U} \right]^3}. \quad (2.29)$$

The type II effective potential takes a form similar to the heterotic one, as in Eq. (2.26), where now

$$v(\mathcal{U}) = \sum_{\substack{\epsilon_4 \tilde{g}_4 + \epsilon_5 \tilde{g}_5 = 1 \pmod{2} \\ \bar{\epsilon}_4 \tilde{g}_4 + \bar{\epsilon}_5 \tilde{g}_5 = 1 \pmod{2}}} n_{\tilde{g}_4 \tilde{g}_5} v_{\tilde{g}_4 \tilde{g}_5}(\mathcal{U}), \quad (2.30)$$

with $n_{\tilde{g}_4 \tilde{g}_5} = n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$, given in (2.25) (or (2.24)).

2.4 Non thermal cosmologies

In the zero temperature limit, the 1-loop effective action takes the form:

$$S = \int d^4x \sqrt{-\det g} \left\{ \frac{1}{2} R - g^{\mu\nu} \left(\frac{\partial_\mu S \partial_\nu \bar{S}}{(S + \bar{S})^2} + \frac{\partial_\mu T_1 \partial_\nu \bar{T}_1}{(T_1 + \bar{T}_1)^2} + \frac{\partial_\mu U_1 \partial_\nu \bar{U}_1}{(U_1 + \bar{U}_1)^2} \right) - \mathcal{V} \right\}. \quad (2.31)$$

All other moduli can be frozen since they do not appear in the effective potential. More precisely, their contributions to the effective potential are exponentially suppressed. Freezing further $\text{Im}S$, $\text{Im}T_1$ and $\text{Im}U_1$, we obtain for the Lagrangian

$$\mathbf{L} = \frac{1}{2} R - \frac{1}{2} \left((\partial\phi_S)^2 + (\partial\phi_T)^2 + (\partial\phi_U)^2 \right) - e^{-2\sqrt{2}(\phi_S + \phi_T)} \frac{v(\mathcal{U})}{(2\pi)^4}, \quad (2.32)$$

where

$$\text{Re}S := e^{\sqrt{2}\phi_S}, \quad \mathcal{T} := e^{\sqrt{2}\phi_{\mathcal{T}}}, \quad \mathcal{U} := e^{\sqrt{2}\phi_{\mathcal{U}}}. \quad (2.33)$$

It is useful to redefine the fields as follows

$$\begin{pmatrix} \phi \\ \phi_- \end{pmatrix} := \begin{pmatrix} -1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix} \begin{pmatrix} \phi_S \\ \phi_{\mathcal{T}} \end{pmatrix} \quad (2.34)$$

since the field ϕ_- does not appear in the potential:

$$\mathbb{L} = \frac{1}{2}R - \frac{1}{2}((\partial\phi)^2 + (\partial\phi_-)^2 + (\partial\phi_{\mathcal{U}})^2) - M^4 v(\mathcal{U}) \quad \text{with} \quad M = \frac{e^\phi}{2\pi}. \quad (2.35)$$

We look for homogeneous and isotropic solutions where the metric is of the Friedmann-Robertson-Walker form, with vanishing spatial curvature:

$$ds^2 = -N(t)^2 dt^2 + a(t)^2 dx^i dx^i, \quad H \equiv \left(\frac{\dot{a}}{a} \right). \quad (2.36)$$

Here N is the laps function, $a(t)$ the scale factor and H the Hubble parameter. In the gauge choice $N = 1$, the gravitational field equations are

$$3H^2 = \frac{1}{2}\dot{\phi}^2 + \frac{1}{2}\dot{\phi}_-^2 + \frac{1}{2}\dot{\phi}_{\mathcal{U}}^2 + \mathcal{V}, \quad (2.37)$$

$$2\dot{H} + 3H^2 = -\frac{1}{2}\dot{\phi}^2 - \frac{1}{2}\dot{\phi}_-^2 - \frac{1}{2}\dot{\phi}_{\mathcal{U}}^2 + \mathcal{V}. \quad (2.38)$$

Their linear sum is independent of the fields kinetic terms:

$$\dot{H} + 3H^2 = \mathcal{V} = M^4 v(\mathcal{U}). \quad (2.39)$$

Eq. (2.38) follows by differentiating the Friedmann-Hubble Eq. (2.37), once the following moduli field equations are satisfied:

$$\ddot{\phi} + 3H\dot{\phi} = -\frac{\partial\mathcal{V}}{\partial\phi} = -4M^4 v(\mathcal{U}), \quad (2.40)$$

$$\ddot{\phi}_{\mathcal{U}} + 3H\dot{\phi}_{\mathcal{U}} = -\frac{\partial\mathcal{V}}{\partial\phi_{\mathcal{U}}} = -\sqrt{2}M^4 \mathcal{U} v'(\mathcal{U}), \quad (2.41)$$

$$\ddot{\phi}_- + 3H\dot{\phi}_- = 0, \quad (2.42)$$

where prime derivatives are with respect to \mathcal{U} . The last equation can be integrated giving

$$\frac{1}{2}\dot{\phi}_-^2 = \frac{c_-}{a^6}, \quad (2.43)$$

where c_- is a positive constant. Eq. (2.41) can be satisfied for a constant \mathcal{U} , if there exists a solution to

$$v'(\mathcal{U}) = 0. \quad (2.44)$$

We will look for models for which this extremum is a local minimum so that the complex structure modulus \mathcal{U} is stabilized. The compatibility of Eqs (2.39) and (2.40) requires that there exists a constant c_ϕ such that:

$$\dot{\phi} = -4H + \frac{c_\phi}{a^3}. \quad (2.45)$$

In [22], it is shown that the solution for $c_\phi = 0$ is an attractor. Thus, we concentrate on the case $c_\phi = 0$, so that

$$M \equiv \frac{e^\phi}{2\pi} = M_0 \left(\frac{a_0}{a}\right)^4, \quad (2.46)$$

where $M_0 a_0^4$ is a positive integration constant.

The Hubble equation takes the form:

$$3H^2 = -\frac{c_m}{a^6} + \frac{c}{a^{16}} \quad \text{where} \quad c_m = \frac{3}{5} c_- > 0, \quad c = -v(\mathcal{U}) \times \frac{3}{5} M_0^4 a_0^{16}. \quad (2.47)$$

If $c > 0$, one has for $c_m = 0$

$$a(t) = A t^{1/8} \quad \text{where} \quad A = 2^{3/8} \left(\frac{c}{3}\right)^{1/16}. \quad (2.48)$$

When the kinetic energy for ϕ_- is switched on, i.e. when $c_m > 0$, one has a big bang/big crunch cosmology. The solution $t(a)$ is given by

$$t(a) = \pm t_0 \int_{a/A}^1 \frac{x^7 dx}{\sqrt{1-x^{10}}}, \quad 0 \leq a \leq A \quad \text{where} \quad A = \left(\frac{c}{c_m}\right)^{1/10}, \quad t_0 = \sqrt{3} c^{3/10} c^{-4/5}. \quad (2.49)$$

We investigate whether some of the heterotic and type II models we considered satisfy the minimization condition (2.44), which fixes the modulus \mathcal{U} , and the positivity of the parameter c , Eq. (2.47), which allows for real time solutions:

$$\textit{Extremum} : v'(\mathcal{U}) = 0, \quad \textit{Stability} : v''(\mathcal{U}) > 0, \quad \textit{Real time} : v(\mathcal{U}) < 0. \quad (2.50)$$

The shape of the potential as a function of \mathcal{U} depends on the R-symmetry breaking charges which define the coefficients $n_{\tilde{g}_4 \tilde{g}_5}$. Since the functions $v_{\tilde{g}_4 \tilde{g}_5}$ in Eq. (2.29) are negative, we need some of the $n_{\tilde{g}_4 \tilde{g}_5}$ to be positive.

In the heterotic models, the functions v_{10} and v_{01} defining the effective potential, Eq. (2.27), are monotonic functions of \mathcal{U} , the former decreasing and the latter increasing, (see Fig. 1a). To have an extremum, n_{10} and n_{01} must be of the same sign. The real time condition requires




Figure 1: The functions $v_{\tilde{g}_4 \tilde{g}_5}$ versus $\ln \mathcal{U}$.

this sign to be positive, which implies that the solution is unstable under fluctuations of \mathcal{U} . Stable cosmological solutions exist only at non-zero temperature, as we will see in the next section. We may have, however, stationary domain wall solutions.

We now study the effective potential given in Eq. (2.30) for the different type II cases.

-*Case 1* : The potential is proportional to the function v_{11} , which is invariant under $\mathcal{U} \rightarrow 1/\mathcal{U}$. This implies that a stationary point, which is a minimum, occurs at $\mathcal{U} = 1$, (see Fig. 1b). The conditions (2.50) are simultaneously satisfied if $n_{11} > 0$. This can be realized by choosing $Q_4 = \bar{Q}_5 = 0$ or H .

-*Case 2* : Since the potential is proportional to the monotonic function $v_{01}(\mathcal{U})$, there is always a runaway behavior for this model: $\mathcal{U} \rightarrow 0$ *i.e.* $R_4 \rightarrow +\infty$ for $n_{01} > 0$ and $\mathcal{U} \rightarrow +\infty$ *i.e.* $R_5 \rightarrow +\infty$ for $n_{01} < 0$. Thus, the dynamics drives the system out of the scope of our analysis, and one should look for a solution in five dimensions, where supersymmetry is spontaneously broken by the remaining finite size internal radius.

-*Case 3* : This type II model has an effective potential of the form encountered in the heterotic case.

-*Case 1'* : The model is supersymmetric with a flat effective potential.

At finite temperature, we are going to see that the situations in the type II cases also change drastically.

3 Thermal cosmologies

3.1 Equations of motion and thermodynamics

Using the field redefinitions (2.33)–(2.34) and the FRW ansatz (2.36), the thermal effective action is

$$S = -\frac{1}{6} \int dt N a^3 \left(\frac{3}{N^2} H^2 - \frac{1}{2N^2} \dot{\phi}^2 - \frac{1}{2N^2} \dot{\phi}_-^2 - \frac{1}{2N^2} \dot{\phi}_u^2 - P \right), \quad (3.1)$$

where the pressure can be expressed in terms of the supersymmetry breaking scale $M(\phi)$ as

$$P = M^4 \frac{p(u, \mathcal{U})}{u^4}, \quad M = \frac{e^\phi}{2\pi}, \quad u = \frac{M}{T}. \quad (3.2)$$

The gravity equations are obtained by varying with respect to N and a ,

$$\frac{3}{N^2} H^2 = \frac{1}{2N^2} \dot{\phi}^2 + \frac{1}{2N^2} \dot{\phi}_-^2 + \frac{1}{2N^2} \dot{\phi}_u^2 + \rho \quad \text{with} \quad \rho := -P - N \frac{\partial P}{\partial N}, \quad (3.3)$$

$$\frac{2}{N^2} \dot{H} + \frac{3}{N^2} H^2 - \frac{2}{N^3} H \dot{N} = -\frac{1}{2N^2} \dot{\phi}^2 - \frac{1}{2N^2} \dot{\phi}_-^2 - \frac{1}{2N^2} \dot{\phi}_u^2 - P - \frac{1}{3} a \frac{\partial P}{\partial a}, \quad (3.4)$$

where in general P can depend on N and a . To determine the dependence, we recall in which specific frame we computed P :

$$P = \frac{Z}{(\text{Re}S)^2 V_4} = \frac{Z}{N a^3} \quad \Longrightarrow \quad N = 2\pi R_0 \sqrt{\text{Re}S} = \frac{1}{T}, \quad a = 2\pi R \sqrt{\text{Re}S}, \quad (3.5)$$

where $R_1 = R_2 = R_3 \equiv R$ are the radii of the large three spatial directions (before the large volume limit $R \rightarrow +\infty$ is taken). Since $P = T^4 p(u, \mathcal{U})$ and T is identified with the inverse of the laps function, we have

$$\rho = T \frac{\partial P}{\partial T} - P \quad \text{and} \quad \frac{\partial P}{\partial a} = 0. \quad (3.6)$$

It is remarkable that these expressions are identical to the ones derived from thermodynamics. We thus show that the variational principle is in perfect agreement with thermodynamics.

Since P (in the action (3.1)) and ρ (in Eq. (3.3)) are scalars under time reparameterizations, we can write the gravitational equations in the simple gauge $N = 1$ as follows:

$$3H^2 = \frac{1}{2} \dot{\phi}^2 + \frac{1}{2} \dot{\phi}_-^2 + \frac{1}{2} \dot{\phi}_u^2 + \rho, \quad (3.7)$$

$$2\dot{H} + 3H^2 = -\frac{1}{2} \dot{\phi}^2 - \frac{1}{2} \dot{\phi}_-^2 - \frac{1}{2} \dot{\phi}_u^2 - P. \quad (3.8)$$

Combining Eqs (2.12) and (3.6), we obtain

$$\rho = T^4 r(u, \mathcal{U}) \quad \text{where} \quad r = 3p - u \partial_u p. \quad (3.9)$$

The equations of motion for the moduli fields can be written as follows:

$$\ddot{\phi} + 3H\dot{\phi} = \frac{\partial P}{\partial \phi} \equiv T^4 \{3p(u, \mathcal{U}) - r(u, \mathcal{U})\}, \quad (3.10)$$

$$\ddot{\phi}_{\mathcal{U}} + 3H\dot{\phi}_{\mathcal{U}} = \frac{\partial P}{\partial \phi_{\mathcal{U}}} \equiv T^4 \sqrt{2} \mathcal{U} \partial_{\mathcal{U}} p(u, \mathcal{U}), \quad (3.11)$$

$$\ddot{\phi}_- + 3H\dot{\phi}_- = 0, \quad (3.12)$$

where the r.h.s. of Eq. (3.10) follows from Eqs (3.2) and (3.9).

The last equation gives (2.43). We would like to find solutions to the remaining system of equations, with \mathcal{U} stabilized. Then, Eq. (3.11) amounts to the following algebraic equation

$$D(u, \mathcal{U}) := \mathcal{U} \partial_{\mathcal{U}} p = 0, \quad (3.13)$$

requiring that u is also a constant. It follows that the time dependence of ρ and P arises from the T^4 pre-factors only. From the relations in (3.2), we have that

$$M(\phi) = uT \quad \Longrightarrow \quad \dot{\phi} = \frac{\dot{T}}{T}. \quad (3.14)$$

Instead of solving the scale factor Eq. (3.8), we choose to solve the equation that arises from the conservation of the energy-momentum tensor:

$$\frac{d}{dt} \left(\frac{1}{2} \dot{\phi}^2 + \frac{1}{2} \dot{\phi}_{\mathcal{U}}^2 + \frac{1}{2} \dot{\phi}_-^2 + \rho \right) + 3H \left(\dot{\phi}^2 + \dot{\phi}_{\mathcal{U}}^2 + \dot{\phi}_-^2 + \rho + P \right) = 0. \quad (3.15)$$

Using the equations of motion for the scalar fields and (3.14), this gives

$$\dot{\phi} = \frac{\dot{T}}{T} = -H \quad \Longrightarrow \quad aT = a_0 T_0, \quad (3.16)$$

where $a_0 T_0$ is a positive integration constant.

Next, we consider the linear sum of Eqs (3.8) and (3.7),

$$\dot{H} + 3H^2 = \frac{1}{2}(\rho - P) = \frac{1}{2} T^4 \{r(u, \mathcal{U}) - p(u, \mathcal{U})\}. \quad (3.17)$$

Using (3.16), the compatibility between this equation and (3.10) implies the following thermal equation of state:

$$\rho = 5P. \quad (3.18)$$

This is one of the main results of this paper. It reminds us of the analogous equation, $\rho = 4P$, derived in [5, 28], when a single modulus was participating in the spontaneous supersymmetry breaking mechanism. These results are very suggestive, and we conjecture that they will be generalized to the cases when more moduli participate in the supersymmetry breaking. When n such fields are involved, we expect the equation of state to take the form:

$$\rho = (3 + n)P. \quad (3.19)$$

In the two moduli case, which we are considering here, Eq. (3.18) can also be written as

$$C(u, \mathcal{U}) := (2 + u\partial_u)p = 0. \quad (3.20)$$

As a result, the complex structure ratios (u, \mathcal{U}) are determined by the equations $D = C = 0$. It is interesting that along this critical trajectory, complex structure moduli participating in the breaking of supersymmetry are stabilized, and thus the cosmology is characterized by a single running scale.

The time dependence of the scale factor is dictated by the Friedmann-Hubble Eq. (3.7). The latter takes the form

$$3H^2 = \frac{c_r}{a^4} + \frac{c_m}{a^6} \quad \text{where} \quad c_r = 6(a_0 T_0)^4 p(u, \mathcal{U}), \quad c_m = \frac{6}{5} c_- > 0. \quad (3.21)$$

When $c_m = 0$, the universe is effectively radiation dominated and a cosmological solution exists if the constant $p(u, \mathcal{U})$ is positive:

$$a(t) = B\sqrt{t} \quad \text{with} \quad B = \sqrt{2} \left(\frac{c_r}{3} \right)^{1/4}. \quad (3.22)$$

When c_m is non-trivial, the time t can be expressed as a function of the scale factor as follows:

$$\begin{aligned} \text{if } c_r > 0 : \quad t(a) &= t_0 \int_0^{a/B} \frac{x^2 dx}{\sqrt{1+x^2}}, \quad \forall a \geq 0, \\ \text{if } c_r < 0 : \quad t(a) &= \pm t_0 \int_{a/B}^1 \frac{x^2 dx}{\sqrt{1-x^2}}, \quad 0 \leq a \leq B, \end{aligned} \quad (3.23)$$

where

$$B = \sqrt{\frac{c_m}{|c_r|}} \quad \text{and} \quad t_0 = \sqrt{3} c_m |c_r|^{-3/2}. \quad (3.24)$$

In the explicit examples presented in the following section, we always find $c_r > 0$. It would be interesting to find if models with “negative effective radiation energy density”, c_r/a^4 , are allowed.

As noticed in [5], the fact that the cosmological evolution we have found behaves effectively like a four dimensional universe filled with thermal radiation is not in contradiction with the state equation $\rho = 5P$. The reason is that the total energy density and pressure contain the “cold” part associated to the kinetic energy of ϕ . When $c_m = 0$, one has $\dot{\phi}^2/2 = \rho/5$, so that

$$\begin{aligned}\rho_{\text{tot}} &= \frac{1}{2}\dot{\phi}^2 + \rho = \frac{6}{5}\rho \\ P_{\text{tot}} &= \frac{1}{2}\dot{\phi}^2 + P = \frac{2}{5}\rho,\end{aligned}\tag{3.25}$$

in agreement with the expected state equation $\rho_{\text{tot}} = 3P_{\text{tot}}$.

3.2 Some stringy examples

We examine whether the extremization condition (3.13) and the compatibility condition (3.20) are simultaneously satisfied in the various heterotic and type II models under consideration:

$$\textit{Extremum} : D(u, \mathcal{U}) = 0, \quad \textit{Compatibility} : C(u, \mathcal{U}) = 0, \quad \textit{Stability} : \partial_{\mathcal{U}}^2 p(u, \mathcal{U}) < 0. \tag{3.26}$$

As in the non-thermal situation, the shape of the potential depends on the R-symmetry breaking charges. Their choices determine the coefficients $n_{\tilde{g}_0\tilde{g}_4\tilde{g}_5}$ that satisfy $-n_{100} \leq n_{\tilde{g}_0\tilde{g}_4\tilde{g}_5} \leq n_{100}$.

However, in the heterotic cases, n_{111} , n_{010} and n_{001} are not totally arbitrary. It is convenient to parameterize the a priori allowed models by separating the n_{100} states into 4 groups, depending on their parity under the operators $(-)^{\tilde{Q}_4}$ and $(-)^{\tilde{Q}_5}$, as shown in Table 3.27.

	$n_{100}\xi_1$ states	$n_{100}\xi_2$ states	$n_{100}\xi_3$ states	$n_{100}(1 - \xi_1 - \xi_2 - \xi_3)$ states	
$(-)^{Q_4}$	+	+	-	-	(3.27)
$(-)^{Q_5}$	+	-	+	-	

We observe that the parameter space of models is the tetrahedron:

$$\left\{ (\xi_1, \xi_2, \xi_3) \in [0, 1]^3 \quad \text{such that} \quad \xi_1 + \xi_2 + \xi_3 \leq 1 \right\}, \tag{3.28}$$

which constrains the ratios:

$$r_{010} := \frac{n_{010}}{n_{100}} = 2(\xi_1 + \xi_2) - 1, \quad r_{001} := \frac{n_{001}}{n_{100}} = 2(\xi_1 + \xi_3) - 1, \quad r_{111} := \frac{n_{111}}{n_{100}} = 1 - 2(\xi_2 + \xi_3). \tag{3.29}$$

The conditions (3.26) and the $\xi_2 \leftrightarrow \xi_3$ duality symmetry can be visualized geometrically in terms of the tetrahedron representation. Some type II models are also characterized in terms of this representation.

In the large/small u and \mathcal{U} regimes, p contains exponentially suppressed contributions that we have to neglect by consistency. The dominant contributions take the form of a linear sum of a finite number of monomials $u^a \mathcal{U}^b$. The $(\ln u, \ln \mathcal{U})$ -plane is divided into 6 sectors inside of which a power expansion is defined, (see Fig. 2). The boundaries of these sectors are the lines $\mathcal{U} = \lambda u^\omega$, where $\omega = -2, 0, 2$. We show in the Appendix the power expansion of




Figure 2: The $(\ln u, \ln \mathcal{U})$ -complex plane can be divided in 6 sectors, I, \dots, VI , separated by 6 edges of slope $-2, 0$ or 2 . The power expansions of $p(u, \mathcal{U})$, which are listed in the Appendix are well defined in each sector.

$p(u, \mathcal{U})$ in each sector. More accurate expressions are also given along the lines $\mathcal{U} = \lambda u^{\pm 2, 0}$ that are useful to connect the power expansions on each sides. $D(u, \mathcal{U}) = 0$ and $C(u, \mathcal{U}) = 0$ define curves which asymptote to the lines $\mathcal{U} = \lambda u^{\pm 2, 0}$, for which we have determined the intercept λ as a function of the ratios $r_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$. The constraints are simultaneously satisfied if these curves meet at a point (u_c, \mathcal{U}_c) . Then, the stability condition, $\partial_{\mathcal{U}}^2 p(u_c, \mathcal{U}_c) < 0$, and the sign of the radiation density are determined. The $C(u, \mathcal{U}) = 0$ constraint requires to have at least one negative $r_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5}$.

- Our analysis shows the existence of non-trivial thermal cosmological solutions in heterotic models with $\bar{Q}_4 \equiv \bar{Q}_5$, and type II *Case 3* models with $Q_4 + \bar{Q}_4 \equiv Q_5 + \bar{Q}_5$, where the pressure takes the general form

$$p(u, \mathcal{U}) = n_{100} [p_{100} + r(p_{010} + p_{001}) + p_{111}], \quad (3.30)$$

with $-1 \leq r \leq 1$. These models lie along the edge $\xi_2 = \xi_3 = 0$ of the tetrahedron. The duality symmetry $\mathcal{U} \rightarrow \mathcal{U}^{-1}$ implies that along the axis $\mathcal{U} \equiv 1$, $D(u, \mathcal{U}) = 0$. As r varies, we

find 4 distinct patterns, specified by

$$r_{c3} = -\frac{f_{5/2}^{ee}(1) + f_{5/2}^{oo}(1)}{2f_{5/2}^{oe}(1)} \simeq -0.215, \quad r_{c2} = -\frac{S_5^e}{S_5^o} = -\frac{1}{31} \quad \text{and} \quad r_{c1} = 0, \quad (3.31)$$

where the functions in the definition of r_{c3} can be found in Eq. (A.12).

- It turns out that there are no cosmological solutions with constant u and \mathcal{U} when $r < r_{c3}$ or $r > 0$, since then $C(u, \mathcal{U}) \neq 0$ everywhere.

- A stable cosmological solution exists when $r_{c3} < r < r_{c2}$, with $c_r > 0$, (see Fig. 3a). It corresponds to a global minimum of the thermal effective potential.




Figure 3: For a pressure of the form (3.30), the curves defined by $C(u, \mathcal{U}) = 0$ (straight lines) and $D(u, \mathcal{U}) = 0$ (dashed lines) are represented. When $r_{c3} < r < r_{c2}$ (Fig. a), there is a stable cosmological solution ($u_c, \mathcal{U}_c = 1$). It corresponds to a global minimum of the thermal effective potential. When $r_{c2} \leq r < 0$ (Fig. b), there is a stable cosmological solution ($u_c, \mathcal{U}_c = 1$) that corresponds to a local minimum. Two runaway behaviors that bring the system to five dimensions are also allowed.

- When $r_{c2} \leq r < 0$, a stable cosmological solution with $\mathcal{U}_c = 1$ still exists, with $c_r > 0$, but this corresponds to a local minimum of the thermal effective potential. Actually, new branches of the locus $D = 0$ are present and converge exponentially towards the curve $C = 0$, (see Fig. 3b). Formally, their common asymptotes define flat directions. However, since these “solutions” imply R_5 (or R_4) to be very large, they are out of the scope of our analysis. They are better understood in terms of runaway behaviors that decompactify the system to five dimensions, where supersymmetry is broken by the remaining finite size internal radius R_4 (or R_5) and thermal effects.²

² A similar runaway behavior can be realized in type II *Case 2* models, with generic operator \bar{Q}_5 . They involve a pressure $p(u, \mathcal{U}) = n_{100}[p_{100} + rp_{001}]$, where $-1 \leq r \leq 1$. Their representative points in the tetrahedron satisfy $\xi_1 = \xi_3 = \frac{1}{2} - \xi_2$, where $0 \leq \xi_2 \leq \frac{1}{2}$. When r varies, there is a phase where the curves $D = 0$ and $C = 0$ are non-trivial and asymptotic to one another. The situation is similar to what is observed in the lower half plane of Fig. 3b.

For the heterotic model with the choice $\bar{Q}_4 = \bar{Q}_5 = \gamma + \gamma'$, one has

$$\begin{aligned} n_{100} &= n_{111} = 2^3 \times 504, \\ n_{010} &= n_{001} = 2^3 [[2]_{X_{2,3}} + [6]_{T^6} + [120 - 128]_{E_8} + [120 - 128]_{E'_8}] = -2^3 \times 8, \end{aligned} \quad (3.32)$$

so that $r_{c2} < r = -1/63 < r_{c1}$. In this specific case, the cosmological solution corresponds to $(u_c, \mathcal{U}_c) \simeq (1.649, 1)$, where $c_r \simeq 0.0708 \times 6(a_0 T_0)^4$.

Another model considered in [5] is based on the heterotic T^4/\mathbb{Z}_2 orbifold, with non-Abelian gauge group $E_8 \times E_7 \times SU(2)$. In that case, one has $r_{c2} < r = -1/127 < r_{c1}$ and the corresponding cosmological solution fixes $(u_c, \mathcal{U}_c) \simeq (1.996, 1)$, with $c_r \simeq 0.0762 \times 6(a_0 T_0)^4$.

• We can treat in a similar way type II *Case 1*' models with arbitrary $Q_4 + Q_5$. The pressure is of the form

$$p(u, \mathcal{U}) = n_{100}[p_{100} + r p_{111}], \quad (3.33)$$

where $-1 \leq r \leq 1$. This class of models belongs to a segment in the interior of the tetrahedron, $\xi_2 = \xi_3 = \frac{1}{2} - \xi_1$, $0 \leq \xi_1 \leq \frac{1}{2}$. The symmetry $\mathcal{U} \rightarrow \mathcal{U}^{-1}$ implies $D(u, \mathcal{U}) = 0$ along the axis $\mathcal{U} \equiv 1$. Also, $p(u, \mathcal{U})$ is constant in sectors *III* and *IV* (and their common edge (4)), implying that $D(u, \mathcal{U})$ is vanishing. For $r < 0$, there is no other solutions to $D = 0$. When r varies, the set of solutions to $C(u, \mathcal{U}) = 0$ is divided into 3 classes characterized by

$$r_{c4} \simeq -0.77, \quad r'_{c3} = -\frac{f_{5/2}^{ee}(1)}{f_{5/2}^{oo}(1)} \simeq -0.215. \quad (3.34)$$

- For $r < r_{c4}$, the right boundary of the locus $C \leq 0$ is asymptotic to the edges (3) and (5), where D is not vanishing yet, (see Fig. 4a). The only solution to $D = C = 0$ arises at $\mathcal{U} = 1$, but the corresponding cosmological evolution is unstable to small fluctuations of \mathcal{U} .

- For $r_{c4} < r < r'_{c3}$, a cosmological solution with constant u and \mathcal{U} exists, (see Fig. 4b), but is again unstable to small fluctuations of \mathcal{U} .

- For $r'_{c3} < r$, one has $C(u, \mathcal{U}) > 0$ everywhere: There is no cosmological solution with constant complex structures.

• The last class of systems we analyze corresponds to type II *Case 1* models with arbitrary $Q_4 + \bar{Q}_5$. The pressure is

$$p(u, \mathcal{U}) = n_{100}[p_{100} + r p_{011}], \quad (3.35)$$

where $-1 \leq r \leq 1$. In contrast to the previous cases, these models are not represented by points in the tetrahedron. For $r < 0$, the only solutions to $D(u, \mathcal{U}) = 0$ are along the axis




Figure 4: For a pressure of the form (3.33), the curves defined by $C(u, \mathcal{U}) = 0$ (straight lines) are represented. The locus $D = 0$ is composed of the axis $\mathcal{U} = 1$, and the sectors III and IV, (see Fig. 2). When $r < r_{c4}$ (Fig. a) or $r_{c4} \leq r < r'_{c3}$ (Fig. b), there is an unstable cosmological solution $(u_c, \mathcal{U}_c = 1)$.

$\mathcal{U} = 1$. The set of solutions to $C(u, \mathcal{U}) = 0$ is characterized by the point r'_{c3} of Eq. (3.34) and $r_{c1} = 0$.

- For $r < r'_{c3}$, the axis $\mathcal{U} = 1$ is entirely inside the region $C(u, \mathcal{U}) < 0$ that has two distinct boundaries, (see Fig. 5a), and so there is no cosmological solution with constant u and \mathcal{U} .

- For $r'_{c3} \leq r < 0$, the region $C(u, \mathcal{U}) < 0$ has now a connected boundary, (see Fig. 5b). The latter crosses the axis $\mathcal{U} = 1$, so that a cosmological solution $(u_c, \mathcal{U}_c = 1)$ exists. It is however unstable to small fluctuations of \mathcal{U} .

- For $0 < r$, one has $C(u, \mathcal{U}) > 0$ everywhere: There is no cosmological solution with constant u and \mathcal{U} .




Figure 5: For a pressure of the form (3.35), the curves defined by $C(u, \mathcal{U}) = 0$ (straight lines) and $D(u, \mathcal{U}) = 0$ (dashed lines) are represented. When $r < r'_{c3}$ (Fig. a), there is no cosmological solution with constant complex structures. When $r'_{c3} \leq r < 0$ (Fig. b), there is an unstable cosmological solution $(u_c, \mathcal{U}_c = 1)$.

3.3 Non-thermal vs thermal cosmologies

The cosmological evolutions found in the non-thermal and thermal cases, Eqs (2.47) and (3.21), have drastically different properties we would like to comment on.

First, the contributions of the kinetic energy of the field ϕ_- to the “effective” Friedmann-Hubble equation are of *opposite sign*. This phenomenon is due to the different compatibility relations $\dot{\phi} = -4H$ for the non-thermal cosmology and $\dot{\phi} = -H$ for the thermal one, once they are inserted in the respective Friedmann-Hubble Eqs (2.37) and (3.7). This is also the reason why these equations have $1/a^{16}$ and $1/a^4$ monomial contributions, respectively. The intermediate cosmological region we are considering is described by the thermal cosmologies. At late times however, the temperature, being proportional to the inverse of the scale factor, tends to zero, and a paradox seems to arise as the cosmological evolution is never described by the non-thermal solutions.

To better understand this point, we rewrite the pressure and energy density as follows, (see the relations (3.2) and (3.9)):

$$P = -M^4 v_{\text{th}}, \quad \rho = M^4 (v_{\text{th}} + u \partial_u v_{\text{th}}) \quad \text{where} \quad v_{\text{th}}(u, \mathcal{U}) = -\frac{p(u, \mathcal{U})}{u^4}. \quad (3.36)$$

When the pressure is of the general form (A.1), one can use the expansion valid for $u \gg 1$ (\mathcal{U} fixed), Eqs (A.11) and (A.15), to decompose v_{th} into two pieces:

$$v_{\text{th}}(u, \mathcal{U}) = -\{n_{010} k_3^{oe}(\mathcal{U}^{-1}) + n_{001} k_3^{oe}(\mathcal{U}) + n_{011} k_3^{oo}(\mathcal{U})\} + \hat{v}_{\text{th}}(u, \mathcal{U}), \quad (3.37)$$

where

$$\hat{v}_{\text{th}}(u, \mathcal{U}) = -\frac{1}{u^4} \times \{(n_{100} + n_{111})S_4^o + (n_{010} + n_{001} + n_{011})S_4^e\} + \dots, \quad (3.38)$$

up to exponentially suppressed terms as $u \rightarrow +\infty$. Clearly, $\hat{v}_{\text{th}}(u, \mathcal{U}) \rightarrow 0$ in this limit. Since $n_{\tilde{g}_0 \tilde{g}_4 \tilde{g}_5} = n_{\tilde{g}_4 \tilde{g}_5}$, the u -independent terms of the r.h.s. of Eq. (3.37) are equal to the zero temperature effective potential $v(\mathcal{U})$ found for the heterotic or type II cases, Eqs (2.27) and (2.30). Thus, P and ρ are only converging to their $T = 0$ counterparts when $u \rightarrow +\infty$. However, the thermal cosmologies we have considered have stabilized $u \equiv u_c$, implying that the finite temperature corrections $\hat{v}_{\text{th}}(u, \mathcal{U})$, even if the temperature is small, are never negligible. In fact, the condition $u = M(t)/T(t) \rightarrow +\infty$ for the thermal system to be correctly approximated by the non-thermal one implies that the thermal corrections should be screened by radiative corrections, and not that they would necessarily vanish. Explicit

cosmological evolutions with $u \rightarrow +\infty$ are analyzed in [22]. They describe Big Crunch cosmologies, where the temperature is proportional to $1/a$. So, T is large in absolute value (but still negligible compared to M).

Finally, we note that in the present work, each time a stable cosmological solution at finite temperature is found, its counterpart at $T = 0$ is unstable, and vice versa. As explained in [22], this is a consequence of the fact that for arbitrary initial conditions, the cosmological evolutions are always converging to an attractor. Depending on the model, the latter can be the radiation dominated universes studied in [5] and the present paper, or Big Crunch cosmologies where the thermal effects are screened by radiative corrections.

4 Conclusions

The main result of this work is to show the existence of the critical stringy cosmologies of [5], even in the cases where more than one modulus participate in the supersymmetry breaking mechanism. They correspond to a radiation dominated era with constant complex structures. A thorough analysis was done for several string models with $N_4 \geq 2$ initial supersymmetry. Depending on the pattern of supersymmetry breaking, the critical values of the thermal effective potential for the complex structure moduli are either minima or maxima. Run away behaviors that bring the system to higher dimensions can also occur. When the radiation era is stable, we explicitly show that even if the temperature tends to zero at late times, thermal corrections to the dynamics are never negligible.

Our approach was to separate the cosmological evolution in distinct regions, according to the value of the temperature. The stringy non-geometrical region, where the temperature is of order the Hagedorn temperature, is much harder to understand. Stringy phenomena occur where conventional field theoretic notions concerning the geometry and topology are breaking down. Some interesting proposals to understand this early time region have been put forward in [11, 19]. The intermediate region, just after the Hagedorn era is shown to be under control. The free energy is free of any infrared and ultraviolet ambiguities, allowing us to follow the backreaction on the geometry and determine the time evolution of several moduli fields. We show that the only relevant moduli are the ones participating in the supersymmetry breaking mechanism. The others are either frozen by receiving a soft-

breaking mass, or they remain flat directions with exponentially suppressed contribution to the free energy.

A very interesting result is that the critical cosmological solutions are governed by a higher dimensional thermal state equation, $\rho = (d - 1) P$, or in the four dimensional effective description, this equation becomes $\rho = (3 + n) P$ where n is the number of moduli participating in the breaking of supersymmetry. In [5, 28], we analyzed the case $n = 1$. Here, we generalize the result to the $n = 2$ case. The overall scale evolves in time such that the state equation is valid, while complex structure -like ratios of scales are frozen.

Although our work covers adequately and unambiguously the intermediate cosmological region, it is incomplete to describe the very early Hagedorn era, where non-geometric stringy phenomena are important. It is incomplete as well to describe relatively late time phenomena, like the radiative breaking of the electroweak gauge symmetry and QCD confinement, where non-perturbative transmutation scales, like for instance Λ_{QCD} , are relevant. The extension to the late time era requires to consider at least models with $N_4 = 1$ initial supersymmetry. Progress concerning the very early era can be made provided that we understand better the resolution of the Hagedorn instabilities and the stringy non-geometrical structure of the early universe.

Acknowledgements

We are grateful to C. Angelantonj, C. Bachas, J. Estes, J. Troost and especially F. Bourliot for useful discussions. H.P. and N.T. thank the Ecole Normale Supérieure and C.K. and H.P. the University of Cyprus for hospitality.

The work of C.K. and H.P. is partially supported by the EU contract MRTN-CT-2004-005104 and the ANR (CNRS-USAR) contract 05-BLAN-0079-01 (01/12/05). N.T. and C.K. are supported by the EU contract MRTN-CT-2004-512194. H.P. is also supported by the EU contracts MRTN-CT-2004-503369 and MEXT-CT-2003-509661, INTAS grant 03-51-6346, and CNRS PICS 2530, 3059 and 3747, while N.T. is also supported by an INTERREG IIIA Crete/Cyprus program.

Appendix A

We list in this Appendix the large and small complex structure expansions of an arbitrary linear combination of functions $p_{\tilde{g}_0\tilde{g}_4\tilde{g}_5}$ defined in Eq. (2.15),

$$p(u, \mathcal{U}) = n_{100}(p_{100} + r_{010}p_{010} + r_{001}p_{001} + r_{111}p_{111} + r_{011}p_{011}). \quad (\text{A.1})$$

For fixed ω and λ , we determine the power expansion along the lines $\ln \mathcal{U} = \omega \ln u + \ln \lambda$ when $\ln u$ and/or $\ln \mathcal{U}$ are large, by making an extensive use of the approximations

$$\begin{aligned} \sum_m \frac{1}{((2m+1)^2x+a)^3} &= \frac{3\pi}{16} \frac{1}{a^{5/2}} \frac{1}{\sqrt{x}} + \dots, \\ \sum_m \frac{1}{((2m)^2x+a)^3} &= \frac{3\pi}{16} \frac{1}{a^{5/2}} \frac{1}{\sqrt{x}} + \dots, \end{aligned} \quad (\text{A.2})$$

where $a > 0$, $x \rightarrow 0_+$ and the dots stand for $\mathcal{O}\left(e^{-\pi\sqrt{a/x}}/x^{3/2}\right)$ terms. Similarly, we use

$$\begin{aligned} \sum_m \frac{1}{((2m+1)^2x+a)^{5/2}} &= \frac{2}{3} \frac{1}{a^2} \frac{1}{\sqrt{x}} + \dots, \\ \sum_m \frac{1}{((2m)^2x+a)^{5/2}} &= \frac{2}{3} \frac{1}{a^2} \frac{1}{\sqrt{x}} + \dots, \end{aligned} \quad (\text{A.3})$$

where again we neglect exponentially suppressed terms. The $(\ln u, \ln \mathcal{U})$ -plane can then be divided in 6 sectors, I, II, \dots, VI , where the expansions are independent of ω (and λ). The boundaries of these sectors are the lines whose slopes are $\omega = -2, 0$ or 2 , (see Fig. 2). In each sector, we find,

$$\begin{aligned} \frac{p^I}{n_{100}} &= u^{-2} S_6^o \\ &\quad + u^3 \mathcal{U}^{-5/2} (S_5^e + r_{001} S_5^o) \\ &\quad + u^3 \mathcal{U}^{3/2} ((1 + r_{010} + r_{001}) S_4^e + (r_{111} + r_{011}) S_4^o), \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned} \frac{p^{II}}{n_{100}} &= u^4 \mathcal{U}^{-3} r_{001} S_6^o \\ &\quad + u^{-1} \mathcal{U}^{-1/2} (S_5^o + r_{001} S_5^e) \\ &\quad + u^3 \mathcal{U}^{3/2} ((1 + r_{001}) S_4^e + (r_{010} + r_{111} + r_{011}) S_4^o), \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned} \frac{p^{III}}{n_{100}} &= u^4 \mathcal{U}^{-3} r_{001} S_6^o \\ &\quad + u^4 \mathcal{U}^2 ((r_{010} + r_{011}) S_5^o + r_{001} S_5^e) \\ &\quad + (1 + r_{111}) S_4^o + (r_{010} + r_{001} + r_{011}) S_4^e, \end{aligned} \quad (\text{A.6})$$

$$\begin{aligned} \frac{p^{IV}}{n_{100}} &= u^4 \mathcal{U}^3 r_{010} S_6^o \\ &\quad + u^4 \mathcal{U}^{-2} ((r_{001} + r_{011}) S_5^o + r_{010} S_5^e) \\ &\quad + (1 + r_{111}) S_4^o + (r_{010} + r_{001} + r_{011}) S_4^e, \end{aligned} \quad (\text{A.7})$$

$$\begin{aligned} \frac{p^V}{n_{100}} &= u^4 \mathcal{U}^3 r_{010} S_6^o \\ &\quad + u^{-1} \mathcal{U}^{1/2} (S_5^o + r_{010} S_5^e) \\ &\quad + u^3 \mathcal{U}^{-3/2} ((1 + r_{010}) S_4^e + (r_{001} + r_{111} + r_{011}) S_4^o), \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} \frac{p^{VI}}{n_{100}} &= u^{-2} S_6^o \\ &\quad + u^3 \mathcal{U}^{5/2} (S_5^e + r_{010} S_5^o) \\ &\quad + u^3 \mathcal{U}^{-3/2} ((1 + r_{010} + r_{001}) S_4^e + (r_{111} + r_{011}) S_4^o), \end{aligned} \quad (\text{A.9})$$

where we have defined

$$\begin{aligned} S_6^o &= \frac{2}{\pi^3} \sum_m \frac{1}{(2m+1)^6} = \frac{\pi^3}{240}, \\ S_5^o &= \frac{2}{\pi^3} \frac{3\pi}{16} \sum_m \frac{1}{|2m+1|^5} = \frac{93}{128} \frac{\zeta(5)}{\pi^2}, \quad S_5^e = \frac{2}{\pi^3} \frac{3\pi}{16} \sum_{m \neq 0} \frac{1}{|2m|^5} = \frac{3}{128} \frac{\zeta(5)}{\pi^2}, \\ S_4^o &= \frac{2}{\pi^3} \frac{3\pi}{16} \frac{2}{3} \sum_m \frac{1}{(2m+1)^4} = \frac{\pi^2}{192}, \quad S_4^e = \frac{2}{\pi^3} \frac{3\pi}{16} \frac{2}{3} \sum_{m \neq 0} \frac{1}{(2m)^4} = \frac{\pi^2}{2880}. \end{aligned} \quad (\text{A.10})$$

The previous sectors are separated by edges, (1), ..., (6), in the neighborhood of which some terms we neglected in the interior of the adjacent sectors are not exponentially suppressed anymore. Along these edges, one has $\mathcal{U} = \lambda u^{\pm 2, 0}$, where $\lambda \simeq 1$, and

$$\begin{aligned} \frac{p^{(1)}}{n_{100}} &= u^{-2} S_6^o \\ &\quad + u^3 (f_{5/2}^{ee}(\lambda) + r_{010} f_{5/2}^{oe}(\lambda^{-1}) + r_{001} f_{5/2}^{oe}(\lambda) + (r_{111} + r_{011}) f_{5/2}^{oo}(\lambda)), \quad \lambda = \mathcal{U}, \end{aligned}$$

$$\begin{aligned} \frac{p^{(2)}}{n_{100}} &= u^3 \mathcal{U}^{3/2} ((1 + r_{001}) S_4^e + (r_{010} + r_{111} + r_{011}) S_4^o) \\ &\quad + u^{-2} (g_3^{eo}(\lambda) + r_{001} g_3^{oe}(\lambda)), \quad \lambda = \mathcal{U} u^{-2}, \end{aligned}$$

$$\begin{aligned} \frac{p^{(3)}}{n_{100}} &= u^4 \mathcal{U}^{-3} r_{001} S_6^o \\ &\quad + h_{5/2}^{oe}(\lambda) + (r_{010} + r_{011}) h_{5/2}^{eo}(\lambda) + r_{001} h_{5/2}^{ee}(\lambda) + r_{111} h_{5/2}^{oo}(\lambda), \quad \lambda = \mathcal{U} u^2, \end{aligned}$$

$$\begin{aligned} \frac{p^{(4)}}{n_{100}} &= (1 + r_{111}) S_4^o + (r_{010} + r_{001} + r_{011}) S_4^e \\ &\quad + u^4 (r_{010} k_3^{oe}(\lambda^{-1}) + r_{001} k_3^{oe}(\lambda) + r_{011} k_3^{oo}(\lambda)), \quad \lambda = \mathcal{U}, \end{aligned}$$

$$\begin{aligned} \frac{p^{(5)}}{n_{100}} &= u^4 \mathcal{U}^3 r_{001} S_6^o \\ &\quad + h_{5/2}^{oe}(\lambda^{-1}) + r_{010} h_{5/2}^{ee}(\lambda^{-1}) + (r_{001} + r_{011}) h_{5/2}^{eo}(\lambda^{-1}) + r_{111} h_{5/2}^{oo}(\lambda^{-1}), \quad \lambda = \mathcal{U} u^{-2}, \end{aligned}$$

$$\begin{aligned} \frac{p^{(6)}}{n_{100}} &= u^3 \mathcal{U}^{-3/2} \left((1 + r_{010}) S_4^e + (r_{001} + r_{111} + r_{011}) S_4^o \right) \\ &\quad + u^{-2} \left(g_3^{eo}(\lambda^{-1}) + r_{010} g_3^{oe}(\lambda^{-1}) \right), \quad \lambda = \mathcal{U} u^2, \end{aligned} \tag{A.11}$$

where we have introduced the functions

$$\begin{aligned} f_{5/2}^{ee}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \sum_{(m,n) \neq (0,0)} \frac{1}{((2m)^2 \lambda + (2n)^2 \lambda^{-1})^{5/2}}, \\ f_{5/2}^{oe}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n)^2 \lambda^{-1})^{5/2}}, \\ f_{5/2}^{oo}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n+1)^2 \lambda^{-1})^{5/2}}, \end{aligned} \tag{A.12}$$

$$g_3^{eo}(\lambda) = \frac{2}{\pi^3} \sum_{m,n} \frac{1}{((2m)^2 \lambda + (2n+1)^2)^3}, \quad g_3^{oe}(\lambda) = \frac{2}{\pi^3} \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n)^2)^3}, \tag{A.13}$$

$$\begin{aligned} h_{5/2}^{oe}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \lambda^2 \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n)^2)^{5/2}}, \\ h_{5/2}^{eo}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \lambda^2 \sum_{m,n} \frac{1}{((2m)^2 \lambda + (2n+1)^2)^{5/2}}, \\ h_{5/2}^{ee}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \lambda^2 \sum_{(m,n) \neq (0,0)} \frac{1}{((2m)^2 \lambda + (2n)^2)^{5/2}}, \\ h_{5/2}^{oo}(\lambda) &= \frac{2}{\pi^3} \frac{3\pi}{16} \lambda^2 \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n+1)^2)^{5/2}}, \end{aligned} \tag{A.14}$$

$$k_3^{oe}(\lambda) = \frac{2}{\pi^3} \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n)^2 \lambda^{-1})^3}, \quad k_3^{oo}(\lambda) = \frac{2}{\pi^3} \sum_{m,n} \frac{1}{((2m+1)^2 \lambda + (2n+1)^2 \lambda^{-1})^3}. \tag{A.15}$$

References

- [1] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], “Measurements of Ω and Λ from 42 high-redshift supernovae,” *Astrophys. J.* **517**, 565 (1999) [arXiv:astro-ph/9812133].
- [2] C. Kounnas, N. Toumbas and J. Troost, “A wave-function for stringy universes,” *JHEP* **0708** (2007) 018 [arXiv:0704.1996 [hep-th]].
- [3] C. Kounnas and H. Partouche, “Instanton transition in thermal and moduli deformed de Sitter cosmology,” *Nucl. Phys. B* **793** (2008) 131 [arXiv:0705.3206 [hep-th]].

- [4] C. Kounnas and H. Partouche, “Inflationary de Sitter solutions from superstrings,” Nucl. Phys. B **795** (2008) 334 [arXiv:0706.0728 [hep-th]].
- [5] T. Catelin-Jullien, C. Kounnas, H. Partouche and N. Toumbas, “Thermal/quantum effects and induced superstring cosmologies,” Nucl. Phys. B **797** (2008) 137 [arXiv:0710.3895 [hep-th]].
- T. Catelin-Jullien, C. Kounnas, H. Partouche and N. Toumbas, “Thermal and quantum superstring cosmologies,” Fortsch. Phys. **56** (2008) 792 [arXiv:0803.2674 [hep-th]].
- [6] M. Gasperini and G. Veneziano, “Pre-big bang in string cosmology,” Astropart. Phys. **1** (1993) 317 [arXiv:hep-th/9211021].
- M. Gasperini, M. Maggiore and G. Veneziano, “Towards a non-singular pre-big bang cosmology,” Nucl. Phys. B **494**, 315 (1997) [arXiv:hep-th/9611039].
- [7] E. Kiritsis and C. Kounnas, “Dynamical topology change, compactification and waves in a stringy early universe,” arXiv:hep-th/9407005.
- E. Kiritsis and C. Kounnas, “Dynamical topology change in string theory,” Phys. Lett. B **331** (1994) 51 [arXiv:hep-th/9404092].
- E. Kiritsis and C. Kounnas, “Dynamical topology change, compactification and waves in string cosmology,” Nucl. Phys. Proc. Suppl. **41** (1995) 311 [arXiv:gr-qc/9701005].
- E. Kiritsis and C. Kounnas, “String gravity and cosmology: Some new ideas,” arXiv:gr-qc/9509017.
- [8] J. Khoury, B. A. Ovrut, N. Seiberg, P. J. Steinhardt and N. Turok, “From big crunch to big bang,” Phys. Rev. D **65**, 086007 (2002) [arXiv:hep-th/0108187].
- [9] H. Liu, G. W. Moore and N. Seiberg, “The challenging cosmic singularity,” arXiv:gr-qc/0301001.
- [10] B. Craps, S. Sethi and E. P. Verlinde, “A matrix big bang,” JHEP **0510**, 005 (2005) [arXiv:hep-th/0506180].
- [11] C. Kounnas, “Massive boson-fermion degeneracy and the early structure of the universe,” arXiv:0808.1340 [hep-th].

- [12] C. Angelantonj, M. Cardella and N. Irges, “An alternative for moduli stabilisation,” *Phys. Lett. B* **641**, 474 (2006) [arXiv:hep-th/0608022].
- [13] J. Atick and E. Witten, “The Hagedorn transition and the number of degrees of freedom of string theory,” *Nucl. Phys. B* **310**, 291 (1988).
- [14] C. Kounnas and B. Rostand, “Coordinate dependent compactifications and discrete symmetries,” *Nucl. Phys. B* **341** (1990) 641.
- [15] I. Antoniadis and C. Kounnas, “Superstring phase transition at high temperature,” *Phys. Lett. B* **261** (1991) 369.
- I. Antoniadis, J. P. Derendinger and C. Kounnas, “Non-perturbative supersymmetry breaking and finite temperature instabilities in $N = 4$ superstrings,” arXiv:hep-th/9908137.
- C. Kounnas, “Universal thermal instabilities and the high-temperature phase of the $N = 4$ superstrings,” arXiv:hep-th/9902072.
- [16] R. H. Brandenberger and C. Vafa, “Superstrings in the Early Universe,” *Nucl. Phys. B* **316**, 391 (1989).
- [17] R. Easther, B. R. Greene, M. G. Jackson and D. N. Kabat, “String windings in the early universe,” *JCAP* **0502**, 009 (2005) [arXiv:hep-th/0409121].
- B. Greene, D. Kabat and S. Marnerides, “Bouncing and cyclic string gas cosmologies,” arXiv:0809.1704 [hep-th].
- [18] E. Cremmer, S. Ferrara, C. Kounnas and D. V. Nanopoulos, “Naturally vanishing cosmological constant in $N = 1$ supergravity,” *Phys. Lett. B* **133** (1983) 61.
- J. R. Ellis, C. Kounnas and D. V. Nanopoulos, “No scale supersymmetric GUTs,” *Nucl. Phys. B* **247** (1984) 373.
- J. R. Ellis, C. Kounnas and D. V. Nanopoulos, “Phenomenological $SU(1, 1)$ supergravity,” *Nucl. Phys. B* **241** (1984) 406.
- J. R. Ellis, A. B. Lahanas, D. V. Nanopoulos and K. Tamvakis, “No-scale supersymmetric standard model,” *Phys. Lett. B* **134**, 429 (1984).

- [19] C. Angelantonj, C. Kounnas, H. Partouche and N. Toumbas, “Resolution of Hagedorn singularity in superstrings with gravito-magnetic fluxes,” Nucl. Phys. B **809** (2009) 291 [arXiv:0808.1357 [hep-th]].
- [20] J. P. Derendinger, C. Kounnas, P. M. Petropoulos and F. Zwirner, “Superpotentials in IIA compactifications with general fluxes,” Nucl. Phys. B **715** (2005) 211 [arXiv:hep-th/0411276].
- J. P. Derendinger, C. Kounnas, P. M. Petropoulos and F. Zwirner, “Fluxes and gaugings: $N = 1$ effective superpotentials,” Fortsch. Phys. **53** (2005) 926 [arXiv:hep-th/0503229].
- G. Villadoro and F. Zwirner, “ $N = 1$ effective potential from dual type-IIA D6/O6 orientifolds with general fluxes,” JHEP **0506**, 047 (2005) [arXiv:hep-th/0503169].
- G. Villadoro and F. Zwirner, “D terms from D-branes, gauge invariance and moduli stabilization in flux compactifications,” JHEP **0603**, 087 (2006) [arXiv:hep-th/0602120].
- L. Andrianopoli, M. A. Lledo and M. Trigiante, “The Scherk-Schwarz mechanism as a flux compactification with internal torsion,” JHEP **0505** (2005) 051 [arXiv:hep-th/0502083].
- G. Dall’Agata and N. Prezas, “Scherk-Schwarz reduction of M-theory on G_2 -manifolds with fluxes,” JHEP **0510** (2005) 103 [arXiv:hep-th/0509052].
- J. P. Derendinger, P. M. Petropoulos and N. Prezas, “Axionic symmetry gaugings in $N = 4$ supergravities and their higher-dimensional origin,” Nucl. Phys. B **785**, 115 (2007) [arXiv:0705.0008 [hep-th]].
- [21] J. Scherk and J. H. Schwarz, “Spontaneous breaking of supersymmetry through dimensional reduction,” Phys. Lett. B **82** (1979) 60.
- R. Rohm, “Spontaneous supersymmetry breaking in supersymmetric string theories,” Nucl. Phys. B **237** (1984) 553.
- C. Kounnas and M. Porrati, “Spontaneous supersymmetry breaking in string theory,” Nucl. Phys. B **310** (1988) 355.
- S. Ferrara, C. Kounnas and M. Porrati, “ $N = 1$ superstrings with spontaneously broken symmetries,” Phys. Lett. B **206** (1988) 25.

- S. Ferrara, C. Kounnas, M. Porrati and F. Zwirner, “Effective superhiggs and Str M^2 from four-dimensional strings,” Phys. Lett. B **194** (1987) 366.
- S. Ferrara, C. Kounnas and M. Porrati, “Superstring solutions with spontaneously broken four-dimensional supersymmetry,” Nucl. Phys. B **304** (1988) 500.
- [22] F. Bourliot, C. Kounnas and H. Partouche, “Attraction to a radiation era in early superstring cosmology”, CPHT-RR 089.1208, LPTENS - 08/64 preprint.
- [23] L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, “Strings on orbifolds,” Nucl. Phys. B **261**, 678 (1985).
- L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, “Strings on orbifolds. 2,” Nucl. Phys. B **274**, 285 (1986).
- K. S. Narain, M. H. Sarmadi and C. Vafa, “Asymmetric orbifolds,” Nucl. Phys. B **288**, 551 (1987).
- K. S. Narain, M. H. Sarmadi and C. Vafa, “Asymmetric orbifolds: Path integral and operator formulations,” Nucl. Phys. B **356**, 163 (1991).
- [24] I. Antoniadis, C. P. Bachas and C. Kounnas, “Four-dimensional superstrings,” Nucl. Phys. B **289** (1987) 87.
- H. Kawai, D. C. Lewellen and S. H. H. Tye, “Construction of fermionic string models in four-dimensions,” Nucl. Phys. B **288**, 1 (1987).
- [25] C. Kounnas, “BPS states in superstrings with spontaneously broken SUSY,” Nucl. Phys. Proc. Suppl. **58** (1997) 57 [arXiv:hep-th/9703198].
- E. Kiritsis and C. Kounnas, “Perturbative and non-perturbative partial supersymmetry breaking: $N = 4 \rightarrow N = 2 \rightarrow N = 1$,” Nucl. Phys. B **503** (1997) 117 [arXiv:hep-th/9703059].
- E. Kiritsis, C. Kounnas, P. M. Petropoulos and J. Rizos, “String threshold corrections in models with spontaneously broken supersymmetry,” Nucl. Phys. B **540** (1999) 87 [arXiv:hep-th/9807067].
- [26] I. Antoniadis, C. Bachas, C. Fabre, H. Partouche and T. R. Taylor, “Aspects of type I - type II - heterotic triality in four dimensions,” Nucl. Phys. B **489** (1997) 160 [arXiv:hep-th/9608012].

- I. Antoniadis, H. Partouche and T. R. Taylor, “Duality of $N = 2$ heterotic - type I compactifications in four dimensions,” Nucl. Phys. B **499** (1997) 29 [arXiv:hep-th/9703076].
- I. Antoniadis, H. Partouche and T. R. Taylor, “Lectures on heterotic - type I duality,” Nucl. Phys. Proc. Suppl. **61A** (1998) 58 [Nucl. Phys. Proc. Suppl. **67** (1998) 3] [arXiv:hep-th/9706211].
- [27] A. Gregori, C. Kounnas and J. Rizos, “Classification of the $N = 2$, $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetric type II orbifolds and their type II asymmetric duals,” Nucl. Phys. B **549**, 16 (1999) [arXiv:hep-th/9901123].
- [28] I. Antoniadis and C. Kounnas, “The dilaton classical solution and the supersymmetry breaking evolution in an expanding universe,” Nucl. Phys. B **284**, 729 (1987).