

RATIONAL POINTS IN ARITHMETIC PROGRESSION
ON $y^2 = x^n + k$

MACIEJ ULAS

ABSTRACT. Let C be a hyperelliptic curve given by the equation $y^2 = f(x)$, where $f \in \mathbb{Z}[x]$ and f hasn't multiple roots. We say that points $P_i = (x_i, y_i) \in C(\mathbb{Q})$ for $i = 1, 2, \dots, n$ are in arithmetic progression if the numbers x_i for $i = 1, 2, \dots, n$ are in arithmetic progression.

In this paper we show that there exists a polynomial $k \in \mathbb{Z}[t]$ with such a property that on the elliptic curve $\mathcal{E} : y^2 = x^3 + k(t)$ (defined over the field $\mathbb{Q}(t)$) we can find four points in arithmetic progression which are independent in the group of all $\mathbb{Q}(t)$ -rational points on the curve \mathcal{E} . In particular this result generalizes some earlier results of Lee and Vélez from [5]. We also show that if $n \in \mathbb{N}$ is odd then there are infinitely many k 's with such a property that on the curves $y^2 = x^n + k$ there are four rational points in arithmetic progressions. In the case when n is even we can find infinitely many k 's such that on the curves $y^2 = x^n + k$ there are six rational points in arithmetic progression.

1. INTRODUCTION

Many problems in number theory are equivalent to the problem of solution of certain equation or system of equations in integers or in rational numbers. Problems of this type are called *diophantine problems*. In the case when we are able to show that our problem has infinitely many solutions a natural question arises whether it is possible to show the existence of rational parametric solutions i.e. solutions in polynomials or in rational functions. In general this kind of problems are difficult and we don't have any general theory which can give even partially answer to such kind questions. For example, N. Elkies showed in [3] that the set of rational points on the surface $x^4 + y^4 + z^4 = t^4$ is dense in the set of all real points on this surface. However we still haven't known if this equation has rational parametric solutions.

In this paper we meet with the problem of similar type. Our question is related to the construction of integers k with such a property that on the elliptic curve $E_k : y^2 = x^3 + k$ there are four rational points in arithmetic progression. Let us recall that if the curve $C : f(x, y) = 0$ is defined over \mathbb{Q} and we have the rational points $P_i = (x_i, y_i)$ on C then we say that P_i 's are in arithmetic progression if the numbers x_i for $i = 1, 2, \dots, n$ are in arithmetic progression.

In connection with this problem Lee and Vélez in [5] showed that each rational point on the elliptic curve

$$E : y^2 = x^3 - 39x - 173$$

gives an integer k with such a property that on the elliptic curve $E_k : y^2 = x^3 + k$ there are four rational points in arithmetic progression. Due to the fact that the

2000 *Mathematics Subject Classification.* 11G05.

Key words and phrases. arithmetic progressions, elliptic curves, rational points.

set $E(\mathbb{Q})$ of all rational points on E is generated by the point $(11, 27)$ of infinite order we get infinitely many k 's which satisfy demanded conditions.

Now, it should be noted that S. P. Mohanty stated the following

Conjecture 1.1 (S. P. Mohanty [6]). *Let $k \in \mathbb{Z}$ and suppose that the rational points $P_i = (x_i, y_i)$ for $i = 1, \dots, n$ are in arithmetic progression on the elliptic curve $y^2 = x^3 + k$. Then $n \leq 4$.*

We think that the above conjecture is not true. It is clear that in order to find counterexample we should have plenty of k 's for which on the curve $E_k : y^2 = x^3 + k$ we have four points in arithmetic progression. Integer number which satisfy this condition will be called *number of AP4 type*.

Main aim of this paper is to construct parametric families of numbers of AP4 type. We should note that the method employed in [5] cannot be used in order to find families of these kinds.

In Section 2 we show that each rational point on the surface

$$\mathcal{S}_1 : (p^2 - 3q^2 + 3r^2 - s^2)(11p^2 - 18q^2 + 9r^2 - 2s^2) = 3(2p^2 - 5q^2 + 4r^2 - s^2)^2$$

give us an integer k of AP4 type. In particular we prove that on the \mathcal{S}_1 there are infinitely many rational curves. Using this result we show that there exists a polynomial $k \in \mathbb{Z}[t]$ with such a property that on the corresponding elliptic curve $\mathcal{E} : y^2 = x^3 + k(t)$ defined over $\mathbb{Q}(t)$ there are four $\mathbb{Q}(t)$ -rational points and these points are independent in the set $\mathcal{E}(\mathbb{Q}(t))$. Using this result we deduce that the set of rational points on the surface \mathcal{S}_1 is dense in the set of all real points on \mathcal{S}_1 in the Euclidean topology.

In Section 3 we consider natural generalizations of the problem of construction of rational points in arithmetic progressions on hyperelliptic curves of the form $y^2 = x^n + k$.

In Section 4 we give a special sextic hypersurface which is connected with the problem of construction of integer k with such a property that on the curve $y^3 = x^3 + k$ there are five points in arithmetic progression.

2. RATIONAL POINTS ON \mathcal{S}_1

In this section we are interested in the construction of integer numbers of AP4 type. Let $f(x) = ax^3 + bx^2 + cx + d \in \mathbb{Q}[a, b, c, d][x]$ and consider the curve $C : y^2 = f(x)$. Using now change of coordinates

$$(x, y) = \left(\frac{X - 3b}{9a}, \frac{Y}{27a} \right) \text{ with inverse } (X, Y) = (9ax + 3b, 27ay),$$

we can see that the curve C birationally equivalent to the curve

$$E : Y^2 = X^3 + 27(3ac - b^2)X + 27(27a^2d - 9abc + 2b^3).$$

Let p, q, r, s be rational parameters and let us put

$$(1) \quad \begin{aligned} a &= -(p^2 - 3q^2 + 3r^2 - s^2)/6, & b &= (2p^2 - 5q^2 + 4r^2 - s^2)/2, \\ c &= -(11p^2 - 18q^2 + 9r^2 - 2s^2)/6, & d &= p^2. \end{aligned}$$

For a, b, c, d defined in this way we have

$$f(0) = p^2, \quad f(1) = q^2, \quad f(2) = r^2, \quad f(3) = s^2.$$

We can see that in order to prove that the set of integers of AP4 type contains image of certain polynomial we must consider the surface $3ac = b^2$, where a, b, c, d are define by the (1).

Indeed, the points $(0, p), (1, q), (2, r), (3, s)$ are in arithmetic progression on the curve C , and due to the fact that our mapping from C to E given by (2) is affine we deduce that the images of these points will be in arithmetic progression on the curve E . So we consider the surface in \mathbb{P}^3 given by the equation

$$(2) \mathcal{S}_1 : (p^2 - 3q^2 + 3r^2 - s^2)(11p^2 - 18q^2 + 9r^2 - 2s^2) = 3(2p^2 - 5q^2 + 4r^2 - s^2)^2.$$

It is easy to see that the surface \mathcal{S}_1 is singular and that the set $(\pm 1, \pm 1, \pm 1, \pm 1)$ is the set of all singular points (the sign $+$ and $-$ are independent of each other). The surface \mathcal{S}_1 has sixteen singular points which is maximal number of singular points for surfaces of degree four. Surfaces with this property are known under the name of *Kummer surfaces*.

We will show the following

Theorem 2.1. *The set of rational curve on the surface \mathcal{S}_1 is infinite. In particular the set $\mathcal{S}_1(\mathbb{Q})$ is dense in the set of all real points $\mathcal{S}_1(\mathbb{R})$ in the Euclidean topology.*

Proof. In order to show that on the surface \mathcal{S}_1 there are infinitely many rational curves defined over \mathbb{Q} let us consider the following system of equations

$$\begin{cases} t(p^2 - 3q^2 + 3r^2 - s^2) = (2p^2 - 5q^2 + 4r^2 - s^2), \\ (11p^2 - 18q^2 + 9r^2 - 2s^2) = 3t(2p^2 - 5q^2 + 4r^2 - s^2), \end{cases}$$

or equivalently

$$(3) \quad \begin{cases} (3t^2 + 3t + 1)r^2 = -(3t^2 + 9t + 7)p^2 + 2(3t^2 + 6t + 4)q^2, \\ (3t^2 + 3t + 1)s^2 = -2(3t^2 + 12t + 13)p^2 + 9(t^2 + 3t + 3)q^2, \end{cases}$$

where t is an indeterminate parameter. It is easy to see that each rational solution of the system (3) leads us to the rational point on the surface \mathcal{S}_1 . From geometric point of view the system (3) as intersection of two quadratic surfaces with rational points $(p, q, r, s) = (1, 1, 1, 1)$ is birationally equivalent with an elliptic curve defined over the field $\mathbb{Q}(t)$. Now, we show the construction of appropriate mapping.

Using standard substitution $(p, q, r) = (u + r, v + r, r)$ we can parametrize all rational points on the first equation of the system (3) in the following way

$$(4) \quad \begin{cases} p = (3t^2 + 9t + 7)u^2 - 4(3t^2 + 6t + 4)uv + 2(3t^2 + 6t + 4)v^2, \\ q = (3t^2 + 9t + 7)u^2 - 2(3t^2 + 9t + 7)uv + 2(3t^2 + 6t + 4)v^2, \\ r = (3t^2 + 9t + 7)u^2 - 2(3t^2 + 9t + 4)v^2. \end{cases}$$

Without lose of generality we can assume that $u = 1$ and let us substitute the parametrization we have obtained into the second equation in the system (3). We get the curve defined over the field $\mathbb{Q}(t)$ with the equation

$$\begin{aligned} \mathcal{C}_1 : s^2 &= 4(3t^2 + 6t + 4)^2v^4 + 8(3t^2 + 6t + 4)(3t^2 + 15t + 19)v^3 + \\ &\quad - 4(36t^4 + 243t^3 + 618t^2 + 702t + 313)v^2 + \\ &\quad + 4(3t^2 + 9t + 7)(3t^2 + 15t + 19)v + (3t^2 + 9t + 7)^2 \end{aligned}$$

with $\mathbb{Q}(t)$ -rational point $Q = (0, 3t^2 + 9t + 7)$. Let us define the following quantities

$$\begin{aligned} C(t) &= (99t^4 + 756t^3 + 2253t^2 + 3114t + 1709)/3, \\ D(t) &= 36(t^2 + 3t + 3)(3t^2 + 12t + 13)(3t^2 + 15t + 19). \end{aligned}$$

If we treat Q as a point at infinity on the curve \mathcal{C}_1 and use the method described in [7, page 77] we conclude that \mathcal{C}_1 is birationally equivalent over $\mathbb{Q}(t)$ to the elliptic curve with the Weierstrass equation

$$\mathcal{E}_1 : Y^2 = X^3 + f(t^2 + 3t)X + g(t^2 + 3t),$$

where

$$\begin{aligned} f(u) &= -27(1053u^4 + 10152u^3 + 37530u^2 + 62616u + 39673), \\ g(u) &= 54(9u^2 + 60u + 85)(45u^2 + 192u + 227)(63u^2 + 312u + 397). \end{aligned}$$

The mapping $\varphi : \mathcal{C}_1 \ni (v, s) \mapsto (X, Y) \in \mathcal{E}_1$ is given by

$$\begin{aligned} v &= \frac{2Y - 27d(t) - 6(3t^2 + 15t + 19)(X - 9c(t))}{12(X - 9c(t))(3t^2 + 6t + 4)}, \\ s &= \frac{-(2Y - 27d(t))^2 + 4(2X + 9c(t))(X - 9c(t))^2}{72(3t^2 + 6t + 4)(X - 9c(t))^2}. \end{aligned}$$

The discriminant of \mathcal{E}_1 is

$$\begin{aligned} &2^8 3^{16} (3 + 3t + t^2)^2 (1 + 3t + 3t^2)^2 (4 + 6t + 3t^2)^2 \times \\ &(7 + 9t + 3t^2)^2 (13 + 12t + 3t^2)^2 (19 + 15t + 3t^2)^2, \end{aligned}$$

so that $\mathcal{E}_{1,t}$ is singular for the values $t \in \mathcal{A}$, where

$$\mathcal{A} = \left\{ \frac{-15 \pm \sqrt{-3}}{6}, \frac{-9 \pm \sqrt{-3}}{6}, \frac{-3 \pm \sqrt{-3}}{6}, \frac{-6 \pm \sqrt{-3}}{3}, \frac{-3 \pm \sqrt{-3}}{3}, \frac{-3 \pm \sqrt{-3}}{2} \right\}.$$

For $t \in \mathcal{A}$, the decomposition is of Kodaira classification type I_2 . Let us note that \mathcal{E}_1 is a K3-surface. As we know, the Néron-Severi group over \mathbb{C} , denoted by $\text{NS}(\mathcal{E}_1) = \text{NS}(\mathcal{E}_1, \mathbb{C})$, is a finitely generated \mathbb{Z} -module. From Shioda [8], we have

$$\text{rank NS}(\mathcal{E}_1, \mathbb{C}) = \text{rank } \mathcal{E}_1(\mathbb{C}(t)) + 2 + \sum_{\nu} (m_{\nu} - 1),$$

where the sum ranges over all singular fibers of the pencil $\mathcal{E}_{1,t}$, with m_{ν} the number of irreducible components of the fiber. Here, we have

$$\text{rank NS}(\mathcal{E}_1, \mathbb{C}) = \text{rank } \mathcal{E}_1(\mathbb{C}(t)) + 2 + 6 \cdot 2 \cdot (2 - 1).$$

Since the rank of the Néron-Severi group of a K3-surface cannot exceed 20, then the rank $\mathcal{E}_1(\mathbb{C}(t)) \leq 6$. Although it would be interesting to know the rank of $\mathcal{E}_1(\mathbb{C}(t))$ precisely, we are rather interested with the rank of $\mathcal{E}_1(\mathbb{Q}(t))$. We will show that $\text{rank } \mathcal{E}_1(\mathbb{Q}(t)) \geq 1$.

Let us note that the curve \mathcal{E}_1 has three points of order two

$$\begin{aligned} T_1 &= (6(9(t^2 + 3t)^2 + 60(t^2 + 3t) + 85), 0), \\ T_2 &= (3(45(t^2 + 3t)^2 + 192(t^2 + 3t) + 227), 0), \\ T_3 &= (-3(63(t^2 + 3t)^2 + 312(t^2 + 3t) + 397), 0). \end{aligned}$$

On the curve \mathcal{E}_1 we have also the point $P = (X_P, Y_P)$, where

$$\begin{aligned} X_P &= -3(9t^4 + 108t^3 + 357t^2 + 468t + 229), \\ Y_P &= 54(3t^2 + 6t + 4)(3t^2 + 9t + 7)(3t^2 + 15t + 19). \end{aligned}$$

It is easy to see that the point P is of infinite order on the curve \mathcal{E}_1 . In order to prove this let us consider the curve $\mathcal{E}_{1,1}$ which is specialization of the curve \mathcal{E}_1 at $t = 1$. We have $\mathcal{E}_{1,1} : Y^2 = X^3 - 48867651X + 115230640770$. On the curve $\mathcal{E}_{1,1}$ we have the point $P_1 = (-3513, 493506)$ which is the specialization of the point P at $t = 1$. Now, let us note that

$$3P_1 = \left(\frac{3953140143}{1408969}, \frac{24183154596042}{1672446203} \right).$$

As we know, the points of finite order on the elliptic curve $y^2 = x^3 + ax + b$, $a, b \in \mathbb{Z}$ have integer coordinates [9, page 177], while $3P_1$ is not integral point; therefore, P_1 is not a point of finite order on $\mathcal{E}_{1,1}$, which means that P is not a point of finite order on \mathcal{E}_1 . Therefore, \mathcal{E}_1 is a curve of positive rank. In particular we have proved that $1 \leq \text{rank } \mathcal{E}_1(\mathbb{Q}(t)) \leq \text{rank } \mathcal{E}_1(\mathbb{C}(t)) \leq 6$. However, we have not identified the $\mathbb{Q}(t)$ rank of \mathcal{E}_1 exactly. Numerical calculations give that for many specializations of \mathcal{E}_1 at $t \in \mathbb{Q}$ we have that the rank of $\mathcal{E}_{1,t}(\mathbb{Q})$ is equal to one and this suggest that $\text{rank } \mathcal{E}_1(\mathbb{Q}(t)) = 1$.

Before we prove that the set of rational points on the surface \mathcal{S}_1 is dense in Euclidean topology we prove Zariski density of the set of rational points. Because the curve \mathcal{E}_1 is of positive rank over $\mathbb{Q}(t)$, the set of multiplicities of the point P i.e. $mP = (X_m(t), Y_m(t))$ for $m = 1, 2, \dots$, gives infinitely many rational curves on the curve \mathcal{E}_1 . Now, if we look on the curve \mathcal{E}_1 as on the elliptic surface in the space with coordinates (X, Y, t) we can see that each rational curve (X_m, Y_m, t) is included in the Zariski closure, say \mathcal{R} , of the set of rational points on \mathcal{E}_1 . Because this closure consists of only finitely many components, it has dimension two, and as the surface \mathcal{E}_1 is irreducible, \mathcal{R} is the whole surface. Thus the set of rational points on \mathcal{E}_1 is dense in the Zariski topology and the same is true for the surface \mathcal{S}_1 .

To obtain the statement of our theorem, we have to use two beautiful results: Hurwitz Theorem ([10, p. 78]) and the Silverman's Theorem ([9, p. 368]). Let us recall that Hurwitz Theorem states that if an elliptic curve E defined over \mathbb{Q} has positive rank and one torsion point of order two (defined over \mathbb{Q}) then the set $E(\mathbb{Q})$ is dense in $E(\mathbb{R})$. The same result holds if E has three torsion points of order two under assumption that we have rational point of infinite order which lives on bounded branch of the set $E(\mathbb{R})$.

Silverman Theorem states that if \mathcal{E} is an elliptic curve defined over $\mathbb{Q}(t)$ with positive rank, then for all but finitely many $t_0 \in \mathbb{Q}$, the curve \mathcal{E}_{t_0} obtained from the curve \mathcal{E} by the specialization $t = t_0$ has positive rank. From this result we see that for all but finitely many $t \in \mathbb{Q}$ the elliptic curve $\mathcal{E}_{1,t}$ is of a positive rank.

In order to finish the proof of our theorem let us define the polynomial $X_i(t) = X - \text{coordinate of the torsion point } T_i$ for $i = 1, 2, 3$. Tying these two cited theorems together and the fact that we have inequalities $X_3(t) < X_P(t) < X_1(t) < X_2(t)$ for each $t \in \mathbb{R}$ we conclude that for all but finitely many t the set $\mathcal{E}_{1,t}(\mathbb{Q})$ is dense in the set $\mathcal{E}_{1,t}(\mathbb{R})$. This proves that the set $\mathcal{E}_1(\mathbb{Q})$ is dense in the set $\mathcal{E}_1(\mathbb{R})$ in Euclidean topology. Thus, the set $\mathcal{S}_1(\mathbb{Q})$ is dense in the set $\mathcal{S}_1(\mathbb{R})$ in the Euclidean topology. \square

Using the above result we can deduce the following

Theorem 2.2. *There exists a polynomial $k \in \mathbb{Z}[t]$ with such a property that on the elliptic curve $\mathcal{E} : y^2 = x^3 + k(t)$ there are four independent $\mathbb{Q}(t)$ -rational points in arithmetic progression.*

Proof. In order to construct a polynomial $k \in \mathbb{Z}[t]$ with such a property that for all $t \in \mathbb{Z}$ value $k(t)$ is an integer of AP4 type we use the points P and T_1 we have constructed in the proof of Theorem 2.1. Let us note that

$$P+T_1 = (3(99t^4+432t^3+795t^2+684t+251), -486(t^2+3t+3)(3t^2+3t+1)(3t^2+6t+4)).$$

Let us note that this point leads us to the point

$$\varphi(P+T_1) = \left(\frac{3t^2+9t+10}{3(2t+3)}, -\frac{(1+3t+3t^2)(18t^4+162t^3+516t^2+738t+413)}{9(2t+3)^2} \right),$$

which is the point on the curve \mathcal{C}_1 . Performing all necessary simplifications we find that the point $\varphi(P+T_1)$ lead us to the polynomial solution of the equation defining the surface \mathcal{S}_1 in the form

$$\begin{aligned} p &= 18t^4 + 54t^3 + 30t^2 - 72t - 73, & q &= 18t^4 + 90t^3 + 192t^2 + 210t + 107, \\ r &= 18t^4 + 126t^3 + 354t^2 + 456t + 233, & s &= 18t^4 + 162t^3 + 516t^2 + 738t + 413. \end{aligned}$$

Using this parametric solution and the remark on the beginning of our proof we find the curve

$$(5) \quad \mathcal{E} : y^2 = x^3 + k(t),$$

where

$$k(t) = -324^2(2t+3)^2(3t^2+9t+10)^2(6t^2+18t+17)^2h(t),$$

and

$$h(t) = (t^2+3t)^4 + 612(t^2+3t)^3 + 300(t^2+3t)^2 - 3504(t^2+3t) - 5329.$$

On the curve \mathcal{E} we have four points in arithmetic progression

$$\begin{aligned} P_1 &= (tp(t), p(t)(18t^4 + 54t^3 + 30t^2 - 72t - 73)), \\ P_2 &= ((t+1)p(t), p(t)(18t^4 + 90t^3 + 192t^2 + 210t + 107)), \\ P_3 &= ((t+2)p(t), p(t)(18t^4 + 126t^3 + 354t^2 + 456t + 233)), \\ P_4 &= ((t+3)p(t), p(t)(18t^4 + 162t^3 + 516t^2 + 738t + 413)), \end{aligned}$$

where $p(t) = 108(2t+3)(3t^2+9t+10)(6t^2+18t+17)$.

We will show that the above points are independent in the group $\mathcal{E}(\mathbb{Q}(t))$ of all $\mathbb{Q}(t)$ -rational points on the curve \mathcal{E} . We specialize the curve \mathcal{E} at $t = 1$ and we get the elliptic curve \mathcal{E}_1 given by the equation

$$\mathcal{E}_1 : y^2 = x^3 - 111610206808689600.$$

On the curve \mathcal{E}_1 we have the points

$$\begin{aligned} P_{1,1} &= (487080, 62833320) & P_{2,1} &= (974160, 901585080) \\ P_{3,1} &= (1461240, 1734491880) & P_{4,1} &= (1948320, 2698910280), \end{aligned}$$

which are specialization of the points P_1, P_2, P_3, P_4 at $t = 1$. Using now program APECS [2] we obtain that the determinant of the height matrix of the points $P_{1,1}, P_{2,1}, P_{3,1}, P_{4,1}$ is equal to 266.618020487005. This proves that the points $P_{1,1}, P_{2,1}, P_{3,1}, P_{4,1}$ are independent on the curve \mathcal{E}_1 and thus we get that that the points P_1, P_2, P_3, P_4 are independent on the curve \mathcal{E} . \square

Remark 2.3. Let us consider the polynomial $g(t, x) = x^3 + k(t)$, where $k \in \mathbb{Z}[t]$ is defined in the proof of the above theorem. Then in order to find a rational value of t which lead to five points in arithmetic progression on the curve \mathcal{E}_t we must have $g(t, (t-1)p(t)) = \square$ or $g(t, (t+4)p(t)) = \square$. In other words we must be able to find rational point on one of the hyperelliptic curves of genus 3

$$C_1 : v^2 = 324t^8 + 648t^7 - 4428t^6 - 21384t^5 - 34884t^4 + \\ - 17388t^3 + 12828t^2 + 12804t - 791, \\ C_2 : v^2 = 324t^8 + 7128t^7 + 63612t^6 + 309096t^5 + 912816t^4 \\ + 1704132t^3 + 1985988t^2 + 1332624t + 397009.$$

Unfortunately, we are unable to find rational point on any of this curves which lead to nonzero value of $k(t)$.

3. RATIONAL POINTS IN ARITHMETIC PROGRESSIONS ON $y^2 = x^n + k$ FOR $n \geq 4$

In this section we consider a natural generalization of the problem we have considered in Section 1. To be more precise we consider the following

Question 3.1. *Let $n \in \mathbb{N}$ be fixed and suppose that $n \geq 4$. It is possible to find an integer k with such a property that on the hyperelliptic curve $y^2 = x^n + k$ there are at least four rational points in arithmetic progression?*

We show that if n is odd then the answer on the above question is affirmative and it is possible to find infinitely many demanded k 's.

In the case when n is even we show that it is possible to construct infinitely many k 's with such a property that on the curve $y^2 = x^n + k$ there is three term arithmetic progression of rational points with x -coordinates of the form $a, 3a, 5a$, where a is positive. Using the involution $(x, y) \mapsto (-x, y)$ we can see that on these curves we will have six rational points in arithmetic progression with x -coordinates of the form $-5a, -3a, -a, a, 3a, 5a$.

We start with the following

Theorem 3.2. *Let us fix an $n \geq 2$. Then there are infinitely many integers k with such a property that on the hyperelliptic curve $H : y^2 = x^{2n+1} + k$ there are four points in arithmetic progression.*

Proof. In order to prove our theorem let us consider a hyperelliptic curve with the equation $y^2 = ax^{2n+1} + bx^2 + cx + d =: f(x)$, where

$$(6) \quad \begin{aligned} a &= \frac{-p^2 + 3q^2 - 3r^2 + s^2}{2^{2n+1} - 2}, & b &= \frac{p^2 - 2q^2 + r^2}{2}, \\ c &= \frac{-(2^{2n} - 2)p^2 - 3q^2 + (2^{2n} + 2)r^2 - s^2}{2^{2n+1} - 2}, & d &= q^2. \end{aligned}$$

For a, b, c, d defined in this way we have that the points $(-1, p), (0, q), (1, r), (2, s)$ are on the curve and are in arithmetic progression. Now, let us note that if we are able to show that the system of equations in the variables p, q, r, s given by

$$(7) \quad \begin{cases} p^2 - 2q^2 + r^2 = 0, \\ -(2^{2n} - 2)p^2 - 3q^2 + (2^{2n} + 2)r^2 - s^2 = 0, \end{cases}$$

has infinitely many rational solutions, then the points

$$(-a, pa^n), \quad (0, qa^n), \quad (a, ra^n), \quad (2a, sa^n)$$

will be in arithmetic progression on the curve $y^2 = x^{2n+1} + da^{2n}$, where a, b are given by (6).

Now, we show that the system (7) has infinitely many solutions in rational numbers. In order to do this let us parametrize all solutions of the first equation in the system (7). Using standard method we find parametrization given by

$$p = 2u^2 - 4uv + v^2, \quad q = 2u^2 - 2uv + v^2, \quad r = -2u^2 + v^2.$$

Putting now the calculated values for p, q, r into the second equation of the system (7) and taking $u = 1$ we get

$$\mathcal{C}_n^o : s^2 = v^4 + 4(2^{2n+1} - 1)v^3 - 8(3 \cdot 2^{2n} - 1)v^2 + 8(2^{2n+1} - 1)v + 4.$$

The curve \mathcal{C}_n^o is a quartic curve with rational point $Q = (0, -2)$. If we treat Q as a point at infinity on the curve \mathcal{C} and use the method described in [7, page 77] one more time we conclude that \mathcal{C}_n^o is birationally equivalent over \mathbb{Q} to the elliptic curve with the Weierstrass equation

$$\mathcal{E}_n^o : Y^2 = X^3 - 27(3 \cdot 2^{4n+2} + 1)X + 54(9 \cdot 2^{4n+2} - 1).$$

The mapping $\varphi : \mathcal{E}_n^o \ni (X, Y) \mapsto (v, s) \in \mathcal{C}_n^o$ is given by

$$v = \frac{2Y - 27 \cdot 2^{2n+2}(2^{4n+2} - 1)}{6(X - 3(3 \cdot 2^{4n+2} - 1))} - (2^{2n+1} - 1),$$

$$s = -(v + 2^{2n+1} - 1)^2 + \frac{2X + 3(3 \cdot 2^{4n+2} - 1)}{9}.$$

Inverse mapping $\psi : \mathcal{C}_n^o \ni (v, s) \mapsto (X, Y) \in \mathcal{E}_n^o$ has the form

$$X = \frac{3}{2}(3v^2 + 6(2^{2n+1} - 1)v + 3s - 4(3 \cdot 2^{2n} - 1)),$$

$$Y = \frac{27}{2}(v^3 + 3(2^{2n+1} - 1)v^2 - 4(3 \cdot 2^{2n} - 1)v + (2^{2n+1} - 1)s + 2(2^{2n+1} - 1)).$$

Let us note that the curve \mathcal{E}_n^o has three points of order two

$$T_1 = (6, 0), \quad T_2 = (3(3 \cdot 2^{2n+1} - 1), 0), \quad T_3 = (-3(3 \cdot 2^{2n+1} + 1), 0).$$

On the curve \mathcal{E}_n^o we have also the point P_n given by

$$P_n = (-3(3 \cdot 2^{2n+1} - 5), 54(2^{2n+1} - 1)).$$

It is easy to see that the point P_n is of infinite order on the curve \mathcal{E}_n^o . In order to prove this we compute the point $4P_n = (X, Y)$. We have where

$$X = \frac{3 \cdot 2^{-4n-2}(3 + 3 \cdot 2^{16n} - 2^{4n+4} + 13 \cdot 2^{8n+1} + 2^{12n+5})}{(2^{4n} - 1)^2},$$

$$Y = \frac{3 \cdot 2^{-2n-1}(3 \cdot 2^{8n} + 1)}{2^{4n} - 1}X + 27 \cdot 2^{2n}(2^{4n} - 1).$$

It is easy to see that under our assumption $n \geq 2$ the X -coordinate of the point $4P_n$ is not an integer. Thus, theorem of Nagell and Lutz implies that the point P_n is of infinite order. This implies that the set of rational solutions of the system (7) is infinite and thus our theorem is proved. \square

Remark 3.3. Using `mwrnk` program we calculated the rank r_n of elliptic curve \mathcal{E}_n^o and generators for the free part of the group $\mathcal{E}_n^o(\mathbb{Q})$ for $2 \leq n \leq 8$. Results of our computations are given below.

n	r_n	Generators for the free part of the group $\mathcal{E}_n^o(\mathbb{Q})$
2	1	(303, 17820)
3	1	(1167, 6966)
4	2	(4623, 27702), (11773, 1175552)
5	2	(18447, 110646), (350011167/6241, 6184493104374/493039)
6	1	(73743, 442422)
7	3	(294927, 1769526), (153394089/400, 1214402001813/800), (124356529/256, 1101957449705/4096)
8	1	(1179663, 7077942)

Form the above theorem we get an interesting

Corollary 3.4. *Let $n \geq 2$ and consider the family of hyperelliptic curves given by the equation $C_k : y^2 = x^{2n+1} + k$. Then the set \mathcal{A} of integers k with such a property that on the curve C_k there are at least 8 rational points, is infinite. Moreover, we can construct the set \mathcal{A} that for each pair $k_1, k_2 \in \mathcal{A}$ the curves C_{k_1}, C_{k_2} are not isomorphic.*

Proof. First part of our theorem is an immediate consequence of the previous theorem. Second part of our theorem is a simple consequence of the following reasoning. Curves C_{k_1} and C_{k_2} are isomorphic if and only if $k_1/k_2 \in \mathbb{Q}^{10}$. From the previous theorem we can take $k = q^2 = (v^2 - 2v + 2)^2$ for some $v \in \mathbb{Q}$ which is calculated from the point which lies on the elliptic curve C_n^o . Let us suppose that we constructed the integers k_1, k_2, \dots, k_n such that the curves C_{k_i} are pairwise non-isomorphic. We have that $k_i = q_i^2 = (v_i^2 - 2v_i + 2)^2$ for $i = 1, 2, \dots, n$. Then the n curves $(2v^2 - 2v + 2)^2 = (v_i^2 - 2v_i + 2)^2 w^{10}$ for $i = 1, 2, \dots, n$ are all of genus ≥ 2 , thus the set of $\mathbb{C}_1(\mathbb{Q}) \cup \dots \cup C_n(\mathbb{Q})$ is finite (Faltings Theorem [4]). Because the elliptic curve C_n^o has infinitely many rational points we can find v_{n+1} such that the curve $C_{k_{n+1}}$ with $k_{n+1} = (v_{n+1}^2 - 2v_{n+1} + 2)^2$ is not isomorphic to any of the curves C_i for $i = 1, 2, \dots, n$. Using now the presented reasoning we can construct an infinite set \mathcal{A} with demanded property. \square

The above corollary and the Corollary 3.10 gives a generalization A. Bremner result from [1] (Theorem 2.1 and Theorem 3.1) .

Theorem 3.2 we have proved suggest the following

Question 3.5. *Let us fix an integer $n \geq 1$. What is the smallest value $|k_{2n+1}| \in \mathbb{N}$, say M_{2n+1} , with such a property that on the curve $y^2 = x^{2n+1} + k_{2n+1}$ there are at least four rational points in arithmetic progression?*

Example 3.6. If we take $n = 2$ then from the existence of rational point of infinite order on the curve \mathcal{E}_2^o we get that

$$M_5 \leq 3391541395170708368688169980^4 \cdot 2609^2 \cdot 127165689041^2.$$

Moreover in the light of Corollary 3.4 it is natural to sate the following

Question 3.7. *Let us fix an integer $n \geq 2$ and consider the hyperelliptic curve $H_{2n+1} : y^2 = x^{2n+1} + k_{2n+1}$ where k_{2n+1} is constructed with the use method we have presented in the proof of the previous theorem. In particular on the curve H_{2n+1} we have four points in arithmetic progression, say P_i for $i = 1, 2, 3, 4$. Are the divisors $(P_i) - (\infty)$ independent in the jacobian variety associated with the curve H_{2n+1} ?*

Now, we prove the following

Theorem 3.8. *Let us fix an $n \geq 2$. Then there are infinitely many integers k with such a property that on the hyperelliptic curve $H : y^2 = x^{2n} + k$ there are six points in arithmetic progression.*

Proof. In order to prove our theorem let us consider a hyperelliptic curve with the equation $y^2 = x^{2n} + ax^2 + bx + c =: f(x)$, where

$$\begin{aligned} a &= \frac{p^2 - 2q^2 + r^2 - 5^{2n} + 2 \cdot 3^{2n} - 1}{8}, \\ b &= \frac{-2p^2 + 3q^2 - r^2 + 5^{2n} - 3^{2n+1} + 2}{2}, \\ c &= \frac{15p^2 - 10q^2 + 3r^2 - 3 \cdot 5^{2n} + 10 \cdot 3^{2n} - 15}{8}. \end{aligned}$$

Then we have that $f(1) = p^2$, $f(3) = q^2$, $f(5) = r^2$. It is easy to see that in order to prove our theorem it is enough to find infinitely many solutions of the system of equations $a = b = 0$ in rational numbers p, q, r . Indeed, if $a = b = 0$, then on the curve $y^2 = x^{2n} + c$ we will have six points in arithmetic progression which x -coordinates belong to the set $\{-5, -3, -1, 1, 3, 5\}$.

System $a = b = 0$ is equivalent with the system of equations

$$(8) \quad \begin{cases} q^2 = p^2 + 3^{2n} - 1, \\ r^2 = p^2 + 5^{2n} - 1. \end{cases}$$

By putting $p = u + 1$, $q = tu + 3^n$ we find that all rational solutions of the equation $q^2 = p^2 + 3^{2n} - 1$ are contained in the formulas

$$p = \frac{t^2 - 2 \cdot 3^n t + 1}{t^2 - 1}, \quad q = -\frac{3^n t^2 - 2t + 3^n}{t^2 - 1}.$$

Putting the value of p we have calculated into the second equation of the system (8) we get the equation of the quartic curve

$$\mathcal{C}_n^e : s^2 = 5^{2n} t^4 - 4 \cdot 3^n t^3 - 2(5^{2n} - 2 \cdot 3^{2n} - 2)t^2 - 4 \cdot 3^n t + 5^{2n} =: g(t),$$

where $s = (t^2 - 1)r$. For convenience of our calculations let us put $u = 3^n$ and $v = 5^n$. Then, the polynomial g takes the form $g_{u,v}(t) = v^2 t^4 - 4ut^3 - 2(v^2 - 2u^2 - 2)t^2 - 4u + v^2$.

The curve $\mathcal{C}_n^e : s^2 = g_{u,v}(t)$ is a quartic curve with rational point $Q = (0, v)$. If we treat Q as a point at infinity on the curve \mathcal{C}_n^e and use the method described in [7, page 77] one more time we conclude that \mathcal{C}_n^e is birationally equivalent over \mathbb{Q} to the elliptic curve with the Weierstrass equation

$$\begin{aligned} \mathcal{E}_n^e : Y^2 = X^3 - 27(v^4 - (u^2 + 1)v^2 + u^4 - u^2 + 1)X + \\ + 27(1 + u^2 - 2v^2)(2u^2 - v^2 - 1)(u^2 + v^2 - 2). \end{aligned}$$

The mapping $\varphi : \mathcal{E}_n^e \ni (X, Y) \mapsto (t, x) \in \mathcal{C}_n^e$ is given by

$$\begin{aligned} t &= \frac{v^3 Y - 27u(u^2 - v^2)(v^2 - 1)}{3v^2(v^2 X - 3(3u^2 - 2v^2 + v^4 - 2u^2 v^2))} + \frac{u}{v^2}, \\ s &= -\frac{1}{v^3} \left(v^2 t - \frac{u}{v^2} \right)^2 + \frac{v^2 X + 9u^2 - 6(u^2 + 1)v^2 + 3v^4}{9v^3}. \end{aligned}$$

Inverse mapping $\psi : \mathcal{C}_n^e \ni (t, s) \mapsto (X, Y) \in \mathcal{E}_n^e$ has the form

$$X = \frac{2 - 6tu + 2u^2 + 3sv + (3t^2 - 1)v^2}{2},$$

$$Y = -\frac{27}{2}(su + ((3t^2 + 1)u - 2t(u^2 + 1))v - stv^2 - (t^3 - t)v^3).$$

Let us note that the curve \mathcal{E}_n^e has three points of order two

$$T_1 = (3(1 + u^2 - 2v^2), 0), \quad T_2 = (3(u^2 + v^2 - 2), 0), \quad T_3 = (-3(1 - 2u^2 + v^2), 0).$$

On the curve \mathcal{E}_n^e we have also the point P_n given by

$$P_n = (3(u^2 + v^2 + 1), -27uv).$$

It is easy to see that the point P is of infinite order on the curve \mathcal{E}_n^e . Indeed, if we calculate $2P_n = (X_2, Y_2)$ then we have

$$X_2 = \frac{3(3u^4 - 2(u^2 + 1)u^2v^2 + (3 - 2u^2 + 3u^4)v^4)}{4u^2v^2}.$$

Due to the fact that $u = 3^n$ and $v = 5^n$ it is easy to see that for any choice for $n \geq 2$ the above fraction is not an integer. From the Nagell-Lutz theorem we get that the point P_n is not of finite order on the curve \mathcal{E}_n^e . This implies that the set of rational solutions of the system (8) is infinite. As a consequence we get that for any $n \geq 2$ we can construct infinitely many k 's with such a property that on the curve $y^2 = x^{2n} + k$ we have three points in arithmetic progression with x -coordinates belonging to the set $\{m, 3m, 5m\}$ with $m > 0$. Note that on this curve we have also rational points with x -coordinates belonging to the set $\{-m, -3m, -5m\}$. This observation finishes the proof of our theorem. \square

Remark 3.9. Using `mwrnk` program we calculated the rank r_n of elliptic curve \mathcal{E}_n^e and generators for the free part of the group $\mathcal{E}_n^e(\mathbb{Q})$ for $2 \leq n \leq 10$. Results of our computations are given below.

n	r_n	Generators for the free part of the group $\mathcal{E}_n^e(\mathbb{Q})$
2	1	(3840, 176256)
3	2	(80808, 14478912), (130704, 40007520)
4	2	(1230432, 116328960), (31769376/25, 24804389376/125)
5	1	(36278088, 68748343488)
6	1	(836384640, 5022400795776)
7	1	(19863352968, 370669722011712)
8	1	(480955252992, 27480236025415680)

Using similar argument as in the proof of the Corollary 3.4 we can prove the following

Corollary 3.10. *Let $n \geq 2$ and consider the family of hyperelliptic curves given by the equation $C_k : y^2 = x^{2n} + k$. Then the set \mathcal{A} of integers k with such a property that on the curve C_k there are at least 12 rational points, is infinite. Moreover, we can construct the set \mathcal{A} that for each pair $k_1, k_2 \in \mathcal{A}$ the curves C_{k_1}, C_{k_2} are not isomorphic.*

Similarly as in the case of odd exponents we can state the following questions.

Question 3.11. *Let us fix an integer $n \geq 2$. What is the smallest value $|k_{2n}| \in \mathbb{N}$, say M_{2n} with such a property that on the curve $y^2 = x^{2n} + k_{2n}$ there are at least three rational points in arithmetic progression?*

Question 3.12. *Let us fix an integer $n \geq 2$ and consider the hyperelliptic curve $H_{2n} : y^2 = x^{2n} + k_{2n}$ where k_{2n} is constructed with the use method we have presented in the proof of the above theorem. In particular, on the curve H_{2n} we have three points in arithmetic progression, say P_i for $i = 1, 2, 3$. Do the divisors $(P_i) - (\infty)$ are independent in the jacobian variety associated with the curve H_{2n} ?*

4. SEXTIC THREEFOLD RELATED TO FIVE RATIONAL POINTS IN ARITHMETIC PROGRESSIONS ON $y^2 = x^3 + k$

In the section 2 we have used very natural reasoning in order to construct quartic surface which is closely related to the problem of existence of numbers of AP4 type. A natural question arises whether we can construct an algebraic variety, say \mathcal{T} , with such a property that each (nontrivial) rational point on \mathcal{T} gives an integer k of AP5 type, so an integer k such that on the curve $y^2 = x^3 + k$ we have five rational points in arithmetic progression. In order to construct demanded hypersurface we will use similar method as in section 2.

Let $f(x, y) = y^2 + ay - (bx^3 + cx^2 + dx + e) \in \mathbb{Q}[a, b, c, d, e][x]$ and consider the curve $C : f(x, y) = 0$. Using now change of coordinates

$$(x, y) = \left(\frac{X - 12c}{36b}, \frac{Y - 108ab}{216b} \right) \text{ with inverse } (X, Y) = (12(c + 3bx), 108b(a + 2y)),$$

we can see that the curve C is birationally equivalent to the curve

$$E : Y^2 = X^3 - 432(c^2 - 3bd)X + 432(27a^2b^2 + 8c^3 - 36bcd + 108b^2e).$$

Now, let p, q, r, s, t be free parameters and consider the system of equations

$$f(-2, p) = f(-1, q) = f(0, r) = f(1, s) = f(2, t) = 0.$$

This system has exactly one solution in respect to a, b, c, d, e . This solution belong to the field $\mathbb{Q}(p, q, r, s, t)$ and has the form

$$a = \frac{A}{6H}, \quad b = \frac{B}{6H}, \quad c = \frac{C}{6H}, \quad D = \frac{D}{6H}, \quad e = \frac{E}{6H},$$

where $A, \dots, E \in \mathbb{Z}[p, q, r, s, t]$ are homogeneous, A is of degree two and B, \dots, E are of degree three. Moreover, we have $H = p - 4q + 6r - 4s + t$. From this computations we can see that in order to find an integer k with such a property that on the curve $y^2 = x^3 + k$ there are five points in arithmetic progression it is enough to find rational points on the sextic threefold given by the equation

$$\mathcal{T} : C(p, q, r, s, t)^2 = 3B(p, q, r, s, t)D(p, q, r, s, t),$$

where

$$B = (p - 3q + 3r - s)t^2 - (p^2 - 3q^2 + 3r^2 - s^2)t + (q - 3r + 3s)p^2 + \\ - (q^2 - 3r^2 + 3s^2)p + 2(q - s)(3qr - 3r^2 - 4qs + 3rs),$$

$$C = -3((t^2 + p^2)(q - 2r + s) - (t + p)(q^2 - 2r^2 + s^2) + 2r(q^2 - qr - rs + s^2)),$$

$$D = -(p - 6q + 3r + 2s)t^2 + (p^2 - 6q^2 + 3r^2 + 2s^2)t + (2q + 3r - 6s)p^2 + \\ - (2q^2 + 3r^2 - 6s^2)p - 8(q - s)(3qr - 3r^2 - 4qs + 3rs).$$

We have obvious automorphisms of order two which act on the \mathcal{T} given by

$$(p, q, r, s, t) \mapsto (t, s, r, q, p), \quad (p, q, r, s, t) \mapsto (-p, -q, -r, -s, -t).$$

By a trivial rational point on the hypersurface \mathcal{T} we will understand the point (p, q, r, s, t) which lies on one of the lines

$$L_1 : p = q = r, s = t, \quad L_2 : p = q = s, r = t, \quad \dots, \quad L_{10} : r = s = t, p = q,$$

or on the hyperplane $H : p - 4q + 6r - 4s + t = 0$.

We perform small numerical calculations in order to find a non-trivial rational point on \mathcal{T} . We compute all integer solutions of the equation which define the hypersurface \mathcal{T} under assumption $\max\{|p|, |q|, |r|, |s|, |t|\} \leq 10^2$. Unfortunately in this range all solutions are trivial. This suggest the following

Question 4.1. *Is the set of non-trivial rational points on the hypersurface \mathcal{T} non-empty?*

REFERENCES

- [1] A. Bremner, *On the Equation $Y^2 = X^5 + k$* , Experiment. Math. **17**(3) (2008), 371-374.
- [2] I. Connell, APECS: Arithmetic of Plane Elliptic Curves, available from <ftp.math.mcgill.ca/pub/apecs/>.
- [3] N. Elkies, *On $A^4 + B^4 + C^4 = D^4$* , Math. of Comp. **51** (1988), 825-835.
- [4] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, *Invent. Math.* **73** (1983), 349-366.
- [5] J.-B. Lee, W. Y. Vélez, *Integral solutions in arithmetic progression for $y^2 = x^3 + k$* , Periodica Mathematica Hungarica **25** (1992), 31-49.
- [6] S. P. Mohanty, *Integral solutions in arithmetic progression for $y^2 = x^3 + k$* , Acta Math. Acad. Sci. Hung. **34** (3-4) (1980), 261-265.
- [7] L. J. Mordell, *Diophantine equations*, Academic Press, London, 1969.
- [8] T. Shioda, *On elliptic modular surfaces*, J. Math. Soc. Japan **24** (1972), 20-59.
- [9] J. Silverman, *The Arithmetic of Elliptic Curves*, Springer-Verlag, New York, 1986.
- [10] T. Skolem, *Diophantische Gleichungen*, New York : Chelsea Publishing Company, 1950.

Maciej Ulas
 Jagiellonian University
 Institute of Mathematics
 Łojasiewicza 6
 30 - 348 Kraków, Poland
 e-mail: Maciej.Ulas@im.uj.edu.pl
 e-mail: maciej.ulas@gmail.com