Two-step General Linear Methods for Retarded Functional Differential Equations

Anton Tuzov *

Abstract

This paper presents a class of Two-Step General Linear Methods for the numerical solution of Retarded Functional Differential Equations. Explicit methods up to order five are constructed. To avoid order reduction for mildly stiff problems the uniform stage order of the methods is chosen to be close to uniform order.

1 Two-step General Linear Methods for Ordinary Differential Equations

For the numerical approximation of the solution y(t) of a system of Ordinary Differential Equations

$$y'(t) = f(t, y), \quad t \in [t_0, T],$$

 $y(t_0) = y_0,$
(1)

where $f : \mathbb{R} \times \mathbb{R}^d \longrightarrow \mathbb{R}^d$, $y_0 \in \mathbb{R}^d$, we consider the class of General Linear Methods [5]

$$Y_{i}^{[n]} = \sum_{j=1}^{s} a_{ij} h F_{j}^{[n]} + \sum_{j=1}^{r} u_{ij} y_{j}^{[n-1]}, \qquad i = 1, \dots, s,$$

$$y_{i}^{[n]} = \sum_{j=1}^{s} b_{ij} h F_{j}^{[n]} + \sum_{j=1}^{r} v_{ij} y_{j}^{[n-1]}, \qquad i = 1, \dots, r,$$

$$F_{i}^{[n]} = f(t_{n-1} + c_{i}h, Y_{i}^{[n]}), \qquad i = 1, \dots, s,$$
(2)

where $y_1^{[n-1]}, \ldots, y_r^{[n-1]}$ — input vectors, available at step number n, $Y_1^{[n]}, \ldots, Y_s^{[n]}$ — stage values, $F_1^{[n]}, \ldots, F_s^{[n]}$ — derivative values, $a_{ij}, u_{ij}, b_{ij}, v_{ij}$. — coefficients of the method.

^{*}Department of Control systems, Siberian State Aerospace University, Krasnoyarsk, Russia, e-mail: tuzov@sibsau.ru.

Let us restrict ourselves to two-step GLMs and choose r = s + 2,

$$y_1^{[n-1]} \approx y(t_{n-1}), \quad y_2^{[n-1]} \approx y(t_{n-2}), \quad y_{2+i}^{[n-1]} \approx hy'(t_{n-2} + c_i h), \ i = 1, \dots, s.$$
 Then
$$y_1^{[n-1]} = y_{n-1}, \qquad y_2^{[n-1]} = y_{n-2}, \qquad y_{2+i}^{[n-1]} = hf(t_{n-2} + c_i h, Y_i^{[n-1]}) = hF_i^{[n-1]}, \ i = 1, \dots, s.$$

and (2) takes the form

$$Y_{i}^{[n]} = h \sum_{j=1}^{s} a_{ij} F_{j}^{[n]} + u_{i1}y_{n-1} + u_{i2}y_{n-2} + h \sum_{j=1}^{s} u_{i,2+j}F_{j}^{[n-1]}, \qquad i = 1, \dots, s,$$

$$y_{n} = h \sum_{j=1}^{s} b_{1j} F_{j}^{[n]} + v_{11}y_{n-1} + v_{12}y_{n-2} + h \sum_{j=1}^{s} v_{1,2+j}F_{j}^{[n-1]}, \qquad (3)$$

$$F_{i}^{[n]} = f(t_{n-1} + c_{i}h, Y_{i}^{[n]}), \qquad i = 1, \dots, s.$$

In the construction of GLMs it is assumed that $y_i^{[n-1]} = u_i y(t_{n-1}) + v_i h y'(t_{n-1}) + O(h^2)$ and 'preconsistensy conditions' holds

$$\begin{aligned}
Vu &= u, \\
Uu &= \mathbf{1}.
\end{aligned}$$
(4)

For (3) we have $u_1 = 1$, $v_1 = 0$, $u_2 = 1$, $v_2 = -1$, $u_{2+i} = 0$, $v_{2+i} = 1$, $i = 1, \ldots, s$. It follows from (4) that

$$u_{i2} = 1 - u_{i1}, \quad i = 1, \dots, s,$$

 $v_{12} = 1 - v_{11}.$

Let us denote

$$\begin{aligned} K_i^{[n]} &:= F_i^{[n]} \quad \widetilde{a}_{ij} := u_{i,2+j}, \quad u_i := u_{i1}, \quad \Rightarrow u_{i2} = 1 - u_i, \quad j = 1, \dots, s, \ i = 1, \dots, s, \\ b_j &:= b_{1j}, \quad \widetilde{b_j} := v_{1,2+j}, \quad v := v_{11}, \quad \Rightarrow v_{12} = 1 - v, \quad j = 1, \dots, s, \end{aligned}$$

then the method (3) satisfying 'preconsistency conditions' (4) takes the form

$$y_{n} = (1-v)y_{n-2} + vy_{n-1} + h\sum_{j=1}^{s} \widetilde{b}_{j}K_{j}^{[n-1]} + h\sum_{j=1}^{s} b_{j}K_{j}^{[n]},$$

$$K_{i}^{[n]} = f(t_{n-1} + c_{i}h, Y_{i}^{[n]}), \qquad i = 1, \dots, s,$$

$$Y_{i}^{[n]} = (1-u_{i})y_{n-2} + u_{i}y_{n-1} + h\sum_{j=1}^{s} \widetilde{a}_{ij}K_{j}^{[n-1]} + h\sum_{j=1}^{s} a_{ij}K_{j}^{[n]}, \quad i = 1, \dots, s.$$
(5)

2 Two-step General Linear Methods for Retarded Functional Differential Equations

We begin with notations introduced in [1].

Let $r \in [0, +\infty)$, and \mathcal{C} be the space of continuous functions $[-r, 0] \longrightarrow \mathbb{R}^d$, equipped with the maximum(uniform) norm $\|\phi\| = \max_{\theta \in [-r,0]} |\phi(\theta)|$, $\phi \in \mathcal{C}$, where $|\cdot|$ is an arbitrary norm on \mathbb{R}^d . Let u be continuous function $[a - r, b) \longrightarrow \mathbb{R}^d$, where a < b. Then $\forall t \in [a, b)$ shift function is well defined by $u_t(\theta) = u(t + \theta)$, $\theta \in [-r, 0]$, and $u_t \in \mathcal{C}$.

Let us consider a system of Retarded Functional Differential Equations

$$y'(t) = f(t, y_t), \quad t \in [t_0, T],$$

$$y_{t_0}(\theta) = \phi(\theta), \quad \theta \in [-r, 0].$$
(6)

where $(t_0, \phi) \in \Omega$, $f: \Omega \longrightarrow \mathbb{R}^d$, $\Omega \subset \mathbb{R} \times C$, Ω is open set. It is assumed that there exists a unique solution of (6).

We introduce the class of two-step GLMs for RFDEs on the base of approach proposed in [1]. We can reformulate the method (5) for RFDEs (6) as follows

$$\eta^{[n]}(\alpha h) = (1 - v(\alpha))\eta^{[n-1]}(0) + v(\alpha)\eta^{[n-1]}(h) + h\sum_{j=1}^{s} \widetilde{b_{j}}(\alpha)K_{j}^{[n-1]} + h\sum_{j=1}^{s} b_{j}(\alpha)K_{j}^{[n]},$$

$$\alpha \in [0, 1],$$

$$K_{i}^{[n]} = f(t_{n-1} + c_{i}h, Y_{c_{i}h}^{[n]}), \qquad i = 1, \dots, s,$$

$$Y^{[n]i}(\alpha h) = (1 - u_{i}(\alpha))\eta^{[n-1]}(0) + u_{i}(\alpha)\eta^{[n-1]}(h) + h\sum_{j=1}^{s} \widetilde{a}_{ij}(\alpha)K_{j}^{[n-1]} + h\sum_{j=1}^{s} a_{ij}(\alpha)K_{j}^{[n]}, \qquad (7)$$

$$\alpha \in [0, c_{i}], \qquad i = 1, \dots, s,$$

$$\eta^{[n]}(\theta) = \eta_{h}^{[n-1]}(\theta), \qquad \theta \in [-r, 0],$$

$$Y^{[n]i}(\theta) = \eta_{h}^{[n-1]}(\theta), \qquad \theta \in [-r, 0], \qquad i = 1, \dots, s,$$

where

$\eta^{[n-1]}: \ [-r,h] \longrightarrow \mathbb{R}$	$A^d, K_i^{[n-1]} \in \mathbb{R}^d$	are available as approximations
		computed in the step $n-1$,
$Y^{[n] \ i}$		stage functions,
$K_i^{[n]}$		stage values,
$u_i(\cdot), \ \widetilde{a}_{ij}(\cdot), a_{ij}(\cdot), \ a_{ij$	$w(\cdot), \ \widetilde{b_j}(\cdot), \ b_j(\cdot) =$	coefficients of the method.



Thus $\eta^{[n]}(\alpha h) \approx y(t_{n-1} + \alpha h), \ \alpha \in [0, 1], \quad \eta^{[n]}(\theta) \approx y(t_{n-1} + \theta), \ \theta \in [-r, 0],$ hence $\eta_h^{[n]} \approx y_{t_n}$ on [-r - h, 0], where $t_n = t_{n-1} + h$.

Remark 2.1. We chose two-step methods among multi-step methods $(k \ge 2)$ for the following reasons.

- For multistep methods, the local error $E(h, t_{n-1}, y_{t_{n-1}})$ has the required order only if exact solution y(t) is sufficiently smooth on $[t_{n-k}, t_n]$. This is rather severe assumption for many problems (6). For the case of two-step methods (k = 2) this condition imposes the weakest restriction on stepsize.
- Furthermore, in the case of two-step methods, starting procedure and stepsize strategy seem to be simplest ones.

Let us denote

$$\eta := \eta^{[n]}, \ K_i := K_i^{[n]}, \quad \overline{\eta} := \eta^{[n-1]}, \ \overline{K}_i := K_i^{[n-1]}, \quad \sigma := t_{n-1},$$

then the method (7) can be reformulated in Stefano Maset's notations as follows



3 Two-step GLMs for RFDEs in Stefano Maset's notations

Let us consider a system of Retarded Functional Differential Equations

$$x'(t) = f(t, x_t), \quad t \in [t_0, T],$$

$$x_{t_0}(\theta) = \phi(\theta), \qquad \theta \in [-r, 0].$$
(9)

where $(t_0, \phi) \in \Omega, f: \Omega \longrightarrow \mathbb{R}^d, \quad \Omega \subset \mathbb{R} \times \mathcal{C}, \Omega \text{ is open set.}$

It is assumed that conditions of existence and uniqueness theorem for the (9) hold.

When s-stage Two-Step General Linear Method for RFDEs (TSGLM) with coefficients $(a_{ij}(\cdot), b_j(\cdot), c_i, \tilde{a}_{ij}(\cdot), \tilde{b}_j(\cdot), u_i(\cdot), v(\cdot))_{i,j=1,\dots,s}$ is applied with stepsize h to (9) for the computation of the solution x(t), it yields, as an approximation on [-r, h] of the shift function $y := x(\sigma + \cdot)$, the function

$$\eta(\alpha h) = (1 - v(\alpha))\overline{\eta}(0) + v(\alpha)\overline{\eta}(h) + h\sum_{j=1}^{s}\widetilde{b_j}(\alpha)\overline{K_j} + h\sum_{j=1}^{s}b_j(\alpha)K_j, \qquad \alpha \in [0, 1],$$
(10)
$$\eta(\theta) = \overline{\eta}_h(\theta), \qquad \qquad \theta \in [-r, 0],$$

where

function $\overline{\eta}_h \approx y$ on [-r - h, 0] and stage values \overline{K}_i , are available as approximations computed in the previous step,

$$K_i = f(\sigma + c_i h, Y^i_{c_i h}),$$
 $i = 1, \dots, s,$ (11)

and $Y^i: [-r, c_i h] \longrightarrow \mathbb{R}^d$ is a stage function given by

$$Y^{i}(\alpha h) = (1 - u_{i}(\alpha))\overline{\eta}(0) + u_{i}(\alpha)\overline{\eta}(h) + h\sum_{j=1}^{s} \widetilde{a}_{ij}(\alpha)\overline{K}_{j} + h\sum_{j=1}^{s} a_{ij}(\alpha)K_{j}, \quad \alpha \in [0, c_{i}],$$

$$Y^{i}(\theta) = \overline{\eta}_{h}(\theta). \qquad \qquad \theta \in [-r, 0],$$

$$(12)$$

It is assumed that coefficients $(a_{ij}(\cdot), b_j(\cdot), c_i, \tilde{a}_{ij}(\cdot), \tilde{b}_j(\cdot), u_i(\cdot), v(\cdot))_{i,j=1,\dots,s}$ of TSGLMs satisfy the following conditions:

• $a_{ij}(\cdot), \ \widetilde{a}_{ij}(\cdot), \ u_i(\cdot), \quad \text{are polynomial functions } [0, c_i] \longrightarrow \mathbb{R}, \quad i, \ j = 1, \dots, s.$ (13) $b_j(\cdot), \ \widetilde{b}_j(\cdot), \ v(\cdot), \quad \text{are polynomial functions } [0, 1] \longrightarrow \mathbb{R}, \quad j = 1, \dots, s.$

•
$$c_i \in \mathbb{R}, \ c_i \ge 0,$$
 $i = 1, \dots, s.$ (14)

•
$$a_{ij}(0) = \widetilde{a}_{ij}(0) = 0, \quad u_i(0) = 1,$$
 (15)

•
$$b_j(0) = b_j(0) = 0, \quad v(0) = 1,$$
 (16)

The last two conditions correspondingly gurantee continuity of the stage functions $Y_{c_ih}^i \in \mathcal{C}$ and the approximate solution $\eta_h \in \mathcal{C}$ provided that approximate solution computed in the previous step is continuous function $\overline{\eta}_h \in \mathcal{C}$.

Remark 3.1. If the conditions

$$u_{i}(\cdot) = 1, \quad \tilde{a}_{ij}(\cdot) = 0, \quad i, \ j = 1, \dots, s,$$

$$v(\cdot) = 1, \quad \tilde{b}_{j}(\cdot) = 0, \qquad j = 1, \dots, s,$$
(17)

hold, the two-step method (8) becomes the one-step RK method for RFDEs introduced in [1], where initial function $\phi := \overline{\eta}_h$.

Definition 3.2. TSGLM with coefficients $(a_{ij}(\cdot), b_j(\cdot), c_i, \tilde{a}_{ij}(\cdot), \tilde{b}_j(\cdot), u_i(\cdot), v(\cdot))_{i,j=1,\dots,s}$ is called explicit if $a_{ij}(\cdot) = 0$ for all $j: j \ge i, i, j = 1,\dots,s$.

Definition 3.3. The function $E = \eta - y$: $[0, h] \longrightarrow \mathbb{R}^d$ computed under the assumption that $\overline{\eta}_h = y$ on [-r - h, 0] is called the local error of TSGLM (8).

Definition 3.4. The function $E^i = Y^i - y$: $[0, c_i h] \longrightarrow \mathbb{R}^d$ computed under the assumption that $\overline{\eta}_h = y$ on [-r - h, 0] is called the local stage error of TSGLM (8).

By analogy with [7] we define **stage order** for RFDEs.

Definition 3.5. Let functions $\widetilde{E}^i = Y^i - y$: $[0, c_i h] \longrightarrow \mathbb{R}^d$, $i = 1, \ldots, s$, $\widetilde{E}^{s+1} = \eta - y$: $[0, h] \longrightarrow \mathbb{R}^d$ are computed under the assumption that $\overline{\eta}_h = y$ on [-r - h, 0], $\overline{K}_j = y'(-h + c_j h)$, $K_j = y'(c_j h)$, $j = 1, \ldots, s$, that is

$$\widetilde{E}^{i}(\alpha h) = (1 - u_{i}(\alpha))y(-h) + u_{i}(\alpha)y(0) + h\sum_{j=1}^{s} \widetilde{a}_{ij}(\alpha)y'(-h + c_{j}h) + h\sum_{j=1}^{s} a_{ij}(\alpha)y'(c_{j}h) - y(\alpha h), \qquad \alpha \in [0, c_{i}], \qquad i = 1, \dots, s, \qquad (18)$$

$$\widetilde{E}^{s+1}(\alpha h) = (1 - v(\alpha))y(-h) + v(\alpha)y(0) + h\sum_{j=1}^{s} \widetilde{b}_{j}(\alpha)y'(-h + c_{j}h) + h\sum_{j=1}^{s} b_{j}(\alpha)y'(c_{j}h) - y(\alpha h), \qquad \alpha \in [0, 1].$$

Denote $c_{s+1} := 1$. If there are positive integers \widetilde{p}_i and reals $D_i > 0$, H > 0 such that

$$\max_{\alpha \in [0,c_i]} |\widetilde{E}^i(\alpha h)| \le D_i h^{\widetilde{p}_i + 1}, \qquad h \in [0,H], \quad i = 1, \dots, s + 1,$$
(19)

then positive integer $\tilde{p} = \min\{\tilde{p}_1, \ldots, \tilde{p}_{s+1}\}$ is called uniform stage order of TSGLM (8).

4 Order conditions

Assume that f is of class C^l with respect to the second argument for a sufficiently large l and solution x(t) of (9) is of piecewise class C^m for a sufficiently large m.

We introduce the polynomial functions $\Gamma_k : [0,1] \longrightarrow \mathbb{R}$ and $\Gamma_{ik} : [0,c_i] \longrightarrow \mathbb{R}$ given by

$$\Gamma_{k}(\alpha) = \frac{1}{(k-1)!} \left[\frac{(1-v(\alpha))(-1)^{k}}{k} + \sum_{j=1}^{s} \widetilde{b}_{j}(\alpha)(-(1-c_{j}))^{k-1} + \sum_{j=1}^{s} b_{j}(\alpha)c_{j}^{k-1} - \frac{\alpha^{k}}{k} \right],$$

$$\alpha \in [0,1],$$
(20)

$$\Gamma_{ik}(\alpha) = \frac{1}{(k-1)!} \left[\frac{(1-u_i(\alpha))(-1)^k}{k} + \sum_{j=1}^s \widetilde{a}_{ij}(\alpha)(-(1-c_j))^{k-1} + \sum_{j=1}^s a_{ij}(\alpha)c_j^{k-1} - \frac{\alpha^k}{k} \right],$$

$$\alpha \in [0, c_i], \qquad i = 1, \dots, s.$$

Remark 4.1. If the conditions (17) hold the Γ_{ik} , Γ_k are the same as for the one-step RK method [1].

Let $c_1^*, \ldots, c_{s^*}^*$ such that $c_1^* < c_2^* < \cdots < c_{s^*}^*$ and $\{c_1^*, \ldots, c_{s^*}^*\} = \{c_1, \ldots, c_s\}$, i.e. c_i^* are distinct c_i in increasing order.

Lemma 4.2. Let p be a positive integer. If x is of piecewise class C^{p+1} and the local stage errors computed in the previous step are $\overline{E}^i = O(h^p)$, i = 1, ..., s, then the local error E and the local stage errors E^i satisfy

$$E(\alpha h) = h \sum_{j=1}^{s} b_j(\alpha) D_j + \sum_{k=1}^{p} y^{(k)}(0) h^k \Gamma_k(\alpha) + O(h^{p+1}), \quad \alpha \in [0, 1],$$
(21)

$$E^{i}(\alpha h) = h \sum_{j=1}^{s} a_{ij}(\alpha) D_{j} + \sum_{k=1}^{p} y^{(k)}(0) h^{k} \Gamma_{ik}(\alpha) + O(h^{p+1}), \quad \alpha \in [0, c_{i}], \quad i = 1, \dots, s,$$
(22)

where

$$D_{i} = f(\sigma + c_{i}h, y_{c_{i}h} + E_{c_{i}h}^{i}) - f(\sigma + c_{i}h, y_{c_{i}h}), \qquad i = 1, \dots, s.$$
(23)

The Lemma 4.2 has been proved, but its proof is omitted here for brevity.

In the following we assume that the TSGLM satisfies the conditions $\Gamma_1 = 0$ and $\Gamma_{i1} = 0, \ i = 1, \ldots, s$, that is

$$v(\alpha) - 1 + \sum_{j=1}^{s} \widetilde{b}_j(\alpha) + \sum_{j=1}^{s} b_j(\alpha) = \alpha, \qquad \alpha \in [0, 1],$$
(24)

$$u_i(\alpha) - 1 + \sum_{j=1}^s \tilde{a}_{ij}(\alpha) + \sum_{j=1}^s a_{ij}(\alpha) = \alpha, \qquad \alpha \in [0, c_i], \quad i = 1, \dots, s.$$

The above condition is an equivalent form of uniform stage order **one** condition.

Theorem 4.3. A TSGLM satisfying (24) has uniform order two iff $\Gamma_2 = 0$.

The theorem 4.3 has been proved, but its proof is omitted here for brevity.

Theorem 4.4. Let TSGLM satisfy (24) and has uniform order two. If $\Gamma_3 = 0$ and $\sum_{\substack{i=1\\c_i=c_m^*}}^{s} b_i(\alpha)\Gamma_{i2}(\beta) = 0$, $\alpha \in [0,1]$, $\beta \in [0,c_m^*]$, $m = 1,\ldots,s^*$ then the method has uniform order three.

The theorem 4.4 has been proved, but its proof is omitted here for brevity.

Theorem 4.5. Let TSGLM satisfy (24) and has uniform order three.

$$If \ \Gamma_4 = 0, \\ \sum_{\substack{i=1\\c_i = c_m^*}}^s b_i(\alpha) \Gamma_{i3}(\beta) = 0, \qquad \alpha \in [0, 1], \quad \beta \in [0, c_m^*], \qquad m = 1, \dots, s^*,$$
(25)

$$\sum_{\substack{i=1\\c_i=c_m^*}}^{s} \sum_{\substack{j=1\\c_j=c_l^*}}^{s} b_i(\alpha) a_{ij}(\beta) \Gamma_{j2}(\gamma) = 0, \quad \alpha \in [0,1], \quad \beta \in [0,c_m^*], \quad \gamma \in [0,c_l^*], \quad l,m = 1,\dots,s^*.$$

then the method has uniform order four.

The theorem 4.5 has been proved, but its proof is omitted here for brevity.

Theorem 4.6. TSGLM has uniform stage order \tilde{p} iff $\Gamma_{ik} = 0, \ \Gamma_k = 0, \ i = 1, \dots, s, \ k = 1, \dots, \tilde{p}.$

Proof. Follows by Taylor series expansion of functions \widetilde{E}^i given by (18).

The following results can be obtained as corollary of theorems (4.6) and (4.4), (4.5).

Corollary 4.7. Let TSGLM has uniform stage order two. If $\Gamma_3 = 0$ then the method has uniform order three.

Corollary 4.8. Let TSGLM has uniform stage order three. If $\Gamma_4 = 0$ then the method has uniform order four.

The results of Corollary (4.7) and (4.8) can be easily generalized as follows.

Theorem 4.9. Let TSGLM has uniform stage order \tilde{p} . It has uniform order $p = \tilde{p} + 1$ iff $\Gamma_{\tilde{p}+1} = 0$.

The theorem 4.9 has been proved, but its proof is omitted here for brevity.

5 Construction of explicit two-stage GLMs of uniform stage order four and five

Consider two-stage explicit TSGLM satisfying (24). It's Butcher tableau is

TABLE 5.1. Butcher tableau for 2-stage explicit TSGLMs

c_1	$u_1(\alpha)$	$\widetilde{a}_{11}(\alpha)$	$\widetilde{a}_{12}(\alpha)$	0	0
c_2	$u_2(\alpha)$	$\widetilde{a}_{21}(\alpha)$	$\widetilde{a}_{22}(\alpha)$	$a_{21}(\alpha)$	0
	$v(\alpha)$	$\widetilde{b}_1(lpha)$	$\widetilde{b}_2(lpha)$	$b_1(\alpha)$	$b_2(\alpha)$

A natural choice will be to space out the abscissae c_i , i = 1, ..., s uniformly in the interval [0, 1] so that [6] $c_1 = 0$, $c_2 = \frac{1}{s-1}, ..., c_{s-1} = \frac{s-2}{s-1}, c_s = 1$. In the case of s = 2 we have $c_s = 0$, $c_s = 1$.

have $c_1 = 0, c_2 = 1$.

Since $c_1 = 0$, conditions $\Gamma_{1k}(\alpha) = 0$, $\alpha \in [0, c_1]$, $k = 1, 2, \ldots$ reduce to $\Gamma_{1k}(0) = 0$, $k = 1, 2, \ldots$ that follows from (15). It also follows that $u_1(\cdot) = 1$, $\tilde{a}_{11}(\cdot) = 0$, $\tilde{a}_{12}(\cdot) = 0$.

For brevity we omit the argument α of the method coefficient functions. By theorem (4.9), the method has uniform order four and uniform stage order three if $\Gamma_k = 0$, k = 1, 2, 3, 4 and $\Gamma_{2k} = 0$, k = 1, 2, 3, that is

$$-(1-v) + \tilde{b}_{1} + \tilde{b}_{2} + b_{1} + b_{2} = \alpha,$$

$$\frac{1-v}{2} - \tilde{b}_{1} + b_{2} = \frac{\alpha^{2}}{2}$$

$$-\frac{1-v}{3} + \tilde{b}_{1} + b_{2} = \frac{\alpha^{3}}{3}$$

$$\frac{1-v}{4} - \tilde{b}_{1} + b_{2} = \frac{\alpha^{4}}{4}$$

$$-(1-u_{2}) + \tilde{a}_{21} + \tilde{a}_{22} + a_{21} = \alpha,$$

$$\frac{1-u_{2}}{2} - \tilde{a}_{21} = \frac{\alpha^{2}}{2}$$

$$-\frac{1-u_{2}}{3} + \tilde{a}_{21} = \frac{\alpha^{3}}{3}$$

(26)

where $\alpha \in [0, 1]$.

The coefficients are defined by

$$u_{2} = -(2 \alpha - 1) (\alpha + 1)^{2},$$

$$v = (\alpha - 1)^{2} (\alpha + 1)^{2},$$

$$\tilde{a}_{2,1} = \alpha^{2} (\alpha + 1),$$

$$\tilde{b}_{1} = -\frac{1}{12} \alpha^{2} (\alpha + 1) (5 \alpha - 7),$$

$$a_{2,1} = \alpha (\alpha + 1)^{2} - \tilde{a}_{2,2},$$

$$b_{1} = -\frac{1}{3} \alpha (2 \alpha - 3) (\alpha + 1)^{2} - \tilde{b}_{2},$$

$$b_{2} = \frac{1}{12} \alpha^{2} (\alpha + 1)^{2},$$

(27)

where $\tilde{a}_{2,2}$, \tilde{b}_2 remain free. The relation $\Gamma_5(1) = \frac{4}{15} \neq 0$ implies that it is impossible to attain discrete order five.

The uniform order and the uniform stage order can be increased by finding a suitable value for c_2 . Assume that $c_1 = 0$, $c_2 \neq 0$ (in general case $c_2 \neq 1$). By theorem (4.9), the method has uniform order five and uniform stage order four if

 $\Gamma_k = 0, \quad k = 1, 2, 3, 4, 5 \text{ and } \Gamma_{2k} = 0, \quad k = 1, 2, 3, 4, \text{ that is}$

$$-(1-v) + \tilde{b}_{1} + \tilde{b}_{2} + b_{1} + b_{2} = \alpha,$$

$$\frac{1-v}{2} - \tilde{b}_{1} - (1-c_{2})\tilde{b}_{2} + c_{2}b_{2} = \frac{\alpha^{2}}{2},$$

$$-\frac{1-v}{3} + \tilde{b}_{1} + (1-c_{2})^{2}\tilde{b}_{2} + c_{2}^{2}b_{2} = \frac{\alpha^{3}}{3},$$

$$\frac{1-v}{4} - \tilde{b}_{1} - (1-c_{2})^{3}\tilde{b}_{2} + c_{2}^{3}b_{2} = \frac{\alpha^{4}}{4},$$

$$-\frac{1-v}{5} + \tilde{b}_{1} + (1-c_{2})^{4}\tilde{b}_{2} + c_{2}^{4}b_{2} = \frac{\alpha^{5}}{5},$$

$$-(1-u_{2}) + \tilde{a}_{21} + \tilde{a}_{22} + a_{21} = \alpha,$$

$$\frac{1-u_{2}}{2} - \tilde{a}_{21} - (1-c_{2})\tilde{a}_{22} = \frac{\alpha^{2}}{2},$$

$$-\frac{1-u_{2}}{3} + \tilde{a}_{21} + (1-c_{2})^{2}\tilde{a}_{22} = \frac{\alpha^{3}}{3},$$

$$\frac{1-u_{2}}{4} - \tilde{a}_{21} - (1-c_{2})^{3}\tilde{a}_{22} = \frac{\alpha^{4}}{4},$$
(28)

where $\alpha \in [0, 1]$ in the first five equations (28) and $\alpha \in [0, c_2]$ in other ones.

The coefficients are defined by

$$\begin{aligned} u_2 &= (\alpha+1)^2 \left(1-2\alpha + \frac{3\alpha^2}{2c_2-1}\right), \\ v &= -\frac{(\alpha+1)^2 \left((10\alpha-5)c_2^2 - 15c_2\alpha^2 + (\alpha+1)\left(6\alpha^2 - 3\alpha + 1\right)\right)}{5c_2^2 - 1}, \\ \tilde{a}_{21} &= \alpha^2 \left(\alpha+1\right) - \frac{\alpha^2 \left(\alpha+1\right)^2 \left(3c_2 - 1\right)}{2c_2 \left(2c_2 - 1\right)}, \\ \tilde{a}_{22} &= \frac{\alpha^2 \left(\alpha+1\right)^2}{2c_2 \left(c_2 - 1\right) \left(2c_2 - 1\right)}, \\ \tilde{b}_1 &= \frac{\alpha^2 \left(\alpha+1\right) \left(20c_2^4 - \left(30\alpha + 10\right)c_2^3 + \left(12\alpha^2 + 3\alpha - 13\right)c_2^2 + \left(4\alpha^2 + 11\alpha + 3\right)c_2 - 2\alpha \left(\alpha + 1\right)\right)}{4c_2 \left(5c_2^2 - 1\right) \left(c_2 + 1\right)}, \\ \tilde{b}_2 &= \frac{\alpha^2 \left(\alpha+1\right)^2 \left(5c_2^2 - \left(4\alpha - 3\right)c_2 - 2\alpha\right)}{4c_2 \left(5c_2^2 - 1\right) \left(c_2 - 1\right)} \end{aligned}$$
(29)
$$a_{21} &= \alpha \left(\alpha+1\right)^2 \left(1 - \frac{\alpha \left(3c_2 - 2\right)}{2 \left(2c_2 - 1\right) \left(c_2 - 1\right)}\right), \\ b_1 &= \frac{\alpha \left(\alpha+1\right)^2 \left(20c_2^4 - \left(30\alpha + 20\right)c_2^3 + \left(12\alpha^2 + 21\alpha - 4\right)c_2^2 + \left(-4\alpha^2 + 3\alpha + 4\right)c_2 - 2\alpha \left(\alpha + 1\right)\right)}{4c_2 \left(5c_2^2 - 1\right) \left(c_2 - 1\right)} \\ b_2 &= -\frac{\alpha^2 \left(\alpha+1\right)^2 \left(5c_2^2 - \left(4\alpha + 7\right)c_2 + 2\alpha + 2\right)}{4c_2 \left(5c_2^2 - 1\right) \left(c_2 + 1\right)}. \end{aligned}$$

To attain the discrete stage order five, we determine c_2 from $\Gamma_{25}(1) = 0$. We have

$$c_2 = \frac{11 - \sqrt{41}}{10}.\tag{30}$$

The relation $\Gamma_6(1) = -\frac{16(17 - 2\sqrt{41})}{75(71 - 11\sqrt{41})} \neq 0$ implies that it is impossible to attain discrete order six.

So we construct explicit TSGLM of uniform order five, uniform stage order four and discrete stage order five.

Remark 5.1. There is not a method of uniform stage order two in a class of explicit onestep RK methods for RFDEs. Indeed, for explicit one-step RK methods $c_1 = 0$, $c_2 \neq 0$ and $a_{2,j} = 0$, j = 2, ..., s, hence $\Gamma_{2k} = -\frac{\alpha^k}{k!} \neq 0$, $\alpha \in (0, c_2]$, k = 2, 3,

It is known [7] that methods with low stage order suffer from the order reduction phenomenon when applied to stiff ODEs. Hence, explicit TSGLMs may be more appropriate for some mildly stiff RFDEs (of course, if the smoothness conditions in Remark 2.1 and in the beginning of section 4 hold).

References

- S. Maset, L. Torelli, R. Vermiglio Runge-Kutta methods for Retarded Functional Differential Equations Mathematical Models and Methods in Applied Sciences, 8 (2005), 1203-1251.
- [2] L. Tavernini, One-step methods for the numerical solution of Volterra functional differential equations, SIAM J. Numer. Anal., 8 (1971), 786-795.
- [3] C. Cryer and L. Tavernini, The numerical solution of Volterra functional differential equations by Euler's method, *SIAM J. Numer. Anal.*, **9** (1972), 105–129.
- [4] A. Bellen and M. Zennaro, Numerical Methods for Delay Differential Equations, Oxford Univ. Press 2003.
- [5] J.C. Butcher, Numerical Methods for Ordinary Differential Equations, John Wiley&Sons, Chichester 2003.
- [6] J.C. Butcher and Z. Jackiewicz Construction of diagonally implicit general linear methods of type 1 and 2 for ordinary differential equations, *Applied Numerical Mathematics*, 21 (1996), 385-415.
- [7] K. Dekker and J.G. Verwer, Stability of Runge-Kutta methods for stiff nonlinear differential equations, North-Holland, Amsterdam-New-York-Oxford 1984.