Dark states of dressed spinor Bose-Einstein condensates of spin-1 atoms

Ma Luo, Chengguang Bao, Zhibing Li*

State Key Laboratory of Optoelectronic Materials and Technologies School of Physics and Engineering

Sun Yat-Sen University, Guangzhou, 510275, P.R. China

Dressed states of spinor Bose-Einstein condensates of spin-1 atoms located in optical cavity of near resonance frequency of excited state of the atom are investigated. The wave function of dark states are given.

PACS numbers: 03.75. Kk, 03.65. Fd

Implementation of optical trap of spin-1 to spin-3 bosonic atoms, which librate the freedom of spin of the atoms, has given a great chance to investigate more prolific physics of cold atomic cluster [1, 2, 3, 4, 5]. When temperature is low, spinor Bose-Einstein condensate (spinor BEC) is form, whose eigenstate conserve the total spin and its Z-component [6]. If the system is coupled to a high Q-factor optical cavity mode, the eigenstate will be modified. This is a system of cavity quantum electrodynamics (cavity QED) with a spinor BEC.

Cavity QED investigate the interaction between atom and optical cavity mode [7, 8]. In the parameter region that the interaction strength between atom and optical mode is much larger than decay rate of cavity mode and excited state of the atom, the system is in strong coupling region. In this region, energy transfer between atom and cavity mode by absorbing and emitting photon periodically. The eigen state of the coupling system form dressed state described by Jaynes-Cummings model [9]. When there are more than one atom, the coupling effect will be enhanced. Several previous research has investigated cavity QED with a BEC theoretically and experimentally [10, 11, 12, 13, 14, 15, 16]. Dressed state of BEC is confirmed [15]. For a multicomponent condensates, it is shown that dark states of dressed BEC exist [16]. When the dressed BEC is in dark state, no atom is state at excited state, but oscillate between two ground states of Λ type energy level atom. The research of dressed BEC with has potential in understanding of quantum state [17] and quantum information.

This paper consider the dark states of dressed spinor BEC of spin-1 atoms. At first, coupling Hamiltonian between spinor BEC and optical cavity mode is given. And then, we calculate the matrix elements of the coupling Hamiltonian, where we make use of fractional parentage coefficients (FPCs) to get a simple analytic expression. At last, wave function of the dark states is deduced and value scope of quantum number for the dark states is fix.

The system we are going to investigate consist of $N^{87}Rb$ atoms optically trapped to form a spinor BEC, located inside a high-Q optical cavity. In the strong cou-

pling region, the photon decay rate of the cavity and the non-resonant decay rate of the atoms is neglected. The optical trap can be modeled by a harmonic trap potential. The frequency of the cavity mode is near resonance to the transition frequency of D_2 line of ⁸⁷Rb : $5^2S_{1/2}, F = 1 \longrightarrow 5^2P_{3/2}, F = 0$, so that only the two levels is considered in the dressed states. The cavity consist of two parallel high reflection mirror, which has two optical modes at the resonance frequency: one mode is left circular polarization, the other mode is right circular polarization. The two modes have opposite angular momentum along the axis of the cavity. The Hamiltonian of the atoms and cavity system can be written, respective, as:

$$H_A = \sum_{i=1}^{N} \left(\frac{\mathbf{p}_i^2}{2m} + U(\mathbf{r}_i) + \hbar\omega_e |e_i\rangle\langle e_i|\right) + \sum_{i< j} V(\mathbf{r}_i - \mathbf{r}_j) \quad (1)$$

$$H_C = \sum_{\mu=\pm 1} \hbar \omega_c \hat{a}^+_\mu \hat{a}_\mu \tag{2}$$

where \mathbf{p}_i is momentum of the ith atom, m is mass of the atoms, $U(\mathbf{r})$ is harmonic potential of optical trap, $|e_i\rangle$ is the excited state $5^2 P_{3/2}, F = 0$ of the ith atom, $\hbar\omega_e$ is energy level of the excited state, $V(\mathbf{r}_i - \mathbf{r}_i)$ is interaction between two atom that depend on coupling of their spins, and $\hbar\omega_c$ is energy level of cavity mode, \hat{a}_{μ} is annihilate operator of standing wave cavity mode of polarization $\mu = \pm$. $V(\mathbf{r}_i - \mathbf{r}_j)$ can be given as $V_{gg}(\mathbf{r}_i - \mathbf{r}_j)$ $\mathbf{r}_{j}||g_{i}g_{j}\rangle\langle g_{i}g_{j}| + V_{ge}(\mathbf{r}_{i} - \mathbf{r}_{j})(|g_{i}e_{j}\rangle\langle g_{i}e_{j}| + |e_{i}g_{j}\rangle\langle e_{i}g_{j}|),$ where $|g_i\rangle$ is ground state of the ith atom. V_{gg} is spin dependent interaction between two ground state atoms with spin-1, which is $(c_0 + c_2 \mathbf{F}_i \cdot \mathbf{F}_i) \delta(\mathbf{r}_i - \mathbf{r}_i)$, \mathbf{F}_i is spin operator of the ith atom, $c_0(c_2)$ is spin independent (dependent) interaction constant [1]. For V_{qe} , one of the atom is spin-0, the scattering has only one channel, so that the interaction is spin independent, i.e. $V_{ge} = c_4 \delta(\mathbf{r}_i - \mathbf{r}_j)$ where c_4 is a constant.

The direction of quantize axis of the atoms is along the axis of the cavity, so that by absorbing a left (right) circular polarization photon from the cavity, only the σ^+ (σ^-) transition that conserve the angular momentum could occur; the same conservation law of angular momentum apply for emission one photon to the cavity. Therefore,

^{*}Corresponding author: stslzb@mail.sysu.edu.cn

the Hamiltonian of coupling between atoms and cavity mode is,

$$H_{AC} = \sum_{i=1}^{N} [\hbar \Omega_{+} | e_{i} \rangle \langle g_{i}, -1 | \hat{a}_{+} + \hbar \Omega_{-} | e_{i} \rangle \langle g_{i}, +1 | \hat{a}_{-}] + H.c.$$

$$(3)$$

where $\hbar\Omega_{\pm} = d_{\pm}\sqrt{\hbar\omega_c/2\epsilon_0 V}$ is coupling strength of σ^{\pm} transition, d_{\pm} is the corresponding atomic dipole-matrix element that satisfy $d_+ = -d_-$, V is mode volume of the cavity mode, $|g_i, \pm 1\rangle$ is the ground state of atom with spin component ± 1 .

The eigenstate of non-coupling Hamiltonian $H_A + H_C$ can be labeled as $|g, S, M; N_1, N_2\rangle$ and $|e, S', M'; N_1, N_2\rangle$, where g means all N atoms are at ground state, e means one of the atoms is at excited state, S and M is total spin and the Z component, N_1 (N_2) is photon number of $\mu = + (\mu = -)$ polarization photon. For the state that all N atoms are at ground state, N-S is limited to be even, and the eigen energy is approximated to be $E_{g,S}^{N_1,N_2} = J_g S(S+1) + \hbar \omega_c (N_1+N_2)$ [6], where J_q is spinor interaction constant that depend on total spin and spatial wave function. When one atom is at excited state of spin-0, all the other atoms couple to total spin S', with the limitation that N - 1 - S' is even. The atomic part of wave function of this state can be written as

$$|e, S', M'\rangle = \frac{1}{\sqrt{N}} \sum_{i} \{ [\varphi_e(\mathbf{r}_i) \prod_{j \neq i} \varphi_g(\mathbf{r}_j)] \vartheta_{S', M'}^{N-1}(\bar{i}) \} \quad (4)$$

where $\varphi_{e(g)}$ is spatial wave function of excited (ground) state atom, $\vartheta_{S',M'}^{N-1}(\bar{i})$ is total symmetric spin state of all the other atoms without the ith atom with quantum number S' and M'. Spin of the ith atom is zero, and it couple with spin state of the other N-1 atoms, forming total spin state S'. Using variation method to deduce Schrödinger equation for the spatial wave function $\varphi_{e(q)}$ and solving the equation by perturbation approximation, eigen energy of $|e, S', M'\rangle$ plus energy of the cavity mode is given as $E_{e,S'}^{N_1,N_2} = J_e S'(S'+1) + \hbar \omega_e + \hbar \omega_c (N_1 + N_2).$ J_e is spinor interaction constant. When N is small, J_q and J_e is approximated to be equated to each other, noted as J. J is much smaller than $\hbar\omega_c$ and $\hbar\omega_e$, so that for a fix set of quantum number $(g(e), N_1, N_2)$, the energy levels expand to be a narrow band by quantum number S(S') around energy level $(\hbar\omega_e) + \hbar\omega_c(N_1 + N_2)$. When there is no static magnetic field, the energy levels do not depend on M(M'). We consider the dressed states consist of states in the first excited band: $|q, S, M; 1, 0\rangle$, $|g, S, M; 0, 1\rangle$ and $|e, S', M'; 0, 0\rangle$. According to the conservation of angular momentum at quantize axis, the dressed states with total angular momentum along quantize axis $M\hbar$ can be written as

$$|M\rangle = \sum_{S'} c_e^{S',M}(t)|e, S', M; 0, 0\rangle + \sum_{S} c_+^{S,M-1}(t)|g, S, M-1; 1, 0\rangle + \sum_{S} c_-^{S,M+1}(t)|g, S, M+1; 0, 1\rangle$$
(5)

where the summation of $c_e^{S',M}$ run through S' that N - S' - 1 is even and S' >= M, the summation of $c_+^{S,M-1}(c_-^{S,M+1})$ run through S that N - S is even and S >= M - 1(S >= M + 1).

In order to get the equation of motion of probability amplitudes under the full Hamiltonian $H = H_A + H_C + H_{AC}$, we deduce the matrix element of the coupling part of Hamiltonian H_{AC} at the bases of eigen states of non-coupling Hamiltonian. We consider matrix element $\langle e, S', M'; 0, 0|H_{AC}|g, S, M; 1, 0 \rangle$ first. The atomic state $\langle e, S, M|$ is expanded as equation (4). The atomic spin state of $|g, S, M; 1, 0 \rangle$ is total symmetric spin state of Natoms, $\vartheta^N_{S,M}$. Making use of FPCs, the spin state of the ith atom can be extracted, so that the atomic state can be expanded as

$$|g, S, M\rangle = [\prod_{j=1}^{N} \varphi_g(\mathbf{r}_j)] \quad [a_S^N[\chi(i)\vartheta_{S+1,M}^{N-1}(\bar{i})]_S + b_S^N[\chi(i)\vartheta_{S-1,M}^{N-1}(\bar{i})]_S]$$
(6)

where $a_S^N = \{[1 + (-1)^{N-S}](N - S)(S + 1)/[2N(2S + 1)]\}^{1/2}$ and $b_S^N = \{[1 + (-1)^{N-S}]S(N + S + 1)/[2N(2S + 1)]\}^{1/2}$ is FPCs given in reference [18], $\chi(i)$ is spin state of the ith spin-1 ground state atom. The coupling between $\chi(i)$ and $\vartheta_{S+1,M}^{N-1}(i)$ can be expanded by using Clebsch-Gordan coefficients [19]. When we calculate the matrix element of operator of the ith atom in H_{AC} , we can extracted spin state of the ith atom and make use of the orthogonal relation of spin states to get the exact expression of the spin part of the element. Under approximation that the spatial wave functions are independent of total spin and internal state, the matrix element can be expressed as

$$\langle e, S', M'; 0, 0 | H_{AC} | g, S, M; 1, 0 \rangle$$

= $\sqrt{N} \hbar \Omega_{+} a_{S}^{N} C_{S',1,M',-1}^{SM} \delta_{S,S'-1} \delta_{M,M'-1}$
+ $\sqrt{N} \hbar \Omega_{+} b_{S}^{N} C_{S',1,M',-1}^{SM} \delta_{S,S'+1} \delta_{M,M'-1}$ (7)

where C_{S_1,S_2,M_1,M_2}^{SM} is Clebsch-Gordan coefficients. Using similar analysis, the matrix element with another photon state can be obtained as

$$\langle e, S', M'; 0, 0 | H_{AC} | g, S, M; 0, 1 \rangle$$

= $\sqrt{N} \hbar \Omega_{-} a_{S}^{N} C_{S',1,M',1}^{SM} \delta_{S,S'-1} \delta_{M,M'+1}$
+ $\sqrt{N} \hbar \Omega_{-} b_{S}^{N} C_{S',1,M',1}^{SM} \delta_{S,S'+1} \delta_{M,M'+1}$ (8)

Inserting the state into Schrodinger equation, and take the inner product with all bases in equation (4), the equation of motion of probability amplitudes are obtained as

$$i\hbar \frac{dc_e^{S',M}}{dt} = [\hbar\omega_e + JS'(S'+1)]c_e^{S',M} + \sqrt{N}\hbar\Omega_+ a_{S'-1}^N C_{S',1,M,-1}^{S'-1,M-1}c_+^{S'-1,M-1} + \sqrt{N}\hbar\Omega_+ b_{S'+1}^N C_{S',1,M,-1}^{S'+1,M-1}c_+^{S'+1,M-1} + \sqrt{N}\hbar\Omega_- a_{S'-1}^N C_{S',1,M,1}^{S'-1,M+1}c_-^{S'-1,M+1} + \sqrt{N}\hbar\Omega_- b_{S'+1}^N C_{S',1,M,1}^{S'+1,M+1}c_-^{S'+1,M+1}$$
(9)

where N - S' - 1 is limited to be even, and

$$i\hbar \frac{dc_{+}^{S,M-1}}{dt} = [\hbar\omega_{c} + JS(S+1)]c_{+}^{S,M-1} + \sqrt{N}\hbar\Omega_{+}a_{S}^{N}C_{S+1,1,M,-1}^{S,M-1}c_{e}^{S+1,M} + \sqrt{N}\hbar\Omega_{+}b_{S}^{N}C_{S-1,1,M,-1}^{S,M-1}c_{e}^{S-1,M}$$
(10)

$$i\hbar \frac{dc_{-}^{S,M+1}}{dt} = [\hbar\omega_c + JS(S+1)]c_{-}^{S,M+1} + \sqrt{N}\hbar\Omega_{-}a_S^N C_{S+1,1,M,1}^{S,M+1}c_e^{S+1,M} + \sqrt{N}\hbar\Omega_{-}b_S^N C_{S-1,1,M,1}^{S-1,M}c_e^{S-1,M}$$
(11)

where N-S is limited to be even. $M\hbar$ is total angular momentum along quantize axis. The exist of dark state require that the probability amplitude $c_e^{S',M}$ equate to zero. From equation (10) and (11), this will result in the total spin dependence of the time factor of $c_{\pm}^{S,M\mp 1}$. From equation (9), this will result in relations between different probability amplitude. Because the time factor of $c_{\pm}^{S,M\mp 1}$ depend on S as $e^{-iJS(S+1)t/\hbar}$, and $\Omega_{+} = -\Omega_{-}$, equation (9) reduce to two equation for dark states,

$$a_{S'-1}^{N}C_{S',1,M,-1}^{S'-1,M-1}c_{+}^{S'-1,M-1}-a_{S'-1}^{N}C_{S',1,M,1}^{S'-1,M+1}c_{-}^{S'-1,M+1} = 0$$
(12)
$$b_{S'+1}^{N}C_{S',1,M,-1}^{S'+1,M-1}c_{+}^{S'+1,M-1}-b_{S'+1}^{N}C_{S',1,M,1}^{S'+1,M+1}c_{-}^{S'+1,M+1} = 0$$
(13)

Thus far, it is easy to find that a dark state consist of two states with the same total angular momentum along quantize axis, $M\hbar$, and the same total spin S. The two states are $|g, S, M - 1; 1, 0\rangle$ and $|g, S, M + 1; 0, 1\rangle$, and their probability amplitude satisfy the equations

$$a_{S}^{N}C_{S+1,1,M,-1}^{S,M-1}c_{+}^{S,M-1} - a_{S}^{N}C_{S+1,1,M,1}^{S,M+1}c_{-}^{S,M+1} = 0$$

$$b_{S}^{N}C_{S-1,1,M,-1}^{S,M-1}c_{+}^{S,M-1} - b_{S}^{N}C_{S-1,1,M,1}^{S,M+1}c_{-}^{S,M+1} = 0 \quad (14)$$

Existence of solution for equation (14) is the determine of coefficient matrix equation to zero, which gives

$$a_{S}^{N}b_{S}^{N}(C_{S+1,1,M,-1}^{S,M+1}C_{S-1,1,M,1}^{S,M+1} - C_{S+1,1,M,1}^{S,M+1}C_{S-1,1,M,-1}^{S,M-1}) = a_{S}^{N}b_{S}^{N}\frac{2M(2S+1)}{\sqrt{4S(2S-1)(S+1)(2S+3)}} = 0$$
(15)

It is obvious that one kind solution is M = 0 and S > 0. When S = 0, M could not be ± 1 ; when S > 0 and M = 0, all pairs of $(|g, S, -1; 1, 0\rangle$, $|g, S, 1; 0, 1\rangle)$ form dark states with $c_{+}^{S,-1} = c_{-}^{S,1}$. Therefore, this kind of dark state is

$$|S, M = 0\rangle_d = \frac{1}{\sqrt{2}}(|g, S, -1; 1, 0\rangle + |g, S, 1; 0, 1\rangle) \quad (16)$$

Another kind of solution is $a_S^N = 0$ when S = N. In this case, there is not atomic excited state with total spin S = N + 1, so that only the second equation in (14) is required. Thus, when S = N, all pairs of $(|g, N, M - 1; 1, 0\rangle$, $|g, N, M + 1; 0, 1\rangle)$ form dark states with $\sqrt{(N - M)(N - M + 1)}c_+^{S=N,M-1} = \sqrt{(N + M)(N + M + 1)}c_-^{S=N,M+1}$. This kind of dark state is

$$|S = N, M\rangle_d = \frac{\sqrt{(N+M)(N+M+1)}}{\sqrt{2(M^2+N+N^2)}} |g, N, M-1; 1, 0\rangle_d + \frac{\sqrt{(N-M)(N-M+1)}}{\sqrt{2(M^2+N+N^2)}} |g, N, M+1; 0, 1\rangle$$
(17)

In equation (16) and (17), the time factor in include into the state itself, because it only depend on S.

In the above analysis, it is assumed that magnetic field is zero. However, it is impossible to have absolute zero magnetic field environment. Therefore, we consider how small the magnetic field is required to achieve dark states. Because the dark states consist of two states with atomic Z-component of angular momentum difference of $2\hbar$, the Zeeman split is 2gB, where B is magnetic field, g = -0.70 MHz/G is Zeeman split factor [20]. The energy difference between state with S + 1 and S - 1total spin is J(4S + 2). For a system with N = 1000atoms and $\omega_{trap} = 1000 Hz$ optical trap frequency, J is estimated to be 1Hz approximately. if Zeeman split is much smaller than energy split due to total spin, .i.e. $2gB \ll J(4S+2)$, the Zeeman split can be neglected and the dark state exist: otherwise, the dark states is damaged. It can be seen that magnetic field is require to be smaller than $10^{-6}G$ so as to mean the requirement of dark states. Since magnetic field on the surface of the earth is about 0.6G, it need very accurate experiment set up the counteract this magnetic field.

In conclusion, the Hamiltonian of dressed spinor BEC regardless of atomic and photon decay is given. The Hamiltonian is diagonalized under bases of eigenstates of non-coupling part of Hamiltonian of spinor BEC and optical cavity mode. Only the matrix elements that conserve total angular momentum along quantize axis is nonzero. As a result, the dressed state with total angular momentum along quantize axis $M\hbar$ consist of three kind of states in non-coupling bases: $|e, S', M; 0, 0\rangle$, $|g, S, M - 1; 1, 0\rangle$ and $|g, S, M + 1; 0, 1\rangle$. Dark state is a state that oscillate between two kind of states,

 $|g, S, M - 1; 1, 0\rangle$ and $|g, S, M + 1; 0, 1\rangle$, while is not excited to the states $|e, S, M; 0, 0\rangle$. It is found that there are two species of dark states, whose wave function is given by equation (16) and equation (17), respectively. It is obvious from the two wave functions that the dark state is entangle state of spinor BEC $|g, S, M \pm 1\rangle$ and optical cavity mode $|\mu = \pm\rangle$. This entanglement could be very useful to the theory and application of quantum information and quantum computer.

- [1] Tin-Lun Ho, Phys. Rev. Lett. 81, 742 (1998).
- [2] D. M. Stamper-Kurn, M. R. Andrews, A. P. Chikkatur, S. Inouye, H.-J. Miesner, J. Stenger, and W. Ketterle, Phys. Rev. Lett. 80, 2027 (1998).
- [3] J. Stenger, S. Inouye, D. M. Stamper-Kurn, H.-J. Miesner, A. P. Chikkatur, and W. Ketterle, Nature (London) 396, 345 (1998).
- [4] A. Gorlitz, T. L. Gustavson, A. E. Leanhardt, R. L?w, A. P. Chikkatur, S. Gupta, S. Inouye, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. 90, 090401 (2003).
- [5] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, Phys. Rev. Lett. 94, 160401 (2005).
- [6] C. G. Bao and Z. B. Li, Phys. Rev. A 70, 043620 (2004).
- [7] A. Boca, R. Miller, K. M. Birnbaum, A. D. Boozer, J. McKeever, and H. J. Kimble, Phys. Rev. Lett. 93, 233603 (2004)
- [8] P. Maunz, T. Puppe, I. Schuster, N. Syassen, P. W. H. Pinkse, and G. Rempe, Phys. Rev. Lett. 94, 033002 (2005)
- [9] E.T.Jaynes and F.W.Cummings, Proc. IEEE 51, 89(1963).
- [10] Ferdinand Brennecke, Tobias Donner, Stephan Ritter, Thomas Bourdel, Michael K??hl, Tilman Esslinger, Nature, 450, 268(2007).
- [11] Christoph Maschler and Helmut Ritsch, Phys. Rev. Lett. 95, 260401 (2005)
- [12] Igor B. Mekhov, Christoph Maschler, Helmut Ritsch, Nature Physics 3, 319 - 323 (01 May 2007)
- [13] Peter Horak, Stephen M. Barnett, and Helmut Ritsch, Phys. Rev. A 61, 033609 (2000)
- [14] Peter Horak and Helmut Ritsch, Phys. Rev. A 63, 023603 (2001)
- [15] Elena V. Goldstein, Ewan M. Wright, and P. Meystre, Phys. Rev. A 57, 1223(1998)
- [16] E. S. Lee, C. Geckeler, J. Heurich, A. Gupta, Kit-Iu Cheong, S. Secrest, and P. Meystre, Phys. Rev. A 60, 4006(1999)
- [17] J. Higbie and D. M. Stamper-Kurn, Phys. Rev. A 69, 053605 (2004).
- [18] C. G. Bao and Z. B. Li, Phys. Rev. A 72, 043614 (2005).
- [19] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, Quantum Theory of Angular Momentum (World Scientific, Singapore, 1988).
- [20] Hans Kleinpoppen, "Atoms" in Ludwig Bergmann and Clemens Schaefer, Constituents of Matter: Atoms, molecules, Nuclei, and Particles, Wilhelm Raith, Ed. (Walter de Gruyter, Berlin, 1997)

Acknowledgments

We appreciate the support from the NSFC under the grants 10574163 and 10674182.