# Dark Energy, Hyperbolic Cosecant Cardassian and Virial Collapse for Power-style Cardassian

Wen-Jie Tian<sup>1</sup>

College of Physics and Information Technology, Shaanxi Normal University, Xi'an, 710062, P.R.China

<sup>1</sup>tianwj1@gmail.com

# Contents

1	Intr	Introductions					
	1.1	Funda	amentals of Cosmology	6			
		1.1.1	the Cosmic Principle	6			
		1.1.2	Cosmos Expansion & Hubble's Law	7			
		1.1.3	Cosmic Microwave Background Radiation	7			
		1.1.4	Global Scenario	7			
		1.1.5	First three Minutesin Fine Cosmology	8			
		1.1.6	Dark Matter	10			
	1.2	Gener	al Relativity & Cosmic Dynamics	11			
		1.2.1	Gravitational Field Equation & FRW metric	11			
		1.2.2	Perfect Fluid Scenario & Conservation Laws	11			
		1.2.3	Friedmann Equation	12			
		1.2.4	EOS in Perfect Fluid Scenario	14			
<b>2</b>	Acc	elerati	ing Expansion of the Universe	16			
	2.1	Accele	erating Expansion and Dark Energy	16			
	2.2	Chara	cteristic Parameters of Cosmos	17			
	2.3	Mechanism for Accelerating Expansion					
		2.3.1	Scalar Fields	18			
		2.3.2	Modified Gravitational Field Equation/Friedmann Equation	19			
		2.3.3	Modern QFT approaches	19			
3	Qui	ntesse	nce & Phantom	<b>21</b>			
	3.1	Quint	essence Scalar Field	21			
		3.1.1	Basic Structure of Quintessence	21			
		3.1.2	Quintessence Potential	22			
		3.1.3	Scalar Fields' problems and Tracker Field	23			
		3.1.4	Typical discussions for Quintessence	24			
	3.2	Phantom Scalar Field					

		3.2.1	basic Structures of Phantom	25				
		3.2.2	Hubble Parameter & Phantom Cosmology	26				
4	Qui	uintom & Hessence						
	4.1	om Scalar Field	28					
		4.1.1	Scalar Field with $w$ crossing -1	28				
		4.1.2	Basic Structure of Double Field Quintom	29				
		4.1.3	Generalized Quintom	30				
	4.2	Hesser	nce Scalar Field	30				
		4.2.1	Basic Structure of Hessence	30				
		4.2.2	Hessence Reconstruction via Hubble Parameter	33				
		4.2.3	Autonomous Equations for Hessence	34				
<b>5</b>	k-es	k-essence & Chaplygin gas						
	5.1	k-esse	nce Mechanism	36				
	5.2	5.2 Chaplygin gas model						
		5.2.1	Special Chaplygin gas	39				
		5.2.2	Generalized Chaplygin gas	41				
		5.2.3	Generalized Chaplygin gas & Hessence	42				
		5.2.4	Chaplygin gas & Superstring QFT	43				
6	Hyj	Hyperbolic Cosecant Cardassian Cosmology 4						
	6.1	Introd	luction	44				
	6.2	Funda	amentals of Cardassian Cosmology	45				
		6.2.1	Fundamentals for Implicit Cardassian term	46				
		6.2.2	Fundamentals for Explicit Cardassian term	46				
		6.2.3	Phase Space, Autonomous Equations for Separate Matter & Ra-					
			diation	47				
		6.2.4	Unitary Treatment of Matter & Radiation	48				
		6.2.5	Constraints for Cardassian & Hyperbolic cosecant Cardassian	49				
	6.3	3 Fundamentals of Hyperbolic Cosecant Function						
	6.4	4 Csch Cardassian for unitary matter and radiation						
	6.5	6.5 Csch Cardassian for separate matter and radiation						
		6.5.1	Dynamical Equations	53				
		6.5.2	Phase Space and Autonomous Equations	54				
		6.5.3	Critical Points	55				
		6.5.4	Stability Matrix for Autonomous System	57				

7	Viri	al Col	lapse of Power-style Cardassian	<b>59</b>
	7.1	Applic	cations of Virialization in Cosmology	59
	7.2	Viriali	zation Power-style Cardassian	60
		7.2.1	Virialization of Matter Term	61
		7.2.2	Virialization GF Fluid/Cardassian Term	61
		7.2.3	Unitary Virialization of Matter Term and GF Fluid Term $\ . \ . \ .$	62
	7.3	Conclu	sions and Evaluation	64

### Abstract

The problem of dark energy(DE) is the greatest challenge for modern physics, and therefore this thesis is dedicated to the modeling establishment of DE. In the first chapter, qualitative and descriptive methods are employed to draw a rough map for the cosmos in its finely analyzed era. Then the dynamical equations of cosmology based on the general relativity are introduced, together with the measuring methods and parameters. The second chapter reviews the results of cosmic microwave background radiation and supernova observations, which conclude the flat topology and accelerating expansion of cosmos, and give rise to existence of DE and the associated dynamical mechanism problem. Taking WMAP data as an example, we introduced the practical parameters. In the end we classify the existing DE models into three categories for further discussion. The first two chapters make up the basis for the whole thesis.

Chapter 3 reviews the fundamental structures and research progress of two scalar fields, Quintessence and Phantom, which correspond to the case of EOS beyond -1 and below -1 respectively. Chapter 4 reviews the fundamental structures and research progress of another two scalar fields, Phantom and Quintom with non-canonical lagrangian, both of which have EOS crossing -1. But Phantom field is real, while Quintom complex. Chapter 5 reviews two non-scalar-field DE formalism of K-essence and Chaplygin gas, which are both rooted in modern quantum field theory. Totally six kinds of DE is analyzed, all of which have positive energy density, negative pressure and could drive the lately accelerating expansion. And they are compared and identified in details.

New work gathers in Chapter 6 and Chapter 7. Firstly we make a comprehensive analysis of the existing work, pay particular attention to Gondolo and Freese's idea of treating Cardassian energy term as relativistic perfect fluid (GF fluid), point out that a potential Cardassian term should meet three conditions, and review three existing Cardassian terms (power style, polytropic style and its modification, exponential style and its modification), and eventually put forward the newly found hyperbolic cosecant Cardassian.

Then the Cardassian dynamical equations are introduced generally and logically under GF fluid scenario, together with the flowing process of constructing phase space and differential dynamical systems from Friedmann equation. Hyperbolic cosecant Cardassian term is employed for concrete computing. The analysis proceeds in two cases, namely a unified description of matter and radiation energy density (case 1) and a separate description of matter and radiation terms (case 2). Formalism of case 2 is more exact at the expense of more complicatedness, and due to the mathematical symmetry of matter term and radiation term in hyperbolic cosecant function, the differential dynamical equations are considerably simplified. Phase space and dynamical systems for both cases are achieved. When we calculate the critical points for case 2, amazingly interesting behaviors of self-consistency and auto-normalization are exhibited, which is a strong support for the new model, along with a forever positive sound speed.

The process of virial collapse in Cardassian cosmos is analyzed. Power-style Cardassian term is employed for its simplicity. Calculation declares that virial collapse of matter alone is forbidden. Yet Cardassian has excellent ability for virial collapse, after the virial collapse ending up with a stable sphere, the ratio of the ultimate radius to the original radius depends on the adjustable parameters in Cardassian term. And, the mixture of GF fluid and matter could conduct virial collapse, the ratio of the ultimate radius to the original radius depends on the adjustable parameters in Cardassian term, too. No singularity is generated.

The creative work in this thesis incudes the introduction of hyperbolic cosecant Cardassian, which is the fourth ever available Cardassian term ever found, after the introduction of exponential style in 2005. This work, along with the calculating of power Cardassian virialization, enriches the research of Cardassian cosmology.

**Key Words**: Friedmann Equation, Dark Energy, Scalar Field, Cardassian, Hyperbolic Cosecant Function, Virial Collapse

### Chapter 1

## Introductions

### **1.1** Fundamentals of Cosmology

Cosmology, which ranks one of the most encouraging realms in natural science, is the scientific branch researching into the cosmos structure, origin and evolution on the base of astronomic data and physics principles. Modern cosmology is basically supported by three events, namely the cosmic principle, the expansion of the universe together with Hubble Law, and the Cosmic Microwave Background Radiation.

### 1.1.1 the Cosmic Principle

Shortly afterwards the establishment of General Relativity, A. Einstein computed the global behaviors of the cosmos. To stay in harmony with Mach Principle and to simplify the calculating process, he made a basic assumption, which is nowadays honored as the Cosmos Principle. It declares that the universe is homogeneous and isotropic on large scales. Under modern considerations, the so-called large scale means covering a distance of  $10^8$  light year or more, which belongs to the cosmic scale. Homogeneous means different points identify with each other and could not be tell apart via physical experiments, while *isotropic* refers to the fact that , when taking observations into arbitrary directions at any coordinates origin, the scenarios are the same. A most significant application of this principle is to simplify the form of the space-time metric quantitatively. At the present days, the Cosmic Principle has already been established firmly on the base of observation data. As a matter of a typical sample, the results of the Willkinson Microwave Anisotropic Probe (WMAP) indicate that the universe is in high homogeneity and isotropy, only with tiny fluctuations at the magnitude of  $10^{-5} \sim 10^{-4}$ , which is in fact essentially required for the generation of galaxies according to the standard model of cosmology.

#### 1.1.2 Cosmos Expansion & Hubble's Law

In 1929, Hubble was engaged in comparing the light spectrum of far distant galaxies. To his surprise, he discovered a systematic redshift with the galaxies beyond our milkyway. As to *redshift*, it means the whole spectrum translates to the red wave interval. According to Doppler Effect, all galaxies are leaving us, and the spatial scale of the universe is undoubtedly expanding. After some algebra with the redshift amount and the distance to the galaxies, Hubble found the routing that the farther they are, the quicker they leaves, at a speed v proportional to the very distance r:

$$v = H_0 r \tag{1.1}$$

where the coefficient  $H_0$  is honored Hubble Constant, which is simply the present-day value of the more generalized dynamical Hubble parameter. Currently, the results mainly lie in the interval  $H_0 \approx 60 \sim 70 Km \cdot s^{-1} \cdot Mpc^{-1}$ .

### 1.1.3 Cosmic Microwave Background Radiation

The background space of galaxies is a radiation field that is highly homogeneous and isotropic, whose power spectrum curve agrees with that of the 2.7K blackbody perfectly. The so-called Cosmic Microwave Background Radiation (CMBR) is a key tool to understand the structure, origin and evolutionary history of the universe. It was researching into CMBR that lead to the era of finely quantitative analysis of cosmology.

CMBR could translate quite a lot of information on the young cosmos. For instance, the perfect homogeneity and isotropy suggest that the cosmos is also homogeneous and isotropic at its youth era, and the scattering surface is spherical, on which the gas are of identical temperature. The background photons are in thermal equilibrium before decoupling, thus the spectra, which describ how the radiation strength depending on the power at arbitrary direction, are all blackbody-like. The temperature fluctuations at present-day represent the medium density fluctuations at early time, which magnify via gravitation and form the clustering structure, or the inhomogeneity on the small scale. CMBR plays a pretty important role in the standard model of modern cosmology.

### 1.1.4 Global Scenario

The standard model of modern cosmology comprises two parts: One is the heat Big Bang theory, which was put forward by G. Gamow at around 1948, the other is the inflation theory, which is born in early 1980s when Guth,Linde, Albrecht and Steinhart employed this mechanism to beat the difficulties that Big Bang ran into at extremely early era.

The heat big bang theory is based on the observational evidences of Hubble Expansion and the Helium-4 amounts, and the theoretical formalism of General Relativity and solar heat nuclear reaction. It succeeds in explaining the genesis of light chemical elements, Hubble redshift and forecast the 5K CMBR (although the measured value being 2.7K, this prediction is still a greatest success of mankind intelligence). The big bang formalism accounts for the process from  $10^{-2}$  second (T ~  $10^{11}K$ ) to current time (~  $10^{17}s$ ,2.7K), but falls in trouble with the so-called singularity problem, the horizon problem, the asymmetry of baryonic numbers, the flat state problem and origin of galaxies, all of which are concerned with the extremely early era of the cosmos.

The inflation theory is based on general relativity, as well as the standard model for high energy particle physics established at that time. It aims at the time scale from  $10^{-2}$  second dating back to  $10^{-43}$  second (Plank time), and solved all the problems the big bang theory come across all at once, except for the singularity problem, which is principally impossible to overcome under the framework of classically general relativity. There seems no avoiding the singularity problem, and only the quantum cosmology theory declares to possess the power to solve the diagnose.

### 1.1.5 First three Minutesin Fine Cosmology

The time scale is generated together with the Creation of Big Bang. When time reaches  $10^{-43}$  second, the scale grows to  $10^{-33}cm$ , the four finds of elementary forces are unified as a grand force. Only energy lives, without any matter, and the world is dominated by quantum uncertainty principle. This era is called the Plank era or the Grand Unified era[3].

when it goes to  $10^{-36}s$ , temperature drops to  $10^{29}K$ , and certain kinds of particles are produced. Although some particles annihilate with their anti-particle companies, the creating rate is faster than the annihilate rate. Temperature slowing down, gravitation and the nuclear strong force decouple from the grand unified force one after one.

Then since  $10^{-35} \sim 10^{-33} s$ , it comes the inflation era. The spatial scale doubles every  $10^{-34}$  second. In a period of  $10^{-32} s$ , the scale expanded by  $10^{100}$  times, ending with a spatial scale of  $3 \times 10^{-25} cm$  and temperature of  $3 \times 10^{28} K$ . By the end of inflation, the electroweak force decouples into electromagnetic force and nuclear weak force.

When it comes to  $10^{-11}$ th second, the corresponding temperature is  $10^{15}K$ . The highest energy produced on the large hadron colliders is able to simulate this situation. And the validity of the numerical calculating and experimental simulating stands testing and win a wide agreement.

When it comes to  $10^{-6}$  second, the temperature decreases to  $10^{12}K$ , and "quarkgluon soup" is generated. Quarks and gluons are in free state originally. With the temperature continuously doing down, a gluon will bound three quarks together to form a nucleon (proton or neutron).

When it comes to 1st second, temperature drops to 10 billion Kelvin. The contains of the cosmos are photons, quarks, electrons, neutrinos and other particles, and of course, proton and neutrons. Temperature decreasing, neutrons begin to decay into steady proton, ending with a proton-neutron number ratio 7 : 1.

With time slipping down, the creation rate drops, and annihilation rate of particleantiparticle pairs exceed that of creation. Large amounts of electron-positrons annihilate to produce more photons. Eventually the annihilation era finished, the bulk being the remaining quarks and electrons (leptons). The cosmos evolves into a new creating era, and above all more protons and neutrons are created.

At about 100 seconds after the big bang, it gets 1 billion Kelvin. Protons and neutrons can never escape from the constraints of nuclear strong force and forms  $H_1^2$ . Then  $H_1^2$  turns to  $He_2^3$ , and  $He_2^3$  to  $He_2^4$ , with large quantities of neutrinos generated meanwhile.

Three minutes after the big bang, the light nuclear synthesis process ends due to low pressure. Now about 25% of protons and neutrons forms Helium, with a few  $H_1^2$ ,  $Li_3$ ,  $Be_4$ , and the remaining neutrons decay into proton. This results in a universe of 75% Hydrogen and 25% Helium.

The standard model of particle physics also believes that, in the extremely early era, some other particles are produced, such as x-boson, Higgs boson, Weakly Interacting heavy particle, cosmic string and monomagneton .But none of these are explored. At three minutes after the big bang, the universe is filled with electron, nude hydrogen, nude helium and dark matter, as well as high energy  $\gamma$ . Because photons, which is the medium particle of electromagnetic force, interact with free electrons and positrons via collision, absorption, emitting frequently, the lifetime and free-diatance of a photon is quite short, and information couldnot travel to the distance. So, the whole cosmos is in dark plasma state.

0.3 million years after the big bang, it has reached a temperature as low as 4000K. Electrons are bounded in atoms, so photons could travel to distant places to compare notes. The cosmos is apparent now, and it expands steadily and silently, and enriching clustering structures are formed. The temperature of background radiation goes on falling ,to the present-day value of 2.7K.

### 1.1.6 Dark Matter

Just before the discovery of dark energy, the so-called *dark matter* is already a remarkable challenge to man's brains. Although Zwicky had suggested the existence of dark matter, it is not until the late 1970 did the idea is highly treasured. Now, it is commonly accepted the existence of dark matter. Lots of experiments lead to this result, the classical representative of which comes from the measured rotating speed curve of spinning galaxies. With the hard efforts of high energy physists, conclusions are drawn that the dark matter, which has hardly any interactions with photons, couldnot be baryonic. Hence it is now generally called cold dark matter (hereafter CDM).

There are several candidates of elementary particles for CDM. The most hopeful one is the weakly interacting massive particles (WIMP), whose mass is probably far beyond a proton. Among several WIMPs, a particular kind is the neutralino, with a typical mass of 102GeV, or  $10^{-22}$  gram. Other candidates include the axion with a typical mass of  $10^{-6}$ eV or  $10^{-39}$  gram, fuzzy CDM, neutrinos, etc[2].

Nevertheless, CDM might also be ancient black wholes that form before nuclearsynthesis, rather than elementary particles. And the lower mass limit of the blackholes is Hawking evaporating limit, namely  $10^{15}$  gram.

### **1.2** General Relativity & Cosmic Dynamics

### 1.2.1 Gravitational Field Equation & FRW metric

The research into cosmic behaviors on large scales is based on Einstein's gravitational field equation[4] [5]:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G T_{\mu\nu}$$
(1.2)

where natural units are employed to set c = h = 1, and  $G_{\mu\nu}$  is Einstein tesor,  $R_{\mu\nu}$  being Ricci tensor, R being Ricci scalar, G being Newton's gravitational constant.

Take another consideration, as for the four-dimensional spacetime that obeys the Cosmological Principle and is therefore homogeneous and isotropic, the metric is Friedmann-Robertson-Walker(FRW) metric:

$$ds^{2} = dt^{2} - a^{2}(t) \left( \frac{dr^{2}}{1 - kr^{2}} + r^{2}d\theta^{2} + r^{2}\sin^{2}\theta d\phi^{2} \right)$$
(1.3)

where a(t) is the scale factor whose evolution is dependent on cosmic time t;  $r, \theta, \phi$  is the set of comoving coordinates fixed on the medium, the medium expanding with the universe while the comoving coordinates remaining static. k refers to the curvature scalar. When k > 0, the spatial topology is limited curved and namely closed universe; when k = 0, the spatial topology is infinitely flat and namely flat universe; when k < 0, the spatial topology is infinitely curved and namely open universe. Moreover, if the unit for r is properly selected, the values of k can be reduced to be k = +1, 0, -1correspondingly. Sometimes, FRW metric (1.3) is transformed to take the formalism below for computing convenience:

$$ds^{2} = dt^{2} - a^{2}(t) \left[ d\chi^{2} + f_{k}^{2}(\chi) (d\theta^{2} + \sin^{2}\theta d\phi^{2}) \right]$$
(1.4)

where

$$f_k(\chi) = \begin{cases} \sin \chi, & k = +1 \\ \chi, & k = 0 \\ \sinh \chi, & k = -1 \end{cases}$$
(1.5)

### 1.2.2 Perfect Fluid Scenario & Conservation Laws

Now let's combine the cosmological principle and the energy-momentum tensor which reflects the general states of matters, and get:

$$T_{\mu\nu} = (\rho + p)U_{\mu}U_{\nu} + pg_{\mu\nu}$$
(1.6)

where the energy density  $\rho$  and pressure p are dependent on t but independent of  $r, \theta, \phi$ , and  $U_{\mu}$  is the four-dimensional velocity vector:

$$U^t = 1 \tag{1.7}$$

$$U^i = 0 \tag{1.8}$$

Nevertheless, the energy-momentum tensor for 4D perfect fluid also takes the form of Eq.(1.6), where  $\rho$  and p are measured in an inertial reference comoving with the fluid. Hence perfect fluid scenarios are generally employed as dynamical mechanism in cosmology.

The spatial media(taken as perfect fluid) are pretty homogeneous and isotropic, which determines the diagonal matrix of its energy-momentum tensor and the identification of the spatial elements.

$$T^{\mu}_{\nu} = diag(-\rho(t), p(t), p(t), p(t)), \qquad (1.9)$$

The total energy-momenata ought to be conserving, hence there being the covariance equation:

$$T^{\mu\nu}_{;\nu} = 0$$
 (1.10)

as to  $\nu = r, \theta, \phi$  the above equation holds obviously; as for  $\nu = 0$ , one gets:

$$d[a^{3}(\rho+p)] = a^{3}dp \tag{1.11}$$

and translate it into the formalism of the first law of thermodynamics:

$$d(\rho a^3) = -pd(a^3) \tag{1.12}$$

which indicates, for a comoving volume element, the multiplication of its pressure and volume (i.e. volume work)  $pd(a^3)$  is the opposite value of the energy change of the mass element  $d(\rho a^3)$ .

### 1.2.3 Friedmann Equation

The expanding rate of the universe is reflected via Hubble parameter H, defined as:

$$H \equiv \frac{\dot{a}}{a} \tag{1.13}$$

Hubble parameter evolves according to observations, and is dynamically treated in theory too. The so-called Hubble Constant  $H_0$  refers to the presentday value of Hubble parameter in particular. Under FRW meric, the non-null component for the Ricci tensor of gravitational field equation (1.2) is:

$$R_{00} = -3\frac{\ddot{a}}{a} \tag{1.14}$$

$$R_{ij} = -\left(\frac{\ddot{a}}{a} + 2\frac{\dot{a}^2}{a^2} + 2\frac{k}{a^2}\right)g_{ij}$$
(1.15)

Ricci scalar being:

$$R = -6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2}\right) \tag{1.16}$$

the 0 - 0 (time) component of the gravitation equation being

$$\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} = \frac{8\pi G\rho}{3} \tag{1.17}$$

This is the very Friedmann equation that represents the dynamical evolution of cosmic scalar factor, which indicates that that the expansion process depends on both energy term and geometrical curvature term. The i - i (spatial) component of gravitational equation (1.2) is:

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} = -8\pi Gp \tag{1.18}$$

Meantime it is also induced to get Raychaudhuri equation which shows the evolution of Hubble parameter:

$$\dot{H} = -4\pi G(p+\rho) + \frac{k}{a^2}$$
(1.19)

Eq(1.12), Eq(1.17), Eq(1.18) is connected via Bianchi identity, hence only two ones are independent. From (1.10), (1.17), (1.18) one could induce the energy-momenta conserving equation for cosmological perfect fluid:

$$\dot{\rho} + 3H(\rho + p) = 0 \tag{1.20}$$

Unite (1.17)(1.18)to deduct  $\frac{k}{a^2}$  and one would get a straight overlook on the cosmic expansion:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p) \tag{1.21}$$

Conclusion is directly drawn that  $\rho + 3p < 0$  holds for accelerating expansion.

Via Hubble parameter, Fredmann equation could be rewritten as:

$$\frac{k}{H^2 a^2} = \frac{\rho}{\frac{3H^2}{8\pi G}} - 1 \tag{1.22}$$

and therefore the cosmic curvature completely depends on the total energy density, which is in casuality agreement with general relativity. If the density is above  $\frac{3H^2}{8\pi G}$ , k will be positive, and results in a limited and closed cosmos; If the density is identical to  $\frac{3H^2}{8\pi G}$ , k will be null, and results in a infinite and flat cosmos; If the density is below  $\frac{3H^2}{8\pi G}$ , k will be negative, and results in a infinite and open cosmos. So in further discussion it is proper to define the unitary quantity  $\frac{3H^2}{8\pi G}$ , which share the dimension of energy density, to be the critical energy density of the universe:

$$\rho_{crit} \equiv \frac{3H^2}{8\pi G} \tag{1.23}$$

together with a non-dimensional quantity of relative energy density:

$$\Omega \equiv \frac{\rho}{\rho_{crit}} \tag{1.24}$$

According to the current theories, it is  $\rho_{crit}$  and  $\Omega$  that determine the ultimate fate of the universe. Hence Friedmann equation take the further form:

$$\frac{k}{H^2 a^2} = \Omega - 1 \tag{1.25}$$

thus a clear graph is finished to reflect how the evolution of the universe depends on its property parameters, which prepares a firm foundation for cosmic observations and cosmological dynamics.

### 1.2.4 EOS in Perfect Fluid Scenario

Actually to resolve cosmological problems under perfect fluid scenario, we still need the equation of state (EOS):

$$p = p(\rho) \tag{1.26}$$

Thus theoretically, the combination of Friedmann Equation (1.17), energy-momenta conservation equation (1.20) and EOS (1.26) is equal to solve the dynamics of cosmos.

Generally, a simplest EOS is employed for cosmological perfect fluid:

$$p = w\rho \tag{1.27}$$

where  $w \in \mathbb{R}$ . For ordinary matter, w = 0. And the accelerating expansion conditions  $\rho + 3p < 0$  can be re-expressed via EOS parameter:

$$w < -1/3 \tag{1.28}$$

Integration by (1.12) give rise to the relation of energy density and cosmic scalar factor:

$$\rho \propto a^{-3(1+\omega)} \tag{1.29}$$

When the universe is dominated by relativistic matter and radiation (i.e. the baby era), w = 1/3, hence

$$a(t) \propto (t - t_0)^{1/2}, \qquad \rho \propto a^{-4}$$
 (1.30)

When the universe is dominated by non-relativistic matter, w = 0:

$$a(t) \propto (t - t_0)^{2/3}, \qquad \rho \propto a^{-3}$$
 (1.31)

All in all, different eras correspond to different domination and evolution.

### Chapter 2

# Accelerating Expansion of the Universe

### 2.1 Accelerating Expansion and Dark Energy

The inflation theory, which is a great success as supplement to Big Bang cosmology, declares that the cosmos is flat in early 1980s. Theorists' insistence lead to annoying disagreement. However, cosmic observations found that there is far less matter than needed for flat topology. Shortly afterwards in early 1990s, COBE is sent by NASA to explore the cosmic microwave background. To theorists' delight, the location of the first peak in angular power spectrum confirms the flatness of the cosmos. This strongly require a new density component to fit the remarkable gap between the matter density and the critical density. Before physists could take a calm breath, the redshift measurement of 1a supernova declared the accelerating expansion of the universe! This milestone is among the greatest ever discoveries, and bring the discoverers the Shaw Awards for Astronomy.

So, CMB exploring proves the flatness, homogeneity and isotropy of the universe, and SN1a confirms the accelerating expansion. These results lead to nondebatable conclusions that, if general relativity and the standard model of particle physics are correct, then

(1) The density of cosmos is  $\Omega = 1$ . Since baryons, non-baryonic CDMs and radiation add up to only around  $\Omega_m + \Omega_\gamma = 0.3$ , the major energy (the gap) comprise of an unknown component, dubbed Dark Energy (DE).

(2) Only limited dynamical characters are known. Dark energy has positive energy,

negative pressure, and anti-gravitation effects that drive the accelerating expansion. DE doesnot condensate at scales smaller than 100Mpc; DE density is almost constant; DE is only dominant recently ( $z \in (0 \sim 1)$  and  $z_0 = 0$  today), so it doesn't bother the nuclear-synthesis process at early era.

And even up till now, we are still quite unfamiliar with DE. DE may be scalar fields, vector fields, tensor fields, or even cosmic strings, etc[7]. And a set of typical data is that the critical density is made up of 72% DE, 24% CDM, 4% ordinary matter.

### 2.2 Characteristic Parameters of Cosmos

The properties of the universe need to be parameterized for quantitative study (this is a common method to handle huge and complicated systems).

Presently, there are 12 parameters that are mostly popular. [40]:

$$P \equiv (\omega_b, \quad \omega_c, \quad \mathcal{K}, \quad H, \tau, \quad w_0, \quad w_1, \sum m_{\nu}, \quad n_s, \quad A_s, \quad \alpha_s, \quad r)$$

where  $\omega_b \equiv \Omega_b h^2$  and  $\omega_c \equiv \Omega_c h^2$  represent the relative energy density of baryonic matter and cold dark matter.  $\mathcal{K}$  represents the presentday scalar curvature ,and cosmic observations support the result of  $\mathcal{K} = 0$ , a flat cosmos H represents Hubble parameter; $\tau$ represents luminous depth  $\sum m_{\nu}$  represents the total mass of the three generation of neutrinos. In general computing process, the EOS parameter is often rewritten as  $w = w_0 + w_1(1-a), w_0$  and  $w_1$  represent the new EOSs respectively. The last four parameters are related to inflation theory. $A_s$  represents the amplitudes of original power $n_s$ represents the power index for scalar spectra $\alpha_s$  represents shift of spectra index  $n_s r$  is the tensor power parameter.

It is of great importance to investigate these patameters , which help constraint the properties of dark energy. Here below we list the data from Willkinson Microwave Anisotropy Probe, which can de downloaded from the WMAP homepage of NASA.

### 2.3 Mechanism for Accelerating Expansion

Ever since the proof that DE is dominant, its components, origin, dynamics are waiting to be determined. With the parameter constraining via cosmic observations, various models have been put forward for a cosmos with accelerating expansion. I would divide these models into three categories for further discussion. Table: Data from WMAP.

Cosmic Parameters	Symbol	Value	+ uncertainty	-uncertainty
total density	$\Omega_{tot}$	1.02	0.02	0.02
Quintessence EOS	ω	< -0.78	95% CL	_
DE density	$\Omega_{\Lambda}$	0.73	0.04	0.04
Baryon Energy Density	$\Omega_b$	0.044	0.004	0.004
Baryon Number Density $(cm^{-3})$	$n_b$	$2.5  imes 10^{-7}$	$0.1  imes 10^{-7}$	$0.1  imes 10^{-7}$
Matter Energy Density	$\Omega_m$	0.27	0.04	0.04
Neutrino Density	$\Omega_{\nu}h^2$	< 0.0076	95% CL	_
CMB Temperature(K)	$T_{cmb}$	0.275	0.002	0.002
Baryon-photon Number	$\eta$	$6.1  imes 10^{-10}$	$0.3\times 10^{-10}$	$0.2\times 10^{-10}$
$8h^{-1}Mpc$ Spherical Fluctuation.	$\delta_8$	0.84	0.04	0.04
Scalar Spectra Index $k_0 = 0.05 M p c^{-1}$ )	$n_s$	0.93	0.03	0.03
Index shift $(k_0 = 0.05 M p c^{-1})$	$dn_s/dlnk$	-0.031	0.016	0.018
Scalar-Tensor Ratio $(k_0 = 0.002 M p c^{-1})$	r	< 90%	95% CL	_
decoupling redshift	$z_{dec}$	1089	1	1
decoupling width(FWHM)	$\Delta z_{dec}$	195	2	2
Hubble parameter(reduced)	h	0.71	0.04	0.03
Cosmos Age(Gyr)	$t_0$	13.7	0.02	0.02
Decoupling time(Kyr)	$t_{dec}$	379	8	7
Heavy Ionic time (Myr,95% $CL$ )	$t_r$	180	220	2580
decoupling interval(Kyr)	$\Delta t_{dec}$	118	3	2
equality redshift	$z_{eq}$	3233	194	210
Baryonic luminous depth	au	0.17	0.04	0.04

### 2.3.1 Scalar Fields

Bulk of these proposals are some kinds of scalar fields with shallow potentials and thus tiny mass. The field rolls slowly, up (canonical) or down (noncanonical), in its own potential. This process takes place only considerable on Hubble time scale. EOS in scalar field models takes the traditionally most simple form  $p = w\rho$ , and EOS parameter w turns the identity of different fields. w > -1 or -1 < w < -1/3 means Quintessence field; w = -1 is Einstein's cosmological constant  $\Lambda$ ; w < -1 reflects Phantom field. These cases are all real fields; and those with w crossing -1 can be real (Quintom) or complex (Hessence).

And the key point is that a scalar field is itself meaningful at all, if not treated together with the accelerating expansion. Firstly, it's generated or associates with many attractive theories, such as superstring quantum field, pseudo-Nambu-Goldstone Bosonic Formulation, Brans-Dicke theory. So it's a natural trial to test the cosmological consequence of a scalar field, typical work like Ratra and Peebles in late 1980s. Secondly, a scalar field could help make clear the CDM spectra. Last but not least, even a light scalar field can generate testable consequences in the standard CDM scenario. For those reasons, scalar fields such as Quintessence has been discussed pioneeringly, after the inflation theory, before the accelerating expansion observations. More discussions are available in this thesis afterwards.

### 2.3.2 Modified Gravitational Field Equation/Friedmann Equation

A most remarkable candidate is Cardassian models, which add a new energy term to the right side of Friedmann equation, or to the energy-momenta tensor of Gravitational Field Equation. This Cardassian term depends on the density of matter and radiation only, and the modified Friedmann equation could produce a flat, matter dominated universe with accelerating expansion at late time, while the early evolution of baby cosmos is not perturbed. The creative work of this thesis is exactly connected with this models.

Modifications to gravitation equation includes operations on the geometric term of Ricci tensor, or energy-momenta tensor. Besides Cardassian, other examples include Randall-Sundrum model, etc.

### 2.3.3 Modern QFT approaches

Such models include k-essence, tachyon and Chaplygin gas. K-essence focuses on modifying the kinetic energy term of Lagrangian, who acts as the first principle. Tachyon plays an important role in standard model of particle physics and superstring guage theory, and its connections with accelerating expansion is established after 1998. Both k-essence and tachyon modeling are deeply rooted in quantum field theory. Chaplygin gas seems less mathematically complicated than k-essence and tachyon formulation. It is a kind of non-relativistic perfect fluid in superstring theory, being the only liquid allowing supersymmetric generalization. Its EOS is quite different from that of scalar fields, and can be reached via several approaches, such as non-relativistic approximation of Born-Infield theory. In (generalized) Chaplygin gas cosmology, the ultimate fate of accelerating expansion is de Sitter Universe. A most interesting property is that (generalized) Chaplygin gas indicates a unified description of dark energy with dark matter.

In the following test, I firstly review some typical existing work, including four scalar fields (Quintessence, Phantom, Quintom, Hessence), k-essence and Chaplygin gas. I put my work in the last two chapters, where I come up with a new Cardassian model and introduce virial collapse to Cardassian cosmology.

### Chapter 3

# Quintessence & Phantom

### 3.1 Quintessence Scalar Field

Quintessence scalar field is the first introduced dark energy model, and has received most attention. At early 1980s, a scalar field is employed in the inflation theory, which encourages people to analyze the consequence of a scalar field rolling in its own potential at the whole Hubble time scales, rather than the tiny time interval at the extremely early universe. This work is done in late 1980s, and is re-introduced as the first non- $\Lambda$ mechanism for dark energy after 1998.

### 3.1.1 Basic Structure of Quintessence

Generally Quintessence is treated as a classical scalar field, with smallest coupling with gravitation field, and its action is:

$$S = \int \mathcal{L}_{\phi} \sqrt{-g} d^4 x \tag{3.1}$$

where g is the determinant of the matrix  $g_{\mu\nu}$ . With metric (+--), phenomenently, the Lagrangian density with positive energy and positive pressure  $\mathcal{L}_{\phi}$  is:

$$\mathcal{L}_{\phi} = \frac{1}{2} g^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi - V(\phi)$$
(3.2)

Take variance operation on (3.1), and one gets the energy-momenta tensor for a scalar field:

$$T^{\phi}_{\mu\nu} = \partial_{\mu}\phi\partial_{\nu}\phi - g_{\mu\nu} \left[\frac{1}{2}\partial_{\mu}\phi\partial_{\nu}\phi - V(\phi)\right]$$
(3.3)

Take FRW metric(1.3) in a flat space into account, via the conservation equation  $T^{\mu\nu}_{\phi}_{\phi}_{;\nu} = 0$ , one can get the dynamical equation for the scalar field  $\phi$ :

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV}{d\phi} = 0 \tag{3.4}$$

Assuming that the scalar field is homogeneous, whose derivative to spatial coordinates is zero and dependent on cosmological time only. Via FRW line elements in such a field, one can get the energy density and pressure of Quintessence field:

$$\rho_{\phi} = \frac{1}{2}\dot{\phi}^2 + V(\phi) \tag{3.5}$$

and

$$p_{\phi} = \frac{1}{2}\dot{\phi}^2 - V(\phi)$$
 (3.6)

Hence EOS parameter of Quintessence field is:

$$w_{\phi} = \frac{p_{\phi}}{\rho_{\phi}} = \frac{\frac{1}{2}\dot{\phi}^2 - V(\phi)}{\frac{1}{2}\dot{\phi}^2 + V(\phi)}$$
(3.7)

The dynamical equations for the evolutionary cosmos is

$$H^{2} = \frac{8\pi G}{3} \left[ \frac{1}{2} \dot{\phi}^{2} + V(\phi) \right]$$
(3.8)

$$\frac{\ddot{a}}{a} = -\frac{8\pi G}{3} \left[ \dot{\phi}^2 - V(\phi) \right] \tag{3.9}$$

From Eq.(3.9) indicates that accelerating expansion is possible if  $\dot{\phi}^2 < V(\phi)$  holds. Insert (3.7) and (1.20) into Quintessence field, then:

$$\rho_{\phi} = \rho_0 exp \left[ -\int 3(1+w_{\phi})\frac{da}{a} \right]$$
(3.10)

where  $\rho_0$  is an integral constant. According to (3.9), the value of  $w_{\phi}$  must be mamong -1 to 1. If the rolling in the potential proceeds very slowly, i.e.  $\dot{\phi}^2 \ll V(\phi)$ , then  $\omega_{\phi} = -1$  and  $\rho_{\phi}$  being a constant; If the rolling in the potential proceeds very sharply, i.e.  $\dot{\phi}^2 \gg V(\phi)$ , then  $\omega_{\phi} = 1$  and  $\rho \propto a^{-6}$ As to the media cases, Quintessence energy density will be

$$\rho_{\phi} \propto a^{-m}, \qquad 0 < m < 6 \tag{3.11}$$

Since  $\omega = -1/3$  is the critical condition for accelerating expansion and decelerating expansion, constraints can be strengthened for 0 < m < 2 for accelerating expansion.

#### 3.1.2 Quintessence Potential

:

In discussions with Quintessence field, three potentials are most popular

(1)Inverse power style potential by supersymmetry models[10]:

$$V(\phi) = V_0 \left(\frac{\phi_0}{\phi}\right)^{a_0} \qquad a_0 > 1 \tag{3.12}$$

(2)Exponential potential by Kaluza-Klein theory[11]:

$$V(\phi) = V_0 e^{\phi - \phi_0} \tag{3.13}$$

(3)Cosine potential by Pseudo-Nambu-Goldstone Bosonic Mechanism[12]:

$$V(\phi) = \frac{1}{2} V_0 \left[ \cos(\frac{\phi - \phi_0}{f}) + 1 \right]$$
(3.14)

Let's take a closer look through a concrete sample: For power style expansion, the driven potential is exponential. The scalar factor expands in power style,  $a(t) \propto t^p$ , where p = 1 is the critical condition to separate accelerating and decelerating expansion. According to (1.18), one has

$$\dot{H} = -4\pi G \dot{\phi}^2 \tag{3.15}$$

So one could employ the observables H and  $\dot{H}$  to represent the potential  $V(\phi)$  and  $\dot{\phi}$ :

$$V = \frac{3H^2}{8\pi G} \left( 1 + \frac{\dot{H}}{3H^2} \right) \tag{3.16}$$

$$\phi = \int dt \left( -\frac{\dot{H}}{4\pi G} \right)^{1/2} \tag{3.17}$$

This operation can be treated as the reconstruction of the model. Hence the driven potential for a power style expanding potential is:

$$V(\phi) = V_0 exp\left(-\sqrt{\frac{16\pi}{p}}\frac{\phi}{m_{pl}}\right)$$
(3.18)

where  $V_0$  is a constant, unitary parameter  $m_{pl}$  being Plank mass. In a word, an exponential Quintessence potential will drive the universe expanding in power style. Moreover, exponential Quintessence could lead to scaling solutions of dynamical phase system, where the ratio of Quintessence energy density  $\rho_{\phi}$  to that of background matter  $\rho_m$  is constant.

### 3.1.3 Scalar Fields' problems and Tracker Field

If one employs a slowly rolling and light scalar field to drive the accelerating expansion, two serious problem will arise. This is the difficulty that all scalar field come across.

### (1)Fine-tuning Problem

The kernel of this problem lies that, why the missing energy density differs so much in comparison with the typical energy scale in particle physics ? According to the current data,  $\Omega_m = 0.3$ , then dark energy density is in the magnitude of  $10^{-47} GeV^4$ , which needs an extra 14 magnitude beyond electromagnetic interactions.

### (2)Coincidence Problem

The kernel of this problem lies at the initial conditions of the cosmos. The evolutionary rate of Quintessce field differs very much with that of energy density, and the process is quite complicated. So, to evolve into the state where the energy density of Quintessence field and matter are of the same magnitude after 137Gyr, the initial conditions must be exactly set.

All scalar fields cosmological models are upset with the two problems, but, the discovery of the so-called Tracker Field, which originate from Quintessence and then spread to all other scalar fields, throws light on beating these problems[15]. It's found that, as for some special potential, such as  $V \sim V_0 \phi^{-n}$ ,  $V \sim V_0 \exp(M/\phi - 1)$ ,  $V \sim V_0 [sinh(\alpha \sqrt{k_0} \Delta \Phi)]^{\beta}$  [16], there are tracker solution for the Quintessence scalar field. Tracker solutions have quite unrestricted initial conditions, and the energy density ratio of Quintessence to matter is permitted to fluctuate for 100 magnitudes to have the same evolutionary results. This means it's almost independent of initial conditions, or perfectly Markovian. This solves the two problems referred to above in some sense. And from then on, the existence of tracker solution, in the differential autonomous dynamical system turn an important judgement for the quality of a scalar field for dark energy, and for the quality of a potential for certain scalar field.

### 3.1.4 Typical discussions for Quintessence

(1)In [17] it's shown that the gravitational field equation and conservation laws put strict constraints to Quintessence with w = const., For constant EOS parameter w and  $w \neq -1$  (simply the case of cosmological constant  $\Lambda$ ):

$$\frac{V'}{V'_0} = \sqrt{\Omega_\phi \left(\frac{V}{V_0}\right)^2 + \Omega_M \left(\frac{V}{V_0}\right)^{\frac{w+2}{w+1}} + \Omega_k \left(\frac{V}{V_0}\right)^{\frac{3w+5}{3w+3}}} \tag{3.19}$$

where V is the scalar potential for Quintessence field,  $V' \equiv \pm 3H_0\sqrt{(1-w^2)V_0/2}, V_0$ being the presentday value.

(2)Carroll made typical evaluation to the strength of Quintessence field[18]. Cosmological observations restrict that the scalar potential is quite shallow, which means the tiny mass in excited states,  $m_{\phi} \equiv \sqrt{V''(\phi)/2} \leq H_0 \sim 10^{-33} eV$ . In order to support the observed energy density, the current value of the potential must be approximately the closure density,  $V(\phi_0) \sim (10^{-3} eV)^4$ , and the field be  $\phi_0 \sim 10^{18} GeV \sim M_{pl} = (8\pi G)^{-1/2}$ , where  $M_{pl}$  is the reduced Plank mass. A light scalar field will generate testable long-range force, and if Quintessence could couple with matter, it would lead to the evolution of physical constants at Hubble time scale.

(3)[19] investigates the origin of Quintessence from breakdown of supersymmetry. [20] investigated the possibility that axions and QCD massless quark comprise the Quintessence. As a matter of fact, in the following years since the discovery of dark energy, attentions are focused on the theoretical formulation, and practical constrains for Quintessence takes fall behind obviously. Various connections with Quitessence from other branches have been established, and large amounts of theses are available, but this thesis wouldnot go further and wider review for Quintessence.

### 3.2 Phantom Scalar Field

According to Riess et al, if the matter density is treated as  $\Omega_m^0 = 0.27 \pm 0.04$ , one would see  $w = -1.02_{-0.19}^{+0.13}$ , and -1.46 < w < -0.78 for 95% confidence level. If one goes further to take the results of CMB and 2dFGRs into consideration, it's  $w = -1.08^{+2.0} - 0.18$ . So, w < -1 is also possible. Caldwell for the first time discussed the cosmological consequence for w < -1, and introduced the second scalar field, Phantom[21], as opposed to Quintessence. Phantom has many amazing properties. Phantom violates the weak energy conditions of gravitation; w < -1 leads to the effective sound speed  $v = \sqrt{|dp/d\rho|}$  beyond light velocity. The most debating result is the so-called Big Rip. The dark energy density is dynamical and increase along with the accelerating expansion; thus the increasing dark energy and the accelerating expansion unite to form a positive feedback pair, and one day in the future, the huge negative pressure will tear out the spacetime and all matter. For a review, see [22] and [24].

### 3.2.1 basic Structures of Phantom

Canonical scalar fields couldnot realize the process for  $\omega < -1$ , so noncanonical ones are employed for Phantom. A noncanonical Lagrangian for free Phantom field, which also has the smallest couplings with gravitation, can be written as [21][25]:

$$\mathcal{L}_{\phi} = -\frac{1}{2}g^{\mu\nu}\partial_{\mu}\phi\partial_{\nu}\phi - V(\phi)$$
(3.20)

Hence the dynamical equation for Phantom field is:

$$\ddot{\phi} + 3H\dot{\phi} - \frac{dV}{d\phi} = 0 \tag{3.21}$$

the minus sign before the potential term makes Phantom climbs up the potential, as opposed to a canonical field's rolling down in its potential (see Quintessence).

Variance of Phantom Lagrangian leads to Phantom energy density and pressure:

$$\rho_{\phi} = -\frac{1}{2}\dot{\phi}^2 + V(\phi)$$
 (3.22)

and

$$p_{\phi} = -\frac{1}{2}\dot{\phi}^2 - V(\phi) \tag{3.23}$$

as well as Phantom EOS (To avoid the increase of Tackyon module, w is always assumed to be constant in Phantom):

$$w_{\phi} = \frac{p_{\phi}}{\rho_{\phi}} = \frac{-\frac{1}{2}\dot{\phi}^2 - V(\phi)}{-\frac{1}{2}\dot{\phi}^2 + V(\phi)}$$
(3.24)

Obviously  $w_{\phi} < -1$  when  $\frac{1}{2}\dot{\phi}^2 < V(\phi)$ . And energy-momenta tensor for Phantom field is:

$$T^{\phi}_{\mu\nu} = -\partial_{\mu}\phi\partial_{\nu}\phi - g_{\mu\nu}\left[\frac{1}{2}\partial_{\mu}\phi\partial_{\nu}\phi - V(\phi)\right]$$
(3.25)

### 3.2.2 Hubble Parameter & Phantom Cosmology

The energy density of matter becomes equal with that of Phantom field at the cosmological time  $t_{eq}$ , and the scalar factor driven by Phantom energy will be:

$$a(t) \simeq a(t_{eq}) \left[ (1+w) \frac{t}{t_{eq}} - w \right]^{\frac{2}{3(1+w)}}, \quad w < -1$$
 (3.26)

So the scalar factor will be divergent one day, i.e.  $t \to t_{BR} = \left(\frac{w}{1+w}\right) t_{eq}$  and  $a(t) \to \infty$ . Insert w < -1 into Friedmann Equation:

$$H(z) = H_0 \left[ \Omega_m (1+z)^3 + \Omega_X (1+z)^{3(1+w)} \right]^{1/2}$$
(3.27)

where  $H_0 = H(z = 0)$  means the presentday value of Hubble parameter, and  $w = w_{DE}$ refers to EOS parameter of dark energy.  $\Omega_m = \frac{8\pi G\rho_{0m}}{3H_0^2}$ ,  $\Omega_X = \frac{8\pi G\rho_{DE}}{3H_0^2}$ , is relative energy density for matter and dark energy respectively, and  $\Omega_m + \Omega_X = 1$ . Thus,

$$H(z) \sim t_{eq}^{-1} \left[ (1+w)\frac{t}{t_{eq}} - w \right]^{-1}$$
(3.28)

When  $t \to t_{BR} = \left(\frac{w}{1+w}\right) t_{eq}$ , the Hubble parameter is also divergent mathematically. This indicates that one day the expanding velocity will approach the extremely upper limit. So Phantom dark energy density  $\rho(t) \propto \left[\Omega_m (1+z)^3 + \Omega_X (1+z)^{3(1+w)}\right]^{-2}$  is divergent at Big Rip, too. At the Big Rip singularity, spacetime and matter are all destroyed.

Phantom is also a kind of scalar field under large amounts of discussions, and its connections with quantum field theory, standard model of particle physics, superstring theory and quantum cosmology have all been established [33] [34] [35] [36], etc. And not all Phantom cosmos of  $w_{\phi} < -1$  ends up with Big Rip, but extra mechanism must be introduced, see [26] [27] [28] [29] [30] [31] [32] and the pioneering work of classification of Big Rips from Qhantom and Quintessence theories [23], as well as alternative attempt to quintom [27].

### Chapter 4

# Quintom & Hessence

### 4.1 Quintom Scalar Field

### 4.1.1 Scalar Field with w crossing -1

Riess et al analysed their SN1a Gold data (with dark energy EOS parameter  $w(z) = w_0 + w'z$ ), and found  $w = -1.02^{+0.13}_{-0.19}$ , which indicates that the presentday EOS parameter is probably crossing -1. Shortly, Alam, Sahni et al combined Gold data with CMB data to examine the behaviors of dark energy, and they drew clear conclusion that EOS parameter evolves continuously, from  $w \simeq 0$  at  $z \simeq 1$ , to  $w \leq -1$  at  $z \simeq 0$  today.

Hubble parameter and luminal distance is connected via:

$$H(z) = \left[\frac{d}{dz}\left(\frac{d_L(z)}{1+z}\right)\right]^{-1}$$
(4.1)

Alam et al selected a formalism in which Hubble parameter is independent of models:

$$H(x) = H_0 [\Omega_{m0} x^3 + A_0 + A_1 x + A_2 x^2]^{\frac{1}{2}}$$
(4.2)

where x = 1 + z, and  $\Omega_{m0}$  being the presentday matter density, and  $A_0$ ,  $A_1$ ,  $A_2$  are constants. Thus EOS parameter is reconstructed tobe:

$$\omega(x) = \frac{(2x/3)H'/H - 1}{1 - (H_0/H)^2 \Omega_{m0} x^3}$$
(4.3)

where prime ' refers to derivative with x. Together with modified  $\chi^2$  test, the above conclusions are drawn.

Hunterer confirmed the result of crossing -1, via data fitting of piecewise function and cosmic obsevations, and to be more exact, the cosmological redshift z < 0.2 for w < -1, and z > 0.2 for z > -1. Corasaniti went even further to make combined constraints with CMB, SN1a, and large scale exploring data, to get the result of w crossing -1.

As it is put above,  $-1 < w < -\frac{1}{3}$  for Quintessence and w < -1 for Phantom, and this constraints is principly inviolated, so, both Quintessence and Phantom are powerless to realize the crossing of the state w = -1. And therefore, the so-called Quintom scalar field is put forward, which aims at crossing EOS -1.

However, to build such a Quintom model, one runs into the No-Go theorem (see A. Vikman[37]): (1)Under the frame of four dimensional FRW metric and general relativity, (2)if dark energy is a single perfect fluid, or single scalar field  $\phi$  with Lagrangian  $\mathcal{L} = \mathcal{L}(\phi, \partial_{\mu}\phi\partial^{\mu}\phi)$ , (3)with smallest coupling to gravitation, then the EOS parameter w couldnot cross -1. This means, to make w cross -1, at least one of the three conditions should be violated, such as Quintom with double field[38], or Quintom based on Lagrangian with high order modification term[39].

### 4.1.2 Basic Structure of Double Field Quintom

Quintom field includes two real scalar fields, one being canonical Quintessence style field  $\phi$ , the other being Phantom style field  $\varphi$ .

Lagrangian deensity of Pantom field is:

$$\mathcal{L} = \frac{1}{2} \partial^{\mu} \phi_Q \partial_{\mu} \phi_Q - \frac{1}{2} \partial^{\mu} \phi_P \partial_{\mu} \phi_P - V(\phi_Q, \phi_P)$$
(4.4)

thus Quitom dark energy density is:

$$\rho_q = \frac{\dot{\phi}^2}{2} - \frac{\dot{\varphi}^2}{2} + V(\phi, \varphi)$$
(4.5)

and with pressure of:

$$p_q = \frac{\dot{\phi}^2}{2} - \frac{\dot{\varphi}^2}{2} - V(\phi, \varphi)$$
(4.6)

With the dark energy density and pressure, EOS parameter of Phantom dark energy is:

$$w_q = \frac{p_q}{\rho_q} = \frac{\dot{\phi}^2 - \dot{\varphi}^2 - 2V(\phi, \varphi)}{\dot{\phi}^2 - \dot{\varphi}^2 + 2V(\phi, \varphi)}$$
(4.7)

From Eq(4.7) we can see if  $\dot{\phi}^2 > \dot{\varphi}^2$ ,  $\omega > -1$ ; if  $\dot{\phi}^2 < \dot{\varphi}^2$ ,  $\omega < -1$ . EOS of Quintom crosses -1.

At first these two componential potential have no interactions with each other, Quintom potential is the simple addition of those two potentials:

$$V(\phi,\varphi) = V_1(\phi) + V_2(\varphi) \tag{4.8}$$

take a simple potential  $V(\phi_Q, \phi_P)$  for example:

$$V(\phi_Q, \phi_P) = V_0 \left[ exp(-\frac{\lambda}{M_P}\phi_Q) + exp(-\frac{\lambda}{M_P}\phi_P) \right]$$
(4.9)

For these cases, the steady attractor solution is in Phantom style, or the ultimate state is Phantom, where one has to face the nightmare of Big Rip.

### 4.1.3 Generalized Quintom

There are various methods to make expansion, a simplest and natural way is to set weights for the addition of Quintessence and Phantom.

### 4.2 Hessence Scalar Field

Quintessence, Phantom, Quintom are all real, and the earliest attempt to describe dark energy is Spintessence model. However, Spintessence falls into deadly trouble when dealing with the steadiness of Q-balls and is generally abandoned as a dark energy model(but Spintessence turns out to be a good candidate for dark matter).

### 4.2.1 Basic Structure of Hessence

Considering a single complex scalar field with EOS parameter crossing -1[41][42]:

$$\Phi = \phi_1 + i\phi_2 \tag{4.10}$$

As for complex field, its canonical Lagrangian is:

$$\mathcal{L} = \frac{1}{2} (\partial_{\mu} \phi_1)^2 + \frac{1}{2} (\partial_{\mu} \phi_2)^2 - V(|\Phi|)$$
(4.11)

But, Dark energy based on canonical Lagrangian is not steady (Spintessence). Thus, the modified noncanonical Lagrangian is introduced as the first principle for Hessence complex model:

$$\mathcal{L}_{he} = \frac{1}{4} [(\partial_{\mu} \Phi)^{2} + (\partial_{\mu} \Phi^{*})^{2}] - U(\Psi^{2} + \Psi^{*2})$$
$$= \frac{1}{2} [(\partial_{\mu} \phi)^{2} - \phi^{2} (\partial_{\mu} \theta)^{2}] - V(\phi)$$
(4.12)

the action of Hessence field is:

$$S = \int d^4x \sqrt{-g} \left( -\frac{R}{16\pi G} + \mathcal{L}_{he} + \mathcal{L}_m \right)$$
(4.13)

where  $(\psi, \theta)$  are two newly defined variables for Hessence:

$$\phi_1 = \phi \cosh \theta, \qquad \phi_2 = \phi \sinh \theta \tag{4.14}$$

thus

$$\phi^2 = \phi_1^2 - \phi_2^2, \qquad \coth \theta = \frac{\phi_1}{\phi_2}$$
 (4.15)

This is why Hessence receives its name, and H here stands for *hyperbolic*.

In flat FRW spacetime with scalar factor a(t), and supposing  $\phi$  and  $\theta$  is isotropic, from Hessence Lagrangian (4.12), one gets Hessence dynamical equation which depends on  $\phi$  and  $\theta$ :

$$\ddot{\phi} + 3H\dot{\phi} + \phi\dot{\theta}^2 + \frac{\partial V}{\partial\phi} = 0 \tag{4.16}$$

$$\phi^2 \ddot{\theta} + (2\phi \dot{\phi} + 3H\phi^2) \dot{\theta} = 0 \tag{4.17}$$

Thus Hessence energy density and pressure are:

$$\rho_{he} = \frac{1}{2} (\dot{\phi}^2 - \phi^2 \dot{\theta}^2) + V(\Psi)$$
(4.18)

$$p_{he} = \frac{1}{2}(\dot{\phi}^2 - \phi^2 \dot{\theta}^2) - V(\Psi)$$
(4.19)

Based on(4.17), there's

$$Q = a^3 \Psi^2 \dot{\theta} = const. \tag{4.20}$$

where Q is the conservational quantity in certain volume, which is generated due to inner symmetry of complex scalar fields. In details, the Hessence Lagrangian (4.12) is conserving under local gauge transformations, or remain a same mathematical form after  $\phi \rightarrow \phi$  and  $\theta \rightarrow \theta - i\alpha$ . According to Nöether theorem, conserving fluid and quantity associates (noted Q). On the other hand, Hessence potential  $V(\Phi, \Phi^*)$  or  $V(\phi_1, \phi_2)$ should only depend on  $\Phi^2 + \Phi^{*2}$  or  $\phi_1^2 - \phi_2^2$  (or noted  $V(\phi)$ ), to keep the symmetry holding. Now turn attention to the comparison of Hessence Lagrangian with Quintom Lagrangian, which shows that it is  $\theta$  term or Q term that plays the role of Phantom field. When Q = 0, Hessence reduces to Quintessence, and there is no more necessity for crossing w = -1. And there's the pity that physical meaning of Q is still uncertain.

It should be stressed that Hessence differs from Quintom and is an independent model. An important difference is that  $\phi_1^2 - \phi_2^2$  is treated as a whole part to be the independent variable, while the two componential fields are generally independent to each other in Phantom.

With Eq(4.20) one can rewrite the dynamical equation, energy density and pressure of Hessence as:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{Q^2}{a^6\phi^3} + \frac{\partial V}{\partial\phi} = 0$$
(4.21)

$$\rho_{he} = \frac{1}{2}\dot{\phi}^2 - \frac{Q^2}{2a^6\phi^2} + V(\phi) \tag{4.22}$$

$$p_{he} = \frac{1}{2}\dot{\phi}^2 - \frac{Q^2}{2a^6\phi^2} - V(\Psi)$$
(4.23)

I t is notable that (4.21) is equal to Hessence energy conserving equation:

$$\dot{\rho}_{he} + 3H(\rho_{he} + p_{he}) = 0 \tag{4.24}$$

while Friedmann and Raychaudhuri equations turn to be

$$H^{2} = \frac{8\pi G}{3}(\rho_{he} + \rho_{m})$$
(4.25)

$$\dot{H} = -4\pi G(\rho_{he} + \rho_m + p_{he} + p_m)$$
(4.26)

where  $\rho_m$  and  $p_m$  are the energy density and pressure for background matter. Ultimately, EOS parameter of Hessence is:

$$w_{he} = \frac{\dot{\phi}^2 - \frac{Q^2}{a^6\phi^2} - 2V(\phi)}{\dot{\phi}^2 - \frac{Q^2}{a^6\phi^2} + 2V(\phi)}$$
(4.27)

And apparently when  $\dot{\phi}^2 \ge \frac{Q^2}{a^6 \Psi^2}$ , Hessence EOS parameter  $w_{he} \ge -1$  when  $\dot{\phi}^2 \le \frac{Q^2}{a^6 \Psi^2}$ ,  $w_{he} \le -1$ . Via the evolution of  $\phi$  field, EOS parameter w crosses -1 when  $\dot{\phi}^2 \ge \frac{Q^2}{a^6 \Psi^2}$ . Eqs 4.20-4.27 all depends on  $Q^2$  straightly rather than Q, which means +Q and -Q share the same dynamical behaviors, and leads to the assumption of existence of Hessence DE and anti-Hessence DE.

Canonical and complex fields could easily generate the so-called Q - ball structure. Q-ball is a kind of non-topological soliton, whose steadiness is related to the corresponding Q conserving quantity. As to complex fields with canonical Lagrangian, such as Spintessence, once the energy density fluctuates, the perturbation would magnify nonlinearly at an exponential speed to form Q-balls condensation. This a a big trouble for DE, because DE doesn't condensate at the scales smaller than 100Mpc. Once come into being, Q-balls act like matter and its energy density decay into other particles at the speed proportional to  $a^{-3}$ . However, as for Hessence DE, whose Lagrangian is noncanonical, the formation of Q-balls are easily avoided. Although the Hessence Lagrangian never occures before in quantum field theory, due to its steadiness consequences, Hessence is treated highly as a reasonable complex scalar field candidate for dark energy [45] [46].

### 4.2.2 Hessence Reconstruction via Hubble Parameter

Firstly, denote  $M_{pl}^2 \equiv (8\pi G)^{-1/2}$ , and  $M_{pl}^2$  is the reduced Plank mass. Then (4.25), (4.26) could be rewritten as [43]:

$$H^{2} = \frac{1}{3M_{pl}^{2}}(\rho_{he} + \rho_{m})$$
(4.28)

$$\dot{H} = \frac{1}{2M_{pl}^2} (\rho_{he} + \rho_m + p_{he} + p_m)$$
(4.29)

This is an interesting formulation. According to the two equations above, one gets:

$$V(\phi) = 3M_{pl}^2 H^2 + M_{pl}^2 \dot{H} - \frac{1}{2}\rho_m$$
(4.30)

$$\dot{\phi}^2 - \frac{Q^2}{a^6 \phi^2} = -2M_{pl}^2 \dot{H} - \rho_m \tag{4.31}$$

and the redshift being  $z = a^{-1} - 1$  (supposing  $z_0 = 1$  and subscript 0 refers to today), then for arbitrary f,

$$\dot{f} = -(1+z)H\frac{df}{dz} \tag{4.32}$$

and (4.30), (4.31) could be expressed as:

$$V(z) = 3M_{pl}^2 H^2 - M_{pl}^2 (1+z) H \frac{dH}{dz} - \frac{1}{2}\rho_{m0} (1+z)^3$$
(4.33)

$$\left(\frac{d\phi}{dz}\right)^2 - \frac{Q^2}{\phi^2}(1+z)^4 H^{-2} = 2M_{pl}^2(1+z)^{-1}H_{-1}\frac{dH}{dz} - \rho_{m0}(1+z)H^{-2}$$
(4.34)

now introduce such non-dimensional quantities as below:

$$\tilde{V} \equiv \frac{V}{M_{pl}^2 H_0^2}, \qquad \tilde{\phi} \equiv \frac{\phi}{M_{pl}}$$
(4.35)

$$\tilde{H} \equiv \frac{H}{H_0}, \qquad \tilde{Q} \equiv Q M_{pl}^2 H_0 \tag{4.36}$$

thus (4.33), (4.34) turn to:

$$\tilde{V}(z) = 3\tilde{H}^2 - (1+z)\tilde{H}\frac{d\tilde{H}}{dz} - \frac{3}{2}\Omega_{m0}(1+z)^3$$
(4.37)

$$\left(\frac{d\tilde{\phi}}{dz}\right) - \tilde{Q}^2 \tilde{\phi}^{-2} (1+z)^4 \tilde{H}^{-2} = 2(1+z)^{-1} \tilde{H} \frac{\tilde{H}}{dz} - 3\Omega_{m0} (1+z) \tilde{H}^{-2}$$
(4.38)

where  $\Omega_{m0} = \rho_0/(3M_{pl}^2H_0^2)$  refers to the presentday energy density of matter. Once  $\tilde{H}(z)$  or H(z) is known, we could reconstruct V(z) and  $\phi(z)$  via (4.37) and (4.38), and reconstruct  $V(\phi)$  via V(z) and  $\phi(z)$ . From (4.31) (4.37) one gets the reconstruction of Hessence EOS parameter:

$$w_{he}(z) \equiv \frac{p_{he}}{\rho_{he}} = \frac{-1 + \frac{2}{3}(1+z)\frac{dlnH}{dz}}{1 - \Omega_{m0}\tilde{H}^{-2}(1+z)^3}$$
(4.39)

Kinetic Energy could be induced from reconstruction of (4.31):

$$K \equiv \frac{\dot{\phi}^2}{2} - \frac{Q^2}{2a^6\phi^2}$$
(4.40)

thus,

$$\tilde{K}(z) \equiv \frac{K}{M_{pl}^2 H_0^2} = (1+z)\tilde{(}H)\frac{dH}{dz} - \frac{3}{2}\Omega_{m0}(1+z)^3$$
(4.41)

And this reconstruction depends on testable H(z) rather than certain models.

### 4.2.3 Autonomous Equations for Hessence

Firstly, to express the background matter via EOS of perfect fluid:

$$p_m = w_m \rho_m \equiv (w - 1)\rho_m \tag{4.42}$$

where  $0 < \gamma < 2$ . Particularly,  $\gamma = 1$  and  $\gamma = 4/3$  represent the matter dominated state and radiation dominated state respectively. Introduce the interaction term C of Hessence and matter at energy equilibrium, and when energy conservation conditon  $\rho_{tot} + 3H(\rho_{tot} + p_{tot}) = 0$  holds[42][44], there's:

$$\dot{\rho}_{he} + 3H(\rho_{he} + p_{he}) = -C \tag{4.43}$$

$$\dot{\rho}_m + 3H(\rho_m + p_m) = C \tag{4.44}$$

If there's no interaction between Hessence and background matter, C = 0, and interaction for  $C \neq 0$ .

To establish the differential autonomous equations, one takes the following transformations,  $\kappa^2 = 8\pi G, \kappa > 0$ , and introduce the non-dimensional variables:

$$x \equiv \frac{\kappa \dot{\phi}}{\sqrt{6}H}, \quad y \equiv \frac{\kappa \sqrt{V}}{\sqrt{3}H}, \quad z \equiv \frac{\kappa \sqrt{\rho_m}}{\sqrt{3}H}, \quad u \equiv \frac{\sqrt{6}}{\kappa \phi}, \quad v \equiv \frac{\kappa}{\sqrt{6}H} \frac{Q}{a^3 \phi}$$
(4.45)

They forms the Hessence phase space, and now Hessence dynamical problems can be treated as geometrical problems in Hessence phase space. And according to basic Hessence equations one gets

$$x' = 3x(x^2 - v^2 + \frac{\gamma}{2}z^2 - 1) - uv^2 - \sqrt{\frac{3}{2}}y^2f - C_1$$
(4.46)

$$y' = 3y(x^2 - v^2 + \frac{\gamma}{2}z^2) + \sqrt{\frac{3}{2}xyf}$$
(4.47)

$$z' = 3z(x^2 - v^2 + \frac{\gamma}{2}z^2 - \frac{\gamma}{2}) + C_2$$
(4.48)

$$u' = -xu^2 \tag{4.49}$$

$$v' = 3v(x^2 - v^2 + \frac{\gamma}{2}z^2 - 1) - xuv$$
(4.50)

where prime operator (') refers to derivative to e-folding time  $N \equiv lna$ , and

$$f \equiv \frac{1}{\kappa V} \frac{\partial V}{\partial \phi} \tag{4.51}$$

$$C_1 \equiv \frac{\kappa C}{\sqrt{6}H^2\dot{\phi}}, \qquad C_2 \equiv \frac{\kappa C}{2\sqrt{3}H^2\sqrt{\rho_m}} = \frac{x}{z}C_1 \tag{4.52}$$

Thus Friedmann Equation reads:

$$x^2 + y^2 + z^2 - v^2 = 1 (4.53)$$

And the relative energy density for each component reads:

$$\Omega_{he} = \frac{\rho_{he}}{\rho_{crit}} = x^2 + y^2 - v^2, \qquad \Omega_m = \frac{\rho_m}{\rho_{crit}} = z^2 \tag{4.54}$$

and the effective EOS parameters for Hessence and the whole cosmos are:

$$w_{he} = \frac{p_{he}}{\rho_{he}} = \frac{x^2 - v^2 - y^2}{x^2 - v^2 + y^2}$$
(4.55)

$$w_{eff} = \frac{p_{he} + p_m}{\rho_{he} + \rho_m} = x^2 - v^2 - y^2 + (\gamma - 1)z^2$$
(4.56)

the critical points  $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v})$  is gained via setting  $\bar{x}' = \bar{y}' = \bar{z}' = \bar{u}' = \bar{v} = 0$ . And this is the foundation for further analysis. Hessence has perfect mathematical structure, so it is employed to show the normal process of dynamical analysis here.

It is shown in [42] that, as for exponential potential and power potential, all late attractor solutions meet  $\omega_{he} \geq -1$  and  $\omega_{eff} \geq -1$  with 4-form interaction term C. Nevertheless, Phantom style solutions with  $\omega_{he} < -1$  or  $\omega_{eff} < -1$  is not stable, and will evolve into Quintessence attractor with  $\omega_{he} \geq -1$  or  $\omega_{eff} \geq -1$ , or de Sitter attractor with  $\omega_{he} = -1$  or  $\omega_{eff} = -1$ . So Big Rip is avoided in Hessence cosmology.

In [44], it is shown that in 4 dimensional phase space, the existence of stable late attractor solutions is independent of the forms of  $V(\phi)$  and C, but the stability of the solutions depends on the second order derivative of Hessence potential  $V(\phi)$ .
## Chapter 5

# k-essence & Chaplygin gas

## 5.1 k-essence Mechanism

The Scalar field dark energy models employ slowly rolling potentials to drive the accelerating expansion. Yet, such a cosmos scenario can be gained via modifying the kinetic energy term. In 1999, Armendariz-Picon realized that the kinetic energy term could lead to cosmic inflation in the baby era that is governed by high energy mechanism, and introduced k-inflation. Afterwards Chiba utilized the same mechanism to describe dark energy that drive the accelerating expansion, and Armendariz-Picon improved this work into k-essence. Fork-essence, high order modifications appears at the kinetic terms to drive the acceleration. And Bohn-Infeld field is one form of k-essence. If the form of pressure is properly selected, the fine-tuning problem and coincidence problem can also be solved[47] [48] [49].

k-essence has noncanonical kinetic energy. As for a scalar field with  $\phi$  and kinetic energy  $X \equiv -(1/2)(\nabla \phi)^2$ , the action with the most generalized form is:

$$S = \int d^4x \sqrt{-g} p(\phi, X) \tag{5.1}$$

where Lagrangian density is the same with pressure density  $p(\phi, X)$ . and the action is also used in generalized Quintessence models.

Usually, Lagrangian density of k-essence obeys the following constraints:

$$p(\phi, X) = f(\phi)\hat{p}(X) \tag{5.2}$$

This constraints originates from superstring theory, in which low energy effect will leads to higher order derivative terms from a' and loop modifications (the relation of a' and string length scalar  $\lambda_s$  is  $a' = \lambda_s/2\pi$ ). The four dimensional effective string action is:

$$S = \int d^4x \sqrt{-\tilde{g}} \{ B_g(\phi) \tilde{R} + B_{\phi}^{(0)}(\phi) (\tilde{\nabla}\phi)^2 - a' [c_1^{(1)} B_{\phi}^{(1)}(\phi) (\tilde{\nabla}\phi)^4 + \ldots] + O(a'^2) \}$$
(5.3)

where  $k^2 = 8\pi G = 1$ . Here  $\phi$  is Dilaton field, which controls the string coupling strength  $g_s^2$  via  $g_s^2 = e^{\phi}$ . For weak couplings  $e^{\phi} \ll 1$ , the coupling equation is  $B_g \approx B_{\phi}^{(0)} \approx B_{\phi}^{(1)} \approx e^{-\phi}$ . If the coupling is one order, it will take a more complicated form. Via conformal transformation  $g_{\mu\nu} = B_g(\phi)\tilde{g}_{\mu\nu}$ , the string action will transforms into Einsteinian action form:

$$S_E = \int d^4x \sqrt{-g} \left[\frac{1}{2}R + K(\phi)X + L(\phi)X^2 + \dots\right]$$
(5.4)

with the component

$$K(\phi) = \frac{3}{2} \left(\frac{1}{B} \frac{dB_g}{d\phi}\right)^2 - \frac{B^{(0)}}{B_g}$$
(5.5)

$$L(\phi) = 2c_1^{(1)} a' B_{\phi}^{(1)}(\phi)$$
(5.6)

Thus we could get the Lagrangian with noncanonical kinetic energy term:

$$p(\phi, X) = K(\phi)X + L(\phi)X^2$$
(5.7)

and redefine the field  $\phi_{new}$  as:

$$\phi_{new} = \int^{\phi_{old}} d\phi \sqrt{\frac{L}{|K|}} \tag{5.8}$$

Thus Lagrangian transforms into:

$$p(\phi, X) = f(\phi)(-X + X^2)$$
(5.9)

where  $\phi = \phi_{new}, X \equiv X_{new} = (L/|K|)X_{old}$ , and  $f(\phi) = K^2(\phi_{old})/L(\phi_{old})$ . This is the k-essence model induced from (5.2)via  $\hat{p}(X) = -X + X^2$ . Thus the pressure is also known, thus the energy density of the scalar field  $\phi$  is:

$$\rho = 2X \frac{\partial p}{\partial X} - p = f(\phi)(-X + 3X^2)$$
(5.10)

With energy density and pressure, one immediately gets EOS parameter for  $\phi$ :

$$w_{\phi} = \frac{p}{\rho} = \frac{1 - X}{1 - 3X} \tag{5.11}$$

and as we can see, for certain X,  $w_{\phi}$  is also a constant. For example, when X = 1/2 we could get a cosmological constant style EOS parameter  $w_{\phi} = -1$ ; to get an accelerating expansion w < -1/3, one gets X < 2/3.

According to energy density  $\rho$ , one gets the continuity equation (1.20). When matter and radiation is dominant and EOS parameter for background fluid is  $w_m$ , Eq(1.18) gives rise to the Hubble parameter:

$$H = \frac{2}{3(1+w_m)(t-t_0)}$$
(5.12)

and the energy density being

$$\dot{\rho} = -\frac{2(1+w_{\phi})}{(1+w_m)(t-t_0)}\rho \tag{5.13}$$

For certain X, or certain  $w_{\phi}$ ,  $f(\phi)$  is constraint for

$$f(\phi) \propto (\phi - \phi_0)^{-\alpha}, \quad \alpha = \frac{2(1 + w_\phi)}{1 + w_m}$$
 (5.14)

when  $w_{\phi} = w_m$ ,  $f(\phi) \propto (\phi - \phi_0)^{-2}$  for radiation dominated or matter dominated era, and this agrees with the scaling solutions. Thus, as for dark energy, one must take fine tuning of  $f(\phi)$  to satisfy the presentday cosmic density. And one should note that the density of the scalar field must be significantly less than that of the background fluid, i.e.  $\rho \ll \rho_m$ , which contradicts with a DE dominated universe. For example,  $f(\phi) \propto (\phi - \phi_0)^2$ , except for a late scaling solution, we has another solution of accelerating expansion. Actually this is the divide of accelerating expansion with decelerating expansion.

From (5.11) we could know that, the kinetic term x plays the key role to influent the EOS. If 1/2 < X < 2/3 holds, the scalar field  $\phi$  will acts as dark energy with  $0 \leq \alpha \leq 2$ . Armendariz-Picon et al have made more generalized analysis with  $\hat{p}(X)$  to avoid the fine-tuning problem.

## 5.2 Chaplygin gas model

In the discussions with Quintessence, the background perfect fluid always has a EOS of  $p = w\rho$ , and change different potentials and Lagrangian. What if employing some other form of EOS ? It is very coincident that, when three superstring experts were studying the stability of black holes in brane world, they found they have to add a background matter called Chaplygin gas, to ensure the stability. Chaplygin gas is a perfect fluid with negative pressure in superstring theory. They proceeded to calculate the cosmological consequence for a FRW universe comprised of matter, CDM and chaplygin gas, and found that a flat universe with accelerating expansion is gained, and chaplygin gas not only acts as dark energy, but strongly indicates a unified description of dark energy and dark matter. Hence, Chaplygin gas is introduced as a DE candidate

in a quite natural manner.

#### 5.2.1 Special Chaplygin gas

EOS of Chaplygin gas is (all components depend on comoving coordinates):

$$p_{CG} = -\frac{A}{\rho_{CG}} \tag{5.15}$$

where  $\rho_{CG} > 0$ ,  $p_{CG} > 0$ , A is a positive constant. Chaplygin gas obeys the energymomenta equation (1.20), (1.12). Insert the EOS of Chaplygin gas into (1.12), and the integral induces:

$$\rho_{CG} = \sqrt{A + \frac{B}{a^6}} \tag{5.16}$$

Where B is an integral constant, and is selected to be positive in the following discussions; a is the scalar factor. Eq(5.16) show the behaviors:

$$a \ll \left(\frac{B}{A}\right)^{\frac{1}{6}} \longrightarrow \rho_{GC} \sim \frac{\sqrt{B}}{a^3}$$
 (5.17)

$$a \gg (\frac{B}{A})^{\frac{1}{6}} \longrightarrow \rho_{CG} \sim -p_{CG} \sim \sqrt{A}$$
 (5.18)

And we can easily draw the conclusions that, at early epoches, a is small enough to ensure  $a \ll (\frac{B}{A})^{\frac{1}{6}}$ , and density of Chaplygin gas  $\rho_{CG} \sim a^{-3}$ , which is like non-relativistic matters; lately a is large enough to ensure  $a \gg \frac{B}{A})^{\frac{1}{6}}$ , thus  $\rho_{CG} \sim -p_{CG} \sim \sqrt{A}$  and acts like the cosmological constant, which has negative pressure to drive the accelerating expansion, and the time needed to arrive at this state is:

$$t = \frac{1}{6\sqrt[4]{A}} \left( \ln \frac{\sqrt[4]{A + \frac{B}{a^6}} + \sqrt[4]{A}}{\sqrt[4]{A + \frac{B}{a^6}} - \sqrt[4]{A}} - 2 \arctan \sqrt[4]{1 + \frac{B}{Aa^6}} \right)$$
(5.19)

Because Chaplygin gas acts like dark matter and dark energy at different epoches, it indicates a unification of CDM and DE.

A more detailed perturbative form of (5.18) is:

$$\rho \sim \sqrt{A} + \sqrt{\frac{B}{4A}}a^6 \tag{5.20}$$

$$p \sim -\sqrt{A} + \sqrt{\frac{B}{4A}}a^6 \tag{5.21}$$

(5.20), (5.21) show that the expanding universe is made up of two components, the major part acts like the cosmological constant with  $p = -\rho$ ; the minor part acts as

 $p=\rho,$  whose energy density decrease sharply with the expansion.

The cosmological consequences of Chaplygin gas could be treated as a homogeneous scalar field  $\phi(t)$  together with a potential  $V(\phi)$ . Consider the Lagrangian:

$$\mathcal{L}_{GC}(\phi) = \frac{1}{2}\dot{\phi}^2 - V(\phi) \tag{5.22}$$

and set the density of the field equal to that of Chaplygin gas:

$$\rho_{\phi} = \frac{1}{2}\dot{\phi}^2 + V(\phi) = \sqrt{A + \frac{B}{a^6}}$$
(5.23)

the pressure is identical to Lagrangian:

$$p_{\phi} = \frac{1}{2}\dot{\phi}^2 - V(\phi) = -\frac{A}{\sqrt{A + \frac{B}{a^6}}}$$
(5.24)

After some algebra, the reasonable potential is:

$$V(\phi) = \frac{1}{2}\sqrt{A}\left(\cosh 3\phi + \frac{1}{\cosh 3\phi}\right)$$
(5.25)

The potential is independent of B, and it reflects the properties EOS of Chaplygin gas (5.15) only.

There are several approaches to EOS of Chaplygin gas, including:

(1) Action and Potential of Quintessence

$$V(\phi) = \frac{\sqrt{a}}{2} (\cosh 3\phi + \frac{1}{\cosh 3\phi}) \tag{5.26}$$

and in return, with the factors A, B and scale factor a in Chaplygin gas, one can re-express the kinetic energy and potential in Quintessence:

$$\dot{\phi}^2 = \frac{B}{a^6 \sqrt{A + B/a^6}}$$
(5.27)

and

$$V(\phi) = \frac{2a^6(A + B/a^6) - B}{2a^6\sqrt{A + B/a^6}}$$
(5.28)

(2)Born-Infeld Lagrangian

$$\mathcal{L} = -V_0 \sqrt{1 - \partial_\mu \partial^\mu} \tag{5.29}$$

has the induction that:

$$\rho = \frac{V_0}{\sqrt{1 - \partial_\mu \partial^\mu}} \tag{5.30}$$

$$p = -V_0 \sqrt{1 - \partial_\mu \partial^\mu} \tag{5.31}$$

which end up with Chaplygin gas EOS.

#### 5.2.2 Generalized Chaplygin gas

Firstly to generalize EOS of Chaplygin gas(GCG) into:

$$p = -\frac{A}{\rho^{\beta}} \tag{5.32}$$

where  $0 < \beta \leq 1$ .(further constraints see mage in [24] to get  $0 \leq \beta \leq 0.5$ . The following task is to find out the scalar field corresponding to (5.32)

For a complex field with non-zero mass, supposing:

$$\Phi = \left(\frac{\phi}{\sqrt{2}}m\right)\exp\left(-im\theta\right) \tag{5.33}$$

with Lagrangian of:

$$\mathcal{L}_{GCG} = g^{\mu\nu} \left( \phi^2 \theta_{,\mu} \theta_{,\nu} \right) - V(\phi^2/2) \tag{5.34}$$

where ,  $\mu$  refers to variance with spatial coordinates  $x^{\mu}$ . The condition  $\phi_{,\mu} \ll m\phi$  is used to ensure the fluctuations of energy density in spacetime, and together with the generalized EOS (5.32) and Thomas-Fermi, to rewrite its Lagrangian:

$$\mathcal{L}_{TF} = \frac{\phi^2}{2} g^{\mu\nu} \theta_{,\mu} \theta_{,\nu} \tag{5.35}$$

This acts as first principle for generalized Chaplygin gas cosmology.

From (5.32), one has a generalized (5.16):

$$\rho = \left(A + \frac{B}{a^{3(1+\beta)}}\right)^{\frac{1}{1+\beta}} \tag{5.36}$$

Energy density for generalized Chaplygin gas being:

$$\rho_{GCG} = \left(A + \frac{B}{a^{3(1+\beta)}}\right)^{\frac{1}{1+a}} \tag{5.37}$$

the effective EOS is:

$$\omega(a) = -\frac{|\omega_0|}{|\omega_0| + \frac{1 - |\omega_0|}{a^{3(1+\beta)}}}$$
(5.38)

and the perturbations (5.20), (5.21) turn to be:

$$\rho \sim A^{\frac{1}{1+\beta}} + \left(\frac{1}{1+\beta}\right) \frac{B}{A^{\frac{\beta}{1+\beta}}} a^{-3(1+\beta)}$$
(5.39)

$$p \sim -A^{\frac{1}{1+\beta}} + \left(\frac{1}{1+\beta}\right) \frac{B}{A^{\frac{\beta}{1+\beta}}} a^{-3(1+\beta)}$$
 (5.40)

Thus, as for the generalized Chaplygin gas, the expanding cosmos is also made up of two components. The major part is dark energy with  $p = \rho$  and  $A^{\frac{1}{1+\beta}}$ . The minor part

is of EOS  $p = \beta \rho$  now, and the potential for generalized Chaplygin gas with Lagrangian (5.35) is:

$$V = \frac{1}{2} \left( \Psi^{2/\beta} + \frac{A}{\Psi^2} \right) \tag{5.41}$$

where  $\psi \equiv B^{(1-\beta/1+\beta)}a^{3(1-\beta)}\phi^2$ .

There's some other generalized form, for example

$$p = -\frac{A(a)}{\rho} \tag{5.42}$$

where A(a) depends on the scale factor a, and is set positive. A modified or generalized version can be more flexible and can supply better cosmic scenario.

### 5.2.3 Generalized Chaplygin gas & Hessence

If the field is properly selected, the late behaviors of Hessence complex scalar field could be expressed via generalized Chaplygin gas. Firstly, Hessence should obey EOS (5.32):

$$p_{he} = -\frac{A}{\rho_{he}^{\beta}} \tag{5.43}$$

Via EOS of Hessence (4.27):

$$2V(\phi) = \rho_{he} - p_{he} = \rho_{he} + \frac{A}{\rho_{he}^{\beta}}$$
(5.44)

$$\dot{\phi} - \frac{Q^2}{a^6 \phi^2} = \rho_{he} + p_{he} = \rho_{he} - \frac{A}{\rho_{he}^\beta}$$
(5.45)

Friedmann Equation(1.17) for Hessence dominated

$$\left(\frac{\dot{a}}{a}\right)^2 = H^2 = \frac{8\pi G}{3}\rho_{he} \tag{5.46}$$

If the above condition holds,

$$\dot{\phi}^2 \ll \frac{Q^2}{a^6 \phi^2}, \qquad \beta = 1$$
 (5.47)

Analyzable solutions are achieved:

$$\rho_{he} = \frac{B\phi^2}{Q^2}, \qquad a = \left(\frac{BQ^4}{AQ^4 - B^2\phi^4}\right)^{1/6}$$
(5.48)

$$\dot{\phi}^2 = \frac{6\pi G B Q^2}{a^1 2 (A - B/a^6)} \tag{5.49}$$

## 5.2.4 Chaplygin gas & Superstring QFT

The origin of Chaplygin gas cosmology has shown its connections with superstring quantum field theory. Chaplygin gas is the non-relativistic limit of Born-Infeld mechanism. For excellent details, see Jackiw[52].

## Chapter 6

# Hyperbolic Cosecant Cardassian Cosmology

## 6.1 Introduction

In 2002, K. Freese and M.Lewis from Michigan University came up with a new mechanism for dark energy, where dark energy doesn't exist at all<sup>[53]</sup>! They modified Friedmann equation, or the gravitational field equation, which is of fundamental significance in cosmological dynamics, to establish a scenario where the universe is flat, matter dominated, and has accelerating expansion recently. They treated matter and radiation together (denoted as a matter term), and introduced a new term, Cardassian energy density, which depends quantitatively only on the density of the matter term, to the right side of Friedmann equation (the original discussion takes power function style Cardassian term as an example). When dated back to the baby era, the Friedmann equation would return to its normal form with no appearance of Cardassian term, and the process such as nuclearsynthesis and galaxies formation are not disturbed. Cardassian term turns dominant only recently, with redshift  $z \sim O(1)$ . In Cardassian cosmos, the critical density is remarkably less than the normal value, and can be exactly  $\Omega_{crit} = \Omega_m$  via tuning parameters. So, in Cardassian world, matter itself determines the flatness of the universe, and Cardassian term takes over to drive an accelerating expansion recently. There is no room for dark energy any more. Cardassian cosmology has survived several observational test, such as the age of the universe, CMB, etc. Initially, the origin of Cardassian term is attributed to the natural consequence when we insert the observational cosmos into a world with higher dimension.

But the work of K. Freese and M. Lewis didn't raised much attention originally. So, 7 months later Freese expressed the idea again and generalize the density term in Friedmann equation to a general form  $g(\rho)$  by the way [54]. Shortly, Gondolo and Freese put forward the relativistic fluid interpretation of Cardassian dynamics [55](denoted as GF fluid hereafter). Cardassian cosmology is raised to the hight of theory from phenomenology, and the dynamical mechanism is analysed comprehensively. This work is a milestone in the study of Cardassian cosmology, which lay the foundation for future research on this subject, and up to now, nearly all the discussions with Cardassian are based on GF fluid scenario. To ensure the positiveness of sound speed, they analysed a new kind of Cardassian term, the polytropic and a better modified polytropic style. They also come up with a proposal that the negative pressure arise from the interactions between dark matter. From this thesis on, Cardassian cosmology wins large attention and hot discussions.

In [56] it is shown that power style Cardassian violates the weak energy conditions of gravitation on small scales. Although suffering this problem, power style Cardassian, due to its simplest form, is still widely employed when new ideas are introduced. Two other important work attribute to [57](M Szydlowski and W Czaja) and [58](R Lazkoz, G Leon). The former provides an excellent example for autonomous dynamics and numerical analysis for Cardassian. The latter divides the Cardassian discussion of autonomous dynamical system into high energy limit and low energy limit two categories for the first time. The typical work to generalize Cardassian model includes the introduction of exponential Cardassian term[61]; Statefinder diagnose for modified polytropic Cardassian[62]. Generally, computations with Cardassian cosmology require the background to be homogeneous and isotropic, and thesis [63] makes an attempt to cancell the restriction of homogeneity. Meantime, numerical analysis and observational constraints for Cardassian take step [64] [65] [66] [67] [68] [69] [70] [71] [72] [73] [74] [75] [76] [77], mainly with CMB, the cosmic age, gravitational lensing and large scale exploring.

Freese and her cooperators have written several reviews on Cardassian cosmology, which does good for a glance of Cardassian, see [54] [59] [60]. And in this chapter, we will do a job similar to [61], to introduce a new model via the introduction of hyperbolic cosecant Cardassian term.

## 6.2 Fundamentals of Cardassian Cosmology

As expressed above, the following analysis is based on GF fluid scenario. In the original work, matter and radiation are treated together for a matter energy density term and the independent variable of Cardassian term. The following analysis will change into two cases. This treatment is reasonable, because Cardassian term is effective only recently  $(z \sim O(1))$  when matter is far more enriching than radiation, and radiation is considerable and makes a difference only in the early era, when the Cardassian modification is neglected. But for a clearer scenario or as a generalization, to treat matter and radiation separately is also a good choice. So in this text two cases are discussed, namely treating matter and radiation together as a matter term and treating them separately.

#### 6.2.1 Fundamentals for Implicit Cardassian term

In Cardassian cosmology, the total effective energy density is [53][54][55]:

$$\rho' = g(\rho) \tag{6.1}$$

where  $\rho = \rho_m + \rho_\gamma$  reflects the summary of energy density of matter  $\rho_m$  and energy density of radiation  $\rho_\gamma$ , which are testable parameters. From the first law and second law of thermaldynamics, we have  $Td(sV) = d(\rho V) + pdV$ ; for adiabatic expanding d(sV) = 0; conservation of particles' number  $d(\rho_m V) = 0$ ; conservation of radiation  $d(\rho_\gamma V^{\frac{4}{3}}) = 0$ . Insert these relations into (6.1) and we could get the total pressure:

$$p' = -\rho' + \rho_m \frac{\partial \rho'}{\partial \rho_m} + \frac{4}{3}\rho_\gamma \frac{\partial \rho'}{\partial \rho_\gamma}$$
(6.2)

In Cardassian cosmos, Friedmann equation reads:

$$H^{2} = \frac{8\pi G}{3}g(\rho)$$
(6.3)

and Raychaudhuri equation reads:

$$\dot{H} = -4\pi G \left(\rho_m \frac{\partial \rho'}{\partial \rho_m} + \frac{4}{3} \rho_\gamma \frac{\partial \rho'}{\partial \rho_\gamma}\right) \tag{6.4}$$

the energy-momentum conservation equation:

$$\dot{\rho}' = -3H(\rho' + p') \tag{6.5}$$

The above equations are the fundamental equations for Cardassian cosmology.

#### 6.2.2 Fundamentals for Explicit Cardassian term

Firstly dividing  $\rho' = g(\rho)$  into two terms:

$$\rho' = \rho + f(\rho) \tag{6.6}$$

The first term  $\rho$  is the matter and radiation term, and  $f(\rho) \equiv g(\rho) - \rho$  is the explicit Cardassian term, which is the modification term of Friedmann equation. When matter and radiation are treated separately:

$$\rho' = \rho_m + \rho_\gamma + f(\rho_m, \rho_\gamma) \tag{6.7}$$

Thus from (6.2) one gets the total energy:

$$p' = \frac{\rho_{\gamma}}{3} + \rho_m \frac{\partial f}{\partial \rho_m} + \frac{4}{3}\rho_{\gamma} \frac{\partial f}{\partial \rho_{\gamma}} - f$$
(6.8)

and from this equation, we can easily see that only radiation and Cardassian term contribute to the total pressure, or EOS parameter for matter is w = 0. This conclusion will be quoted in next chapter for virilization. Friedmann equation and Raychaudhuri equation are

$$H^{2} = \frac{8\pi G}{3} [\rho_{m} + \rho_{\gamma} + f(\rho_{m}, \rho_{\gamma})]$$
(6.9)

and

$$\dot{H} = -4\pi G(\rho_m + \frac{4}{3}\rho_\gamma + \rho_m \frac{\partial f}{\partial \rho_m} + \frac{4}{3}\rho_\gamma \frac{\partial f}{\partial \rho_\gamma})$$
(6.10)

And energy conservation is equal to the conservation of particles' number and conservation of radiation:

$$\dot{\rho}_m = -3H\rho_m \tag{6.11}$$

$$\dot{\rho}_{\gamma} = -4H\rho_{\gamma} \tag{6.12}$$

## 6.2.3 Phase Space, Autonomous Equations for Separate Matter & Radiation

Based on Friedmann equation, we define

$$1 = \frac{8\pi G}{3H^2} [\rho_m + \rho_\gamma + f(\rho_m, \rho_\gamma)]$$

$$\equiv \Omega_m + \Omega_\gamma + \Omega_{card}$$
(6.13)

where  $\Omega_m$ ,  $\Omega_\gamma$ ,  $\Omega_{card}$  are the relative density parameter of matter, radiation, Cardassian term respectively. Now we introduce the non-dimensional parameters (with  $\kappa^2 = 8\pi G$ ):

$$x = \frac{\kappa \sqrt{\rho_m}}{\sqrt{3}H}, \qquad y = \frac{\kappa \sqrt{\rho_\gamma}}{\sqrt{3}H}, \qquad z = \frac{\kappa \sqrt{f}}{\sqrt{3}H}, \qquad N = \ln a$$
(6.14)

where N is e-folding time. Apparently N keeps the same pace with the cosmological time in our flat, expanding FRW universe. Those newly introduced variables form the 4 dimensional phase space, and when x,y,z, N are inserted into the basic dynamical equations, they will turn to the geometric curves in the phase space by:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}x^3 + 2xy^2 + \frac{3}{2}x^3\frac{\partial f}{\partial\rho_m} + 2xy^2\frac{\partial f}{\partial\rho_\gamma}$$
(6.15a)

$$\frac{dy}{dN} = -2y + \frac{3}{2}yx^2 + 2y^3 + \frac{3}{2}yx^2\frac{\partial f}{\partial\rho_m} + 2y^3\frac{\partial f}{\partial\rho_\gamma}$$
(6.15b)

$$\frac{dz}{dN} = \frac{3}{2}x^2\left(z - \frac{1}{z}\right)\frac{\partial f}{\partial\rho_m} + 2y^2\left(z - \frac{1}{z}\right)\frac{\partial f}{\partial\rho_\gamma} + \frac{3}{2}x^2z + 2y^2z \tag{6.15c}$$

These curves are called the differential autonomous equations of the system. For further discussion, we treat the cosmos as flat (the confirmed result by CMB etc), then there will be the restriction  $x^2 + y^2 + z^2 = 1$ , which simplify the equations (6.15) above into:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}x^3 + 2xy^2 + \frac{3}{2}x^3\frac{\partial f}{\partial\rho_m} + 2xy^2\frac{\partial f}{\partial\rho_\gamma}$$
(6.16a)

$$\frac{dy}{dN} = -2y + \frac{3}{2}yx^2 + 2y^3 + \frac{3}{2}yx^2\frac{\partial f}{\partial\rho_m} + 2y^3\frac{\partial f}{\partial\rho_\gamma}$$
(6.16b)

Particularly, if the Cardassian term has the symmetry  $f(\rho_m, \rho_\gamma) = f(\rho_m + \rho_\gamma)$ , the equations (6.16) can be further reduced into:

$$\frac{dx}{dN} = -\frac{3}{2}x + \left(\frac{3}{2}x^3 + 2xy^2\right)\left(1 + \frac{\partial f}{\partial\rho_m}\right)$$
(6.17a)

$$\frac{dy}{dN} = -2y + \left(\frac{3}{2}x^2y + 2y^3\right)\left(1 + \frac{\partial f}{\partial\rho_m}\right)$$
(6.17b)

This case is important, because the hyperbolic cosecant Cardassian term, as we will put forward later, just has such a lovely character.

So, once  $f(\rho_m, \rho_\gamma)$  is known, we will master the dynamics of the Cardassian universe, and the proximation behaviors of the corresponding autonomous equations in phase space.

#### 6.2.4 Unitary Treatment of Matter & Radiation

If matter and radiation are treated together as a matter term, or only take matter into account for recent universe, the energy total density will be

$$\rho' = \rho_m + f(\rho_m) \tag{6.18}$$

From(6.2) one gets:

$$p' = \rho_m \frac{\partial f}{\partial \rho_m} - f \tag{6.19}$$

Friedmann equation and Raychaudhuri are separately:

$$H^{2} = \frac{8\pi G}{3} [\rho_{m} + f(\rho_{m})]$$
(6.20)

and

$$\dot{H} = -4\pi G\rho_m (1 + \frac{\partial f}{\partial \rho_m}) \tag{6.21}$$

Now the energy conservation equation  $\dot{\rho}' = -3H(\rho' + p')$  is equal to the conservation of particles' number  $\dot{\rho_m} = -3H\rho_m$ . When matter and radiation is treated together, there's convenient form for the speed of sound, which should be positive:

$$c_s^2 = \frac{\partial p'}{\partial \rho'} = \rho_m \frac{\frac{\partial^2 \rho'}{\partial \rho_m^2}}{\frac{\partial \rho'}{\partial \rho_m}}$$
(6.22)

Based on Friedmann equation, we define

$$1 = \frac{8\pi G}{3H^2} [\rho_m + f(\rho_m)]$$

$$\equiv \Omega_m + \Omega_{card}$$
(6.23)

and introduce the new variables  $(\kappa^2 = 8\pi G)$ :

$$x = \frac{\kappa \sqrt{\rho_m}}{\sqrt{3}H}, \qquad y = \frac{\kappa \sqrt{f}}{\sqrt{3}H}$$
 (6.24)

Together with e-folding time this leads to a 3-dimensional phase space. Insert x, y into the equations above to get:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}\left(1 + \frac{\partial f}{\partial \rho_m}\right) \tag{6.25a}$$

$$\frac{dy}{dN} = -\frac{3}{2}\frac{x^2}{y}\frac{\partial f}{\partial \rho_m} + \frac{3}{2}x^2y\left(1 + \frac{\partial f}{\partial \rho_m}\right)$$
(6.25b)

with constraint  $x^2 + y^2 = 1$  for flat FRW space, the number of independent equations reduces to one:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}\left(1 + \frac{\partial f}{\partial\rho_m}\right) \tag{6.26}$$

Also, we know the universe if we know  $f(\rho_m)$ . And for simple Cardassian terms, we can even get the exact and analyzable solution.

#### 6.2.5 Constraints for Cardassian & Hyperbolic cosecant Cardassian

Based on the general discussion above, we can make detailed analysis with certain form of Cardassian term. Yet the Cardassian term should not be arbitrary, as has been shown in the above algebra, and a potential candidate must obey at least three constraints [61]:

(1) As for the baby universe with  $\rho_{crit,0} \ll \rho$ , total energy density  $g(\rho)$  must reduces to  $\rho$ ; or Cardassian term  $f(\rho)$  vanishes.

(2)Cardassian term is dominant recently with redshift  $z \sim O(1)$ , when  $g(\rho)$  differs from  $\rho$  significantly, or  $f(\rho)$  differs from null significantly.

(3)The speed of sound  $c_s^2 > 0$  must be positive.

The constraints is in fact powerful. Only those that satisfies the first condition is worth further algebra during which the second and third condition proceeds to be tested. Three set of reasonable Cardassian terms have been introduced before:

(1)Power-style Cardassian[53]

$$\rho' = \rho + b\rho^n \tag{6.27}$$

where b and n is tuning parameters, and n < 2/3. (2)Polytropic and Modified Cardassian[55]

$$\rho' = \rho_{card} \left[ 1 + \left( \frac{\rho_m}{\rho_{card}} \right)^q \right]^{\frac{1}{q}}$$
(6.28a)

$$\rho' = \rho_{card} \left[ 1 + \left( \frac{\rho_{card}}{\rho_m} \right)^{qv} \right]^{\frac{1}{q}}$$
(6.28b)

where q and v is tuning parameters;  $\rho_{card}$  a constant density value for reference. (3)Exponential and Modified Cardassian[61]

$$\rho' = \rho \exp\left(\frac{\rho_{card}}{\rho}\right)^n \tag{6.29a}$$

$$\rho' = (\rho + \rho_{card}) \exp\left(\frac{q\rho_{card}}{\rho + \rho_{card}}\right)^n \tag{6.29b}$$

where q and n is tuning parameters.

Next we will discuss a new Cardassian term, namely hyperbolic cosecant Cardassian term:

$$\rho' = \rho \left( 1 + b \cdot csch \left( \frac{\rho}{\rho_{card}} \right)^n \right)$$
(6.30)

where b, n are tuning parameters and

$$f(\rho) = \rho b \cdot csch \left(\frac{\rho}{\rho_{card}}\right)^n \tag{6.31}$$

Hyperbolic cosecant function differs very much from double exponential potential function. Before further discussion, it's a convenience to review the properties of hyperbolic cosecant function.

### 6.3 Fundamentals of Hyperbolic Cosecant Function

Hyperbolic sine sh(x) and hyperbolic cosine ch(x) are the two basic functions which are orthogonal to each other in the hyperbolic function space, and are defined as:

$$sh(x) = \frac{e^x - e^{-x}}{2}$$
  $ch(x) = \frac{e^x + e^{-x}}{2}$  (6.32)

And definition of hyperbolic cosecant function is:

$$csch(x) = \frac{1}{sh(x)} = \frac{2}{e^x - e^{-x}}$$
 (6.33)

Its one order derivative is:

$$\frac{dcsch(x)}{dx} = -csch(x)cth(x) \tag{6.34}$$

Where cth(x) is the hyperbolic cotangent function:

$$cth(x) = \frac{ch(x)}{sh(x)} = \frac{e^x + e^{-x}}{e^x - e^{-x}}$$
(6.35)

The hyperbolic cotangent function can be expressed with hyperbolic cosecant function:

$$cth(x) = \sqrt{1 + csch^2(x)} \tag{6.36}$$

The inverse function of hyperbolic cosecant is denoted as Arcsch(x), and certainly csch(Arcsch(x)) = x.

## 6.4 Csch Cardassian for unitary matter and radiation

We treat matter and radiation together for the matter term, denoted as  $\rho_m$ , and the total energy density of hyperbolic cosecant (hereafter donated as Csch) Cardassian universe is, as above:

$$\rho' = \rho_m \left( 1 + b \cdot csch \left( \frac{\rho_m}{\rho_{card}} \right)^n \right)$$
(6.37)

with the Cardassian term:

$$f(\rho) = \rho_m b \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n \tag{6.38}$$

n and b are tuning parameters, which need observational constraints.  $\rho_{card}$  is a constant density value for reference, as above in exponential Cardassian. when  $\rho_m \to \infty$ ,  $\rho' \to \rho_m$ , thus the modified Friedmann equation returns to the normal form, and

$$\frac{\partial f}{\partial \rho_m} = b \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n - nb \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n \cdot cth \left(\frac{\rho_m}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m}{\rho_{card}}\right)^n \tag{6.39}$$

Hence from (6.18) we get the pressure that:

$$p = \rho_m \frac{\partial f}{\partial \rho_m} - f$$

$$= \rho_m b \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n - n\rho_m b \cdot csch \left(\frac{\rho_m}{\rho_{crit}}\right)^n \cdot cth \left(\frac{\rho_m}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m}{\rho_{card}}\right)^n - \rho_m b \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n$$

$$= -bn\rho_m \cdot csch \left(\frac{\rho_m}{\rho_{crit}}\right)^n \cdot cth \left(\frac{\rho_m}{\rho_{crit}}\right)^n \cdot \left(\frac{\rho_m}{\rho_{crit}}\right)^n$$
(6.40)

the Friedmann quation and Raychaudhuri equation in Csch cosmos is:

$$H^{2} = \frac{8\pi G}{3} \left[ \rho_{m} + \rho_{m} b \cdot csch \left(\frac{\rho_{m}}{\rho_{card}}\right)^{n} \right]$$
(6.41)

and

$$\dot{H} = -4\pi G \rho_m \left(1 + \frac{\partial f}{\partial \rho_m}\right)$$
$$= -4\pi G \rho_m \left[1 + b \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n - nb \cdot csch \left(\frac{\rho_m}{\rho_{card}}\right)^n cth \left(\frac{\rho_m}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m}{\rho_{card}}\right)^n\right]$$
(6.42)

And the energy conservation equation  $\dot{\rho'} = -3H(\rho' + p')$  is equal to the conservation of particles' number  $\dot{\rho}_m = -3H\rho_m$ , After the algebra above, we have known the kinetic and dynamical equations of Csch Cardassian cosmos when matter and radiation are treated together. Then we will research into its phase space and autonomous equations. We define:

$$1 = \frac{8\pi G}{3H^2} [\rho_m + f(\rho_m)]$$

$$\equiv \Omega_m + \Omega_{card}$$
(6.43)

and introduce new variables  $\kappa^2 = 8\pi G$ :

$$x = \frac{\kappa \sqrt{\rho_m}}{\sqrt{3}H}, \qquad y = \frac{\kappa \sqrt{f}}{\sqrt{3}H}$$
 (6.44)

thus (6.39) can be rewritten as:

$$\frac{\partial f}{\partial \rho} = \frac{y^2}{x^2} - n\frac{y^2}{x^2} \cdot \sqrt{1 + \left(\frac{y^2}{bx^2}\right)^2} \cdot Arcsch\frac{y^2}{bx^2} \tag{6.45}$$

According to (6.25), we get the autonomous equations:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}\left(1 + \frac{\partial f}{\partial\rho_m}\right)$$
$$= -\frac{3}{2}x + \frac{3}{2}\left[\frac{y^2}{x^2} - n\frac{y^2}{x^2} \cdot \sqrt{1 + \left(\frac{y^2}{bx^2}\right)^2} \cdot Arcsch\frac{y^2}{bx^2}\right]$$
(6.46a)

$$\frac{dy}{dN} = -\frac{3}{2} \frac{x^2}{y} \frac{\partial f}{\partial \rho_m} + \frac{3}{2} x^2 y \left(1 + \frac{\partial f}{\partial \rho_m}\right)$$
$$= -\frac{3}{2} \left[\frac{y^2}{x^2} - n\frac{y^2}{x^2} \cdot \sqrt{1 + \left(\frac{y^2}{bx^2}\right)^2} \cdot Arcsch\frac{y^2}{bx^2}\right]$$
$$+ \frac{3}{2} x^2 y \left[1 + \frac{y^2}{x^2} - n\frac{y^2}{x^2} \cdot \sqrt{1 + \left(\frac{y^2}{bx^2}\right)^2} \cdot Arcsch\frac{y^2}{bx^2}\right]$$
(6.46b)

For flat FRW universe,  $\Omega_m + \Omega_{card} = 1$ , and there's the constraining equation  $x^2 + y^2 = 1$ . So(6.46) can be reduced into:

$$\frac{dx}{dN} = -\frac{3}{2}x + \frac{3}{2}\left[\frac{1}{x^2} - 1 - \left(\frac{1}{x^2} - 1\right) \cdot \sqrt{1 + \left(\frac{1 - x^2}{bx^2}\right)^2} \cdot Arcsch\frac{1 - x^2}{bx^2}\right]$$
$$= -\frac{3}{2}x^3 - \frac{3}{2}x(1 - x^2) \cdot \sqrt{1 + \left(\frac{1 - x^2}{bx^2}\right)^2} \cdot Arcsch\frac{1 - x^2}{bx^2} \tag{6.47}$$

This is the very differential autonomous equation for Csch Cardassian cosmos with unitary matter and radiation.

## 6.5 Csch Cardassian for separate matter and radiation

#### 6.5.1 Dynamical Equations

In this section the energy density of matter and radiation will be treat separately, so the total energy density is:

$$\rho' = (\rho_m + \rho_\gamma) \left[ 1 + b \cdot csch \left( \frac{\rho_m + \rho_\gamma}{\rho_{card}} \right)^n \right]$$
(6.48)

thus the Csch Cardassian term is:

$$f(\rho_m, \rho_\gamma) = f(\rho_m + \rho_\gamma) = b(\rho_m + \rho_\gamma) \cdot csch\left(\frac{\rho_m + \rho_\gamma}{\rho_{card}}\right)^n \tag{6.49}$$

As for Csch function, the symmetric property  $csch(\rho_m, \rho_\gamma) = csch(\rho_m + \rho_\gamma)$  holds, so the matter term and radiation term are symmetric mathematically to each other.

$$\frac{\partial f}{\partial \rho_m} = \frac{\partial f}{\partial \rho_\gamma} = b \cdot csch \left(\frac{\rho_m + \rho_\gamma}{\rho_{card}}\right)^n - nb \cdot csch \left(\frac{\rho_m + \rho_\gamma}{\rho_{card}}\right)^n \cdot cth \left(\frac{\rho_m + \rho_\gamma}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m + \rho_\gamma}{\rho_{card}}\right)^n \tag{6.50}$$

According to 6.8 the total pressure is:

$$p' = \frac{\rho_{\gamma}}{3} + \rho_m \frac{\partial f}{\partial \rho_m} + \frac{4}{3}\rho_{\gamma} \frac{\partial f}{\partial \rho_{\gamma}} - f$$

$$= \frac{\rho_{\gamma}}{3} + (\rho_m + \frac{4}{3}\rho_{\gamma})\frac{\partial f}{\partial\rho_m} - f$$

$$= \frac{\rho_{\gamma}}{3} + (\rho_m + \frac{4}{3}\rho_{\gamma})b \cdot \operatorname{csch}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n$$

$$-(\rho_m + \frac{4}{3}\rho_{\gamma})nb \cdot \operatorname{csch}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n \cdot \operatorname{cth}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n$$

$$-(\rho_m + \rho_{\gamma})b(\rho_m + \rho_{\gamma}) \cdot \operatorname{csch}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n$$

$$= \frac{\rho_{\gamma}}{3} + \frac{\rho_{\gamma}}{3}b \cdot \operatorname{csch}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n$$

$$-(\rho_m + \frac{4}{3}\rho_{\gamma}) \cdot nb \cdot \operatorname{csch}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n \cdot \operatorname{cth}\left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n \cdot \left(\frac{\rho_m + \rho_{\gamma}}{\rho_{card}}\right)^n$$
(6.51)

With Friedmann equation and Raychaudhuri equation being:

$$H^{2} = \frac{8\pi G}{3} [\rho_{m} + \rho_{\gamma} + f(\rho_{m}, \rho_{\gamma})]$$
(6.52)

and

$$\dot{H} = -4\pi G(\rho_m + \frac{4}{3}\rho_\gamma + \rho_m \frac{\partial f}{\partial \rho_m} + \frac{4}{3}\rho_\gamma \frac{\partial f}{\partial \rho_\gamma})$$
(6.53)

Here the energy conservation means the conservation of particles' number and the radiation conservation:

$$\dot{\rho}_m = -3H\rho_m \tag{6.54}$$

$$\dot{\rho}_{\gamma} = -4H\rho_{\gamma} \tag{6.55}$$

Now we have got the dynamical equations for Csch Cardassian cosmos with the radiation term and matter term treated separately. Then we'll investigate the autonomous equations in its phase space.

## 6.5.2 Phase Space and Autonomous Equations

From Friedmann equations, we define:

$$1 = \frac{8\pi G}{3H^2} [\rho_m + \rho_\gamma + f(\rho_m, \rho_\gamma)]$$

$$\equiv \Omega_m + \Omega_\gamma + \Omega_{card}$$
(6.56)

where  $\Omega_m, \Omega_\gamma, \Omega_{card}$  is the relative energy density for matter, radiation and Csch Cardassian term respectively. Further more we introduce new variables with  $\kappa^2 = 8\pi G$ :

$$x = \frac{\kappa \sqrt{\rho_m}}{\sqrt{3H}}, \qquad y = \frac{\kappa \sqrt{\rho_\gamma}}{\sqrt{3H}}, \qquad z = \frac{\kappa \sqrt{f}}{\sqrt{3H}}, \qquad N = \ln a$$
(6.57)

Thus (6.50) reads:

$$\frac{\partial f}{\partial \rho_m} = \frac{\partial f}{\partial \rho_\gamma} = \frac{z^2}{x^2 + y^2} - n \cdot \frac{z^2}{x^2 + y^2} \cdot \sqrt{1 + \left(\frac{z^2}{b(x^2 + y^2)}\right)^2} \cdot \operatorname{Arcsch} \frac{z^2}{b(x^2 + y^2)} \tag{6.58}$$

For flat FRW cosmos with  $\Omega_m + \Omega_{\gamma} + \Omega_{card}$ , or  $x^2 + y^2 + z^2 = 1$ , combined with the symmetry,  $f(\rho_m, \rho_{\gamma}) = f(\rho_m + \rho_{\gamma})$ , according to (6.17), we have the autonomous equations:

$$\frac{dx}{dN} = -\frac{3}{2}x + \left(\frac{3}{2}x^3 + 2xy^2\right) \left(1 + \frac{z^2}{x^2 + y^2}\right)$$
(6.59a)  
+  $\left(\frac{3}{2}x^3 + 2xy^2\right) \left[-n\frac{z^2}{x^2 + y^2}\sqrt{1 + \left(\frac{1 - x^2 - y^2}{b(x^2 + y^2)}\right)^2} \cdot \operatorname{Arcsch}\frac{1 - x^2 - y^2}{b(x^2 + y^2)}\right]$   
 $\frac{dy}{dN} = -2y + \left(\frac{3}{2}x^2y + 2y^3\right)\frac{1 - x^2 - y^2}{x^2 + y^2}$ (6.59b)  
-  $\left(\frac{3}{2}x^2y + 2y^3\right) \left[n \cdot \frac{1 - x^2 - y^2}{x^2 + y^2} \cdot \sqrt{1 + \left(\frac{1 - x^2 - y^2}{b(x^2 + y^2)}\right)^2} \cdot \operatorname{Arcsch}\frac{1 - x^2 - y^2}{b(x^2 + y^2)}\right]$ 

#### 6.5.3 Critical Points

From the above analysis, we get the autonomous equations in Csch Cardassian for separate treatment of matter and energy density. The following work is to get the critical points. and we will show there are three and only three critical points, namely:(0,0)(1,0)(0,1).

Obviously, 1 > x > 0, 1 > y > 0 and  $1 > x^2 + y^2$  hold, firstly let's introduce a new variable:

$$\phi = \frac{1 - (x^2 + y^2)}{x^2 + y^2} > 0 \tag{6.60}$$

and formulate a new function of:

$$\Psi = \phi - n\phi\sqrt{1 + \phi^2/b^2} \cdot Arcsch\frac{\phi}{b}$$
(6.61)

For (6.61), when  $(x, y) \to (0, 0)$ ,  $\phi \to +\infty$ ,  $n\phi\sqrt{1+\phi^2/b^2} \to \infty$ ,  $Arcsch\frac{\phi}{b} \to 0$ ; and when  $(x, y) \to (1, 0)$  or (0, 1),  $\phi \to 0$ ,  $n\phi\sqrt{1+\phi^2/b^2} \to 0$ ,  $Arcsch\frac{\phi}{b} \to \infty$ , The limit of  $\Psi$  is impossible to be read directly in both approximation cases. So we transform  $\Psi$ and employ the familiar Los'pital theorem (prime operator (')refers to derivative of the dependent variable to the independent variable):

$$\Psi = \phi - \frac{Arcsch\frac{\phi}{b}}{\frac{1}{n\phi\sqrt{1+\phi^2/b^2}}}$$

$$= \phi - \frac{\left(Arcsch\frac{\phi}{b}\right)'}{\left(\frac{1}{n\phi\sqrt{1+\phi^{2}/b^{2}}}\right)'} \\ = \phi - \frac{\frac{n^{2}\phi^{4}}{b^{2}} + n^{2}\phi^{2}}{\frac{n\phi^{3}}{b^{3}} + \frac{n\phi^{2}}{b^{3}} + \frac{n\phi}{b}}$$
(6.62)

Based on this equation, let's consider two approximation cases: (1) When  $(x, y) \to (0, 0), \phi \to \infty$ :

$$\Psi = \phi - \phi \frac{\frac{n^2 \phi^3}{b^2} + n^2 \phi}{\frac{n \phi^3}{b^3} + \frac{n \phi^2}{b^3} + \frac{n \phi}{b}}$$

$$\sim \phi - nb \cdot \phi$$
(6.63)

So the infinitesimal  $\operatorname{Arcsch}^{\phi}_{\overline{b}}$  is magnified by the infinity  $n\phi\sqrt{1+\phi^2/b^2}$  to get an unitary infinity  $n\phi\sqrt{1+\phi^2/b^2}\cdot\operatorname{Arcsch}^{\phi}_{\overline{b}}$ , which is coincidently of the same order with the infinity  $\phi$ , who will conduct a minus operation with it. The minus of two infinity of the same order can leads to conservance ! And we put  $\phi$  back to the original equation, and when  $(x, y) \to (0, 0)$ , we have:

$$\frac{dx}{dN} \sim \left[\frac{3}{2}x^3 + 2xy^2\right] \cdot (1 - nb)\frac{1 - (x^2 + y^2)}{x^2 + y^2}$$

$$> \left[\frac{3}{2}x^3 + \frac{3}{2}xy^2\right] \cdot (1 - nb)\frac{1 - (x^2 + y^2)}{x^2 + y^2}$$

$$\sim \frac{3}{2}x \cdot (1 - \frac{b^2}{2}n)[1 - (x^2 + y^2)] \sim 0$$
(6.64)

It's notable that  $\Psi$  is conserving itself if we put 1 - nb = 0. However, this constraint is too strong and we will have much less room left for parameter tuning. Considering there is an infinitesimal factor waiting to multiply  $\Psi$ , we'd better realize the conservation in the next step, rather than set 1 - nb = 0 in a hurry in advance. Hence (0, 0) is a critical point of this autonomous system.

(2)When  $(x, y) \to (1, 0)$  or  $(0, 1), \phi \to 0$ , in

$$\frac{\frac{n^2\phi^4}{b^2} + n^2\phi^2}{\frac{n\phi^3}{b^3} + \frac{n\phi^2}{b^3} + \frac{n\phi}{b}}$$

the numerator is higher order infinitesimal compared to the denominator, so the limit is zero, and:

$$\Psi = \phi - \frac{\frac{n^2 \phi^4}{b^2} + n^2 \phi^2}{\frac{n \phi^3}{b^3} + \frac{n \phi^2}{b^3} + \frac{n \phi}{b}}$$

whose limit is also zero. Now let's go back to analyze the original equation:

$$\frac{dy}{dN} \sim -2y + \left[\frac{3}{2}x^2y + 2y^3\right] \left[1 + \phi - \frac{\frac{n^2\phi^4}{b^2} + n^2\phi^2}{\frac{n\phi^3}{b^3} + \frac{n\phi^2}{b^3} + \frac{n\phi}{b}}\right]$$
$$\sim -2y + \frac{3}{2}x^2y + 2y^3 \sim 0 \tag{6.65}$$

Hence (1,0),(0,1) are both critical points of the autonomous system.

All in all, (0,0), (1,0), (0,1) are three critical points of this autonomous system. And because the critical points are numerically achieved via solving the group of two three-order algebraic equations, according to Gauss's basic theorem of arithmetics, we could have only three roots in the complex domain. So the three critical points achieved via approximation analyzing are the only ones.

#### 6.5.4 Stability Matrix for Autonomous System

Now we'll make phenomenological analysis of the stability of the critical points. Set  $X = \frac{dx}{dN}$ ,  $Y = \frac{dy}{dN}$ , and the perturbation matrix will be:

$$\mathcal{M} = \begin{pmatrix} \frac{\partial X}{\partial x} & \frac{\partial X}{\partial y} \\ \frac{\partial Y}{\partial x} & \frac{\partial Y}{\partial y} \end{pmatrix}$$
(6.66)

The four elements of this Jacobian matrix is:

$$\frac{\partial X}{\partial x} = -\frac{3}{2} + \left(\frac{9}{2}x^2 + 2y^2\right) \left[1 + \frac{1 - (x^2 + y^2)}{x^2 + y^2}\right]$$
$$- \left(\frac{9}{2}x^2 + 2y^2\right) \left[n\frac{1 - (x^2 + y^2)}{x^2 + y^2}\sqrt{1 + \left(\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}\right)^2}Arcsch\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}\right]$$
$$+ \left(\frac{3}{2}x^3 + 2xy^2\right)n\frac{2x}{(x^2 + y^2)^2}\sqrt{1 + \left(\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}\right)^2}Arcsch\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}$$
$$+ n\left(\frac{3}{2}x^3 + 2xy^2\right)\frac{(1 - (x^2 + y^2))^2}{(x^2 + y^2)^4}\frac{2x}{b^2\sqrt{1 + \frac{1}{b^2}(\frac{1}{x^2 + y^2} - 1)^2}}Arcsch\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}$$
$$- \frac{2x}{(x^2 + y^2)^2} \cdot \left(\frac{3}{2}x^3 + 2xy^2\right) + \left(\frac{3}{2}x^3 + 2xy^2\right)n\frac{2x}{(x^2 + y^2)^2}$$
(6.67)

and

$$\frac{\partial X}{\partial y} = 4xy \left[ 1 + \frac{1 - (x^2 + y^2)}{x^2 + y^2} \right]$$
$$-4xy \cdot n \frac{1 - (x^2 + y^2)}{x^2 + y^2} \sqrt{1 + \left(\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}\right)^2} Arcsch \frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}$$

$$+\left(\frac{3}{2}x^{3}+2xy^{2}\right)n\frac{2y}{(x^{2}+y^{2})^{2}}\sqrt{1+\left(\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}\right)^{2}}Arcsch\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}$$

$$+\left(\frac{3}{2}x^{3}+2xy^{2}\right)n\frac{(1-(x^{2}+y^{2}))^{2}}{(x^{2}+y^{2})^{4}}\frac{2y}{b^{2}\sqrt{1+\frac{1}{b^{2}}(\frac{1}{x^{2}+y^{2}}-1)^{2}}}Arcsch\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}$$

$$-\frac{2y}{(x^{2}+y^{2})^{2}}\left(\frac{3}{2}x^{3}+2xy^{2}\right)+\left(\frac{3}{2}x^{3}+2xy^{2}\right)n\frac{2y}{(x^{2}+y^{2})^{2}}$$
(6.68)
and
$$\frac{\partial Y}{\partial x}=3xy\left[1+\frac{1-(x^{2}+y^{2})}{x^{2}+x^{2}}\right]$$

$$\frac{\partial x}{\partial x} = \left[ \frac{1}{x^2 + y^2} \right]$$
$$-3xy \cdot n \frac{1 - (x^2 + y^2)}{x^2 + y^2} \sqrt{1 + \left(\frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}\right)^2} Arcsch \frac{1 - (x^2 + y^2)}{b(x^2 + y^2)}$$

$$+\left(\frac{3}{2}x^{2}y+2y^{3}\right)n\frac{2x}{(x^{2}+y^{2})^{2}}\sqrt{1+\left(\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}\right)^{2}}Arcsch\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}$$

$$+\left(\frac{3}{2}x^{2}y+2y^{3}\right)n\frac{(1-(x^{2}+y^{2}))^{2}}{(x^{2}+y^{2})^{4}}\frac{2x}{b^{2}\sqrt{1+\frac{1}{b^{2}}(\frac{1}{x^{2}+y^{2}}-1)^{2}}}Arcsch\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}$$
$$-\frac{2x}{(x^{2}+y^{2})^{2}}\left(\frac{3}{2}x^{2}y+2y^{3}\right)+n\left(\frac{3}{2}x^{2}y+2y^{3}\right)\frac{2x}{(x^{2}+y^{2})^{2}}\tag{6.69}$$

and

$$\frac{\partial Y}{\partial y} = -2 + \left(-\frac{3}{2}x^2 + 6y^2\right) \left[1 + \frac{1 - (x^2 + y^2)}{x^2 + y^2}\right]$$

$$-\left(-\frac{3}{2}x^2+6y^2\right)n\frac{1-(x^2+y^2)}{x^2+y^2}\sqrt{1+\left(\frac{1-(x^2+y^2)}{b(x^2+y^2)}\right)^2}Arcsch\frac{1-(x^2+y^2)}{b(x^2+y^2)}$$

$$+\left(\frac{3}{2}x^{2}y+2y^{3}\right)n\frac{2y}{(x^{2}+y^{2})^{2}}\sqrt{1+\left(\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}\right)^{2}}Arcsch\frac{1-(x^{2}+y^{2})}{b(x^{2}+y^{2})}$$

$$+ \left(\frac{3}{2}x^{2}y + 2y^{3}\right)n\frac{(1 - (x^{2} + y^{2}))^{2}}{(x^{2} + y^{2})^{4}}\frac{2x}{b^{2}\sqrt{1 + \frac{1}{b^{2}}(\frac{1}{x^{2} + y^{2}} - 1)^{2}}}Arcsch\frac{1 - (x^{2} + y^{2})}{b(x^{2} + y^{2})}$$
$$- \frac{2y}{(x^{2} + y^{2})^{2}}\left(\frac{3}{2}x^{2}y + 2y^{3}\right) + \left(\frac{3}{2}x^{2}y + 2y^{3}\right)n\frac{2y}{(x^{2} + y^{2})^{2}}\tag{6.70}$$

## Chapter 7

# Virial Collapse of Power-style Cardassian

## 7.1 Applications of Virialization in Cosmology

The spherical collapse mechanism, which was put forward by Gunn and Gott [82] [83], is an effective tool in understanding the consequence of the density inhomogeneity. An overdensed volume will slow down from the co-expansion, until the expanding velocity vanishes. Then it starts to collapse over rotating, and potential energy translates into rotating kinetic energy. The special state when the expanding just stops and collapse is to start is dubbed turn around state. Ultimately the collapse ends up with stable virial structure.

We can apply the virial theorem to the above process in the light of energy conservation. A general statement of virial theorem says that, during a long period of time, the opposite value of the average kinetic energy of a particle group equals to the virial force acting on it:

$$\bar{T} = -\frac{1}{2} \sum_{i} \vec{F_i} \cdot \vec{r_i} \tag{7.1}$$

where  $\overline{T}$  is the total rotating kinetic energy,  $\vec{F_i}$  and  $\vec{r_i}$  refer to the external forces and location vector of the *i*th particle. For conserving system,

$$\bar{T} = \frac{1}{2} \sum_{i} (\bar{\nabla_i V}) \cdot \vec{\eta}_i \tag{7.2}$$

where V is the potential of the particle group,  $\nabla$  is the vectoring gradient operator:

$$\nabla_i = \frac{\partial}{\partial x_i}\hat{i} + \frac{\partial}{\partial y_i}\hat{j} + \frac{\partial}{\partial z_i}\hat{k}$$
(7.3)

The total potential energy, the total virial potential energy and the static radius of the virialized state at the turn around (when the expansion stops, and collapse and rotating takes place) are denoted as  $U_{ta}$ ,  $U_{vir}$  and  $R_{vir}$  respectively. Supposing the system's potential energy takes the analytic form of U, then the kinetic energy after virialization, which depends on U, is:

$$T_{vir} = \left(\frac{R}{2}\frac{\partial U}{\partial R}\right)_{vir} \tag{7.4}$$

The energy conserves, so after the virial collapse(T=0 when collapse starts):

$$\left[U + \frac{R}{2}\frac{\partial U}{\partial R}\right]_{vir} = U_{ta} \tag{7.5}$$

The analytic expression of the potential energy term U can be deduced as follows: Supposing  $\rho_Q$  is the energy density for the virial component,  $\Phi_Q$  is the total potential acts on a certain point in the sphere, and the total potential energy will be

$$U = \frac{1}{2} \int \rho_Q \Phi_Q dV$$

$$= \frac{1}{2} \int \rho_Q \Phi_Q(r) d(\frac{4}{3}\pi r^3)$$

$$= \frac{1}{2} \int \rho_Q \Phi_Q(r) \cdot 4\pi r^2 dr$$
(7.6)

In a homogeneous sphere,  $\rho_Q$  satisfies:

$$\dot{\rho}_Q + 3(1+w_Q)\frac{\dot{r}}{r}\rho_Q = 0 \tag{7.7}$$

And the total potential in the sphere for a certain point is:

$$\Phi_Q(r) = -2\pi G(1+3w_Q)\rho_Q\left(R^2 - \frac{r^2}{3}\right)$$
(7.8)

Where R is the radius of the sphere, r is the distance between a point and the spherical center.

## 7.2 Virialization Power-style Cardassian

Now let's consider the virialization of power-style Cardassian[53]. Supposing that Cardassian term and matter term both participate the virialization process. i.e.

$$\rho = \rho_m + b\rho_m^n, \qquad n < 2/3 \tag{7.9}$$

#### 7.2.1 Virialization of Matter Term

EOS parameter for matter term is w = 0, and the distribution of potential in the sphere is:

$$\Phi(r) = -2\pi G (1+3\times 0)\rho_m \left(R^2 - \frac{r^2}{3}\right)$$

$$= -2\pi G \rho_m \left(R^2 - \frac{r^2}{3}\right)$$
(7.10)

The total potential energy in the sphere of radius R is:

$$U_m = -\int_0^R \rho_m^2 \cdot 2\pi G \cdot 2\pi r^2 \cdot \left(R^2 - \frac{r^2}{3}\right) dr$$
(7.11)  
=  $-\frac{16}{15} \pi^2 G \rho_m^2 R^5$ 

The rotating kinetic energy associating  $U_m$  is:

$$T_m = \frac{R}{2} \frac{\partial U_m}{\partial R} = -\frac{8}{2} \pi^2 G \rho_m^2 R^5$$
(7.12)

Obviously  $T_m < 0$  in (7.12). Kinetic energy cannot be negative, so the virial collapse of matter term alone is forbidden.

### 7.2.2 Virialization GF Fluid/Cardassian Term

Now let's have a look at the behaviors of Cardassian energy. Power-style Cardassian is perfect fluid with EOS parameter w = n - 1 and the distribution of potential in a virializing sphere is:

$$\Phi(r) = -2\pi G(3n-2)(b\rho_m^n) \left(R^2 - \frac{r^2}{3}\right)$$
(7.13)

So the total potential energy in the sphere is:

$$U = \frac{1}{2} \int \rho \Phi(r) \cdot 4\pi r^2 dr$$

$$= -\frac{1}{2} \int_0^R \rho_m \cdot 2\pi G (3n-2) (b\rho_m^n) \left(R^2 - \frac{r^2}{3}\right) \cdot 4\pi r^2 dr$$
(7.14)

$$=\frac{16}{15}G\pi^2 R^5(2-3n)b\rho_m^{2n}$$

and the associated virial rotating energy is:

$$T_{vir} = \frac{R_{vir}}{2} \frac{\partial U}{\partial R} = \frac{8}{3} G \pi^2 R_{vir}^5 (2 - 3n) (\rho_{m,vir} + b\rho_{m,vir}^n)^2 = \frac{5}{2} U_{vir}$$
(7.15)

Energy is conserving during the virialization process.

$$\frac{7}{2} \times \frac{16}{15} G \pi^2 R_{vir}^5 (2-3n) (b\rho_{m,vir}^{2n}) = \frac{16}{15} G \pi^2 R_{ta}^5 (2-3n) (b\rho_{m,ta}^{2n})$$
(7.16)

reduced to be:

$$\frac{7}{2}R_{vir}^5\rho_{m,vir}^{2n} = R_{ta}^5\rho_{m,ta}^{2n}$$
(7.17)

The mass does't lose, so

$$\frac{\rho_{m,vir}}{\rho_{m,ta}} = \left(\frac{R_{ta}}{R_{vir}}\right)^3 \tag{7.18}$$

Put 7.18 into7.17 to get:

$$\left(\frac{R_{vir}}{R_{ta}}\right)^5 \left(\frac{\rho_{m,vir}}{\rho_{m,ta}}\right)^{2n} = \left(\frac{R_{vir}}{R_{ta}}\right)^5 \left(\left(\frac{R_{ta}}{R_{vir}}\right)^3\right)^{2n} = \left(\frac{R_{vir}}{R_{ta}}\right)^{5-6n} = \frac{2}{7}$$
(7.19)

Hence the ratio of the stable virial radius with turn around radius is:

$$\frac{R_{vir}}{R_{ta}} = \sqrt[5-6n]{\frac{2}{7}}$$
(7.20)

The upper calculations constrains parameter n for twice, namely kinetic energy being positive and collapse radius becoming smaller than initial:

$$T > 0 \sim 2 - 3n > 0$$
 (7.21a)

$$\frac{R_{vir}}{R_{ta}} = \sqrt[5-6n]{\frac{2}{7}} < 1 \tag{7.21b}$$

So, after virialization GF fluids collapse to a steady sphere with certain radius which depends on the initial radius and the Cardassian parameter n. And no singularity is generated.

#### 7.2.3 Unitary Virialization of Matter Term and GF Fluid Term

Although matter term  $\rho_m$  cannot virialize on itself, howerever, as it is shown below, due to the excellent virialization property of GF fluid, the matter term  $\rho_m$  can virialize together with GF fluid  $\rho_{card}$ . This is pretty similar to the fact that high energy  $\gamma$  photon cannot decay into electron-positron pairs when isolated, but will decay if located in Column field nearby the nucleus.

As analyzed in chapter 6, that the total pressure of  $\rho_m$  and  $\rho_{card}$  is  $p = (n-1)b\rho_m^n$ , so the EOS parameter is:

$$w = \frac{(n-1)b\rho_m^n}{\rho_m + b\rho_m^n} \tag{7.22}$$

The distribution of potential generated by  $\rho_m$  and  $\rho_{card}$  in the virial sphere is:

$$\Phi(r) = -2\pi G \left( 1 + \frac{3(n-1)b\rho_m^n}{\rho_m^n} \right) \left(\rho_m + b\rho_m^n\right) \left( R^2 - \frac{r^2}{3} \right)$$
(7.23)  
$$-2\pi G \left(\rho_m + 3(n-2)b\rho_m^n\right)$$

 $\rho_m$  and  $\rho_{card}$  leads to the potential:

$$U = \frac{1}{2} \int_{m}^{r} (-2\pi G) \left(\rho_{m} + 3(n-2)b\rho_{m}^{n}\right) dr$$

$$= \frac{16}{15} \pi^{2} G R^{5} (\rho_{m} + b\rho_{m}^{n}) \left((2-3n)b\rho_{m}^{n} - \rho_{m}\right)$$
(7.24)

the associating rotating energy:

$$T = \frac{8}{3}\pi^2 G R^5 (\rho_m + b\rho_m^n) \left( (2 - 3n)b\rho_m^n - \rho_m \right) = \frac{5}{2}U$$
(7.25)

Thus, T > 0 is possible if b, n are properly tuned. Due to energy conserving:

$$\frac{7}{2} \times \frac{16}{15} \pi^2 G R_{vir}^5 (\rho_{m,vir} + b\rho_{m,vir}^n) ((2-3n)b\rho_m^n - \rho_m)$$

$$= \frac{16}{15} \pi^2 G R_{ta}^5 (\rho_{m,ta} + b\rho_{m,ta}^n) ((2-3n)b\rho_m^n - \rho_m)$$
(7.26)

The mass is conserving

$$\frac{\rho_{m,vir}}{\rho_{m,ta}} = \left(\frac{R_{ta}}{R_{vir}}\right)^3 \tag{7.27}$$

Combine it with energy conserving equation:

$$\frac{7}{2} \left(\frac{R_{vir}}{R_{ta}}\right)^5 = \frac{(\rho_{m,ta} + b\rho_{m,ta}^n)((2-3n)b\rho_m^n - \rho_m)}{(\rho_{m,vir} + b\rho_{m,vir}^n)((2-3n)b\rho_m^n - \rho_m)}$$

$$= \frac{(2-3n)b^2\rho_{m,ta}^{2n} + (1-3n)b\rho_{m,ta}^{n+1} - \rho_{m,ta}^2}{(2-3n)b^2\rho_{m,vir}^{2n} + (1-3n)b\rho_{m,vir}^{n+1} - \rho_{m,vir}^2}$$
(7.28)

Set the ratio  $\frac{R_{vir}}{R_{ta}} = x$ , and via the mass conservation,

$$\frac{7}{2}x^5 = \frac{(2-3n)b^2\rho_{m,vir}^{2n}x^{6n} + (1-3n)b\rho_{m,vir}^{n+1}x^{3n+3} - \rho_{m,vir}^2x^6}{(2-3n)b^2\rho_{m,vir}^{2n} + (1-3n)b\rho_{m,vir}^{n+1} - \rho_{m,vir}^2}$$
(7.29)

Make deduction of  $x^5$  to (7.29) to get

$$\frac{7}{2} \left\{ (2-3n)b^2 \rho_{m,vir}^{2n} + (1-3n)b \rho_{m,vir}^{n+1} - \rho_{m,vir}^2 \right\}$$
$$= (2-3n)b^2 \rho_{m,vir}^{2n} x^{6n-5} + (1-3n)b \rho_{m,vir}^{n+1} x^{3n-2} - \rho_{m,vir}^2 x$$
(7.30)

Introduce a new variable  $q \equiv \frac{\rho_{card,vir}}{\rho_{m,vir}} = \frac{b\rho_{m,vir}^n}{\rho_{m,vir}}$ , and

$$b^2 \rho_{m,vir}^{2n} = q^2 \rho_{m,vir}^2 \tag{7.31a}$$

$$b^2 \rho_{m,vir}^{n+1} = q \rho_{m,vir}^2$$
 (7.31b)

Insert (7.31) into (7.30) to get

$$\frac{7}{2}\left\{(2-3n)q^2\rho_{m,vir}^2 + (1-3n)q\rho_{m,vir} - \rho_{m,vir}^2\right\}$$

$$= (2-3n)q^2\rho_{m,vir}^2 x^{6n-5} + (1-3n)q\rho_{m,vir}^2 x^{3n-2} - \rho_{m,vir}^2 x$$
(7.32)

Obviously  $\rho_{m,vir}^2 > 0$  holds, and deduct it from both sides of the equation (7.32), and we get the relation of x and q:

$$\frac{7}{2} \left\{ (2-3n)q^2 + (1-3n)q - 1 \right\}$$
$$= (2-3n)q^2 x^{6n-5} + (1-3n)q x^{3n-2} - x$$
(7.33)

We can also get the relation  $z_{vir}$  and x via (7.33). Currently, Friedmann equation reads:

$$H_0^2 = \frac{8\pi G}{3} (\rho_{m,0} + b\rho_{m,0}^2)$$
(7.34)

i.e

$$\rho_{crit,0} \equiv \frac{3H_0^2}{8\pi G} = \rho_{m,0} + b\rho_{m,0}^2 \tag{7.35}$$

i.e

$$\frac{\rho_{crit,0}}{\rho_{m,0}} = 1 + \frac{b\rho_{m,0}^n}{\rho_{m,0}} \tag{7.36}$$

i.e

$$\frac{1}{\Omega_{m,0}} - 1 = q \tag{7.37}$$

insert it into x - q relation (7.33) and one will get the  $q - z_{vir}$  relation.

## 7.3 Conclusions and Evaluation

New work gathers in Chapter 6 and Chapter 7. Firstly we make a comprehensive analysis of the existing work, pay particular attention to Gondolo and Freese's idea of treating Cardassian energy term as relativistic perfect fluid (GF fluid), point out that a potential Cardassian term should meet three conditions, and review three existing Cardassian terms (power style, polytropic style and its modification, exponential style and its modification), and eventually put forward the newly found hyperbolic cosecant Cardassian.

Then the Cardassian dynamical equations are introduced generally and logically under GF fluid scenario, together with the flowing process of constructing phase space and differential dynamical systems from Friedmann equation. Hyperbolic cosecant Cardassian term is employed for concrete computing. The analysis proceeds in two cases, namely a unified description of matter and radiation energy density (case 1) and a separate description of matter and radiation terms (case 2). Formalism of case 2 is more exact at the expense of more complicatedness, and due to the mathematical symmetry of matter term and radiation term in hyperbolic cosecant function, the differential dynamical equations are considerably simplified. Phase space and dynamical systems for both cases are achieved. When we calculate the critical points for case 2, amazingly interesting behaviors of self-consistency and auto-normalization are exhibited, which is a strong support for the new model, along with a forever positive sound speed.

The process of virial collapse in Cardassian cosmos is analyzed. Power-style Cardassian term is employed for its simplicity. Calculation declares that virial collapse of matter alone is forbidden. Yet Cardassian has excellent ability for virial collapse, after the virial collapse ending up with a stable sphere, the ratio of the ultimate radius to the original radius depends on the adjustable parameters in Cardassian term. And, the mixture of GF fluid and matter could conduct virial collapse, the ratio of the ultimate radius to the original radius depends on the adjustable parameters in Cardassian term, too.

The creative work in this thesis incudes the introduction of hyperbolic cosecant Cardassian, which is the fourth ever available Cardassian term ever found, after the introduction of exponential style in 2005. This work, along with the calculating of power Cardassian virialization, enriches the research of Cardassian cosmology.

# Bibliography

- Liao LIU, Yuan-Fang Jiang, Zhen-Hua Qian. Inflation Cosmology- 10<sup>-35</sup> second after Big Bang. Progress in Physics, Vol 9(2), 121-186, 1986.
- [2] Bo QIN, Dark Matter Problem in the Fine Cosmology Era, Knowledge of Modern Physics (in Chinese). Vol 19(5). P17-24.
- [3] Scientific American, Section 2, 2007.
- [4] Charles W. Misner, Kip S. Thorne, John Archibald Wheeler. Gravitation. San Francisco, W. H. Freeman 1973.
- [5] Stephani, Hans. General relativity: an introduction to the theory of the gravitational field / Hans Stephani; edited by John Stewart; translated by Martin Pollock and John Stewart. Cambridge [Cambridgeshire]; New York: Cambridge University Press, 1982.
- [6] J. P. Ostriker, P. J. Steinhardt, Nature 377, 600 (1995).
- [7] David Spergel, Ue-Li Pen, Astrophys. J. 491, 67, (1997).
- [8] B. Ratra and P. J. E. Peebles, Phys. Rev. D 37, 3406 (1988);
- [9] P. J. E. Peebles and B. Ratra, Ap. J. 325, L17 (1988).
- [10] A. B. Burd and J. D. Barrow, Nucl. Phys. B 308, 929 (1988);
- [11] E. J. Copeland, A. Lahiri and D. Wands, Phys. Rev. D bf 50, 4868 (1994);
- [12] E. J. Copeland, A. R. Liddle and D. Wands, Phys. Rev. D bf 57, 5064 (1998)
- [13] K. Coble, S. Dodelson and J. A. Frieman, Phys. Rev. D 55, 1851 (1997);
- [14] J. A. Frieman et al., Phys. Rev. Lett. 75, 2077 (1995).
- [15] Ivaylo Zlatev, Limin Wang and Paul J. Steinhardt. Quintessence, Cosmic Coincidence, and the Cosmological Constant. astro-ph/9807002

- [16] L. Arturo Urena-Lopez, Tonatiuh Matos. A new cosmological tracker solution for Quintessence. astro-ph/0003364
- [17] Elisa Di Pietro, Jacques Demaret. A constant equation of state for quintessence? gr-qc/9908071
- [18] Sean M. Carroll. Quintessence and the Rest of the World. astro-ph/9806099
- [19] Models of Dynamical Supersymmetry Breaking and Quintessence.
   P.Binetruya.hep-ph/9810553
- [20] Jihn E. Kim. Axion and Almost Massless Quark as Ingredients of Quintessence. hep-ph/9811509
- [21] R. R. Caldwell. A Phantom Menace? Cosmological Consequences of a Dark Energy Component with Super-Negative Equation of State. astro-ph/9908168
- [22] L.P. Chimento, Ruth Lazkoz. On big rip singularities. gr-qc/0405020
- [23] S Nojiri, S D Odintsov and S Tsujikawa, hep-th 0501025, PRD 71,2005,063004.
- [24] Celine Cattoen, Matt Visser. Necessary and sufficient conditions for big bangs, bounces, crunches, rips, sudden singularities, and extremality events. gr-qc/0508045
- [25] A. E. Schulz, Martin White. The Tensor to Scalar Ratio of Phantom Dark Energy Models. astro-ph/0104112
- [26] A.V. Yurov. Phantom scalar fields result in inflation rather than Big Rip. astro-ph/0305019
- [27] E Elizalde, S Nojiri and S D Odintsov, hep-th/0405034, PRD 70, 2004, 043539.
- [28] S Nojiri and S D Odintsov, hep-th/0405078, PLB 595,1 (2004)
- [29] Puxun Wu , Hongwei Yu . Avoidance of Big Rip In Phantom Cosmology by Gravitational Back Reaction. astro-ph/0407424
- [30] Ruben Curbelo, Tame Gonzalez and Israel Quiros. Interacting Phantom Energy and Avoidance of the Big Rip Singularity. astro-ph/0502141
- [31] Alexander A. Andrianov, Francesco Cannata, Alexander Y. Kamenshchik, Daniele Regoli. Phantom without phantom or how the PT symmetry saves us from the Big Rip. gr-qc/0810.5076v1

- [32] Julio C. Fabris , Diego Pavon. Big rip avoidance via black holes production. gr-qc/0806.2521v1
- [33] Dao-jun Liu, Xin-zhou Li. Born-Infeld-type phantom on the brane world. hep-th/0307239
- [34] Emilio Elizalde, John Quiroga Hurtado . Phantom and quantum matter in an Anti-de Sitter Universe. gr-qc/0310128
- [35] Brett McInnes. The Phantom Divide in String Gas Cosmology. hep-th/0502209
- [36] V. Berezin, V. Dokuchaev, Yu. Eroshenko, A. Smirnov. Phantom shell around black hole and global geometry. gr-qc/0503068
- [37] Alexander Virkman. Can dark energy evolve to the phantom? Phys.Rev.D71:023515,2005. astro-ph/0407107
- [38] Feng B., Wang X., Zhang X. M. Phys. Lett. B. 2005, 607: 35.
- [39] Li M., Feng B., Zhang X. M. JCAP 2005, 0512: 002.
- [40] Jun-Qing Xia, Xin-Min Zhang, Progress in Dark Energy Research, Progress in Physics(chinese), Vol.28, No.3, 263-273, 2008.
- [41] Hao Wei, Rong-Gen Cai, Ding-Fang Zeng. Hessence: A New View of Quintom Dark Energy. hep-th/0501160
- [42] Hao Wei, Rong-Gen Cai. Cosmological Evolution of Hessence Dark Energy and Avoidance of the Big Rip. astro-ph/0509328
- [43] Hao Wei, Ningning Tang, Shuang Nan Zhang. Reconstruction of Hessence Dark Energy and the Latest Type 1a Supernovae Gold Dataset. astro-ph/0612726.
- [44] M. Alimohammadi, H. Mohseni Sadjadi. Attractor solutions for general hessence dark energy. hep-th/0602268.
- [45] Wen Zhao. Holographic hessence models. astro-ph/0706.2211.
- [46] Hao Wei, Shuang Nan Zhang, Dynamics of Quintom and Hessence Energies in Loop Quantum Cosmology. gr-qc/0705.4002
- [47] Armendariz-Picon C., Mukhanov V F, Steinhardt P J. Phys. Rev. Lett., 2000, 85: 4438. astro-ph/0004134.
- [48] Armendariz-Picon C., Mukhanov V F, Steinhardt P J. Phys. Rev. D63:103510, 2001, astro-ph/0006373.

- [49] Chiba T, Okabe T, Yamaguchi M. Phys. Rev. D62:023511. astro-ph/9912463.
- [50] Alexander Kamenschik, Ugo Moschella, Vincent Pasquier. An Alternative to Quintessence. gr-th/0103004.
- [51] M. C. Bento, O. Bertolami, A. A. Sen. Generalized Chaplygin Gas, Accelerated Expansion and Dark Energy-Matter Unification. gr-qc/0202064.
- [52] R. Jackiw. (A Particle Field Theorist's) Lecture on (Supersymmetric, Non-Abelian) Fluid Mechanics (and d-Brane). physics.flu-dyn/0010042.
- [53] Katherine Freese, Matthew Lewis. Cardassian Expansian: a Model in which the Universe is Flat, Matter Dominated, and Accelerating. astro-ph/0201229.
- [54] Katherine Freese. Generalized Cardassian Expansian: a Model in which the Universe is Flat, Matter Dominated, and Accelerating. hep-ph/0208264
- [55] Paolo Gondolo, Katherine Freese. Fluid Interpretation of Cardassian Expansian. hep-ph/0209322
- [56] James M Cline, Jeremie Vinet. Problems with Time-Varying Extra Dimensions or "Cardassian Expansion" as alternative to Dark Energy. hep-ph/0211284
- [57] Marek Szydlowski, Wojciech Czaja. Modified Friedmann Cosmologies-theory & observations. astr-ph/0402510
- [58] Ruth Lazkoz, Genly Leon. Asymptotic behavior of Cardassian cosmologies with exponential potentials. astro-ph/0503478
- [59] Paolo Gondolo, Katherine Freese. An Accelerating Universe from Dark Matter Interactions with Negative Pressure. hep-th/0211397
- [60] Katherine Freese. Cardassian Expansion: Dark Energy Density from Modified Friedmann Equations. astro-ph/0501675.
- [61] Dao-Jun Liu, Chang-Bo Sun, Xin-Zhou Li. Exponential Cardassian Universe. astr-ph/0512355
- [62] Ze-Long Yi, Tong-Jie Zhang. Statefinder Diagnostic for the Modified Polytropic Cardassian Universe. astr-ph/07-03630
- [63] Nikolaus Berndt, Alan Rendall. Isotropization in the Approach to Big Rip Sigularities for Cardassian Models. gr-qc/0804.3693

- [64] S. Sen, AA Sen. Observational Constraints on Cardassian Expansion. astro-ph/0211634.
- [65] Zong-Hong Zhu, Masa-Katsu Fujimoto. Cardassian Expansion: Constraints from Compact Radio Source Angular Size versus Redshift Data.astro-ph/0212192
- [66] Yun Wang, Katherine Freese, Paolo Gondolo, Matthew Lewis. Future Type 1a Supernova Data as tests of Dark Energy from Modified Friedmann Equations. astro-ph/0302064
- [67] Zong-Hong Zhu, Masa-Katsu Fujimoto. Constraints on Cardassian Expansion Expansion from Distant type 1a Supernova. astro-ph/0303021
- [68] A.A. Sen, S. Sen. WMAP constraints on Cardassian model. astro-ph/0303383
- [69] Wlodzimierz Godlowski, Marek Szydlowski. Constraints on a Cardassian model from SN1a data-rivisited. astro-ph/0309569
- [70] W.J. Frith. Constraints on Cardassian Expansion. astr-ph/0311211
- [71] Zong-Hong Zhu, Masa-Katsu Fujimoto. Constraints bon Cardassian Scenario from the Expansion Turnaround Redshift and the Sunyaev-Zeldovich/X-Ray Data. astro-ph/0312022.
- [72] Chris Savage, Noriyuki Sugiyama, Katherine Freese. Age of the Universe in Cardassian Models. astr-ph/0403196
- [73] Tomi Koivisto, Hannu Kurki-Suonio, Finn Ravndal. The CMB Spectrum in Cardassian Models. astro-ph/0409163
- [74] M Amarzguioui, /O. Elgar/oy, T Multamäki. Combined constraints on Cardassian models from supernovae, CMB, and large-scale structure observations. astr-ph/0410408
- [75] J.S. Alcaniz, Abha Dev, Deepak Jain. Constraints on the Cardassian Expansion from the Cosmic Lens All-Sky Survey Gravitational Lens Statistics. astro-ph/0501026
- [76] Stephane Fay, Morad Amarzguioui. Large-scale structure and the Cardassian Fluid. gr-qc/0608126
- [77] Herman J. Mosquera Cuesta, Habib Dumet M., Cristina Furlanetto. Confronting the Hubble Diagram of Gamma-Ray Brusts with Cardassian Cosmology. astroph/0708.1355

- [78] SUN Chang-bo. Modified Friedmann Equation and the Expanding Universe. Ms.D these is. Shanghai Normal University.
- [79] Varun Sahni, Tarun Deep Saini, A. A. Starobinsky, Ujjaini Alam. Statefinder a new geometrical diagnostic of dark energy. astro-ph/0201498. JETPLett.77:201-206,2003
- [80] Varun Sahni. Exploring dark energy using the Statefinder. astro-ph/0211084. Talk at the XVIII'th IAP Colloquium 'On the Nature of Dark Energy', IAP Paris, July 1 - 5, 2002; 4 pages.
- [81] Ujjaini Alam, Varun Sahni, Tarun Deep Saini, A. A. Starobinsky. Exploring the Expanding Universe and Dark Energy using the Statefinder Diagnostic. astro-ph/0303009. Mon.Not.Roy.Astron.Soc.344:1057,2003.
- [82] J.E.Gunn, J.R.1.Gott. AstroPhys.J.176,l(1972)
- [83] Irit Maor, Ofer Lahav. On Virialization with Dark Energy. astro-ph/0505308.