

Phys. Teacher **48**,4(2006)
A SHORT HISTORY OF HINDU ASTRONOMY &
EPHEMERIS

P.Rudra

Formerly of Department of Physics, University of Kalyani,
Kalyani, WB, 741-235, India

e-mail: p.rudra@saha.ac.in prasanta.rudra@gmail.com

PACS no. 01.65.+g 95.10.Km

Abstract

We have summarized here the astronomical knowledge of the ancient Hindu astronomers. This knowledge was accumulated from before 1500 B.C. up to around 1200 A.D. In Section II we have correlated terms used by the Hindu astronomers and their equivalents in modern astronomy. In Section III we have defined the different astronomical coordinate systems and their transformation relations. In Sections IV and V we have collected the main features of solar and lunar motions in terms of modern astronomical terminology. In Section VI we have given the names of the stars mentioned by the Hindu astronomers and their modern names with the present astronomical coordinates. In Section VII we have given a short survey of Indian history with emphasis to Hindu astronomy. In Sections VIII and IX we have given short descriptions of the main sources of Hindu astronomy. In Section X the important features of Hindu astronomy have been described. All through the article we have tried to present the content in tabular forms for ready and unambiguous perception.

I Introduction

Every civilization from the antiquity to the present time has been fascinated by astronomy, study of motion of the heavenly bodies across the sky. To-day astronomy has many tools of investigation: γ -ray, X-ray, ultraviolet, visible, infra-red, microwave and radio-wave radiations emanating from the heavenly bodies. This has opened up a horizon that was not available to the ancients. They only had the visible range of the spectrum. Their studies were thus restricted to what we now call positional astronomy and did not extend to investigations of the nature of these bodies. However, extensive data about positional astronomy exist in ancient Egyptian, Babylonian, Indian and Chinese records.

Here we have summarized the knowledge that ancient Hindu Astronomers had arrived at. The time span is from circa 1500 B.C. to 1200 A.D. This is a very long span of time. The technical terms used by these people have to be collated to present astronomical vocabularies. These data were collected at places distributed over a vast region of the earth. These facts complicate the process of evaluation of correctness of these data.

In Table 2 we have given the Sanskrit names of different astronomical terms and the corresponding technical names of present-day astronomy. In writing Sanskrit words in Roman letters of alphabet we have followed the rule of transliteration recommended by the International Alphabet of Sanskrit Transliteration (IAST) given in Table 1.

In Section III we explain the different coordinate systems used to describe the position of the heavenly bodies on the Celestial Sphere. In Section IV we have described present-day knowledge of the motion of the sun and the moon, the two heavenly bodies most important to terrestrial beings. In Section V we have mentioned the different lunar and solar time periods that are of importance to astronomy. In Section VI we tabulate the fixed stars on the Celestial Sphere that were mentioned by Hindu Astronomers in their studies. In Section VII we describe the salient points of history of the Indians, their interaction with Western Asia. A map (Figure 5) gives us the idea of the region that is of importance to the development of Hindu Astronomy. The Vedas and the Siddhāntas are the main texts that tell us about the range of knowledge of the Hindu astronomers. In Sections VIII and IX we describe the salient features of these sources. In Section X the principal features of Hindu astronomy over the ages are summarized.

Table 1 Rule of transliteration from Sanskrit to Roman letters of alphabet according to International Alphabet of Sanskrit Transliteration (IAST)

अ = a,	आ = ā,	इ = i,	ई = ī,	
उ = u,	ऊ = ū,	ऋ = ṛ,	ॠ = ṝ,	
ए = e,	ऐ = ai,	ओ = o,	औ = au,	
क = k,	ख = kh,	ग = g,	घ = gh,	ङ = ṅ,
च = c,	छ = ch,	ज = j,	झ = jh,	ञ = ñ,
ट = t,	ठ = ṭh,	ड = ḍ,	ढ = ḍh,	ण = ṇ,
त = t,	थ = th,	द = d,	ध = dh,	न = n,
प = p,	फ = ph,	ब = b,	भ = bh,	म = m,
य = y,	र = r,	ल = l,	व = v,	
श = ś,	ष = ṣ,	स = s,	ह = h,	
(◌) = m̐,	: ◌ = ḥ,	◌ = (◌)		

II Equivalence of Astronomical Terms

In all records of Hindu Astronomy we find mentions of different astronomical terms. These are given in Table 2 along with their modern equivalents.

Table 2 Technical terms in Modern Astronomy and their equivalents in Hindu Astronomy.

Modern Astronomical Term	Equivalent Term in Hindu Astronomy
Altitude	Madhyonnati
Apogee	Apabhū
Apse	Apadūrika
Aries, first point of	Meṣādi
Astronomy	Jyotiṣa
Azimuth	Digāṁśa
Declination	Viṣuvalamba
Ecliptic	Krāntivṛtta
Equator, Celestial	Mahāviṣuva
Equinox, Autumnal	Jalaviṣuva Saṁkrānti
Equinox, Vernal	Mahāviṣuva Saṁkrānti
Horizon plane	Digcakra
Hour Angle	Horakoṇa
Libra, first point of	Tulādi
Luminary	Jyotiṣka
Meridian	Madhyarekhā
Moon, Full	Pūrṇimā
Moon, New	Amābasyā
Perigee	Anubhū
Polaris	Dhruvatārā
Right Ascension	Viṣuvāṁśa
Saros	Yuga
Sidereal	Nākṣatra
Solstice, Summer	Karkaṭa Saṁkrānti
Solstice, Winter	Makara Saṁkrānti
Sphere, Celestial	Khagola
Zenith	Khamadhya
Zodiac signs	Rāśicakra

If we go through this table we find that the Hindu Astronomers had known the basic facts of positional astronomy regarding motion of the heavenly bodies through the Celestial Sphere, the different planes of motion of the sun and the moon, and the zodiac signs of the fixed stars.

III Coordinates describing an object X


In positional astronomy, motion of a heavenly body X is described in different coordinate systems. In ordinary 3-dimensional space with spherical symmetry, spherical polar coordinate system is the natural choice. On the Celestial Sphere, the real distance from the observer is not of any importance. Only the two angular coordinates fix the position of a body on the Celestial Sphere. This requires an azimuthal plane and a reference direction on this plane. For the coordinate of a place on the surface of the earth (Cartographic system) we take the terrestrial equatorial plane as the azimuthal plane and Greenwich's meridian supplies the starting position for measuring the azimuthal angle. The topocentric (Horizon) system is suitable for direct measurement by an observer on earth's surface. For this the local horizon plane supplies the azimuthal plane. The geocentric (Equatorial) system, which is independent of the observer's position on earth's surface, uses the Celestial Equator (the plane of intersection of the terrestrial equator with the Celestial Sphere) as the azimuthal plane. Description in terms of the geocentric coordinate system frees the observational data from the dependence on the position of the local observer on earth's surface. Figure 1 shows the topocentric and the geocentric coordinate systems. To describe the motion of the planets and the stars on the Celestial Sphere, rotational and non-uniform yearly motion of earth's motion around the sun has to be avoided. The heliocentric (Ecliptic) system takes Ecliptic (the plane of sun's motion on the Celestial Sphere) as the azimuthal plane. Figure 2 shows the heliocentric coordinate system. In Table 3 we have compared the 3 astronomical coordinate systems and the terrestrial cartographic system. The laws of transformation from one system to another are as follows:

$$\begin{pmatrix} \cos \delta \cos \chi \\ -\cos \delta \sin \chi \\ \sin \delta \end{pmatrix} = \begin{pmatrix} \cos \bar{\phi} & 0 & \sin \bar{\phi} \\ 0 & 1 & 0 \\ -\sin \bar{\phi} & 0 & \cos \bar{\phi} \end{pmatrix} \begin{pmatrix} \cos \zeta \cos \psi \\ -\cos \zeta \sin \psi \\ \sin \zeta \end{pmatrix},$$

$$\begin{pmatrix} \cos \beta \cos \lambda \\ \cos \beta \sin \lambda \\ \sin \beta \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \epsilon_0 & \sin \epsilon_0 \\ 0 & -\sin \epsilon_0 & \cos \epsilon_0 \end{pmatrix} \begin{pmatrix} \cos \delta \cos \alpha \\ \cos \delta \sin \alpha \\ \sin \delta \end{pmatrix}$$


Azimuth ψ_0 of a Star at setting ($\zeta = 0$) is given by $\sin(\psi_0 - 90^\circ) = \sin \delta / \cos \phi$.
For the Sun at *Summer Solstice (S.S.)*, $\delta = 23^\circ 30'$.

Figure 1 Topocentric and Geocentric Coordinates.



- O = Observer, X = Astronomical Object
 Z = Zenith, $SWNE \equiv \text{Plane } \perp OZ = \text{Horizon Plane,}$
(Khamadhya) *(Digcakra)*
 P = Polaris, $S'WN'E \equiv \text{Plane } \perp OP = \text{Celestial Equator,}$
(Dhruvatārā) *(Mahāviṣuva)*
Observer's Meridian = Great Circle through P and Z
Object's Meridian = Great Circle through P and X
 $N \equiv \text{North}$ } = { Points of intersection of the
 $S \equiv \text{South}$ } = { Observer's Meridian with the Horizon Plane,
 $E \equiv \text{East}$ } = { Points of intersection of the
 $W \equiv \text{West}$ } = { Celestial Equator with the Horizon Plane,
 N', S' = Points of intersection of the
Observer's Meridian with the Celestial Equator,
 $\zeta = MX \equiv \text{Altitude (Madhyonnati),}$ $\delta = RX \equiv \text{Declination (Viṣuvalamba)}$
 $\psi = SM \equiv \text{Azimuth (Digāṁśā),}$ $\chi = S'R \equiv \text{Hour Angle}$
Sidereal time $\equiv \tau = \text{Hour Angle of a fixed feature } Q \text{ on the Celestial Sphere}$
Right Ascension (R.A.) of $X \equiv \alpha = \tau - \chi$
 α increases towards E from Q , opposite to the apparent motion of the Celestial Sphere

Figure 2 Heliocentric Coordinates.



Ecliptic = Plane of the motion of the Sun on the *Celestial Sphere*,

K = Pole of the *Ecliptic* on the *Celestial Sphere*,

ϵ_0 \equiv obliquity of the *Ecliptic*
with respect to the *Celestial Equator*

= $\begin{cases} 23^\circ 30' \text{ (in 1950 A.D.)} \\ 23^\circ 51' \text{ (in 1300 B.C.)} \end{cases}$

Υ (*V.E.*) and Ω (*A.E.*) = Points of intersection of the *Ecliptic*
and the *Celestial Equator*

Υ has an involved retrograde motion with respect to the fixed stars. Its *Celestial Longitude* with respect to a given fixed star decreases by about $50''$ per year.

In antiquity Υ was in the *Constellation of Aries* (*Meṣa*), but now Υ has moved to the *Constellation of Pisces* (*Mīna*).

The fixed feature Q is taken to be the *Vernal Equinox* point, Υ .

$$\beta = TX = \text{Celestial Latitude,}$$

$$\lambda = \Upsilon T = \text{Celestial Longitude}$$


Table 3 Coordinates used in different Astronomical Systems.

Cartographic System	Topocentric Horizon system	Geocentric Equatorial system	Heliocentric Ecliptical system
ϕ = <i>Geographical Latitude</i> , measured from the <i>Terrestrial Equator</i> along the <i>Object's Meridian</i> , $\bar{\phi} = 90^\circ - \phi$ = <i>Colatitude</i>	$\zeta = MX$ = <i>Altitude</i> (<i>Madhyonnati</i>), measured from the <i>Horizon plane</i> towards the <i>Zenith</i>	$\delta = RX$ = <i>Declination</i> (<i>Viṣuvalamba</i>), measured from the <i>Celestial Equator</i> along the <i>Object's Meridian</i>	$\beta = TX$ = <i>Celestial Latitude</i> , measured from the <i>Ecliptic</i> towards <i>K</i>
<i>Geographical Longitude</i> , measured from <i>Greenwich's Meridian</i> along the <i>Terrestrial Equator</i>	$\psi = SM$ = <i>Azimuth</i> (<i>Digāṃśa</i>), measured from <i>South</i> towards <i>West</i> along the <i>Horizon plane</i>	$\chi = S'R$ = <i>Hour Angle</i> , measured from the <i>Observer's Meridian</i> towards <i>West</i> along the <i>Celestial Equator</i>	$\lambda = \Upsilon T$ = <i>Celestial Longitude</i> , measured from Υ towards the <i>Summer Solstice</i> along the <i>Ecliptic</i>

IV Sun (*Sol*) - Earth (*Terra*) - Moon (*Luna*) System

To people living on the surface of the earth, the sun and the moon are the two most important heavenly bodies. In Figure 3 we show our present-day knowledge of the motion of the earth around the sun throughout the year. This heliocentric picture was, however, not available to the people of antiquity. What they observed was the Ecliptic, the plane of solar motion on the Celestial Sphere, and the difference of speed of the sun along the Ecliptic at different seasons of the year.

Figure 3 Earth's motion around the Sun.




V.E.	≡ Vernal Equinox (<i>Mahāviṣuva Saṁkrānti</i>)	= 21 March in 1950 A.D.
A.E.	≡ Autumnal Equinox (<i>Jalaviṣuva Saṁkrānti</i>)	= 22 September in 1950 A.D.
W.S.	≡ Winter Solstice (<i>Makara Saṁkrānti</i>)	= 22 December in 1950 A.D.
S.S.	≡ Summer Solstice (<i>Karkaṭa Saṁkrānti</i>)	= 21 June in 1950 A.D.

Exact motion of Earth's axis has a number of periodic terms called *nutations* $\sim 9''$.

Similarly lunar motion around the earth is shown in Figure 4. This motion, of course, tallied with observations from earth's surface.

Figure 4 Moon's motion around the Earth.



Inclination of the plane of Lunar motion to the *Ecliptic* (the plane of Earth's annual motion around the Sun) is approximately 5° .

Eccentricity of Lunar elliptic orbit is approximately $1/18$.

Upper limit of the declination of the Moon has a cycle of approximately 18 years (from $18^\circ 10'$ to $28^\circ 50'$).

Perigee = Point of closest approach,

Apogee = Point of farthest approach

V Lunar and Solar Cycles

The two most important cyclic motions are the lunar and the solar cycles. They provide us with the two most easily observable time periods. Since actual periods are not exactly constant all through the year due to non-uniform motion of the earth around the sun, a number of different associated periods have been used by the astronomers. These are given in Table 4.

X *transits* or *culminates* when the *Meridian* of X coincides with the *Meridian* of the Observer.

<i>Sidereal Day</i> of X	=	from one <i>transit</i> of X to the next <i>transit</i> ,
<i>Mean Solar Day</i> (<i>M.S.D.</i>)	=	<i>Sidereal Day</i> of a fictitious <i>Mean Sun</i> (<i>M.S.</i>) moving with a uniform speed on the <i>Celestial Equator</i> ,
<i>Equation of time</i> <i>Greenwich Mean</i>	=	$\alpha_{M.S.} - \alpha_S$,
<i>Astronomical time</i> (<i>GMAT</i>)	=	<i>Hour Angle</i> of <i>M.S.</i> ,
<i>Universal time</i> (<i>UT</i>)	=	<i>GMAT</i> + 12 <i>hrs</i> ,
Mean midnight	=	0 <i>hr</i> ,
Mean noon	=	12 <i>hr</i>

Table 4 Different Lunar and Solar Time Periods.

Motion	Name (Symbol)	Description	Present value in <i>M.S.D.</i>
Lunar	<i>Tropical Month</i> , T_1	<i>Equinox to Equinox</i>	27.32158
	<i>Anomalistic Month</i> , T_2	<i>Perigee to Perigee</i>	27.53455
	<i>Draconic (Nodical) Month</i> , T_3	<i>Node to Node</i>	27.21222
	<i>Sidereal Month</i>	Return to the same fixed Star	27.32166
	<i>Synodic Month</i> , T	<i>New Moon to New Moon</i> (<i>Amānta</i>)	29.53059
Solar	<i>Tropical Year</i> , T_0 (<i>Sāvāṇa Vatsara</i>)	<i>Equinox to Equinox</i>	365.2422
	<i>Sidereal Year</i>	Return to the same fixed Star	365.257

Hindu astronomers determined transits of heavenly bodies by *Gnomon* (Sun-dial) and time was measured by *Clepsydra* (Water-clock). Names of different subdivisions of time given in Table 5 indicate the possibility that Hindu astronomers were able to measure these short time intervals. In fact in all treatises of Hindu Astronomy accurate directions for constructing water-clocks are given [7].

Table 5 Different subdivisions of time in Hindu Astronomy.

1 Anupala	=	$\frac{1}{150}$ sec,				
60 Anupala	=	1 Bipala	=	0.4 sec,	60 Bipala	= 1 Pala = 24 sec,
60 Pala	=	1 Daṇḍa	=	24 min,	2 Daṇḍa	= 1 Muhūrta = 48 min
7.5 Daṇḍa	=	1 Prahara,	=	3 hr,	8 Prahara	= 1 Divasa = 24 hr

The unit of Muhūrta has a significance in astronomy. It is the time interval by which the time of moonrise is delayed on successive days.

VI Stars and their positions

All astronomical and ephemerical calculations are based on motion of the heavenly bodies with respect to the fixed stars on the Celestial Sphere. Hindu Astronomers also used the same convention. The Celestial Sphere is mentally divided in 27 divisions and each division is assigned to the most important star there. Out of these 27 stars 12 are selected to give the names of the months. Present-day Celestial Longitudes and Latitudes of these stars are given in Table 6. Modern Astronomical names of these stars are also given.

Table 6 Names of *Nakṣatras* in *Vedāṅga Jyotiṣa*, *Sūrya Siddhānta*, corresponding names in *Modern Astronomy* and their *Celestial Longitudes* and *Latitudes* in 1950 A.D. The names of the Lunar months are according to the stars written in bold face letters.

Serial Number	Name of <i>Nakṣatra</i> in		Principal Star	<i>Celestial</i>	
	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>		<i>Longitude</i>	<i>Latitude</i>
1.	Kṛttikā	Kṛttikā	η - Tauri	59° 17' 39"	+ 4° 2' 46"
2.	Rohiṇi	Rohiṇi	α - Tauri	69° 5' 25"	- 5° 28' 14"
3.	Mṛgaśīrṣa	Mṛgaśīrṣa	λ - Orionis	83° 0' 31"	- 13° 22' 32"
4.	Ārdrā	Ārdrā	α - Orionis	88° 3' 22"	- 16° 1' 59"
5.	Punarvasu	Punarvasu	β - Geminorum	112° 31' 29"	+ 6° 40' 51"
6.	Ṭiṣya	Puṣyā	δ - Cancrī	128° 1' 23"	+ 0° 4' 32"
7.	Āśleṣā	Āśleṣā	ϵ - Hydrae	131° 38' 59"	- 11° 6' 25"
8.	Maghā	Maghā	α - Leonis	149° 8' 1"	+ 0° 27' 48"
9.	Pūrva Phalguni	Pūrva Phalguni	δ - Leonis	160° 36' 52"	+ 14° 19' 58"
10.	Uttara Phalguni	Uttara Phalguni	β - Leonis	170° 55' 23"	+ 12° 16' 13"
11.	Hasta	Hasta	δ - Corvi	192° 45' 23"	- 12° 11' 31"
12.	Citrā	Citrā	α - Virginis	203° 8' 37"	- 2° 3' 4"
13.	Svāti	Svāti	α - Bootis	203° 32' 8"	+ 30° 46' 3"
14.	Viśākhā	Viśākhā	α - Librae	224° 23' 7"	+ 0° 20' 19"

Table 6 (Continued)

Serial Number	Name of Nakṣatra in		Principal Star	Celestial	
	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>		<i>Longitude</i>	<i>Latitude</i>
15.	Anurādhā	Anurādhā	δ - Scorpii	241° 52' 23"	– 1° 58' 49"
16.	Rohiṇi, Jyeṣṭhā	Jyeṣṭhā	α - Scorpii	249° 3' 51"	– 33° 33' 50"
17.	Vicṛtau, Mūlabarhaṇī	Mūlā	λ - Scorpii	263° 53' 14"	– 13° 46' 56"
18.	Pūrvāṣādhā	Pūrvāṣādhā	δ - Sagittarii	273° 52' 55"	– 6° 27' 58"
19.	Uttarāṣādhā	Uttarāṣādhā	σ - Sagittarii	281° 41' 11"	– 3° 26' 36"
	Abhijit	Abhijit	α - Lyrae	284° 36' 54"	+ 61° 44' 7"
20.	Śroṇā	Śravaṇā	α - Aquilae	301° 4' 16"	+ 29° 18' 18"
21.	Śraviṣṭhā	Dhaniṣṭhā	β - Delphini	315° 38' 38"	+ 31° 55' 21"
22.	Śatabhiṣak	Śatabhiṣak	λ - Aquarii	340° 52' 38"	– 0° 23' 8"
23.	Pūrvā Proṣṭhapada	Pūrvā Bhādrapada	α - Pegasi	352° 47' 19"	+ 19° 24' 25"
24.	Uttarā Proṣṭhapada	Uttarā Bhādrapada	γ - Pegasi	8° 27' 32"	+ 12° 35' 55"
25.	Revati	Revati	ζ - Piscium	19° 10' 40"	– 0° 12' 52"
26.	Aśvayuja	Aśvini	β - Arietis	33° 16' 18"	+ 8° 29' 7"
27.	Apabharaṇī	Bharaṇī	41 - Arietis	47° 30' 19"	+ 10° 26' 48"

VII Summary of Indian History

Hindu Astronomy developed through about two thousand years and this development took place not at a fixed region. Parallel development of Astronomy also took place at Babylon and Alexandria. Figure 5 shows the part of the world and cities having importance in this historical development. Positions of modern cities near these ancient seats of astronomy are also shown.

A short summary of the History of India and her interaction with Western Asia is given in Table 7. The earliest time in Indian History that is of importance from astronomical interest is the time of Ṛk-vedas. This is the oldest of all the Vedas (Section VIII) and astronomical references found here fixes its time (terminus ad quem) around 1500 B.C., when the Middle Kingdom of Egypt ended. The part of the Vedas known as Vedāṅga was compiled around 1200 B.C. when Egypt had the New Kingdom. Time of Gautama Buddha (end of the 6th century B.C.) is a fixed point in Indian History. Written history is available afterwards. Mahābhārata, the great Indian epic was perhaps compiled around 400 B.C. Alexander's invasion of the Northwestern periphery of India in 323 B.C. and Candragupta's victory over Seleucus in 306 B.C. brought India in contact with Babylonian astronomy through the Greeks. But there is scarcely any evidence of influence of Egyptian or Babylonian astronomy on Hindu astronomy at this time. Advent of the Śakas (Scythians) in Balkh (Bactria) and their victory over the Pāradas (Parthians) in 123 B.C. would later have profound impact on Hindu astronomy and ephemeris. The Old Śaka Era started from this momentous event. With the beginning of the reign of Kaṇiṣka, the Kuṣāṇa king, in 78 A.D. (201 in the Old Śaka Era) the New Śaka Era (omitting the 200 of the Old Śaka Era) began in India. This is the Era *par excellence* in Hindu astronomy and no other Era could take its place in Hindu ephemeris. The Śakas grafted Babylonian system on Hindu astronomy and by 400 A.D. the Siddhānta astronomy (Section IX) replaced the older Vedāṅga astronomy. Āryabhaṭa, perhaps the greatest Hindu mathematician cum astronomer, was born at Kusumapura near Pāṭaliputra (modern Patna) in 476 A.D. and his monumental work Āryabhaṭīya was compiled in 499 A.D. The next important event (6th century A.D.) of Hindu astronomy is Varāhamihira's Pañca-siddhāntikā, second only to the appearance of Āryabhaṭīya. The last important contribution to Hindu astronomy came from Bhāskara-cārya of Devagiri in 1207. After Muhammad Ghouri's conquest of Northern India at the end of the 12th century A.D. all the academic centres of excellence in Northern India were completely destroyed and starved of financial support. This saw the end of ancient Hindu astronomy.

Figure 5 India and her Western neighbors.




Table 7 History of India and her Western neighbors.

<i>Era</i>	<i>Century</i>	<i>Indian</i>	<i>West Asian</i>
A.D.	13-th	<ul style="list-style-type: none"> • 1207 Siddhānta śiromaṇi of Bhāskarācārya of Devagiri, (last Indian contributor of importance) 	
	12-th	<ul style="list-style-type: none"> • 1192 Muhammad Ghouri defeats Pṛthvīrāja in the 2nd battle of Tarain 	
	11-th		
	10-th	<ul style="list-style-type: none"> • 1000 Mahmud of Ghazni invades India • 966 Bhaṭṭapāla's Gaṇitaskandha • 950 Āryabhaṭa II 	
	9-th		
	8-th		
	7-th	<ul style="list-style-type: none"> • 630 Harṣavardhana • 628 Brahmagupta of Khandakhādyā & Brahmasiddhānta 	<ul style="list-style-type: none"> • 700 Beginning of Arab civilization of Baghdad, Damascus, Cordova
	6-th	<ul style="list-style-type: none"> • 587 Death of Varāhamihira of Pañca siddhāntikā • 580 Śriṣeṇa • 523 Death of Āryabhaṭa 	<ul style="list-style-type: none"> • 600 End of Sassanid empire in Iran

Table 7 (Continued)

<i>Era</i>	<i>Century</i>	<i>Indian</i>	<i>West Asian</i>

A.D.	5-th	<ul style="list-style-type: none"> • 500 Viṣṇucandra • 499 Composition of Āryabhaṭiya • 476 Birth of Āryabhaṭa at Pāṭaliputra 	
	4-th	<ul style="list-style-type: none"> • 400 Beginning of Siddhānta Jyotiṣa • 320 Beginning of Gupta Empire 	
	3-rd		<ul style="list-style-type: none"> • 224 Beginning of Sassanid Empire in Iran
	2-nd	<ul style="list-style-type: none"> • 170 Śakas reign at Ujjainī 	<ul style="list-style-type: none"> • 150 Ptolemy of Alexandria
	1-st	<ul style="list-style-type: none"> • 100 Sātabāhana empire ends • 78 New Śaka Era, Kaṇiṣka's reign begins • 50 Kāṇva dynasty at Pāṭaliputra, Śakas rule at Puruṣapura & Mathurā 	<ul style="list-style-type: none"> • 100 Selucid rule at Babylon ends
B.C.	1-st	<ul style="list-style-type: none"> • 70 Śakas capture Takṣaśilā • 85 Sunga reign ends 	<ul style="list-style-type: none"> • 80 Śakas occupy Afghanistan
	2-nd	<ul style="list-style-type: none"> • 175 Menander captures Punjab & Sind • 180 Sunga dynasty at Pāṭaliputra, Selucids capture Takṣaśilā 	<ul style="list-style-type: none"> • 123 Śakas (Scythians) defeat Pāradas (Parthians) & occupy Balkh (Bactria), Old Śaka era begins • 138 End of Parthian empire in Iran

Table 7 (Continued)

<i>Era</i>	<i>Century</i>	<i>Indian</i>	<i>West Asian</i>
B.C.
	3-rd	<ul style="list-style-type: none"> • 220 Beginning of Śātavāhana empire • 226 Death of Aśoka • 268 Aśoka's reign begins 	<ul style="list-style-type: none"> • 248 Beginning of Parthian Empire in Iran • 290 Archimedes • 300 Euclid, End of Achaemenian Empire in Iran
	4-th	<ul style="list-style-type: none"> • 306 Candragupta defeats Seleucus • 323 Alexander invades India • 400 Mahābhārata 	<ul style="list-style-type: none"> • 312 Seleucid Era begins at Babylon
	5-th		
	6-th	<ul style="list-style-type: none"> • 560 Death of Gautama Buddha 	<ul style="list-style-type: none"> • 550 Pythagorus • 560 Cyrus establishes Achaemenian Empire in Iran, End of Assyrian Empire at Babylon, End of Egyptian Empire
	7-th	<ul style="list-style-type: none"> • 640 Birth of Gautama Buddha 	<ul style="list-style-type: none"> • 650 Ashurbanipal rules at Babylon
	8-th		<ul style="list-style-type: none"> • 728 Beginning of Assyrian Empire at Babylon, End of Chaldean Empire at Babylon,
	9-th		
	10-th		
	11-th		<ul style="list-style-type: none"> • 1100 End of New Kingdom at Egypt
	12-th	<ul style="list-style-type: none"> • 1200 Beginning of Vedāᅅga Jyotiᅅa 	

Table 7 (Continued)

<i>Era</i>	<i>Century</i>	<i>Indian</i>	<i>West Asian</i>
B.C.
	13-th		
	14-th		
	15-th	<ul style="list-style-type: none"> • 1500 R̥k – Veda (terminus ad quem) 	<ul style="list-style-type: none"> • 1500 End of Middle Kingdom at Egypt
	16-th		
	17-th		
	18-th		
	19-th		<ul style="list-style-type: none"> • 1900 Hammurabi establishes Chaldean Empire at Babylon
	20-th	Harappan civilization	

VIII The Vedas

The earliest reference to astronomical facts are found in the Vedas, the scripture of the Hindus. There are 4 Vedas, Ṛk, Sāma, Yajur and Atharva. Each of these Vedas are divided in 4 parts, Chanda (or Samhitā), Brāhmaṇa, Āraṇyaka (or Upaniṣad) and Vedāṅga. Vedāṅga deals with Phonetics, Ritualistic literature, Grammar, Etymology, Metrics and Astronomy. The Samhitā part of Ṛk-veda has very scanty astronomical references. That of Yajur has slightly more references. The Vedāṅgas (particularly of the Kṛṣṇa or the Black Yajur-veda), on the other hand, has a full-fledged astronomy. The salient features of the Vedas are given in Table 8. This shows the range of academic activities pursued by the ancient Indians.

Table 8 The Vedas, their subdivisions and salient features.

<i>The Vedas</i>					
↓	↓	↓	↓		
Ṛk	<i>Sāma</i>	<i>Yajuh</i>	<i>Atharva</i>		
divided in <i>10 Maṇḍalas</i>		↓			
		↓	↓		
		<i>Kṛṣṇa</i> (<i>Black</i>)	<i>Śukla</i> (<i>White</i>)		
		↓	↓		
↓	↓	↓	↓		
<i>Chandas</i> or <i>Samhitā</i> Collection of hymns, prayers, incantations, benedictions, sacrificial formulas, litanies	<i>Brāhmaṇa</i> prose texts on theological matters, observations on sacrifices and their mystical significances	<i>Āraṇyaka</i> or <i>Upaniṣad</i> meditations of forest hermits and ascetics on God, the world & mankind	<i>Vedāṅga</i> or <i>Sūtra</i>		
↓	↓	↓	↓	↓	↓
<i>Śikṣā</i> (Phonetics)	<i>Kalpa</i> (Ritualistic literature)	<i>Vyākaraṇa</i> (Grammar) e.g. <i>Pāṇini's</i> <i>Aṣṭādhyāyī</i>	<i>Nirukta</i> (Etymology) e.g. of <i>Yāṣka</i>	<i>Chandas</i> (Metrics) e.g. of <i>Piṅgala</i>	<i>Jyotiṣa</i> (Astro- -nomy) e.g. of <i>Lagadha</i> Ṛk- <i>Jyotiṣa</i> has 36 verses and <i>Yajuh</i> - <i>Jyotiṣa</i> has 43 verses

IX The *Siddhāntas*

Hindu astronomy gradually developed from the Vedāṅga jyotiṣa to what are known as the Siddhāntas. There are 5 Siddhāntas: Paitāmaha, Vāśiṣṭha, Romaka, Pauliśa and Sūrya. The last one is the latest and the most developed. Texts of all the siddhāntas are not available, but Varāhamihira in the 6th century A.D. has described them in his Pañca-Siddhāntikā. Romaka and Pauliśa siddhāntas show influence of Babylonian and Egyptian astronomy. One of the earliest proponent of the Sūrya Siddhānta was Āryabhaṭa. But with time it was upgraded with accumulation of new knowledge. Solar, lunar and stellar motions were accurately described in this Siddhānta. A study of this Siddhānta shows that Hindu astronomers knew about precession of the Equinoxes, but they were lamentably unaware of the difference between the Sidereal and the Tropical years. This is not so amazing, since even European astronomers became aware of this difference only in the 17th century A.D. after Galileo, Kepler and Newton. The 5 Siddhāntas are summarized in Table 9.

Table 9 Different Siddhānta Jyotiṣas and their salient features.

<i>The Siddhāntas</i>				
↓	↓	↓	↓	↓
<p><i>Paitāmaha</i> ascribed to Brahmā and described in 5 stanzas in <i>Pañca–</i> <i>Siddhāntikā</i> of Varāhamihira</p>	<p><i>Vāśiṣṭha</i> ascribed to Vaśiṣṭha and revealed by him to Māṇḍavya and described in 13 couplets by Varāhamihira; method of calculating <i>Tithi</i> and <i>Nakṣatra</i>, mentions <i>Rāśi</i> and <i>Lagna</i></p>	<p><i>Romaka</i> revealed by Viṣṇu to Ṛṣi Romaśa; <i>Yuga</i> = $19 \times 5 \times 30$ yrs, incorporates 19 yr cycle of Babylon and 5 yr <i>Yuga</i> of <i>Vedāṅga</i> <i>Jyotiṣa</i></p>	<p><i>Pauliśa</i> ascribed to the sage Pulastya; refers to <i>Yavanapura</i>, Longitudes of Ujjainī & Benares are given with respect to Alexandria, Lords of the days are mentioned like the Iranian calendar, but the names are Indian</p>	<p><i>Sūrya</i> Revealed by the Sun god to Asura Maya, who propounded it to the Ṛṣis. It has 500 verses in 14 Chapters, does not show any influence of Ptolemy's Almagest, does not mention Alexandria, Longitude of Ujjainī is the standard Meridian, it is continually updated like the Landolt– Bernstein tables, earliest version is by Āryabhaṭa 476 – 523A.D.</p>

X Comparison of different Astronomical systems of India before 1200 A.D.

In Table 10 we have summarized astronomical knowledge of the Hindu astronomers found in the 4 systems: Ṛk-Saṁhitā, Yajuḥ-Saṁhitā, Vedāṅga Jyotiṣa and Sūrya Siddhānta. This knowledge primarily encompasses the following subjects: Lunar asterism, Lunar day (Tithi), Month & Year (both Lunar and Solar), Seasons & Year (Ṛtu), and longer period of time (Yuga).

In the Ṛk-Saṁhitā, scanty though the knowledge is, we find mentions of lunar asterism, some specific fixed stars, lunar zodiac, lunar year of 360 days and solar year of 366 days, 3 Seasons each of 4 months (Cāturmāsya) and a 5 year Yuga. It is interesting to note that in the list of Seasons, the Rainy Season was not mentioned. This is possibly because the Hindu astronomers did not come across this season at the place where they were studying astronomy. This indicates that they were located only in the extreme North-Western corner of the country (Takṣaśilā) and Afghanistan (Balkh).

During the time of the Yajuḥ-Saṁhitā astronomy was more developed. Full list of the Lunar asterism is given though it started with Kṛttikā instead of Aśvinī as in later time. Lunar fortnight (Pakṣa) of 15 lunar days (Pañcadaśī tithi) was mentioned and gradually the concept of lunar month came into vogue. Northern and Southern motions of the Sun and the 6 Seasons connected with the motion of the sun along the Ecliptic was established knowledge. The Spring (Vasanta Ṛtu) consisting of 2 months was symmetrically placed about the Vernal Equinox (V.E.) point and the Solar year started when the Sun was 30° before the V.E.

Vedāṅga Jyotiṣa was a completely developed system with the numerical values of astronomical data comparing favorably with modern values. Evidence of the knowledge of the Sidereal Day, the Solstices, ratio of durations of day & night on the Summer Solstice, Lunar months, Tropical Solar Year, and longer periods of time is found in Vedāṅga Jyotiṣa. During the time of the Vedāṅga Jyotiṣa, Winter Solstice was at the Sraviṣṭhā nakṣatra (Delphini) at the longitude of 270° . Sraviṣṭhā is now at $316^\circ 41'$. This fixes the time of the Vedāṅga Jyotiṣa around 1400 B.C. Similar calculation fixes the time of compilation of the Mahābhārata as 450 B.C.

Sūrya Siddhānta has precise and logical definitions of the Lunar asterism, Tithi (Lunar Day), Solar Month, Tropical (Sāvana) year and the 6 Seasons (Ṛtu). The effect of the precession of the Equinoxes was also noted.

Table 10 Salient features of Hindu Astronomy in different ages.

<i>Astro– nomical features</i>	<i>Ṛk Saṃhitā</i>	<i>Yajuḥ Saṃhitā</i>	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>
Lunar Asterism (Nakṣatra)	<ul style="list-style-type: none"> • Custom of mentioning Nakṣatra perhaps started here • Days were designated by lunar asterism • Aghā & Arjunī were mentioned 	<ul style="list-style-type: none"> • Full list of lunar asterism is given • Lunar asterism starts with Kṛttikā 	<ul style="list-style-type: none"> • Lunar asterism starts at Kṛttikā, most probably Υ • Days were named after Nakṣatra • 1 Nakṣatra day was 1.011608 Solar day, in reality it is 1.011913 Solar day • W.S. was at Śraviṣṭhā at longitude 270°; it is now at $316^\circ 41'$; solstice takes 72 yrs to retrograde through 1°; if α – Delphini, then time is 1413 B.C., if β – Delphini, then time is 1338 B.C. • In Mahābhārata W.S. is at Śravaṇā, time is 450 B.C. 	<ul style="list-style-type: none"> • Lunar asterism starts at Aśvinī (α – or β – Arietis), Υ was at Revati (ζ – Piscium); now Υ has shifted by $\sim 19^\circ$ from ζ – Piscium • At Varāha–mihira’s time S.S. has moved $\frac{1}{2}$ of Aśleṣā + Puṣyā, time is 1500 yrs after Vedāṅga Jyotiṣa • No knowledge of precession of the Equinoxes, but its effect has been observed
...

Table 10 (Continued)

<i>Astro– nomical features</i>	<i>R̥k Saṃhitā</i>	<i>Yajuḥ Saṃhitā</i>	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>
...
Lunar day (Tithi)	<ul style="list-style-type: none"> • No reference • Perhaps individual days were denoted by the Nakṣatra, • Lunar zodiac was known 	<ul style="list-style-type: none"> • Pañcadaśī tithi • Knowledge of Pakṣa • Tithi was from moonrise to moonrise during Kṛṣṇa pakṣa and from moonset to moonset during Śukla pakṣa 	<ul style="list-style-type: none"> • Subdivision of day was measured by Clepsydra • Ratio of day & night on S.S. is $\frac{3}{2}$, this is true for latitude 35° N • Tithi is $\frac{1}{30}$ of lunar month = 0.983871 days, correct value is 0.984353 days • Day is reckoned by tithi 	<ul style="list-style-type: none"> • Tithi is complete when the moon is ahead of the sun by 12° or its multiple, tithi varies from 26 hr 47 min to 19 hr 59 min • If a tithi extends to 2 solar days, then both the days are assigned the same tithi
...

Table 10 (continued)

<i>Astro– nomical features</i>	<i>R̥k Saṁhitā</i>	<i>Yajuḥ Saṁhitā</i>	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>
...
Month & Year	<ul style="list-style-type: none"> • Year has 12 months each of 30 days (12 spoke–boards, & 360 śaṅkus) • Mentions Citrā (α – Virginis) & Aghā (α – Leonis) 	<ul style="list-style-type: none"> • Mentions lunar mansions with their presiding deities • Lunar months came gradually, generally pakṣa is mentioned 	<ul style="list-style-type: none"> • Does not mention zodiac signs • Pūrṇimānta lunar months are named after the Nakṣatras 	<ul style="list-style-type: none"> • Defines only astro–nomical solar month • 1 Solar month = time of passage of the Sun through 30° of its orbit = 30.43823 days; modern value = 30.43685 days • 1 Solar year = 12 Solar months
...

Table 10 (Continued)

<i>Astro–nomical features</i>	<i>R̥k Samhitā</i>	<i>Yajuh̥ Samhitā</i>	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>
...
Seasons & Year	<ul style="list-style-type: none"> • 3 seasons (navels), Cāturmāsya, each of 4 months, Gr̥ṣma (Summer), Śarad (Autumn), Hemanta (Winter); Varṣā (Rainy) is not mentioned • Lunar year of 360 days, 12 lunar months and 1 intercalary month of 12 tithis • Solar year of 366 days 	<ul style="list-style-type: none"> • Mentions Uttarāyana & Dakṣiṇāyana of the Sun • The Solar year has 6 Seasons : <ul style="list-style-type: none"> i) Vasanta (Spring) = Madhu & Mādhava ii) Gr̥ṣma (Summer) = Śukra & Śuci iii) Varṣā (Rains) = Nabha & Nabhasya iv) Śarad (Autumn) = Iṣa & Ūrja v) Hemanta (Early Winter) = Sahas & Sahasya vi) Śīta (Winter) = Tapas & Tapasya; • each Season covers 60° of the Sun's motion along the Ecliptic; solar months are seldom mentioned • Tropical Solar year & Vasanta starts at – 30° of the Ecliptic, 	<ul style="list-style-type: none"> • In Brāhmī inscriptions of the Kuṣāṇas month names are mostly seasonal : Gr̥ṣma, Varṣā, Hemanta • The pakṣa is not mentioned • Astro–nomical year starts when the Sun crosses the V.E., the civil year starts on the next day 	<ul style="list-style-type: none"> • 6 Seasons (R̥tu) in Tropical Year; each season consists of 2 zodiac signs; each zodiac sign covers 30° of the Ecliptic • Tropical Year = 365.25875 days (Varāha–mihira), 365.8756 days (Current S.S), 365.242196 days (Modern Astronomy) • Solar Year starts at Vaiśākha, Lunar Year starts at Caitra

Table 10 (Continued)

<i>Astro– nomical features</i>	<i>R̥k Saṁhitā</i>	<i>Yajuh̥ Saṁhitā</i>	<i>Vedāṅga Jyotiṣa</i>	<i>Sūrya Siddhānta</i>
...
Longer Periods of time	<ul style="list-style-type: none"> • Yuga (≡ Saros of Chaldean Astronomy) • Perhaps a 5 years Yuga 		<ul style="list-style-type: none"> • Yuga = 1830 sāvana days = 1860 tithis = 62 lunar months = 60 solar months = 67 nākṣatra months • Yuga begins at W.S. with the Sun and the moon at Dhaniṣṭhā • Mahāyuga 4.32×10^6 years 	<ul style="list-style-type: none"> • Kalpa 4.32×10^9 years

XI Comments on Astrology

Unlike in Babylonian civilization, astrology had no place in Hindu astronomy. This is perhaps because of Gautama Buddha's strictures against astrological predictions of doom and disorder. This is found in the Buddhist scripture *Dīgha Nikāya*, Vol 1, p. 65, (Pali Text Book Society)

Some brāhmaṇas and śramaṇas earn their livelihood by taking to beastly professions and eating food brought to them out of fear. They say: "There will be a solar eclipse, a lunar eclipse, occultation of the stars, the sun and the moon will move in the correct direction, in the incorrect direction, the nakṣatras will move in the correct path, in the incorrect path, there will be precipitation of meteors, burning of the cardinal directions (?), earthquakes, roar of heavenly war-drums, the sun, the moon and the stars will rise and set wrongly producing wide distress amongst all beings, etc. "

Kautilyīya Arthaśāstra also has the following stricture against astrological predictions.

"The objective (\equiv artha) eludes the foolish man (\equiv bālam) who enquires too much from the stars. The objective should be the nakṣatra of the objective, of what avail are the stars."

The practice of astrological prediction of human fate was perhaps imported from Babylonian civilization by the Śakas in the 1st century B.C. It is also to be noted that in the Hindu books of knowledge Astronomy was called Jyotiṣa whereas Astrology was called Sāmudrika Jyotiṣa. The epithet Sāmudrika indicates that this branch of knowledge came from beyond the boundaries of the country.

References

- [1] al-Bīrūnī, *Kitāb-ul-Hind* and *Āthār-al-Bāqia*, English translation *Alberuni's India* by E. C. Sachau, (London, 1914)
- [2] Āryabhaṭa, *Āryabhaṭiya*, English translation by P. C. Sengupta, (University of Calcutta, Calcutta)
- [3] John Bartholomew, *The Oxford School Atlas*, 17-th Edition, (Oxford University Press, Oxford, 1952)
- [4] Bhāskarācārya, *Siddhānta Śiromaṇi*, Edited by B. D. Sastri, (Benares, 1929)
- [5] Bhāskarācārya, *Golādhyāya*, Edited by G. D. Sastri, (Benares, 1929)
- [6] Bhaṭṭapāla, *Commentary on Br̥hajjātaka of Varāhamihira*, Edited by Sita Ram Jha, (Benares, 1934)
- [7] S. B. Diksīt, *Bharatiya Jyotiḥśāstra*
- [8] P. L. Gangoly (Editor), *Sūrya Siddhānta*, Reprint of the English translation with Notes and Appendix by E. Burgers, (University of Calcutta, Calcutta, 1936)
- [9] *Great World Atlas*, 3-rd Edition, (Reader's Digest Association, London, 1977)
- [10] Martin C. Gutzwiller, *Rev. Mod. Phys.* **70**, 589(1998)
- [11] R. C. Majumdar (General Editor), *The History and Culture of the Indian People*, Vol. III, (*The Classical Age*), 4-th Edition, (Bharatiya Vidya Bhavan, Bombay, 1988)
- [12] R. C. Majumdar (General Editor), *The History and Culture of the Indian People*, Vol. IV, (*The Age of Imperial Kanauj*), 4-th Edition, (Bharatiya Vidya Bhavan, Bombay, 1993)
- [13] R. C. Majumdar (General Editor), *The History and Culture of the Indian People*, Vol. V, (*The Struggle for Empire*), 4-th Edition, (Bharatiya Vidya Bhavan, Bombay, 1989)
- [14] M. N. Saha (Chairman), *Report of the Calendar Reform Committee*, (Government of India, New Delhi, 1955)
- [15] P. C. Sengupta, *Ancient Indian Chronology*

- [16] W. H. Smart, *Text Book on Spherical Astronomy*, (University Press, Cambridge, 1931)
- [17] Varāhamihira, *Bṛhat-saṁhitā*, English translation by Subrahmanya Sastri and M. Ramakrishna Bhat, 2 Vols, (Bangalore, 1947)
- [18] Varāhamihira, *Pañca Siddhāntikā*, Edited by G. Thibaut and S. Dvivedi, (Benares, 1889)
- [19] M. Winternitz, *A Short History of Indian Literature*, Vol. I, (University of Calcutta, Calcutta, 1927)