

On the mutual polarization of two He^4 atoms

Vadim M Loktev¹ and Maksim D Tomchenko²

Bogolyubov Institute for Theoretical Physics, 14b, Metrolohichna Str., 03680 Kiev,
Ukraine

E-mail: vloktev@bitp.kiev.ua; mtomchenko@bitp.kiev.ua

Abstract. We propose a simple method based on the standard quantum-mechanical perturbation theory to calculate the mutual polarization of two atoms He^4 .

PACS numbers: 32.10.Dk

Submitted to: *Paper*

In works [1, 2], it was shown that each of two interacting He^4 atoms induces a dipole moment

$$\mathbf{d} = -D_7 |e| \frac{a_B^8}{R^7} \mathbf{n}, \quad D_7 = 13.2-18.4 \quad (1)$$

on the neighboring atom. Here, $\mathbf{n} = \mathbf{R}/R$ is the unit vector toward the second atom, and $a_B = \hbar^2/me^2 = 0.529 \text{ \AA}$. We will call the polarization mechanism ‘‘tidal’’, since the deformation of electron shells of atoms in this case reminds gravitational tides. The calculation in [1, 2] was executed on the basis of special dispersion relations known mainly to experts in atomic physics. Therefore, we propose to solve the problem by a simpler method which does not leave the frame of the ordinary course of quantum mechanics and, in this case, will verify the result (1). In addition, the presented calculations are useful [3] for the explanation of the recently discovered electric activity of superfluid He II [4, 5].

According to the standard perturbation theory (PT), we will search for the ground-state wave function for two immovable interacting atoms He^4 , A and B, as the expansion in all possible states $\Psi_{\tilde{m}}$ of two free atoms A and B:

$$\Psi_0^{AB} = \sum_{\tilde{m}_A \tilde{m}_B} c_{\tilde{m}_A \tilde{m}_B} \Psi_{\tilde{m}_A}^A \Psi_{\tilde{m}_B}^B, \quad (2)$$

where \tilde{m}_A and \tilde{m}_B are the complete collections of quantum numbers (n, l, m) for these atoms. By $\Psi_{\tilde{l}} \equiv |\tilde{l}\rangle$, we denote the set of states with a given l and all possible corresponding m and n ($m = 0, \pm 1, \dots, \pm l$, $n = l, \dots, \infty$). In what follows, we will omit sometimes the tilde for simplicity: $\Psi_{\tilde{l}} \equiv \Psi_l$, $|\tilde{l}\rangle \equiv |l\rangle$, $c_{\tilde{0}\tilde{0}} \equiv c_{00}$, *etc.* Since the excited s-states (1sns) give the zero contribution to a DM (4), we neglect them, so that Ψ_0 means the ground state 1s1s of an atom He^4 . It is obvious that $c_{ij} = c_{ji}$. According to PT, we have

$$c_{ij} = c_{ij}^{(1)} + c_{ij}^{(2)} + \dots, \quad (3)$$

where the upper index shows the order of PT. The tidal DM of atom A is sought by the general formula

$$\mathbf{d}^A = \int \Psi_0^{*AB} \hat{\mathbf{d}}^A \Psi_0^{AB} d\mathbf{r}_1^A d\mathbf{r}_2^A d\mathbf{r}_1^B d\mathbf{r}_2^B, \quad (4)$$

where \mathbf{r}_j are the coordinates of electrons, and

$$\hat{\mathbf{d}}^A = e(\mathbf{r}_1^A + \mathbf{r}_2^A). \quad (5)$$

The function Ψ_0 describes a parastate. Therefore, the coefficients c_{ij} are nonzero only for parastates Ψ_j . In view of this fact, the spin function is the same for all Ψ_j in (2). Therefore, the spins can be took out in (4). Since the sum over spins gives 1, we omit the spin coordinates.

We note that the quantity $d^A \sim c_{10}(1) \sim (a_B/R)^4$ was determined in [6]. The coefficient c_{10} was found in the first order of PT, but its calculation includes an inaccuracy: in the charge operator \hat{Q} (see formulas (11)–(13) from [6]), the Z -axis was

chosen prior to the differentiation. But this axis should be fixed after all differentiations. In this last case, all the coefficients $c_{10}^{(1)}$ become zero, because $c_{10}^{(1)} \sim \langle 0^B | \hat{Q}_B | 0^B \rangle R^{-1}$ and

$$\langle 0^B | \hat{Q}_B | 0^B \rangle R^{-1} = 0. \quad (6)$$

In particular, $\langle 0^B | \hat{Q}_{B,q} | 0^B \rangle R^{-1} \sim \sum_{\alpha\beta} \delta_{\alpha\beta} \frac{\partial}{\partial R_\alpha} \frac{\partial}{\partial R_\beta} \frac{1}{R} = \Delta \frac{1}{R} \equiv 0$, and we get zero only for the Coulomb potential. Relation (6) is valid, because an helium atom has no moment in the ground state.

In this connection, we will determine c_{10} in the second order of PT:

$$c_{10}^{(2)} = - \frac{\langle 00 | \hat{U} | 00 \rangle \langle 10 | \hat{U} | 00 \rangle}{(E_{0,0}^{(0)} - E_{1,0}^{(0)})^2} + \sum_{\tilde{m} \neq (\tilde{0}, \tilde{0})} \frac{\langle \tilde{m} | \hat{U} | 00 \rangle \langle 10 | \hat{U} | \tilde{m} \rangle}{(E_{0,0}^{(0)} - E_{\tilde{m}}^{(0)})(E_{0,0}^{(0)} - E_{1,0}^{(0)})}, \quad (7)$$

where $\tilde{m} = (\tilde{m}_A, \tilde{m}_B)$, and

$$\langle \tilde{m}_1 | \hat{U} | \tilde{m}_2 \rangle = \int \Psi_{\tilde{m}_1^A}^{*A} \Psi_{\tilde{m}_1^B}^{*B} \hat{U} \Psi_{\tilde{m}_2^A}^A \Psi_{\tilde{m}_2^B}^B d\mathbf{r}_1^A d\mathbf{r}_2^A d\mathbf{r}_1^B d\mathbf{r}_2^B. \quad (8)$$

It follows from $\hat{U} = \hat{Q}_A^+ \hat{Q}_B (1/R)$ [7] and (6) that $\langle 00 | \hat{U} | 00 \rangle = 0$. Therefore, only the second term on the right-hand side in (7) is of importance. The main contribution to this term is given by the states $\tilde{m} = (1, 2)$ and $\tilde{m} = (2, 1)$. We have

$$c_{10}^{(2)} = \sum_{\tilde{m} \neq (\tilde{0}, \tilde{0})} \frac{\langle \tilde{m} | \hat{U} | 00 \rangle \langle 10 | \hat{U} | \tilde{m} \rangle}{\Delta E_{\tilde{m}} \Delta E_{1,0}}. \quad (9)$$

As n_1 and n_2 increase, the terms in (9) decrease. The calculation indicates that \mathbf{d}^A receives a significant contribution from a lot of states at c_{10} with $n > 2$ (in particular, $n = 5$ and 6). Since the number of states is huge, it seems impossible to find and to sum over all the required states in (9), as well as all c_{10} with different n and m .

Nevertheless, this difficulty can be avoided if the series is summed approximately. We note that the difference of energies for an atom He^4 is minimum for those of the first excited level $1s2p$ and the ground state, $\Delta E_1(n=2) = 21.236$ eV, and the maximum difference is represented by the ionization energy $\Delta E_{\text{ion}} = 24.58$ eV [8]. In (9), a significant contribution is given, for example, by the state $1s5p$ $\Delta E_1(n=5) = 24.066$ eV [8]. Thus, the dispersion of values of ΔE for different excited levels of parahelium is approximately $1/7$ of the value of ΔE . Since ΔE are sufficiently close for different n , we set them to be equal to a constant ΔE .

Let us supplement (9) by the term $\tilde{m} = (\tilde{0}, \tilde{0}) \langle 00 | \hat{U} | 00 \rangle \langle 10 | \hat{U} | 00 \rangle / (2(\Delta E)^2)$ (it is equal to zero due to $\langle 00 | \hat{U} | 00 \rangle = 0$). With regard for the fact that $\Delta E_{1,0} \equiv E_{0,0}^{(0)} - E_{1,0}^{(0)} = 2E_0^{(0)} - E_1^{(0)} - E_0^{(0)} = E_0^{(0)} - E_1^{(0)} \approx -\Delta E$ and $\Delta E_{\tilde{m}} \equiv E_{0,0}^{(0)} - E_{\tilde{m}}^{(0)} = 2E_0^{(0)} - E_{\tilde{m}_A}^{(0)} - E_{\tilde{m}_B}^{(0)} \approx -2\Delta E$ in (9), we obtain

$$c_{10}^{(2)} \approx \frac{1}{2(\Delta E)^2} \sum_{\tilde{m}} \langle 10 | \hat{U} | \tilde{m} \rangle \langle \tilde{m} | \hat{U} | 00 \rangle = \frac{1}{2(\Delta E)^2} \langle 10 | \hat{U}^2 | 00 \rangle, \quad (10)$$

where $c_{10}^{(2)}$ depends on n and m in the state $|\tilde{1}\rangle$. By writing (10), we used the well-known property of completeness

$$\sum_{\tilde{m}} |\tilde{m}\rangle \langle \tilde{m}| = 1, \quad (11)$$

where the sum is taken over all states of both atoms. By considering only the terms with c_{00} , c_{10} , and c_{01} in (2), relation (4) yields the first part of the DM of atom A:

$$\begin{aligned} \mathbf{d}_1^A &= \int d\mathbf{r}_1^A d\mathbf{r}_2^A d\mathbf{r}_1^B d\mathbf{r}_2^B \hat{\mathbf{d}}^A |c_{00}\Psi_0^A\Psi_0^B \\ &+ \sum_{\tilde{1}} [c_{10}^{(2)}\Psi_1^A\Psi_0^B + c_{01}^{(2)}\Psi_0^A\Psi_1^B]|^2 \\ &\approx c_{00}^* \sum_{\tilde{1}} c_{10}^{(2)} \langle 00|\hat{\mathbf{d}}^A|10\rangle + \text{c.c.} \\ &= \frac{c_{00}^*}{2(\Delta E)^2} \sum_{\tilde{1}} \langle 00|\hat{\mathbf{d}}^A|10\rangle \langle 10|\hat{U}^2|00\rangle + \text{c.c.} \end{aligned} \quad (12)$$

Here, $\sum_{\tilde{1}}$ is the sum over all states $|\tilde{1}\rangle$, i.e., over $m = 0; \pm 1$ and $n = 2, 3, \dots, \infty$. Besides $|10\rangle$, we include all the remaining states $|\tilde{m}_A\tilde{m}_B\rangle$ in (12): $|1^A\rangle = |1snp\rangle$, and also $|1sns\rangle$, $|1snd\rangle$, $|1snf\rangle$, *etc.*; and, in addition to $|0^B\rangle$, all higher states $|\tilde{m}\rangle$. This is possible to make because $\langle 00|\hat{\mathbf{d}}^A|\tilde{m}_A\tilde{m}_B\rangle = 0$ for them. Then, with regard for (11), we obtain

$$\begin{aligned} \mathbf{d}_1^A &\approx \frac{c_{00}^*}{2(\Delta E)^2} \sum_{\tilde{m}_A\tilde{m}_B} \langle 00|\hat{\mathbf{d}}^A|\tilde{m}_A\tilde{m}_B\rangle \langle \tilde{m}_A\tilde{m}_B|\hat{U}^2|00\rangle + \text{c.c.} \\ &= \frac{c_{00}}{(\Delta E)^2} \langle 00|\hat{\mathbf{d}}^A\hat{U}^2|00\rangle. \end{aligned} \quad (13)$$

Relation (11) and the normalization conditions for Ψ_0^{AB} yield

$$c_{00}^2 = 1 - \sum_{l_1j_1 \neq (0,0)} c_{l_1j_1}^2 \approx 1 - \frac{\langle 00|\hat{U}^2|00\rangle}{4(\Delta E)^2} = 1 + \frac{E_{\text{vdW}}}{2\Delta E}, \quad (14)$$

where $E_{\text{vdW}} \approx -\frac{\langle 00|\hat{U}^2|00\rangle}{2\Delta E} \approx -2.15e^2(a_B^5/R^6)$ is the van der Waals interaction energy for two atoms He^4 . We have $|E_{\text{vdW}}| \ll \Delta E$; therefore, $c_{00} \approx 1$.

With regard for (2) and (4), the total DM of atom A is given by the formula

$$\mathbf{d}^A = \sum_{\tilde{m}_1^A\tilde{m}_2^A\tilde{m}_1^B\tilde{m}_2^B} c_{\tilde{m}_1^A\tilde{m}_1^B}^* c_{\tilde{m}_2^A\tilde{m}_2^B} \langle \tilde{m}_1^A\tilde{m}_1^B|\hat{\mathbf{d}}^A|\tilde{m}_2^A\tilde{m}_2^B\rangle. \quad (15)$$

The above-derived expression for \mathbf{d}_1 is a part of the last relation from the terms $c_{00}c_{10}$ in the second order of PT. The formula for \mathbf{d}^A includes also the nonzero contribution in the first order of PT from terms of the form $c_{11}^{(1)}c_{21}^{(1)}$, $c_{22}^{(1)}c_{21}^{(1)}$, and higher-order ones which behave themselves, respectively, as $\sim (a_B/R)^7$, $\sim (a_B/R)^9$, *etc.* (see below). At l and j which are not equal to zero ($\tilde{0}$) simultaneously, the relation

$$c_{lj}^{(1)} = \frac{\langle l^A j^B|\hat{U}|0^A 0^B\rangle}{E_0^{(0)} - E_j^{(0)} + E_0^{(0)} - E_l^{(0)}} \quad (16)$$

is valid. Since $\langle lj|\hat{U}|00\rangle \neq 0$ only at $l, j \geq 1$, we set $2E_0^{(0)} - E_j^{(0)} - E_l^{(0)} \approx -2\Delta E$. Since the contribution in the DM from c_{00} is taken into account in \mathbf{d}_1 , we redefine c_{00} in

(15) according to (16) with $-2\Delta E$ in the denominator (such c_{00} is equal to zero). Since $c_{10}^{(1)} = 0$, we will write also $-2\Delta E$ in the denominator for it instead of $-\Delta E$. We can now use the relation (11), and relation (15) minus \mathbf{d}_1 is reduced to

$$\begin{aligned}\mathbf{d}_2^A &\approx \frac{1}{4(\Delta E)^2} \sum_{j_1 l_1 j_2 l_2} \langle 00 | \hat{U} | l_1 j_1 \rangle \langle l_1 j_1 | \hat{\mathbf{d}}^A | l_2 j_2 \rangle \langle l_2 j_2 | \hat{U} | 00 \rangle \\ &= \frac{1}{4(\Delta E)^2} \langle 0^A 0^B | \hat{\mathbf{d}}^A \hat{U}^2 | 0^A 0^B \rangle = \frac{\mathbf{d}_1^A}{4}.\end{aligned}\quad (17)$$

Whence we obtain the total DM of atom A in the principal ($\sim (a_B/R)^7$) approximation:

$$\mathbf{d}^A = \mathbf{d}_1^A + \mathbf{d}_2^A \approx \frac{5}{4(\Delta E)^2} \langle 0^A 0^B | \hat{\mathbf{d}}^A \hat{U}^2 | 0^A 0^B \rangle. \quad (18)$$

Using the formulas for the operator of charge \hat{Q} [6], we represent the interatomic potential \hat{U} as the sum

$$\hat{U} = \hat{Q}_A^+ \hat{Q}_B \frac{1}{R} = \hat{U}_{dd} + \hat{U}_{qd} + \hat{U}_{qq} + \dots, \quad (19)$$

where

$$\hat{U}_{dd} = \frac{\hat{\mathbf{d}}^A \hat{\mathbf{d}}^B}{R^3} - \frac{3(\hat{\mathbf{d}}^A \mathbf{R})(\hat{\mathbf{d}}^B \mathbf{R})}{R^5} \quad (20)$$

is the dipole-dipole part of the potential, and

$$\begin{aligned}\hat{U}_{qd} &= \frac{3}{R^4} \sum_{\alpha\beta=1}^3 \left(\hat{Q}_{\alpha\beta}^A \left[\hat{d}_\beta^B n_\alpha + \hat{d}_\alpha^B n_\beta - 5\hat{\mathbf{d}}^B \mathbf{n} n_\alpha n_\beta \right] \right) \\ &+ \frac{3}{R^4} \hat{\mathbf{d}}^B \mathbf{n} \sum_{\alpha=1}^3 \hat{Q}_{\alpha\alpha}^A - (A \leftrightarrow B)\end{aligned}\quad (21)$$

is the dipole-quadrupole part. The quadrupole-quadrupole part, \hat{U}_{qq} , includes a negligible correction to the DM ($\sim (a_B/R)^9$ from $\hat{U}_{qd}\hat{U}_{qq}$). Therefore,

$$\hat{U}^2 \approx \hat{U}_{dd}^2 + 2\hat{U}_{dd}\hat{U}_{qd} + \hat{U}_{qd}^2. \quad (22)$$

In view of the parity relative to the inversion, we preserve only the term $2\hat{U}_{dd}\hat{U}_{qd}$. As a result, we obtain

$$\mathbf{d}^A \approx \frac{5}{2(\Delta E)^2} \langle 0^A 0^B | \hat{\mathbf{d}}^A \hat{U}_{dd} \hat{U}_{qd} | 0^A 0^B \rangle. \quad (23)$$

Let us choose the Z -axis along \mathbf{R} : $\mathbf{R} = R\mathbf{i}_z$. Then a nonzero contribution to \mathbf{d}^A is given by the terms

$$\begin{aligned}\hat{\mathbf{d}}^A \hat{U}_{dd} \hat{U}_{qd} &= \frac{6}{R^7} \left\{ (\hat{d}_x^B)^2 \hat{d}_x^A \hat{d}_z^A \hat{Q}_{xz}^A + (x \leftrightarrow y) \right. \\ &\left. - (\hat{d}_z^B)^2 (\hat{d}_z^A)^2 (\hat{Q}_{xx}^A + \hat{Q}_{yy}^A - 2\hat{Q}_{zz}^A) \right\}.\end{aligned}\quad (24)$$

With regard for the isotropy $\Psi_0 = |0\rangle$, we eventually arrive at the general formula

$$\begin{aligned}\mathbf{d}^A &\approx \frac{30\mathbf{i}_z}{R^7(\Delta E)^2} \langle 0^B | (\hat{d}_z^B)^2 | 0^B \rangle \left(\langle 0^A | \hat{d}_x^A \hat{d}_z^A \hat{Q}_{xz}^A | 0^A \rangle \right. \\ &\left. + \langle 0^A | (\hat{d}_z^A)^2 (\hat{Q}_{zz}^A - \hat{Q}_{xx}^A) | 0^A \rangle \right),\end{aligned}\quad (25)$$

which can be rewritten in the following simple form:

$$\mathbf{d}_A = -D_7 |e| \frac{a_B^8}{R^7} \mathbf{n}, \quad (26)$$

$$D_7 = Z_A Z_B \left(\frac{2Ry}{\Delta E} \right)^2 \langle \tilde{r}_1^2 \rangle_B \langle \tilde{r}_1^4 \rangle_A. \quad (27)$$

Here, $\langle \tilde{r}_1^2 \rangle_B \equiv \langle 0^B | (r_1^B/a_B)^2 | 0^B \rangle$, $\langle \tilde{r}_1^4 \rangle_A \equiv \langle 0^A | (r_1^A/a_B)^4 | 0^A \rangle$, $Ry = e^2/2a_B = 13.6 \text{ eV}$.

The next correction to \mathbf{d}_A is of the order of $\sim (a_B/R)^9$, and it is $\sim (R/a_B)^2$ times less than (26). For He II, $R \sim 3.6 \text{ \AA}$, $(R/a_B)^2 \simeq 50$; therefore, it can be neglected. It is of interest that the separate corrections to D_7 for various n and m in the $(\tilde{1}\tilde{1})$ and $(\tilde{1}\tilde{2})$ states from (2) are small, about 0.1 and less. Moreover, we have an alternating series for \mathbf{d}_A , so that several hundreds of various corrections and about 10 different n significantly contribute to the total DM D_7 (27).

The positive sign of D_7 means that the electron cloud of each of two atoms He^4 shifts to the neighboring atom. That is, the interatomic space acquires an excess of the negative charge.

To calculate the value of D_7 , we need to know the wave function $|0\rangle$ of the ground state of an atom He^4 . The exact analytic solution is not available for it, but some approximations have been proposed. Within the simplest one-parameter one [7, 9]

$$\Psi_0 = \varphi_{1s}(\mathbf{r}_1)\varphi_{1s}(\mathbf{r}_2), \quad \varphi_{1s}(\mathbf{r}) = \sqrt{Z^{*3}/\pi a_B^3} e^{-rZ^*/a_B} \quad (28)$$

($Z^* = Z - 5/16$), we get $\langle \tilde{r}_1^2 \rangle = 3/(Z^*)^2 \approx 1.053$, $\langle \tilde{r}_1^4 \rangle = 45/2(Z^*)^4 \approx 2.775$. In (27) for ΔE , we take the least ($\Delta E_1(n=2)$) and largest (ΔE_{ion}) values, so we obtain $D_7 \approx 14.32-19.18$ (here we take into account also that $Z_A = Z_B = 2$).

The more exact multiparameter Ψ_0 were calculated within various methods. In [10], it was proposed to seek Ψ_0 in the form

$$\Psi_0(s, t, u) = \varphi(ks, kt, ku), \quad (29)$$

where

$$\varphi(s, t, u) = Ae^{-s/2} \sum_{l,n,m=0}^{\infty} c_{n,2l,m} s^n t^{2l} u^m, \quad (30)$$

$$s = r_1 + r_2, \quad t = r_1 - r_2, \quad u = r_{12}.$$

The formalism in the variables s, u, t is given in [11] in detail. For the 3-parameter Ψ_0 [10, 11],

$$\varphi = Ae^{-s/2}(1 + c_1 u + c_2 t^2), \quad (31)$$

$$c_1 \approx 0.081, \quad c_2 \approx 0.01, \quad k \approx 3.63,$$

the calculation yields

$$A = (k^3/4\pi) (4 + 35c_1 + 48c_2 + 96c_1^2 + 576c_2^2 + 308c_1c_2)^{-1/2} \approx 1.325, \quad (32)$$

$$\begin{aligned} \langle \tilde{r}_1^2 \rangle &= \frac{768\pi^2 A^2}{k^8} \left(1 + \frac{189}{16}c_1 + 24c_2 \right. \\ &\quad \left. + 42c_1^2 + 480c_2^2 + 195c_1c_2 \right), \end{aligned} \quad (33)$$

$$\begin{aligned} \langle \tilde{r}_1^4 \rangle &= \frac{23040\pi^2 A^2}{k^{10}} \left(1 + \frac{979}{64}c_1 + 52c_2 \right. \\ &\quad \left. + 68c_1^2 + 1704c_2^2 + \frac{8305}{16}c_1c_2 \right). \end{aligned} \quad (34)$$

We have also performed the numerical calculations for 6-, 10-, and 20-parameter Hylleraas functions [10, 12, 13] and for more general 10-, 39-, and 80-parameter Kinoshita functions [14, 15] of the form (29) with

$$\varphi(s, t, u) = A e^{-s/2} \sum_{l,n,m=0}^{\infty} c_{n,m,l} s^{n+1} (t/u)^{2l} (u/s)^m \quad (35)$$

which include the negative degrees in the expansion. The results are given in the Table I. For the 3-parameter Ψ_0 , the result is obtained numerically and analytically. It is seen from the Table I that the more the number of parameters, the less the variation of the results for $\langle \tilde{r}_1^2 \rangle$, $\langle \tilde{r}_1^4 \rangle$, and D_7 . Moreover, the convergence is quite good. For comparison, we present also a deviation of the theoretical ionization potential J of atoms He^4 from the experimental value, $J^{\text{exp}} \approx 198310.8 \text{ cm}^{-1}$, by the data from [9, 15, 16]. It is seen that the convergence for the energy is faster, which is natural. Indeed, at the determination of Ψ_0 , namely the energy is minimized. However, the quantities $\langle \tilde{r}_1^2 \rangle$ and $\langle \tilde{r}_1^4 \rangle$ are determined also with a high accuracy. In work [16], Ψ_0 was calculated by means of the expansion in Laguerre polynomials. Up to 1078 terms were taken into account, and the determined value of $\langle \tilde{r}_1^2 \rangle$ (see Table I) coincides with accuracy to 4 decimal points with the value obtained by us for the 80-parameter Ψ_0 . This means that the above-determined $\langle \tilde{r}_1^2 \rangle$, $\langle \tilde{r}_1^4 \rangle$, and the upper and lower limits of D_7 for this Ψ_0 have error less than 0.1%. The modern more exact methods of calculations of Ψ_0 are available (see, e.g., [17, 18]), but the too high accuracy is not required for our task, because it will not increase the accuracy of D_7 , since the large error of D_7 remains unremovable due to the ‘‘smearing’’ of ΔE .

Since the main contribution to the DM is given by the states with large n , for which $\Delta E \approx \Delta E_{\text{ion}}$, D_7 should be closer to the lower limit. Using formula (27) and the table of energies of different excited levels [8], it can be seen that the value of D_7 belongs to the interval from $(D_7^{\text{max}} + D_7^{\text{min}})/2$ to D_7^{min} . We can estimate the exact value of D_7 , by determining the arithmetic mean for this interval. Therefore, we define D_7 as

$$\bar{D}_7 \approx D_7^{\text{min}} + (D_7^{\text{max}} - D_7^{\text{min}})/4. \quad (36)$$

The remaining uncertainty will be taken into account in the error term. In view of the result for the 80-parameter Ψ_0 , we finally have

$$D_7 \approx \bar{D}_7 \approx 25.2 \pm 2, \quad (37)$$

where the error covers the whole interval from $(D_7^{\text{max}} + D_7^{\text{min}})/2$ to D_7^{min} .

Table 1. Dependence of various quantities on the number of parameters of Ψ_0 (the wave function of the ground state of He^4 atom).

Number of parameters of Ψ_0	Ref.	$\langle \tilde{r}_1^2 \rangle$	$\langle \tilde{r}_1^4 \rangle$	D_7^{\min}	D_7^{\max}	\bar{D}_7	$ \frac{J^{\text{theor}} - J^{\text{exp}}}{J^{\text{exp}}} $
1	[9, 7]	1.053	2.775	14.32	19.18	15.53	$5 \cdot 10^{-2}$
3	[10, 9]	1.182	3.815	22.1	29.6	23.98	$2.5 \cdot 10^{-3}$
6	[10, 9]	1.188	3.861	22.47	30.1	24.38	$5 \cdot 10^{-4}$
10	[12]	1.249	4.222	25.82	34.6	28.02	$1 \cdot 10^{-4}$
10	[14]	1.19	3.91	22.79	30.54	24.73	$1 \cdot 10^{-6}$
20	[13]	1.19326	3.967	23.19	31.07	25.16	$2 \cdot 10^{-6}$
39	[15]	1.19314	3.961	23.15	31.02	25.12	$1 \cdot 10^{-6}$
80	[15]	1.19346	3.973	23.23	31.12	25.2	$2 \cdot 10^{-7}$
1078	[16]	1.19348	—	—	—	—	$5 \cdot 10^{-7}$

Solution (26), (37) is in agreement with (1), though we have obtained a somewhat larger value for D_7 . We note that the formula different from (27) for D_7 was derived in [1, 2] with the use of dispersion sums. The most exact value, $D_7 = 18.4$, was obtained there on the basis of the 20-parameter Ψ_0 taken from [13]. In this case, certain approximations were used in calculations, but the error arising due to these approximations was not evaluated. For this and more exact functions Ψ_0 , we have calculated a more larger value of D_7 (37). The difference of values of D_7 in (37) and (1) is probably related to the approximations made in both calculations. We note that the above-proposed calculation is much simpler than that in [1, 2].

It is possible that D_7 can be determined more exactly within special numerical methods. But, even in this case, our solution retains its validity, since it was obtained by a relatively simple verifiable analytic method. The calculation error for D_7 is determined by the approximate character of Ψ_0 and by the difference of the energies of excited levels of a He atom. However, at a large number of parameters for Ψ_0 , the first error becomes small (see Table 1), and the second error is approximately equal to the relative dispersion of the values of ΔE , i.e., to within 15%. Thus, the method is rather exact. The main error of the method is related to the change of values of ΔE for different levels in sums of the form (9) by some constant value. Therefore, the method works well only for atoms, for which the energies of excited states are close to one another. One of such atoms is a He^4 atom.

The above-considered (and earlier in [1, 2]) tidal mechanism of the polarization of atoms is, in our opinion, a key to the comprehension of electric properties of superfluid He^4 and other fluids of nonpolar atoms, though we cannot yet exclude the contribution of other mechanisms (for example, the thermoemf).

Acknowledgments

The authors thank A.S. Rybalko, S.I. Shevchenko, G.E. Volovik for the fruitful discussions and Yu.V. Shtanov for the valuable remark concerning the calculation. The work is performed under the financial support of the Division of Physics and Astronomy of the NAS of Ukraine in the frame of a Target program of fundamental studies N 0107U000396.

References

- [1] Byers Brown W and Whisnant D M 1973 *Mol. Phys.* **25** 1385
- [2] Whisnant D M and Byers Brown W 1973 *Mol. Phys.* **26** 1105
- [3] Tomchenko M D 2011 *Phys. Rev. B* **83** 094512 (*Preprint* cond-mat/1003.4389v4)
- [4] Rybalko A S 2004 *Fiz. Nizk. Temp.* **30** 1321 [2004 *Low Temp. Phys.* **30** 994]
- [5] Rybalko A S and Rubets S P 2005 *Fiz. Nizk. Temp.* **31** 820 [2005 *Low Temp. Phys.* **31** 623]
- [6] Loktev V M and Tomchenko M D 2008 *Fiz. Nizk. Temp.* **34** 337 [2008 *Low Temp. Phys.* **34** 262]
- [7] Vakarchuk I O 2004 *Quantum Mechanics* (L'viv:L'viv National University) p 783
- [8] Herzberg G 1948 *Atomic Spectra and Atomic Structure* [Russian translation] (Moscow: IL) p. 279
- [9] Bethe H A and Salpeter E E 1957 *Quantum Mechanics of One- and Two-Electron Atoms* (Berlin: Springer) p. 575
- [10] Hylleraas E 1928 *Z. Phys.* **48** 469; 1929 *Z. Phys.* **54** 347
- [11] Sommerfeld A 1951 *Atombau und Spektrallinien* vol. 2 (Braunschweig: Friedr. Vieweg und Sohn) p. 634
- [12] Chandrasekhar S, Elbert D and Herzberg G 1953 *Phys. Rev.* **91** 1172
- [13] Hart J F and Herzberg G 1957 *Phys. Rev.* **106** 79
- [14] Kinoshita T 1957 *Phys. Rev.* **105** 1490
- [15] Kinoshita T 1959 *Phys. Rev.* **115** 366
- [16] Pekeris C L 1959 *Phys. Rev.* **115** 1216
- [17] Goldman S P 1994 *Phys. Rev. Lett.* **73** 2547
- [18] Mitroy J, Bromley M W J and Ratnavelu K 2007 *Int. J. Quant. Chem.* **107** 907 (*Preprint* physics.atom-ph/0607232)