

Polygamy of Distributed Entanglement

Francesco Buscemi,¹ Gilad Gour,^{2,3} and Jeong San Kim^{2,*}

¹Statistical Laboratory, DPMMS, University of Cambridge, Wilberforce Road, CB3 0WB, UK

²Institute for Quantum Information Science, University of Calgary, Alberta T2N 1N4, Canada

³Department of Mathematics and Statistics, University of Calgary, Alberta T2N 1N4, Canada

(Dated: November 23, 2018)

While quantum entanglement is known to be monogamous (i.e. shared entanglement is restricted in multi-partite settings), here we show that distributed entanglement (or the potential for entanglement) is by nature polygamous. By establishing the concept of one-way unlocalizable entanglement (UE) and investigating its properties, we provide a polygamy inequality of distributed entanglement in tripartite quantum systems of arbitrary dimension. We also provide a polygamy inequality in multi-qubit systems, and several trade offs between UE and other correlation measures.

PACS numbers: 03.67.-a, 03.67.Hk, 03.65.Ud,

I. INTRODUCTION

Quantum entanglement is a non-local quantum correlation providing a lot of useful applications in the field of quantum communications and computations such as quantum teleportation and quantum key distribution [1, 2, 3]. This important role of quantum entanglement has stimulated intensive study in both way of its quantification and qualification.

One of the essential differences of quantum correlations (especially, quantum entanglement) from other classical ones is that it cannot be freely shared among the parties in multipartite quantum systems. In particular, a pair of components that are maximally entangled cannot share entanglement [4, 5] nor classical correlations [6] with any part of the rest of the system, hence the term *Monogamy of Entanglement* (MoE) [7]. Monogamy of entanglement was shown to have a complete mathematical characterization for multi-qubit systems [5] using a certain entanglement measures, the concurrence [8].

Whereas MoE shows the restricted sharability of multi-party quantum entanglement, the distribution of entanglement, or *Entanglement of Assistance* (EoA) [9, 10] in multipartite quantum systems was shown to have a dually monogamous (or *Polygamous*) property. Using *Concurrence of Assistance* (CoA) [11] as the measure of distributed entanglement, it was also shown that whereas monogamy of entanglement inequalities provide an upper bound for bipartite sharability of entanglement in a multipartite system, the same quantity provides a lower bound for distribution of bipartite entanglement in a multipartite system [12]. In this paper, by introducing the concept of *One-way Unlocalizable Entanglement* (UE), we provide a polygamy inequality of entanglement in tripartite quantum systems of arbitrary dimension using entropic entanglement measure. Based on the functional relation between concurrence and entropic measure in

two-qubit systems, we provide a polygamy inequality in multi-qubit systems. We also provide several trade offs between UE and other correlations such as EoA, and localizable entanglement.

The paper is organized as follows. In Sec. II, we provide the definition of UE, and its basic properties. In Sec. III, we provide a polygamy inequality of distributed entanglement in tripartite quantum systems in terms of entropy and EoA. In Sec. IV, we generalize the polygamy inequality of entanglement into multi-qubit systems, and provide a more tight polygamy inequality for three-qubit systems. In Sec. V, we provide several trade offs between UE and other correlations, and we summarize our results, in Sec. VI.

II. ONE-WAY UNLOCALIZABLE ENTANGLEMENT

A. Definition

For any bipartite quantum state ρ_{AB} , its one-way distillable common randomness [13] is defined as

$$C_D^{\leftarrow}(\rho_{AB}) = \lim_{n \rightarrow \infty} \frac{1}{n} I^{\leftarrow}(\rho_{AB}^{\otimes n}), \quad (1)$$

where, the function $I^{\leftarrow}(\rho_{AB})$ [14] is

$$I^{\leftarrow}(\rho_{AB}) = \max_{\{M_x\}} \left[S(\rho_A) - \sum_x p_x S(\rho_A^x) \right], \quad (2)$$

and where the maximum is taken over all the measurements $\{M_x\}$ applied on system B . Here, $S(\rho_A)$ is the von Neumann entropy of $\rho_A \equiv \text{tr}_B(\rho_{AB})$, $p_x \equiv \text{tr}[(I_A \otimes M_x)\rho_{AB}]$ is the probability of the outcome x , and $\rho_A^x \equiv \text{tr}_B[(I_A \otimes M_x)\rho_{AB}]/p_x$ is the state of system A when the outcome was x .

For a tripartite pure state $|\psi\rangle_{ABC}$ with $\rho_A = \text{tr}_{BC}|\psi\rangle_{ABC}\langle\psi|$, $\rho_{AB} = \text{tr}_C|\psi\rangle_{ABC}\langle\psi|$, and $\rho_{AC} = \text{tr}_B|\psi\rangle_{ABC}\langle\psi|$, it was shown that [6]

$$S(\rho_A) = I^{\leftarrow}(\rho_{AB}) + E_f(\rho_{AC}). \quad (3)$$

*Electronic address: jkim@qis.ucalgary.ca

Here, $E_f(\rho_{AC})$ is the *Entanglement of Formation* (EoF) of ρ_{AC} defined as [15]

$$E_f(\rho_{AC}) = \min \sum_i p_i S(\rho_A^i), \quad (4)$$

where the minimization is taken over all pure state decomposition of ρ_{AC} such that,

$$\rho_{AC} = \sum_i p_i |\phi^i\rangle_{AC} \langle \phi^i|, \quad (5)$$

with $\text{tr}_C |\phi^i\rangle_{AC} \langle \phi^i| = \rho_A^i$.

As a dual quantity to EoF, EoA is defined by the maximum average entanglement of ρ_{AC} ,

$$E_a(\rho_{AC}) = \max \sum_i p_i S(\rho_A^i), \quad (6)$$

over all possible pure state decompositions of ρ_{AC} .

Definition 1. The one-way unlocalizable entanglement (UE) of a bipartite state ρ_{AB} is defined as follows:

$$E_u^{\leftarrow}(\rho_{AB}) := S(\rho_A) - E_a(\rho_{AC}), \quad (7)$$

where ρ_{AC} denotes the reduced state of a purification $|\psi\rangle_{ABC}$ of ρ_{AB} .

The one-way unlocalizable entanglement can be equivalently characterized as follows:

Lemma 1. For any given bipartite state ρ_{AB} , its one-way unlocalizable entanglement is given by

$$E_u^{\leftarrow}(\rho_{AB}) = \min_{\{M_x\}} \left[S(\rho_A) - \sum_x p_x S(\rho_A^x) \right], \quad (8)$$

where the minimum is taken over all possible rank-1 measurements $\{M_x\}$ applied on subsystem B .

Proof. Eq. (8) can be rewritten as

$$E_u^{\leftarrow}(\rho_{AB}) = S(\rho_A) - \max_{\{M_x\}} \sum_x p_x S(\rho_A^x), \quad (9)$$

where the maximum is taken over all possible rank-1 measurements $\{M_x\}$ applied on system B .

Since $|\psi\rangle_{ABC}$ is a pure state, all possible pure state decompositions of ρ_{AC} can be realized by rank-1 measurements of subsystem B , and conversely, any rank-1 measurement can be induced from a pure state decomposition of ρ_{AC} . Thus, the second term on the right hand side of Eq. (9) is the maximum average entanglement over all possible pure state decomposition of ρ_{AC} , which is the definition of $E_a(\rho_{AC})$, and this completes the proof. \square

By definition, the UE of ρ_{AB} is the difference between $S(\rho_A)$ and $E_a(\rho_{AC})$. Here, $S(\rho_A)$ quantifies the entanglement of the pure state $|\psi\rangle_{A(BC)}$ with respect to the bipartite cut $A-BC$, whereas $E_a(\rho_{AC})$ measures the maximum average entanglement that can be localized on the subsystem AC with the assistance of B . The terminology used is then clear. Figure 1 graphically illustrates this separation.




Fig. 1: The entanglement of $|\psi\rangle_{A(BC)}$ ((a) in the figure) consists of two different parts: One is the localizable entanglement onto AC represented by $E_a(\rho_{AC})$ ((b) in the figure), and the other part is the unlocalizable entanglement represented by $E_u^{\leftarrow}(\rho_{AB})$ ((c) in the figure).

B. Properties

1. Subadditivity

Lemma 2. For all bipartite states ρ_{AB} and $\sigma_{A'B'}$,

$$E_u^{\leftarrow}(\rho_{AB} \otimes \sigma_{A'B'}) \leq E_u^{\leftarrow}(\rho_{AB}) + E_u^{\leftarrow}(\sigma_{A'B'}), \quad (10)$$

where

$$E_u^{\leftarrow}(\rho_{AB} \otimes \sigma_{A'B'}) = \min_{\{L_z\}} \left[S(\rho_A \otimes \sigma_{A'}) - \sum_z r_z S(\tau_{AA'}^z) \right], \quad (11)$$

with $r_z = \text{tr}[(I_{AA'} \otimes L_z)\rho_{AB} \otimes \sigma_{A'B'}]$, $\tau_{AA'}^z = \text{tr}_{BB'}[(I_{AA'} \otimes L_z)\rho_{AB} \otimes \sigma_{A'B'}]/r_z$, and the minimum is taken over all possible rank-1 measurements $\{L_z\}$ applied on subsystem BB' .

Proof. Let $\{M_x\}$ and $\{N_y\}$ be the optimal rank-1 measurements on subsystems B and B' for $E_u^{\leftarrow}(\rho_{AB})$ and $E_u^{\leftarrow}(\sigma_{A'B'})$ respectively, then, we have

$$\begin{aligned} & E_u^{\leftarrow}(\rho_{AB}) + E_u^{\leftarrow}(\sigma_{A'B'}) \\ &= S(\rho_A) + S(\sigma_{A'}) - \sum_x p_x S(\rho_A^x) - \sum_y q_y S(\sigma_{A'}^y) \\ &= S(\rho_A \otimes \sigma_{A'}) - \sum_{xy} p_x q_y S(\rho_A^x \otimes \sigma_{A'}^y) \\ &\geq E_u^{\leftarrow}(\rho_{AB} \otimes \sigma_{A'B'}), \end{aligned} \quad (12)$$

where $p_x \rho_A^x = \text{tr}_B[(I_A \otimes M_x)\rho_{AB}]$, $q_y \sigma_{A'}^y = \text{tr}_{B'}[(I_{A'} \otimes N_y)\sigma_{A'B'}]$, and the second equality is due to the additivity of von Neumann entropy and the definition of $E_u^{\leftarrow}(\rho_{AB} \otimes \sigma_{A'B'})$. \square

By Lemma 2, we can assure the existence of the *regularized* UE

$$E_{u,\infty}^{\leftarrow}(\rho_{AB}) := \lim_{n \rightarrow \infty} \frac{E_u^{\leftarrow}(\rho_{AB}^{\otimes n})}{n}, \quad (13)$$

which satisfies

$$E_{u,\infty}^{\leftarrow}(\rho_{AB}) \leq E_u^{\leftarrow}(\rho_{AB}). \quad (14)$$

2. Simple Lower Bound

Lemma 3. For any bipartite state ρ_{AB} ,

$$E_u^{\leftarrow}(\rho_{AB}) \geq \max\{I_c^{\leftarrow}(\rho_{AB}), 0\}, \quad (15)$$

where $I_c^{\leftarrow}(\rho_{AB}) := S(\rho_A) - S(\rho_{AB})$ is the coherent information of ρ_{AB} .

Proof. Let $|\psi\rangle_{ABC}$ be a purification of ρ_{AB} , then due to the monotonicity of entanglement, we have

$$E_a(\rho_{AC}) \leq \min\{S(\rho_A), S(\rho_C)\}, \quad (16)$$

where $\rho_{AC} = \text{tr}_B|\psi\rangle_{ABC}\langle\psi|$.

Thus, together with Lemma 1, we have

$$\begin{aligned} E_u^{\leftarrow}(\rho_{AB}) &= S(\rho_A) - E_a(\rho_{AC}) \\ &\geq \max\{S(\rho_A) - S(\rho_C), 0\} \\ &= \max\{I_c^{\leftarrow}(\rho_{AB}), 0\}, \end{aligned} \quad (17)$$

where the last equality is due to the purity of $|\psi\rangle_{ABC}$, that is, $S(\rho_C) = S(\rho_{AB})$. \square

Since $|\psi\rangle_{ABC}^{\otimes n}$ is a purification of both $\rho_{AB}^{\otimes n}$ and $\rho_{BC}^{\otimes n}$, we have

$$E_u^{\leftarrow}(\rho_{AB}^{\otimes n}) + E_a(\rho_{AC}^{\otimes n}) = nS(\rho_A). \quad (18)$$

By taking the limit $n \rightarrow \infty$, and due to the relation [16]

$$\lim_{n \rightarrow \infty} \frac{E_a(\rho_{AC}^{\otimes n})}{n} = \min\{S(\rho_A), S(\rho_C)\}, \quad (19)$$

we have that

$$E_{u,\infty}^{\leftarrow}(\rho_{AB}) = \max\{I_c^{\leftarrow}(\rho_{AB}), 0\}. \quad (20)$$

Eq. (20) implies that, in the asymptotic limit of many copies, separable states do not exhibit quantumness in their correlations, or their correlations are completely erasable. This is a strong evidence that the distinction between separable and entangled states is *operational* only in asymptotic sense, since separable states can exhibit non-zero UE in finite case.

III. POLYGAMY OF ENTANGLEMENT IN TRIPARTITE QUANTUM SYSTEMS

For any bipartite pure state $|\phi\rangle_{AB}$, its concurrence, $\mathcal{C}(|\phi\rangle_{AB})$ is defined as [8]

$$\mathcal{C}(|\phi\rangle_{AB}) = \sqrt{2(1 - \text{tr}\rho_A^2)}, \quad (21)$$

where $\rho_A = \text{tr}_B(|\phi\rangle_{AB}\langle\phi|)$. For any mixed state ρ_{AB} , its concurrence is defined via *convex-roof extension*, that is,

$$\mathcal{C}(\rho_{AB}) = \min \sum_k p_k \mathcal{C}(|\phi_k\rangle_{AB}), \quad (22)$$

where the minimum is taken over all possible pure state decompositions, $\rho_{AB} = \sum_k p_k |\phi_k\rangle_{AB}\langle\phi_k|$.

As a dual value to concurrence, CoA [11] of ρ_{AB} is defined as

$$\mathcal{C}^a(\rho_{AB}) = \max \sum_k p_k \mathcal{C}(|\phi_k\rangle_{AB}), \quad (23)$$

where the maximum is taken over all possible pure state decompositions of ρ_{AB} .

By using concurrence and CoA as the quantification of bipartite entanglement, it was shown that there exists a polygamy relation of entanglement in multi-qubit systems [12]. More precisely, for any pure state $|\psi\rangle_{A_1 \dots A_n}$ in an n -qubit system $\mathcal{H}_{A_1} \otimes \dots \otimes \mathcal{H}_{A_n}$ where $\mathcal{H}_{A_i} \cong \mathbb{C}^2$ for $i = 1, \dots, n$,

$$\mathcal{C}_{A_1(A_2 \dots A_n)}^2 \leq (\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2, \quad (24)$$

where $\mathcal{C}_{A_1(A_2 \dots A_n)}$ is the concurrence of $|\psi\rangle_{A_1 \dots A_n}$ with respect to the bipartite cut A_1 and $A_2 \dots A_n$, and $\mathcal{C}_{A_1 A_i}^a$ is the CoA of $\rho_{A_1 A_i} = \text{tr}_{A_2 \dots A_{i-1} A_{i+1} \dots A_n}(|\psi\rangle_{A_1 \dots A_n}\langle\psi|)$ for $i = 2, \dots, n$.

In this section, we provide an analytic upper bound of UE in Eq. (8), and derive a polygamy inequality of entanglement in terms of von-Neumann entropy and EoA for tripartite quantum systems of arbitrary dimension.

First, for an upper bound of UE, we have the following theorem.

Theorem 4. For any bipartite state ρ_{AB} in a bipartite quantum system $\mathcal{H}_A \otimes \mathcal{H}_B$,

$$E_u^{\leftarrow}(\rho_{AB}) \leq \frac{I(\rho_{AB})}{2}, \quad (25)$$

where $I(\rho_{AB}) = S(\rho_A) + S(\rho_B) - S(\rho_{AB})$ is the mutual information of ρ_{AB} .

Proof. Let $\rho_B = \sum_{i=0}^{d_B-1} \lambda_i |e_i\rangle_B \langle e_i|$ be a spectral decomposition of $\rho_B = \text{tr}_A(\rho_{AB})$ where d_B is the dimension of the subsystem \mathcal{H}_B . The proof method follows the construction used in [17].

For any state $\sigma \in \mathcal{H}_B$, define the channels

$$\begin{aligned} M_0(\sigma) &:= \sum_{i=0}^{d_B-1} |e_i\rangle \langle e_i| \sigma |e_i\rangle \langle e_i| \\ M_1(\sigma) &:= \sum_{i=0}^{d_B-1} |\tilde{e}_i\rangle \langle \tilde{e}_i| \sigma |\tilde{e}_i\rangle \langle \tilde{e}_i|, \end{aligned} \quad (26)$$

where $\{|\tilde{e}_j\rangle\}_j$ is the *Fourier basis* such that,

$$|\tilde{e}_j\rangle = \frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} \omega_d^{jk} |e_k\rangle, \quad j = 0, \dots, d-1, \quad (27)$$

and $\omega_d = e^{\frac{2\pi i}{d}}$ is the d -th root of unity.

Notice that $M_0(\rho_B) = \rho_B$, and $M_1(\rho_B) = \frac{1}{d_B}I_B$, so that $M_1(M_0(\rho_B)) = M_0(M_1(\rho_B)) = \frac{1}{d_B}I_B$. We can also write

$$M_0(\sigma) = \frac{1}{d_B} \sum_{b=0}^{d_B-1} Z^b \sigma Z^{-b}, \quad M_1(\sigma) = \frac{1}{d_B} \sum_{a=0}^{d_B-1} X^a \sigma X^{-a}, \quad (28)$$

where Z and X are generalized d_B -dimensional Pauli operators,

$$Z = \sum_{j=0}^{d_B-1} \omega_d^j |e_j\rangle\langle e_j|, \\ X = \sum_{j=0}^{d_B-1} |e_{j+1}\rangle\langle e_j| = \sum_{j=0}^{d_B-1} \omega_d^{-j} |\tilde{e}_j\rangle\langle \tilde{e}_j|. \quad (29)$$

In the following, we will write

$$(I_A \otimes M_0)(\rho_{AB}) = \sum_{i=0}^{d_B-1} \sigma_A^i \otimes \lambda_i |e_i\rangle_B \langle e_i|, \\ (I_A \otimes M_1)(\rho_{AB}) = \sum_{j=0}^{d_B-1} \tau_A^j \otimes \frac{1}{d_B} |\tilde{e}_j\rangle_B \langle \tilde{e}_j|, \quad (30)$$

where $\lambda_i \sigma_A^i = \text{tr}_B[(I_A \otimes |e_i\rangle_B \langle e_i|)\rho_{AB}]$, and $\tau_A^j/d_B = \text{tr}_B[(I_A \otimes |\tilde{e}_j\rangle_B \langle \tilde{e}_j|)\rho_{AB}]$ for $i, j \in \{0, \dots, d_B-1\}$.

The induced ensembles on A by the channels M_0 and M_1 will be denoted by $\mathcal{E}_0 := \{\lambda_i, \sigma_A^i\}_i$ and $\mathcal{E}_1 := \{\frac{1}{d_B}, \tau_A^j\}_j$, and the entropy defects of the induced ensembles on A will be denoted as

$$\chi(\mathcal{E}_0) = S(\rho_A) - \sum_{i=0}^{d_B-1} \lambda_i S(\sigma_A^i), \\ \chi(\mathcal{E}_1) = S(\rho_A) - \frac{1}{d_B} \sum_{i=0}^{d_B-1} S(\tau_A^i). \quad (31)$$

By defining a four-partite quantum state Ω_{XYAB} in $\mathcal{B}(\mathbb{C}^{d_B} \otimes \mathbb{C}^{d_B} \otimes \mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$ such that

$$\Omega_{XYAB} := \frac{1}{d_B^2} \sum_{x,y=0}^{d_B-1} |x\rangle_X \langle x| \otimes |y\rangle_Y \langle y| \otimes (I_A \otimes X_B^x Z_B^y) \rho_{AB} (I_A \otimes Z_B^{-y} X_B^{-x}), \quad (32)$$

we have

$$\Omega_{XAB} = \frac{1}{d_B} \sum_{x=0}^{d_B-1} |x\rangle_X \langle x| \otimes X_B^x \left(\sum_{i=0}^{d_B-1} \sigma_A^i \otimes \lambda_i |e_i\rangle_B \langle e_i| \right) X_B^{-x}, \\ \Omega_{YAB} = \frac{1}{d_B} \sum_{y=0}^{d_B-1} |y\rangle_Y \langle y| \otimes Z_B^y \left(\sum_{j=0}^{d_B-1} \tau_A^j \otimes \frac{1}{d_B} |\tilde{e}_j\rangle_B \langle \tilde{e}_j| \right) Z_B^{-y}, \quad (33)$$

and

$$\Omega_{AB} = \rho_A \otimes \frac{I_B}{d_B}. \quad (34)$$

By straightforward calculation, we can obtain

$$I(\Omega_{X(AB)}) = S(\Omega_X) + S(\Omega_{AB}) - S(\Omega_{XAB}) \\ = \log d_B + \log d_B + S(\rho_A) - \log d_B \\ - S\left(\sum_i \sigma_A^i \otimes \lambda_i |e_i\rangle_B \langle e_i|\right) \\ = \log d_B + S(\rho_A) - H(\vec{\lambda}) - \sum_i \lambda_i S(\sigma_A^i) \\ = \log d_B - S(\rho_B) + \chi(\mathcal{E}_0), \quad (35)$$

where $I(\Omega_{X(AB)})$ is the mutual information of $\Omega_{X(AB)}$ with respect to the bipartite cut $X-AB$, and the second, third equalities are due to the *joint entropy theorem* [18]. Analogously, we have

$$I(\Omega_{Y(AB)}) = \chi(\mathcal{E}_1), \\ I(\Omega_{(XY)AB}) = \log d_B + S(\rho_A) - S(\rho_{AB}) \\ = \log d_B + I_c^{\leftarrow}(\rho_{AB}). \quad (36)$$

Due to the independence of subsystems X and Y , we have $I(\Omega_{(XY)AB}) \geq I(\Omega_{X(AB)}) + I(\Omega_{Y(AB)})$, which implies

$$\chi(\mathcal{E}_0) + \chi(\mathcal{E}_1) \leq I(\rho_{AB}). \quad (37)$$

Since $\chi(\mathcal{E}_0)$ and $\chi(\mathcal{E}_1)$ of Eq. (37) can be ob-

tained, respectively, from ρ_{AB} by rank-1 measurements $\{|e_i\rangle_B\langle e_i|\}_i$ and $\{|\tilde{e}_j\rangle_B\langle \tilde{e}_j|\}_j$ of subsystem B , by defining a rank-1 measurement

$$\left\{ \frac{|e_i\rangle_B\langle e_i|}{2}, \frac{|\tilde{e}_j\rangle_B\langle \tilde{e}_j|}{2} \right\}_{i,j}, \quad (38)$$

we have

$$E_u^{\leftarrow}(\rho_{AB}) \leq \frac{\chi(\mathcal{E}_0)}{2} + \frac{\chi(\mathcal{E}_1)}{2} \leq \frac{I(\rho_{AB})}{2}, \quad (39)$$

which completes the proof. \square

Corollary 1. *For any tripartite pure state $|\psi\rangle_{ABC}$, we have*

$$S(\rho_A) \leq E_a(\rho_{AB}) + E_a(\rho_{AC}). \quad (40)$$

Proof. By Lemma 7, we have

$$\begin{aligned} E_a(\rho_{AC}) &= S(\rho_A) - E_u^{\leftarrow}(\rho_{AB}), \\ E_a(\rho_{AB}) &= S(\rho_A) - E_u^{\leftarrow}(\rho_{AC}), \end{aligned} \quad (41)$$

and thus,

$$E_a(\rho_{AC}) + E_a(\rho_{AB}) = 2S(\rho_A) - E_u^{\leftarrow}(\rho_{AB}) - E_u^{\leftarrow}(\rho_{AC}). \quad (42)$$

Now, by Theorem 4, we have

$$\begin{aligned} &E_a(\rho_{AC}) + E_a(\rho_{AB}) \\ &\geq 2S(\rho_A) - \frac{I(\rho_{AB})}{2} - \frac{I(\rho_{AC})}{2} \\ &= 2S(\rho_A) - S(\rho_A)/2 - S(\rho_B)/2 + S(\rho_{AB})/2 \\ &\quad - S(\rho_A)/2 - S(\rho_C)/2 + S(\rho_{AC})/2 \\ &= S(\rho_A). \end{aligned} \quad (43)$$

\square

Corollary 1 tells us that for a tripartite pure state of arbitrary dimension, there exists a polygamy relation of entanglement in terms of entropy of entanglement and EoA. Furthermore, this is, we believe, the first result of the polygamous (or dually monogamous) property of distribution of entanglement in multipartite higher-dimensional quantum systems rather than qubits.

IV. POLYGAMY RELATION OF ENTANGLEMENT IN MULTI-QUBIT QUANTUM SYSTEMS

In this section, we show that the polygamy inequality of entanglement in Corollary 1 can be generalized into multipartite quantum systems for the case when each subsystem is a two-level quantum system. By investigating the functional relation between concurrence and EoF in two-qubit systems [8], we show that there exists

a polygamy inequality of entanglement in terms of entropy and EoA in n -qubit systems. We also show that, in three-qubit systems, we have a more tight polygamy inequality than Eq. (40) in Corollary 1.

First, let us consider the functional relation of concurrence with EoF in two-qubit systems. For a 2-qubit mixed state ρ_{AB} (or a pure state $|\psi\rangle_{AB} \in \mathbb{C}^2 \otimes \mathbb{C}^d$), the relation between its concurrence, \mathcal{C}_{AB} and $E_f(\rho_{AB})$ can be given as a monotone increasing, convex function \mathcal{E} [8], such that

$$E_f(\rho_{AB}) = \mathcal{E}(\mathcal{C}_{AB}), \quad (44)$$

where

$$\mathcal{E}(x) = H\left(\frac{1}{2} + \frac{1}{2}\sqrt{1-x^2}\right), \quad \text{for } 0 \leq x \leq 1, \quad (45)$$

and $H(\cdot)$ is the binary entropy function $H(x) = -[x \log_2 x + (1-x) \log_2 (1-x)]$. The same function $\mathcal{E}(x)$ relates also the EoA of a bipartite state ρ_{AB} with its CoA via the equation

$$E_a(\rho_{AB}) \geq \mathcal{E}(\mathcal{C}_{AB}^a), \quad (46)$$

which is due to the convexity of \mathcal{E} and the definition of EoA. The following lemma shows an important property of the function $\mathcal{E}(x)$.

Lemma 5.

$$\mathcal{E}(\sqrt{x^2 + y^2}) \leq \mathcal{E}(x) + \mathcal{E}(y), \quad (47)$$

for $0 \leq x, y \leq 1$ such that $0 \leq x^2 + y^2 \leq 1$.

Proof. By considering

$$f(x, y) = \mathcal{E}(x) + \mathcal{E}(y) - \mathcal{E}(\sqrt{x^2 + y^2}), \quad (48)$$

as a two-variable real-valued function on the domain $D = \{(x, y) | 0 \leq x, y \leq 1, 0 \leq x^2 + y^2 \leq 1\}$, it is enough to show that $f(x, y) \geq 0$ in D .

Since D is a compact subset in \mathbb{R}^2 , whereas f is analytic on the interior of D , and continuous on D , the minimum value of f arises only on the critical points or on the boundary of D . It can be directly checked that f does not have any vanishing gradient on the interior of D , and has non-negative function values on the boundary of D . Thus, f is non-negative on the domain D . \square

A. Three-qubit systems

A direct observation from [4] shows that, for a 3-qubit pure state $|\psi\rangle_{ABC}$,

$$\mathcal{C}_{A(BC)}^2 = \mathcal{C}_{AB}^2 + (\mathcal{C}_{AC}^a)^2, \quad (49)$$

where \mathcal{C}_{AB} and \mathcal{C}_{AC}^a are the concurrence and concurrence of assistance of ρ_{AB} and ρ_{AC} respectively. (Later, Eq. (49) was formally shown in [19].) From Eq. (49) together with Lemma 5, we have the following theorem.

Theorem 6. For a three-qubit pure state $|\psi\rangle_{ABC}$,

$$S(\rho_A) \leq E_f(\rho_{AB}) + E_a(\rho_{AC}). \quad (50)$$

Proof. Since $|\psi\rangle_{ABC}$ is a bipartite pure state in $\mathbb{C}^2 \otimes \mathbb{C}^4$ with respect to the bipartite cut A and BC , we have,

$$S(\rho_A) = E_f(\psi_{A(BC)}) = \mathcal{E}(\mathcal{C}_{A(BC)}). \quad (51)$$

Thus,

$$\begin{aligned} S(\rho_A) &= \mathcal{E}(\mathcal{C}_{A(BC)}) \\ &= \mathcal{E}(\sqrt{\mathcal{C}_{AB}^2 + \mathcal{C}_{AC}^2}) \\ &\leq \mathcal{E}(\mathcal{C}_{AB}) + \mathcal{E}(\mathcal{C}_{AC}) \\ &\leq E_f(\rho_{AB}) + E_a(\rho_{AC}), \end{aligned} \quad (52)$$

where the first inequality is by Lemma 5, and the second inequality is by Eq. (46). \square

Thus, the polygamy relation of distributed entanglement in tripartite quantum systems obtained in Corollary 1 can have a more tight form in three-qubit systems. Furthermore, the result of Theorem 6 together with Eqs. (3) and (7) give us the following corollary.

Corollary 2. For any two-qubit mixed state ρ_{AB} with rank less than or equal to two,

$$I^\leftarrow(\rho_{AB}) \leq E_a(\rho_{AB}), \quad (53)$$

$$E_u^\leftarrow(\rho_{AB}) \leq E_f(\rho_{AB}). \quad (54)$$

Remark 1. Eq. (54) of Corollary 2 implies that any two-qubit separable state ρ_{AB} of rank less than or equal to two has zero UE, $E_u^\leftarrow(\rho_{AB}) = 0$. However, this is not generally true for two-qubit separable states of rank larger than two. Here, we provide an example of two-qubit rank-three separable state with non-zero UE.

Example: Let us consider the following state in $\mathbb{C}^2 \otimes \mathbb{C}^3$ quantum system [20],

$$|\Psi\rangle_{ABC} = \frac{1}{\sqrt{2}}|x\rangle_{AC}|0\rangle_B + \frac{1}{\sqrt{2}}|y\rangle_{AC}|1\rangle_B, \quad (55)$$

where $|x\rangle$ and $|y\rangle$ are two orthogonal states in the $\mathbb{C}^2 \otimes \mathbb{C}^3$ such that

$$\begin{aligned} |x\rangle &= (|02\rangle + \sqrt{2}|10\rangle)/\sqrt{3}, \\ |y\rangle &= (|12\rangle + \sqrt{2}|01\rangle)/\sqrt{3}. \end{aligned} \quad (56)$$

First, since $\rho_A = (|0\rangle_A\langle 0| + |1\rangle_A\langle 1|)/2$, it is clear that $\mathcal{C}_{A(BC)} = \sqrt{2(1 - \text{tr}\rho_A^2)} = 1$, therefore we have $S(\rho_A) = 1$.

Since $\rho_{AC} = (|x\rangle_{AC}\langle x| + |y\rangle_{AC}\langle y|)/2$, Hughston-Jozsa-Wootters (HJW) theorem [21] says that for any decompositions of $\rho_{AC} = \sum_i p_i |\phi_i\rangle_{AC}\langle \phi_i|$, there exists a unitary operator (u_{ij}) such that $\sqrt{p_i}|\phi_i\rangle_{AC} = (u_{i1}|x\rangle_{AC} + u_{i2}|y\rangle_{AC})/\sqrt{2}$ with $2p_i = |u_{i1}|^2 + |u_{i2}|^2$. Thus,

$$\begin{aligned} \text{tr}_C(|\phi_i\rangle_{AC}\langle \phi_i|) &= \frac{1}{6p_i} \begin{pmatrix} |u_{i1}|^2 + 2|u_{i2}|^2 & u_{i1}u_{i2}^* \\ u_{i2}u_{i1}^* & |u_{i2}|^2 + 2|u_{i1}|^2 \end{pmatrix} \\ &= \frac{1}{3}I_A + \frac{1}{3}|\psi_i\rangle_A\langle \psi_i|, \end{aligned} \quad (57)$$

with $|\psi_i\rangle = (u_{i2}^*|0\rangle + u_{i1}^*|1\rangle)/\sqrt{2p_i}$, and we obtain that $\mathcal{C}(|\phi_i\rangle_{AC}) = \frac{2\sqrt{2}}{3}$ for any pure state $|\phi_i\rangle_{AC}$ in any pure state decomposition of ρ_{AC} .

Since $|\phi_i\rangle_{AC}$ is a $2 \otimes 3$ pure state, we have

$$\begin{aligned} E_f(|\phi_i\rangle_{AC}) &= \mathcal{E}(\mathcal{C}(|\phi_i\rangle_{AC})) \\ &= H\left(\frac{2}{3}\right) \\ &= \log_2 3 - \frac{2}{3}, \end{aligned} \quad (58)$$

and thus $E_f(\rho_{AC}) = E_a(\rho_{AC}) = \log_2 3 - \frac{2}{3}$.

Now, we have $E_u^\leftarrow(\rho_{AB}) = S(\rho_A) - E_a(\rho_{AC}) = \frac{5}{3} - \log_2 3 > 0$, whereas, it can be easily seen that ρ_{AB} has a *Positive Partial Transposition* (PPT) which is equivalent to separability for two-qubit states [22]. Thus, ρ_{AB} is a two-qubit, rank-three separable state with non-zero UE.

B. n -qubit systems

The polygamy inequality of entanglement in n -qubit systems in Eq. (24) gives us an inequality

$$\mathcal{C}_{A_1(A_2 \dots A_n)} \leq \sqrt{(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2}. \quad (59)$$

Thus, together with Lemma 5, we have the following theorem.

Theorem 7. For any n -qubit pure state $|\psi\rangle_{A_1 \dots A_n}$,

$$S(\rho_{A_1}) \leq E_a(\rho_{A_1 A_2}) + \dots + E_a(\rho_{A_1 A_n}). \quad (60)$$

Proof. First, let us assume that $(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2 \leq 1$, then we have

$$\begin{aligned} S(\rho_{A_1}) &= \mathcal{E}(\mathcal{C}_{A_1(A_2 \dots A_n)}) \\ &\leq \mathcal{E}\left(\sqrt{(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2}\right) \\ &\leq \mathcal{E}(\mathcal{C}_{A_1 A_2}^a) + \mathcal{E}\left(\sqrt{(\mathcal{C}_{A_1 A_3}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2}\right) \\ &\leq \mathcal{E}(\mathcal{C}_{A_1 A_2}^a) + \mathcal{E}(\mathcal{C}_{A_1 A_3}^a) + \dots + \mathcal{E}(\mathcal{C}_{A_1 A_n}^a) \\ &\leq E_a(\rho_{A_1 A_2}) + \dots + E_a(\rho_{A_1 A_n}), \end{aligned} \quad (61)$$

where the first inequality is due to the monotonicity of the function \mathcal{E} , the second and third inequalities are obtained by iterating Lemma 5, and the last inequality is by Eq. (46).

Now, assume that $(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_n}^a)^2 > 1$. Since $S(\rho_{A_1}) \leq 1$ for any n -qubit pure state $|\psi\rangle_{A_1 \dots A_n}$, it is enough to show that $E_a(\rho_{A_1 A_2}) + \dots + E_a(\rho_{A_1 A_n}) \geq 1$.

Let us first note that there exist $k \in \{2, \dots, n-1\}$ that satisfies

$$\begin{aligned} (\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_k}^a)^2 &\leq 1, \\ (\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_{k+1}}^a)^2 &> 1, \end{aligned} \quad (62)$$

and let

$$T = (\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_{k+1}}^a)^2 - 1. \quad (63)$$

Since, $(\mathcal{C}_{A_1 A_{k+1}}^a)^2 - T = 1 - (\mathcal{C}_{A_1 A_2}^a)^2 - \dots - (\mathcal{C}_{A_1 A_k}^a)^2$, we have

$$0 \leq (\mathcal{C}_{A_1 A_{k+1}}^a)^2 - T \leq 1, \quad (64)$$

and

$$\begin{aligned} 1 &= \mathcal{E} \left(\sqrt{(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_{k+1}}^a)^2 - T} \right) \\ &\leq \mathcal{E} \left(\sqrt{(\mathcal{C}_{A_1 A_2}^a)^2 + \dots + (\mathcal{C}_{A_1 A_k}^a)^2} \right) \\ &\quad + \mathcal{E} \left(\sqrt{(\mathcal{C}_{A_1 A_{k+1}}^a)^2 - T} \right) \\ &\leq \mathcal{E}(\mathcal{C}_{A_1 A_2}^a) + \dots + \mathcal{E}(\mathcal{C}_{A_1 A_k}^a) + \mathcal{E}(\mathcal{C}_{A_1 A_{k+1}}^a) \\ &\leq E_a(\rho_{A_1 A_2}) + \dots + E_a(\rho_{A_1 A_n}), \end{aligned} \quad (65)$$

where the first and second inequalities are by Lemma 5 and by monotonicity of \mathcal{E} , and the last inequality is by Eq. (46). \square

V. UNLOCALIZABLE ENTANGLEMENT VERSUS OTHER MEASURES OF CORRELATION

In this section, we provide some properties of UE concerned with several other correlation measures. By investigating the relation between UE and EoF in $2 \otimes 2 \otimes d$ quantum system, we show that any two-qubit state with zero UE is a separable state. We also provide a quantitative relation among entropy, localizable entanglement, and UE for tripartite mixed states.

A. $2 \otimes 2 \otimes d$ pure state

Let $|\psi\rangle_{ABC}$ be a tripartite pure state in $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^d$.

Theorem 8. For any 2-qubit state ρ_{AB} ,

$$E_u^{\leftarrow}(\rho_{AB}) = 0 \implies \rho_{AB} \text{ separable}, \quad (66)$$

or, equivalently, if $S(\rho_A) = E_a(\rho_{AC})$, then $E_f(\rho_{AB}) = 0$.

Proof. Suppose $S(\rho_A) = E_a(\rho_{AC})$, and let $\rho_{AC} = \sum_i p_i |\phi^i\rangle_{AC} \langle \phi^i|$ be an optimal decomposition such that,

$$\begin{aligned} E_a(\rho_{AC}) &= \sum_i p_i E(|\phi^i\rangle_{AC}) \\ &= \sum_i p_i S(\rho_A^i), \end{aligned} \quad (67)$$

where $\rho_A^i = \text{tr}_C(|\phi^i\rangle_{AC} \langle \phi^i|)$ and $\rho_A = \sum_i p_i \rho_A^i$.

The concavity of von Neumann entropy says, $S(\rho_A) \geq \sum_i p_i S(\rho_A^i)$ and the equality holds if and only if ρ_A^i are identical for all i . So, by the assumption, ρ_A^i are identical for all i .

Since $|\phi^i\rangle_{AC}$ is a pure state in $2 \otimes d$ system and its concurrence is $\mathcal{C}(|\phi^i\rangle_{AC}) = 2\sqrt{\det \rho_A^i}$, we also have that $\mathcal{C}(|\phi^i\rangle_{AC})$ are identical for all i , say $\mathcal{C}(|\phi^i\rangle_{AC}) = \mathcal{C}_{AC}^*$.

Now, we have

$$S(\rho_A^i) = E_f(|\phi^i\rangle_{AC}) = \mathcal{E}(\mathcal{C}(|\phi^i\rangle_{AC})) = \mathcal{E}(\mathcal{C}_{AC}^*), \quad (68)$$

for all i , and

$$\begin{aligned} \mathcal{E}(\mathcal{C}_{A(BC)}) &= E_f(|\psi_{A(BC)}\rangle) \\ &= S(\rho_A) \\ &= \sum_i p_i S(\rho_A^i) \\ &= \sum_i p_i \mathcal{E}(\mathcal{C}(|\phi^i\rangle_{AC})) \\ &= \mathcal{E}(\mathcal{C}_{AC}^*), \end{aligned} \quad (69)$$

where $\mathcal{C}_{A(BC)}$ is the concurrence of $|\psi\rangle_{ABC}$ between subsystems A and BC , and $\mathcal{E}(\cdot)$ is the function in Eq. (45).

Since $\mathcal{E}(\cdot)$ is strictly monotone increasing, (the first derivative $\frac{d}{d\mathcal{C}} \mathcal{E}(\mathcal{C})$ is 0 at $\mathcal{C} = 0$ and positive elsewhere), we have

$$\mathcal{C}_{A(BC)} = \mathcal{C}_{AC}^*, \quad (70)$$

therefore

$$\mathcal{C}_{A(BC)} \geq \mathcal{C}_{AC}^a \geq \sum_i p_i \mathcal{C}(|\phi^i\rangle_{AC}) = \mathcal{C}_{AC}^* = \mathcal{C}_{A(BC)}, \quad (71)$$

and thus,

$$\mathcal{C}_{A(BC)} = \mathcal{C}_{AC}^a. \quad (72)$$

Now, by the Theorem 3 in [20], we have $\mathcal{C}(\rho_{AB}) = 0$ where ρ_{AB} is a 2-qubit state, which implies $E_f(\rho_{AB}) = 0$. \square

Any two-qubit state with zero UE is separable by Theorem 8, and any two-qubit separable state with rank less than or equal to two has zero UE by Corollary 2. However, the converse of Theorem 8 is not generally true, since Remark 1 provides us a two-qubit separable state with non-zero UE.

B. Tripartite Mixed State

Since it is known that the EoA is not a bipartite measure nor an *entanglement monotone* [23], it is not clear yet if there is any quantitative relation between $E_a(\rho_{AB})$ and $E_a(\rho_{A(BD)})$ for a tripartite mixed state ρ_{ABD} . In fact, this is equivalent to the quantitative relation between $E_u^\leftarrow(\rho_{AC})$ and $E_u^\leftarrow(\rho_{A(CD)})$. This is because, if we consider a purification $|\psi\rangle_{ABCD}$ of ρ_{ABD} , then any direction of a quantitative relation between $E_a(\rho_{AB})$ and $E_a(\rho_{A(BD)})$, say $E_a(\rho_{AB}) \leq E_a(\rho_{A(BD)})$, would give us

$$\begin{aligned} S(\rho_A) &= E_u^\leftarrow(\rho_{A(CD)}) + E_a(\rho_{AB}) \\ &= E_u^\leftarrow(\rho_{AC}) + E_a(\rho_{A(BD)}) \\ &\geq E_u^\leftarrow(\rho_{AC}) + E_a(\rho_{AB}), \end{aligned} \quad (73)$$

which implies $E_u^\leftarrow(\rho_{AC}) \leq E_u^\leftarrow(\rho_{A(CD)})$.

In this section, we pay our attentions only to local rank-1 measurements of each subsystems, and we derive a quantitative relation between localizable entanglement, and UE for tripartite mixed states.

For $\rho_{ACD} = \text{tr}_B |\psi\rangle_{ABCD} \langle \psi|$, let us define

$$\tilde{E}_u^\leftarrow(\rho_{A(CD)}) = \min_{\{M_x \otimes N_y\}} \left[S(\rho_A) - \sum_{x,y} p_{xy} S(\rho_A^{xy}) \right], \quad (74)$$

where $p_{xy} \equiv \text{tr}[(I_A \otimes M_x \otimes N_y)\rho_{ACD}]$ is the probability of the outcome x and y on subsystems C and D respectively, and $\rho_A^{xy} \equiv \text{tr}_{CD}[(I_A \otimes M_x \otimes N_y)\rho_{ACD}]/p_{xy}$ is the state of system A when the outcome were x and y . The minimum in Eq. (74) is taken over all possible rank-1 measurements $\{M_x\}$ and $\{N_y\}$ on subsystems C and D respectively. By definition, we have

$$\tilde{E}_u^\leftarrow(\rho_{A(CD)}) \geq E_u^\leftarrow(\rho_{A(CD)}). \quad (75)$$

Furthermore, we have the following lemma.

Lemma 9. *For any tripartite state ρ_{ACD} ,*

$$\tilde{E}_u^\leftarrow(\rho_{A(CD)}) \geq E_u^\leftarrow(\rho_{AC}). \quad (76)$$

Proof. For ρ_{ACD} , let $\{M_x\}$ and $\{N_y\}$ are the optimal rank-1 measurements of C and D respectively, such that

$$\tilde{E}_u^\leftarrow(\rho_{A(CD)}) = S(\rho_A) - \sum_{x,y} p_{xy} S(\rho_A^{xy}). \quad (77)$$

Due to the concavity of von Neumann entropy, we have

$$\begin{aligned} \tilde{E}_u^\leftarrow(\rho_{A(CD)}) &= S(\rho_A) - \sum_{x,y} p_{xy} S(\rho_A^{xy}) \\ &= S(\rho_A) - \sum_x p_x \left[\sum_y \frac{p_{xy}}{p_x} S(\rho_A^{xy}) \right] \\ &\geq S(\rho_A) - \sum_x p_x S\left(\sum_y \frac{p_{xy}}{p_x} \rho_A^{xy}\right) \\ &= S(\rho_A) - \sum_x p_x S(\rho_A^x) \\ &\geq E_u^\leftarrow(\rho_{AC}), \end{aligned} \quad (78)$$

where

$$\begin{aligned} p_x &= \text{tr}_{AC} [(I_A \otimes M_x)\rho_{AC}] \\ &= \text{tr}_{ACD} [(I_A \otimes M_x \otimes I_D)\rho_{ACD}] \\ &= \sum_y \text{tr}_{ACD} [(I_A \otimes M_x \otimes N_y)\rho_{ACD}] \\ &= \sum_y p_{xy}, \end{aligned} \quad (79)$$

$$\begin{aligned} \rho_A^x &= \frac{1}{p_x} \text{tr}_C [(I_A \otimes M_x)\rho_{AC}] \\ &= \frac{1}{p_x} \text{tr}_{CD} [(I_A \otimes M_x \otimes I_D)\rho_{ACD}] \\ &= \sum_y \frac{1}{p_x} \text{tr}_{CD} [(I_A \otimes M_x \otimes N_y)\rho_{ACD}] \\ &= \sum_y \frac{p_{xy}}{p_x} \rho_A^{xy}, \end{aligned} \quad (80)$$

and the second inequality is due to the definition of $E_u^\leftarrow(\rho_{AC})$. \square

Now, we are ready to have the following theorem.

Theorem 10. *For any tripartite mixed state ρ_{ABC} with a purification $|\psi\rangle_{ABCD}$,*

$$S(\rho_A) \geq \tilde{E}_a(\rho_{AB}) + E_u^\leftarrow(\rho_{AC}), \quad (81)$$

where $\tilde{E}_a(\rho_{AB})$ is the localizable entanglement [24] of ρ_{AB} , defined by

$$\tilde{E}_a(\rho_{AB}) = \max_{\{M_x \otimes N_y\}} \sum_{x,y} p_{xy} S(\rho_A^{xy}) \quad (82)$$

over all possible rank-1 measurements $\{M_x\}$ and $\{N_y\}$ on subsystems C and D respectively.

Proof. Eq. (74) can be rewritten as

$$\begin{aligned} \tilde{E}_u^\leftarrow(\rho_{A(CD)}) &= S(\rho_A) - \max_{\{M_x \otimes N_y\}} \sum_{x,y} p_{xy} S(\rho_A^{xy}) \\ &= S(\rho_A) - \tilde{E}_a(\rho_{AB}), \end{aligned} \quad (83)$$

and Lemma 9 completes the proof. \square

Theorem 10 can be considered as an alternative of Lemma 1 for mixed states case. Furthermore, Theorem 10 together with Lemma 1 give us the following simple corollary.

Corollary 3. *For any tripartite mixed state ρ_{ABC} with a purification $|\psi\rangle_{ABCD}$,*

$$E_a(\rho_{A(BC)}) \geq \tilde{E}_a(\rho_{AB}). \quad (84)$$

Proof. By Theorem 10, we have

$$S(\rho_A) \geq \tilde{E}_a(\rho_{AB}) + E_u^+(\rho_{AD}) \quad (85)$$

for any pure state $|\psi\rangle_{ABCD}$, whereas

$$S(\rho_A) = E_a(\rho_{A(BC)}) + E_u^+(\rho_{AD}), \quad (86)$$

for the tripartite partition $A - BC - D$. \square

VI. CONCLUSION

In this paper, we have proposed the concept of UE, and shown that the polygamous nature of distributed quantum entanglement in multipartite systems is strongly due

to this unlocalizable character. As the mathematical interpretation for this polygamous nature of quantum entanglement, we have established polygamy inequalities of entanglement in tripartite quantum systems with arbitrary dimension, and multi-qubit systems. We have also provided several trade offs between UE and other correlations such as EoA, and localizable entanglement.

This is the first result where polygamous property of quantum entanglement in multipartite higher-dimensional quantum systems is provided. Furthermore, the proposed inequalities are in terms of the entropic entanglement measures such as entropy of entanglement for pure states and EoA. In other words, the proposed polygamy inequalities of distributed entanglement have been shown in terms of the actual quantification of entanglement with operational meanings, rather than using other entanglement measures such as concurrence.

Acknowledgments

JSK would like to thank Soojoon Lee for useful discussion, and acknowledges the support from iCORE, MITACS (QIP project) and US Army. GG acknowledges financial support from NSERC and MITACS-QIP.

-
- [1] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres and W. K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 - [2] C. Bennett and G. Brassard, in *Proceedings of IEEE International Conference on Computers, Systems, and Signal Processing* (IEEE Press, New York, Bangalore, India, 1984), p. 175-179.
 - [3] C. H. Bennett, Phys. Rev. Lett. **68**, 3121 (1992).
 - [4] V. Coffman, J. Kundu and W. K. Wootters, Phys. Rev. A **61**, 052306 (2000).
 - [5] T. Osborne and F. Verstraete, Phys. Rev. Lett. **96**, 220503 (2006).
 - [6] M. Koashi and A. Winter, Phys. Rev. A **69**(2) 022309 (2004).
 - [7] B. M. Terhal, IBM J. Research and Development 48, 71 (2004).
 - [8] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [9] D. P. DiVincenzo et al., Lect. Notes Comput. Sci. **1509**, 247 (1999).
 - [10] O. Cohen, Phys. Rev. Lett. **80**, 2493 (1998).
 - [11] T. Laustsen, F. Verstraete and S. J. van Enk, Quantum Inf. Comput. **3**, 64 (2003).
 - [12] G. Gour, S. Bandyopadhyay and B. C. Sanders, J. Math. Phys. **48**, 012108 (2007).
 - [13] I. Devetak, A. Winter, IEEE Transactions on Information Theory **50**(12) pp. 3183-3196 (2004).
 - [14] L. Henderson and V. Vedral, J. Phys. A: Math. Gen. **34**, 6899 (2001).
 - [15] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996).
 - [16] J. A. Smolin, F. Verstraete and A. Winter, Phys. Rev. A **72**, 052317 (2005).
 - [17] M. Christandl and A. Winter, IEEE Trans. Inf. Theory **51**, 3159-3165 (2005).
 - [18] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, U.K., 2000).
 - [19] C-s. Yu and H-s. Song, Phys. Rev. A **76**, 022324 (2007).
 - [20] D. P. Chi, J. W. Choi, K. Jeong, J. S. Kim, T. Kim and S. Lee, J. Math. Phys. **49**, 112102 (2008).
 - [21] L. P. Hughston, R. Jozsa and W. K. Wootters, Phys. Lett. A **183**, 14 (1993).
 - [22] P. Horodecki, Phys. Lett. A **232**, 333 (1997).
 - [23] G. Gour and R. W. Spekkens, Phys. Rev. A **73**, 062331 (2006).
 - [24] M. Popp, F. Verstraete, M. A. Martin-Delgado and J. I. Cirac, Phys. Rev. A **71**, 042306 (2005).