

# QUADRATIC VECTOR FIELDS WITH INVARIANT ALGEBRAIC CURVE OF LARGE DEGREE

RAFAEL RAMIREZ<sup>1</sup> AND NATALIA SADOVSKAIA<sup>2</sup>

ABSTRACT. We construct a polynomial planar vector field of degree two with one invariant algebraic curves of large degree. We exhibit an explicit quadratic vector fields which invariant curves of degree nine , twelve , fifteen and eighteen degree. .

## 1. INTRODUCTION

In the paper [2] the authors present for the first time examples of algebraic limit cycles of degree greater than 4 for planar quadratic vector fields. They also give an example of an invariant algebraic curve of degree 12 and genus one for which the quadratic system has no Darboux integrating factors or first integrals [?].

One of the first example of quadratic planar system with invariant algebraic curve of genus one is the Filipstov differential system [3]

$$\begin{cases} \dot{x} = 16(1+a)x - 6(2+a)x^2 + (2+12x)y \\ \dot{y} = 3a(1+a)x^2 + (15(1+a) - 2(9+5a)x)y + 16y^2 \end{cases}$$

which possesses the irreducible invariant algebraic curve of degree four and genus one

$$g(x, y) \equiv y^3 + \frac{1}{4}(3(1+a) - 6(1+a)x)y^2 + \frac{3}{4}(1+a)ax^2y + \frac{3}{4}(1+a)a^2x^4 = 0.$$

Another example we can find in the paper [2].

This diferential system

$$\dot{x} = 2(1+2x-2ax^2+6xy), \quad \dot{y} = 8-3a-14ax-2axy-8y^2$$

a has the invariant algebraic curve of degree four and genus one

$$x^2y^2 + xy + ax^3 - x^2 + x + 1/4 = 0$$

In the both case we have one algebraic limit cycles.

In [2] we can find a surprisingly simple quadratic system

$$\begin{cases} \dot{x} = xy + x^2 + 1 \\ \dot{y} = 3y^2 - \frac{81}{2}x^2 + \frac{57}{2} \end{cases}$$

---

1991 *Mathematics Subject Classification.* Primary 34C05, 34A34, 34C14.

*Key words and phrases.* limit cycles, algebraic limit cycles, polynomial vector fields.

which has invariant algebraic curve of degree 12 and genus one

$$\begin{cases} -442368 - 7246584x^2 + 71546517x^4 - 97906500x^6 + 41343750x^8 - 23437500x^{10} + \\ 48828125x^{12} + (322272x - 12126312x^3 + 23463000x^5 + 1125000x^7 + 15625000x^9)y - \\ (98784 - 711288x^2 + 5058000x^4 - 375000x^6)y^2 + (32928x - 1124000x^3)y^3 - 5488y^4 = 0 \end{cases}$$

The main result of this paper is to construct a set of quadratic planar vector field with an invariant curves of large degree and genus greater or equal to one. We apply the normal form of the quadratic fields deduced by J.Llibre [?].

## 2. QUADRATIC VECTOR FIELDS WITH A GIVEN INVARIANT ALGEBRAIC CURVES OF DEGREE NINE

In this section we determine three quadratic vector field with invariant curves of degree nine and genus one.

**Corollary 1.** *The following quadratic differential systems have the invariant curve  $g = 0$  of degree nine and genus one and cofactor  $K = 9y$ .*

(i)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{8}{13}qzy - \frac{24}{169}q^2z^2 + q$$

$$g = y^3 - \frac{9}{26}q^2z - \frac{2784}{28561}z^5q^4 + \frac{296}{2197}z^3q^3 + \frac{6144}{371293}z^7q^5 - \frac{4096}{4826809}z^9q^6 + \frac{9}{26}yq - \frac{60}{169}yq^2z^2 \\ + \frac{960}{2197}yz^4q^3 - \frac{1536}{28561}yz^6q^4 - \frac{12}{13}y^2qz - \frac{120}{169}y^2z^3q^2 = 0$$

(ii)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{14}{25}qzy - \frac{84}{125}q^2z^2 + q$$

$$g = y^3 + \frac{772}{3125}z^3q^3 - \frac{39648}{390625}z^5q^4 - \frac{9}{25}q^2z + \frac{9}{25}yq - \frac{312}{625}yq^2z^2 + \\ \frac{2688}{15625}yz^4q^3 - \frac{9408}{78125}yz^6q^4 - \frac{21}{25}y^2qz - \frac{192}{625}y^2z^3q^2 - \frac{87808}{9765625}z^9q^6 + \frac{65856}{1953125}z^7q^5 = 0$$

(iii)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{88}{53}qzy - \frac{3264}{2809}q^2z^2 + q$$

$$g = y^3 - \frac{9}{106}q^2z + \frac{7655424}{7890481}z^5q^4 - \frac{190316544}{418195493}z^7q^5 - \frac{105344}{148877}z^3q^3 - \frac{132}{53}y^2qz + \frac{3840}{2809}y^2z^3q^2 + \\ \frac{1568669696}{22164361129}z^9q^6 + \frac{9}{106}yq + \frac{6240}{2809}yq^2z^2 - \frac{337920}{148877}yz^4q^3 + \frac{4325376}{7890481}yz^6q^4 = 0$$

## 3. QUADRATIC VECTOR FIELDS WITH A GIVEN INVARIANT ALGEBRAIC CURVES OF DEGREE TWELVE

In this section we determine five quadratic vector field with invariant curves of degree twelve and genus one.

**Corollary 2.** *The following quadratic differential systems have the invariant curve  $g = 0$  of degree nine and genus one with cofactor  $K = 12y$ .*

(i)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 + pzy + (-pq - 2p^2)z^2 + q$$

$$\begin{aligned} g = & \frac{1}{36}p^6(q + 2p^2)z^{12} + \frac{1}{2}p(\frac{1}{3}p^5q + \frac{2}{3}p^6)z^{10} + (\frac{1}{3}p^5q + \frac{2}{3}q_{11}^6)y z^9 + \\ & (\frac{1}{2}(\frac{1}{3}qp^3 + \frac{5}{3}p^4)(-q_{11}q - 2p^2) + \frac{1}{4}p^2(\frac{1}{3}qp^3 + \frac{5}{3}p^4) + 3/4p^5q + 3/2p^6)z^8 + \\ & p(\frac{1}{3}qp^3 + \frac{5}{3}p^4)yz^7 + (\frac{17}{12}p(\frac{1}{3}qp^3 + \frac{5}{3}p^4) + (\frac{1}{3}qp^3 + \frac{5}{3}p^4)y^2 + \\ & \frac{1}{4}p^5 + \frac{1}{3}(\frac{1}{3}qp^3 + \frac{5}{3}p^4)q + \frac{5}{4}p^3(-pq - 2p^2))z^6 + \\ & (4p^4 + \frac{1}{2}qq^3 + \frac{3}{2}(-pq - 2p^2)p^2)yz^5 + \\ & (\frac{19}{16}qp^3 + \frac{23}{8}p^4 + \frac{89}{48}(-pq - 2p^2)p^2 + \frac{1}{4}(-pq - 2p^2)^2 + 3p^3y^2)z^4 \\ & + ((qp^2 + 3p^3 + \frac{4}{3}p(-pq - 2p^2))y + 2p^2y^3)z^3 + \\ & (\frac{119}{120}qp^2 + \frac{59}{120}p(-pq - 2p^2) + \frac{59}{60}p^3 + (-pq + p^2)y^2 + \frac{1}{3}(-q_{11}q_{20} - 2q_{11}^2)q_{20})z^2 \\ & + (2y^3p + (\frac{2}{3}pq + \frac{1}{3}p^2)y)z + y^4 + (\frac{2}{3}q + \frac{1}{3}p)y^2 + \frac{1}{9}q^2 + \frac{1}{36}p^2 + \frac{1}{9}pq = 0 \end{aligned}$$

(ii)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{35}{19}qzy - \frac{525}{361}q^2z^2 + q$$

$$\begin{aligned} g = & \frac{16748046875}{271737008656}z^{12}q^8 - \frac{3349609375}{7150973912}q^7z^{10} + \frac{95703125}{188183524}q^6yz^9 \\ & + \frac{1003515625}{752734096}q^6z^8 - \frac{7109375}{2476099}q^5yz^7 + (\frac{203125}{130321}q^4y^2 - \frac{16765625}{9904396}q^5)z^6 \\ & + \frac{1396875}{260642}q^4yz^5 + (\frac{1638125}{2085136}q^4 - \frac{39375}{6859}q^3y^2)z^4 + (\frac{750}{361}q^2y^3 - \frac{22625}{6859}q^3y)z^3 + \\ & (\frac{155}{2888}q^3 + \frac{1875}{361}q^2y^2)z^2 + (\frac{-70}{19}qq^3 - \frac{155}{1444}q^2y)z - \frac{1}{5776}q^2 + y^4 + \frac{1}{19}qy^2 = 0 \end{aligned}$$

(iii)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{10}{59}qzy - \frac{150}{3481}z^2q^2 + q$$

$$\begin{aligned}
g = & -\frac{781250000}{50362840098282103}q^8z^{12} + \frac{312500000}{853607459292917}q^7z^{10} - q_{20}^6 \frac{62500000}{14467923038863}yz^9 - \\
& q_{20}^6 \frac{159375000}{14467923038863}z^8 - q^5 \frac{3750000}{245219034557}yz^7 + (q^4 \frac{375000}{4156254823}y^2 + \frac{230375000}{245219034557}q^5)z^6 - \\
& q_{20}^4 \frac{1425000}{593750689}yz^5 + (q^3 \frac{-4215000}{70444997}y^2 - \frac{82923125}{4156254823}q^4)z^4 + \\
& (\frac{5291500}{70444997}yq^3 + \frac{281000}{1193983}y^3q^2)z^3 + (q^2 \frac{210750}{1193983}y^2 + \frac{64980}{1193983}q^3)z^2 + \\
& (q \frac{-20}{59}y^3 - q^2 \frac{129960}{1193983}y)z + \frac{110592}{1193983}q^2 + y^4 + q \frac{36}{59}y^2 = 0
\end{aligned}$$

(iv)

$$\begin{aligned}
\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{55}{311}qzy + \frac{363}{96721}q^2z^2 + q \\
g = \frac{78460709418025}{22403871730541898055936}q^8z^{12} + \frac{35663958826375}{36019086383507874688}q^7z^{10} - \frac{648435615025}{57908499008855104}yz^9q^6 - \\
\frac{10221703240485}{231633996035420416}q^6z^8 + \frac{16076916075}{46550240360816}yq^5z^7 + (-\frac{292307565}{149679229456}q^4y^2 - \frac{69082021195}{186200961443264}q^5)z^6 + \\
\frac{611188545}{299358458912}q_{20}^4yz^5 + (-\frac{29164418129}{2394867671296}q^4 - \frac{12078825}{240641848}q^3y^2)z^4 + \\
(\frac{73205}{386884}q^2y^3 + \frac{23645215}{481283696}q^3y)z^3 + (\frac{298023}{1547536}q^2y^2 + \frac{689337}{12380288}q^3)z^2 + \\
(-\frac{110}{311}qy^3 - \frac{689337}{6190144}yq^2)z + \frac{189}{311}y^2q + \frac{2278125}{24760576}q^2 + y^4 = 0
\end{aligned}$$

(iv)

$$\begin{aligned}
\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - \frac{187}{107}qzy - \frac{8349}{11449}q^2z^2 + q \\
g = \frac{1227125495297911}{274909788773107216}q^8z^{12} - \frac{82455072806579}{1284625181182744}q^7z^{10} + \\
\frac{440936218217}{6002921407396}yz^9q^6 + \frac{8241027180045}{24011685629584}z^8q^6 - \frac{10932302931}{14025517307}yq^5z^7 + \\
(-\frac{45176577061}{56102069228}q^5 + \frac{58461513}{131079601}y^2q^4)z^6 + \frac{706852839}{262159202}yq^4z^5 + \\
(\frac{1471610833}{2097273616}q^4 - \frac{3733455}{1225043}y^2q^3)z^4 + (\frac{13310}{11449}y^3q^2 - \frac{3587045}{1225043}yq^3)z^3 + \\
(\frac{54087}{11449}q_{20}^2y^2 + \frac{7623}{91592}q^3)z^2 + (-\frac{7623}{45796}yq^2 - \frac{374}{107}y^3q)z + \frac{27}{183184}q^2 + y^4 + \frac{9}{107}qy^2 = 0
\end{aligned}$$

(v)

$$\begin{aligned}
\dot{z} = zy + 1, \quad \dot{y} = 3y^2 + qzy - q^2z^2 - q \\
(\frac{1}{64}q^6p - \frac{1}{288}q^8)z^{12} + (\frac{3}{32}q^5p - \frac{1}{48}q^7)z^{10} + (-\frac{1}{24}yq^6 + \frac{3}{16}yq^4p)z^9 + \\
(-\frac{5}{96}q^6 + \frac{15}{64}q^4p)z^8 + (\frac{3}{4}yq^3p - \frac{1}{6}yq^5)z^7 + (\frac{3}{4}y^2q^2p - \frac{13}{72}q^5 + \frac{5}{16}q^3p - \frac{1}{6}y^2q^4)z^6 + \\
(-\frac{1}{4}q^4 + \frac{9}{8}q^2p)yz^5 + (-\frac{37}{96}q^4 + \frac{15}{64}q^2p + \frac{3}{2}y^2qp)z^4 + (-\frac{5}{6}yq^3 + y^3p + \frac{3}{4}yqp)z^3 + \\
(\frac{3}{32}qp + \frac{3}{4}y^2p - \frac{17}{48}q^3 + \frac{1}{2}y^2q^2)z^2 + (2y^3q + \frac{3}{16}yp - \frac{17}{24}yq^2)z + \frac{1}{64}p + y^4 - \frac{1}{3}y^2q - \frac{1}{288}q^2 = 0
\end{aligned}$$

**Proposition 3.** *The quadratic vector fields with invariant algebraic curve of genus one admits at most two algebraic limit cycles.*

*Proof.* The proof follow from the fact that if the algebraic curve has genus  $G$  then admits at most  $G + 1$  ovals.  $\square$

4. QUADRATIC VECTOR FIELDS WITH A GIVEN INVARIANT ALGEBRAIC CURVES OF DEGREE FIFTEEN

We construct three quadratic vector field with invariant curve of degree fifteen and genus two. In this section we give two of theses quadratic vector fields.

**Corollary 4.** *The following quadratic differential systems have the invariant curve  $g = 0$  of degree fifteen with cofactor  $K = 15y$  and genus two.*

(i)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - 6/17qzy - 8/289q^2z^2 + q$$

$$\begin{aligned} g = & y^5 - 135/1156q^3z + 204600/410338673z^9q^7 - 49546/1419857z^5q^5 - \\ & 31800/24137569z^7q^6 - 365760/6975757441q^8z^{11} - 36864/2015993900449z^{15}q^{10} + \\ & 11915/250563z^3q^4 - 300/289z^3y^4q^2 - 15/17zy^4q - 69120/6975757441yz^{12}q^8 + \\ & 14250/83521yz^4q^4 - 99000/24137569yz^8q^6 - 685/4913yq^3z^2 + 19800/1419857yz^6q^5 + \\ & 190080/410338673yq^7z^{10} - 15780/83521y^2z^5q^4 + 11880/1419857y^2q^5z^7 - 190/289y^2q^2z - \\ & 485/4913y^2q^3z^3 - 31680/24137569y^2z^9q^6 - 145/289y^3q^2z^2 + 3600/4913y^3z^4q^3 - \\ & 1320/83521y^3z^6q^4 + 207360/118587876497q^9z^{13} + 135/1156yq^2 + 35/51y^3q = 0 \end{aligned}$$

(ii)

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - 4/3pzy - 16/3p^2z^2 + p$$

$$\begin{aligned}
g = & y^5 + 2621440/19683yz^{12}p^8 + 2560000/19683yp^6z^8 - 313600/6561yp^5z^6 - \\
& 3768320/19683yp^7z^{10} + 1300/81y^4z^3p^2 - 10/3y^4zp + 265/27y^3p^2z^2 - \\
& 10400/243y^3z^4p^3 + 474880/6561y^3z^6p^4 + 143840/2187y^2p^4z^5 - 949760/6561y^2p^5z^7 \\
& + 35/54y^2p^2z - 9130/729y^2z^3p^3 + 942080/6561y^2z^9p^6 - 695/486yp^3z^2 + \quad cc \\
& 7000/729y^4z^4 + 5/18y^3p + 8388608/177147z^{15}p^{10} - 10/81p^3z + 1015/1458p^4z^3 - \\
& 2512/729p^5z^5 + 10/81yp^2 + 92800/6561p^6z^7 - 7577600/177147p^7z^9 + \\
& 4751360/59049z^{11}p^8 - 5242880/59049p^9z^{13} = 0
\end{aligned}$$

## 5. QUADRATIC VECTOR FIELDS WITH A GIVEN INVARIANT ALGEBRAIC CURVE OF DEGREE EIGHTEEN

We construct a quadratic vector field with invariant curve of degree eighteen and genus two.

**Corollary 5.** *The following quadratic differential systems have the invariant curve  $g = 0$  of degree eighteen with cofactor  $K = 18y$  and genus two.*

$$\dot{z} = zy + 1, \quad \dot{y} = 3y^2 - 6/17qzy - 8/289q^2z^2 + q$$

$$\begin{aligned}
 g = & \frac{16777216}{23298085122481} z^{18} q^{12} - \frac{50331648}{1792160394037} q^{11} z^{16} + \frac{12582912}{137858491849} y z^{15} q^{10} + \\
 & \frac{60555264}{137858491849} z^{14} q^{10} - \frac{26738688}{10604499373} y q^9 z^{13} + \\
 & \left( -\frac{2818048}{815730721} q^9 + \frac{3342336}{815730721} y^2 q^8 \right) z^{12} + \frac{20840448}{815730721} y z^{11} q^8 + \\
 & \left( \frac{11452416}{815730721} q^8 - \frac{1728}{4826809} q^6 p - \frac{4325376}{62748517} y^2 q^7 \right) z^{10} + \\
 & \left( -\frac{540672}{4826809} y q^7 + \frac{360448}{4826809} y^3 q^6 \right) z^9 + \\
 & \left( \frac{2592}{371293} q^5 p + \frac{1695744}{4826809} y^2 q^6 - \frac{1744896}{62748517} q^7 + \frac{288}{28561} y^2 q^4 p \right) z^8 + \\
 & \left( -\frac{196608}{371293} y^3 q^5 + \frac{964608}{4826809} y q^6 - \frac{720}{28561} y q^4 p - \frac{64}{2197} y^3 q^3 p \right) z^7 + \\
 & \left( \frac{4}{169} y^4 p q^2 - \frac{196608}{371293} y^2 q^5 + \frac{132544}{4826809} q^6 - \frac{90}{2197} q^4 p - \frac{264}{2197} y^2 q^3 p + \frac{12288}{28561} y^4 q^4 \right) z^6 + \\
 & \left( \frac{474}{2197} y q^3 p + \frac{18432}{28561} y^3 q^4 - \frac{59136}{371293} y q^5 + \frac{136}{169} y^3 p q^2 \right) z^5 + \\
 & \left( -\frac{5712}{371293} q^5 + \frac{241}{4394} p q^3 + \frac{10896}{28561} y^2 q^4 - \frac{69}{338} y^2 p q^2 - \frac{20}{13} y^4 q p \right) z^4 + \\
 & \left( -\frac{1}{2} y^3 p q - \frac{608}{2197} y^3 q^3 + \frac{1104}{28561} y q^4 + y^5 p - \frac{11}{52} y p q^2 \right) z^3 + \\
 & \left( \frac{177}{2197} q^4 + \frac{3}{26} y^2 p q + \frac{204}{169} y^4 q^2 + \frac{1224}{2197} y^2 q^3 + \frac{3}{4} y^4 p - \frac{23}{832} p q^2 \right) z^2 + \\
 & \left( \frac{3}{16} y^3 p + \frac{23}{416} y p q - \frac{354}{2197} y q^3 - \frac{180}{169} y^3 q^2 - \frac{24}{13} y^5 q \right) z + y^6 + \\
 & \frac{15}{2197} q^3 + \frac{9}{13} y^4 q + \frac{1}{208} p q + \frac{1}{64} y^2 p + \frac{24}{169} y^2 q^2 = 0
 \end{aligned}$$

cc

## REFERENCES

- [1] J. LLIBRE, *Integrability of polynomial differential systems*, Handbook of Differential Equations, Ordinary Differential Equations, Eds. A. Cañada, P. Drabek and A. Fonda, Elsevier (2004), pp. 437–533.
- [2] C. CHRISTOPHER, J. LLIBRE AND G. SWIRSZCZ *Invariant algebraic curves of large degree for quadratic system*, J. of Mathematical Analysis and Applications **303** (2005), 450-461.
- [3] R. RAMIREZ AND N. SADOVSKAIA *Inverse approach in the study of ordinary differential equations*, preprint, 2008.

<sup>1</sup> DEPARTAMENT D'ENGINYERIA INFORMÀTICA I MATEMÀTIQUES, UNIVERSITAT ROVIRA I VIRGILI, AVINGUDA DELS PAÏSOS CATALANS 26, 43007 TARRAGONA, CATALONIA, SPAIN.

*E-mail address:* rafaelorlando.ramirez@urv.cat

<sup>2</sup> DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA, C. PAU GARGALLO 5, 08028 BARCELONA, CATALONIA, SPAIN.

*E-mail address:* natalia.sadovskaia@upc.edu