

MAXIMAL INEQUALITIES AND RIESZ TRANSFORM ESTIMATES ON L^p SPACES FOR MAGNETIC SCHRÖDINGER OPERATORS II

BESMA BEN ALI

ABSTRACT. The paper concerns the magnetic Schrödinger operator $H(\mathbf{a}, V) = \sum_{j=1}^n (\frac{1}{i} \frac{\partial}{\partial x_j} - a_j)^2 + V$ on \mathbb{R}^n . We prove some L^p estimates on the Riesz transforms of H and we establish some related maximal inequalities. The conditions that we arrive at, are essentially based on the control of the magnetic field by the electric potential.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Reverse estimates	8
4. Estimates for weak solution	15
5. Maximal inequalities	21
6. Proof of Theorem 1.2	23
7. Second order Riesz transforms	24
References	27

1. INTRODUCTION

Consider the Schrödinger operator with magnetic field

$$(1.1) \quad H(\mathbf{a}, V) = \sum_{j=1}^n \left(\frac{1}{i} \frac{\partial}{\partial x_j} - a_j \right)^2 + V \text{ in } \mathbb{R}^n, \quad n \geq 2,$$

where $\mathbf{a} = (a_1, a_2, \dots, a_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the magnetic potential and $V : \mathbb{R}^n \rightarrow \mathbb{R}$ is the electric potential. Let

$$(1.2) \quad B(x) = \text{curl } \mathbf{a}(x) = (b_{jk}(x))_{1 \leq j, k \leq n}$$

be the magnetic field generated by \mathbf{a} , where

$$(1.3) \quad b_{jk} = \frac{\partial a_j}{\partial x_k} - \frac{\partial a_k}{\partial x_j}.$$

2000 *Mathematics Subject Classification.* 35J10; 42B20; 58G25; 81Q10; 81V10.

Key words and phrases. Schrödinger operators, maximal inequalities, Riesz transforms, Fefferman-Phong inequality, reverse Hölder estimates, magnetic field.

We will assume that $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in L^1_{loc}(\mathbb{R}^n)$, $V \geq 0$. Let

$$(1.4) \quad L_j = \frac{1}{i} \frac{\partial}{\partial x_j} - a_j \quad \text{for} \quad 1 \leq j \leq n,$$

Set $L = (L_1, \dots, L_n)$ and $|Lu(x)| = (\sum_{j=1}^n |L_j u(x)|^2)^{1/2}$.

Note that $L_j^* = L_j$ for all $1 \leq j \leq n$. Let

$$L^* = (L_1^*, \dots, L_n^*)^T.$$

We define the form \mathcal{Q} by

$$(1.5) \quad \mathcal{Q}(u, v) = \sum_{k=1}^n \int_{\mathbb{R}^n} L_k u \cdot \overline{L_k v} dx + \int_{\mathbb{R}^n} V u \cdot \overline{v} dx,$$

with domain $\mathcal{D}(\mathcal{Q}) = \mathcal{V} \times \mathcal{V}$ where

$$\mathcal{V} = \{u \in L^2, L_k u \in L^2 \text{ for } k = 1, \dots, n \text{ and } \sqrt{V}u \in L^2\}.$$

Let $\dot{\mathcal{V}}$ be the closure of $C_0^\infty \mathbb{R}^n$ under the semi-norm

$$\|f\|_{\dot{\mathcal{V}}} = (\|Lf\|_2^2 + \|V^{1/2}f\|_2^2)^{1/2}.$$

We denote $H(\mathbf{a}, V) = H$, the self-adjoint operator on $L^2(\mathbb{R}^n)$ associated to this symmetric and closed form.

The domain of H is given by:

$$\mathcal{D}(H) = \{u \in \mathcal{D}(\mathcal{Q}), \exists v \in L^2 \text{ so that } \mathcal{Q}(u, \phi) = \int_{\mathbb{R}^n} v \overline{\phi} dx, \forall \phi \in \mathcal{D}(\mathcal{Q})\}.$$

The operators $L_j H(\mathbf{a}, V)^{-1/2}$ are called the Riesz transforms associated with $H(\mathbf{a}, V)$. We know that

$$(1.6) \quad \sum_{j=1}^n \|L_j u\|_2^2 + \|V^{1/2}u\|_2^2 = \|H(\mathbf{a}, V)^{1/2}u\|_2^2, \quad \forall u \in \mathcal{D}(\mathcal{Q}) = \mathcal{D}(H(\mathbf{a}, V)^{1/2}).$$

Hence, the operators $L_j H(\mathbf{a}, V)^{-1/2}$ are bounded on $L^2(\mathbb{R}^n)$, for all $j = 1, \dots, n$.

The aim of this paper is to establish the L^p boundedness of the operators $L_j H(\mathbf{a}, V)^{-1/2}$ and $V^{1/2} H(\mathbf{a}, V)^{-1/2}$.

In the case where the magnetic potential is absent, that is, $H(\mathbf{a}, V) = -\Delta + V$ and $LH(\mathbf{a}, V)^{-1/2} = \nabla(-\Delta + V)^{-1/2}$, many important studies have been established. We mention the works of Helffer-Nourrigat [HNW], Guibourg [Gui2] and Zhong [Z], in which they considered the case of polynomial potentials. A generalization of their results was given by Shen [Sh1], he proved the L^p boundedness of Riesz transforms of Schrödinger operators with electric potential contained in certain reverse Hölder classes. Auscher and I improved this result in [AB], using a different approach based on local estimates. Note that this approach can be extended to more general spaces for instance some Riemannian manifolds and Lie groups (see [BB]).

In the presence of the magnetic field, we know that these operators are of weak type (1.1) and hence, by interpolation, are L^p bounded for all $1 < p \leq 2$. This result was proved by Sikora using the finite speed propagation property [Sik]. Independantly, Duong, Ouhabaz and Yan [DOY] have proved the same result using another method.

The main purpose of this work is to find sufficient conditions on the electric potential and the magnetic field, for which the Riesz transforms of $H(\mathbf{a}, V)$ are L^p bounded for the range $p > 2$. Note that, because of the gauge invariance of the operator $H(\mathbf{a}, V)$ and the nature of the L^p estimates, any such quantitative condition should be imposed on the magnetic field B , not directly on \mathbf{a} .

In a previous paper [Be], many important results about this problem were established. Under certain conditions used by Shen in [Sh4] and given in terms of the reverse Hölder inequality on the magnetic field and the electric potential, we proved that the Riesz transforms of the pure magnetic Schrödinger operator $H(\mathbf{a}, 0)$ are L^p bounded for all $p \geq 2$. We have also extended the results of [AB] about $-\Delta + V$ to the magnetic operator $H(\mathbf{a}, V)$.

The second aim of this article is to establish important maximal inequalities related to the L^p behaviour of $L_j L_k H(\mathbf{a}, V)^{-1}$, $V^{1/2} L H(\mathbf{a}, V)^{-1}$ and other operators called the second order Riesz transforms. Estimates on these operators are of great interest in the study of spectral theory of $H(\mathbf{a}, V)$. There are rather few works around the behaviour of these operators. We cite Guibourg who considered the polynomial case and established an L^2 estimate [Gui1]. Shen [Sh4], generalised [Gui1] and proved under reverse Hölder conditions, the L^p boundedness of $L_j L_k H(\mathbf{a}, V)^{-1}$. Independantly and under the same conditions, we have proved and generalised the results of Shen in [Be]. Note that in [Sh4] and [Be], the contribution of the magnetic field was controlled by introducing an auxiliary function $m(\cdot, \omega)$ defined by Shen [Sh1] for RH_∞ class, he generalizes an early version of a useful auxiliary function for polynomial potentials. In this paper we will use another approach, the contribution of the magnetic field will be controlled by the electric potential and the magnetic Schrödinger operator will be treated as a perturbation of the Laplace operator $-\Delta$.

Here, our assumptions on potentials will be given in terms of reverse Hölder inequality. Let us recall the definition of these weight classes:

Definition 1.1. *Let $\omega \in L^q_{loc}(\mathbb{R}^n)$, $\omega > 0$ almost everywhere, $\omega \in RH_q$, $1 < q \leq \infty$, the class of the reverse Hölder weights with exponent q , if there exists a constant C such that for any cube Q of \mathbb{R}^n ,*

$$(1.7) \quad \left(\int_Q \omega^q(x) dx \right)^{1/q} \leq C \left(\int_Q \omega(x) dx \right).$$

If $q = \infty$, then the left hand side is the essential supremum on Q . The smallest C is called the RH_q constant of ω .

A note about notations: Throughout this paper we will use the following notation $\int_Q \omega = \frac{1}{|Q|} \int_Q \omega$. C and c denote constants. As usual, λQ is the cube co-centered with Q with sidelength λ times that of Q .

We now state our main result:

Theorem 1.2. *Suppose $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in RH_q$, $1 < q \leq +\infty$. Also assume that there exists a constant $C > 0$ such that for any cube Q in \mathbb{R}^n :*

$$(1.8) \quad \begin{cases} \sup_Q |B| \leq C \int_Q V, \\ \sup_Q |\nabla B| \leq C (\int_Q V)^{3/2}, \end{cases}$$

where $|B| = \sum_{j,k} |b_{jk}|$ and $\nabla = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})$. Then, there exists an $\epsilon > 0$ such that for any $1 \leq p < q^* + \epsilon$, there exists a constant $C_p > 0$, depending on V such that

$$(1.9) \quad \|LH(\mathbf{a}, V)^{-\frac{1}{2}}(f)\|_p \leq C_p \|f\|_p,$$

for any $f \in C_0^\infty(\mathbb{R}^n)$ if $p > 1$,
and

$$|\{x \in \mathbb{R}^n; |Lf(x)| > \alpha\}| \leq \frac{C_1}{\alpha} \|H(\mathbf{a}, V)^{\frac{1}{2}} f\|_1,$$

for any $\alpha > 0$ and $f \in C_0^\infty(\mathbb{R}^n)$ if $p = 1$.

Remark 1.3. (1) Note that condition (1.8) implies the following inequalities:

$$(1.10) \quad \begin{cases} |B| \leq CV, \\ |\nabla B| \leq CV^{3/2}, \end{cases}$$

almost everywhere in \mathbb{R}^n . These hypotheses are not sufficient to obtain (1.9).

(2) Condition (1.8) includes the polynomial case.

An important step to prove the previous result is to establish the following reverse estimates that hold an importance of their own :

Theorem 1.4. Suppose $\mathbf{a} \in L_{loc}^2(\mathbb{R}^n)^n$, $V \in A_\infty$. Also assume that there exists a constant $C > 0$ such that for any cube Q in \mathbb{R}^n :

$$(1.11) \quad \sup_Q |B| \leq C \int_Q V.$$

Then, for any $1 \leq p < \infty$, there exists $C_p > 0$, which depends on (1.11) and V , such that

$$(1.12) \quad \|H(\mathbf{a}, V)^{\frac{1}{2}}(f)\|_p \leq C_p (\|Lf\|_p + \|V^{\frac{1}{2}} f\|_p),$$

for any $f \in C_0^\infty(\mathbb{R}^n)$, if $p > 1$,
and

$$(1.13) \quad |\{x \in \mathbb{R}^n; |H(\mathbf{a}, V)^{\frac{1}{2}} f(x)| > \alpha\}| \leq \frac{C_1}{\alpha} \int |Lf| + V^{\frac{1}{2}} |f|,$$

for any $\alpha > 0$ and $f \in C_0^\infty(\mathbb{R}^n)$, if $p = 1$.

Along with the study of Riesz transforms, we will also establish some maximal inequalities:

Theorem 1.5. Suppose $\mathbf{a} \in L_{loc}^2(\mathbb{R}^n)^n$, $V \in L_{loc}^1(\mathbb{R}^n)$ and $V \in RH_q$, $1 < q \leq +\infty$. Also assume that there exists a constant $C > 0$ such that for any cube $Q \subset \mathbb{R}^n$

$$(1.14) \quad \begin{cases} \sup_Q |B| \leq C \int_Q V \\ \sup_Q |\nabla B| \leq C (\int_Q V)^{3/2}. \end{cases}$$

Then, there exists an $\epsilon > 0$ depending on V , such that for every $s = 1, \dots, n$ and $k = 1, \dots, n$, and for any $1p < q + \epsilon$, there exists a constant $C_p > 0$ such that for any $f \in C_0^\infty(\mathbb{R}^n)$,

$$(1.15) \quad \|L_s L_k(f)\|_q \leq C_q \|H(\mathbf{a}, V)f\|_q.$$

The proof of this theorem will use Theorem 1.2 and 1.4. We also need to study the L^p boundedness of different second order Riesz Transforms us $VH(\mathbf{a}, V)^{-1}$, $H(\mathbf{a}, 0)H(\mathbf{a}, V)^{-1}$ (studied in section 5) and especially the behaviour of operator $V^{\frac{1}{2}}LH(\mathbf{a}, V)^{-1}$ described in the following theorem:

Theorem 1.6. *Suppose $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in RH_{n/2}$. Also assume that there exists a constant $C > 0$ such that for any cube Q in \mathbb{R}^n :*

$$(1.16) \quad \begin{cases} \sup_Q |B| \leq C \int_Q V, \\ \sup_Q |\nabla B| \leq C(\int_Q V)^{3/2}. \end{cases}$$

. Then the operator $V^{\frac{1}{2}}LH(\mathbf{a}, V)^{-1}$ is L^p bounded for

$$\begin{cases} 1 \leq p < \frac{2(q+\epsilon)n}{3n-2(q+\epsilon)}, & \text{if } q < n \\ 1 \leq p < 2(q + \epsilon), & \text{if } q \geq n \end{cases}$$

where ϵ depends only on V .

Note that this result was proved by Shen when the magnetic potential is absent (see Theorem 4.13, [Sh1] that can be recovered by [AB] methods under the same hypotheses and for $n \geq 1$ instead of $n \geq 3$).

We mention without proof that our results admit local versions, replacing $V \in RH_q$ by $V \in RH_{q,loc}$ which is defined by the same conditions on cubes with sides less than 1. Then we get the corresponding results and estimates for $H + 1$ instead of H . The results on operator domains are valid under local assumptions.

The arguments are based on local estimates. We briefly sketch the main tools :

- 1) An improved Fefferman-Phong inequality for A_∞ potentials.
- 2) Criteria for proving L^p boundedness of operators in absence of kernels.
- 3) Mean value inequalities for nonnegative subharmonic functions against A_∞ weights.
- 4) Complex interpolation, together with L^p boundedness of imaginary powers of $H(\mathbf{a}, V)$ for $1 < p < \infty$.
- 5) A Calderón-Zygmund decomposition adapted to level sets of the maximal function of $|Lf| + |V^{1/2}f|$.
- 6) A gauge transform adapted to the reverse Hölder conditions on the potentials.
- 7) Reverse Hölder inequalities involving $\mathbb{L}u$, $|B|^{1/2}u$ and $V^{1/2}u$ for weak solutions of $H(\mathbf{a}, V)u = 0$.

The paper is organized as follows. In Section 2 we give the principal tools to prove the theorems mentioned above. We state an improved Fefferman-Phong inequality and we establish an adapted gauge transform. Section 3 is devoted to establish some reverse estimates via a Calderón-Zygmund decomposition. In section 4 we give different estimates for the weak solution of $H(\mathbf{a}, V)u = 0$. We state some useful maximal inequalities in section 5. Section 6 is concerned with the proof of Theorem 1.2. Finally, in section 7, we study the operator $V^{\frac{1}{2}}LH^{-1}$ and give the proof of Theorem 1.5.

2. PRELIMINARIES

We begin by recalling some properties of the reverse Hölder classes.

Proposition 2.1. (Proposition 11.1 [AB]) *Let ω be a nonnegative measurable function. Then the following are equivalent:*

- (1) $\omega \in A_\infty$.
- (2) For all $s \in (0, 1)$, $\omega^s \in B_{1/s}$.
- (3) There exists $s \in (0, 1)$, $\omega^s \in B_{1/s}$.

Remark 2.2. *It is well known that if $\omega \in RH_q$ and $q < +\infty$, then $\omega \in RH_p$ for all $1 < p < q$ and there exists an $\varepsilon > 0$ such that $\omega \in RH_{q+\varepsilon}$.*

We also know that $\omega \in A_\infty$ if and only if there exists $q > 1$ such that $\omega \in RH_q$.

Here A_∞ is the Muckenhoupt weight class, defined as the union of all A_p , $1 \leq p < \infty$.

If $\omega \in A_\infty$ then $\omega(x)dx$ is a doubling measure (see [St],chap V for more details).

The first step of this work is to use the properties of the A_∞ weights to establish some reverse Hölder inequalities with the weak solutions. Then, we apply the following criterion for L^p boundedness [AM1] (A slightly weaker version appears in Shen [Sh2]).

Theorem 2.3. *Let $1 \leq p_0 < q_0 \leq \infty$. Suppose that T is a bounded sublinear operator on $L^{p_0}(\mathbb{R}^n)$. Assume that there exist constants $\alpha_2 > \alpha_1 > 1$, $C > 0$ such that*

$$(2.1) \quad \left(\int_Q |Tf|^{q_0} \right)^{\frac{1}{q_0}} \leq C \left\{ \left(\int_{\alpha_1 Q} |Tf|^{p_0} \right)^{\frac{1}{p_0}} + (S|f|)(x) \right\},$$

for all cube Q , $x \in Q$ and all $f \in L_{comp}^\infty(\mathbb{R}^n)$ with support in $\mathbb{R}^n \setminus \alpha_2 Q$, where S is a positive operator. Let $p_0 < p < q_0$. If S is bounded on $L^p(\mathbb{R}^n)$, then, there is a constant C such that

$$\|Tf\|_p \leq C \|f\|_p$$

for all $f \in L_{comp}^\infty(\mathbb{R}^n)$.

Fix an open set Ω and $f \in L_{comp}^\infty(\mathbb{R}^n)$, the space of compactly supported bounded functions on \mathbb{R}^n . By a weak solution of

$$(2.2) \quad H(\mathbf{a}, V)u = f \text{ in } \Omega,$$

we mean $u \in W(\Omega)$, with

$$W(\Omega) = \{u \in L_{loc}^1(\Omega); V^{1/2}u \text{ and } L_k u \in L_{loc}^2(\Omega) \forall k = 1, \dots, n\}$$

and the equation (2.2) holds in the sense of distribution on Ω . We note that if $u \in W(\Omega)$, then by Poincaré and the diamagnetic inequalities, $u \in L_{loc}^2(\Omega)$.

The weak solution satisfies some important inequalities which will be useful to prove our results:

Lemma 2.4. Caccioppoli type inequality

Let u a weak solution of $H(\mathbf{a}, V)u = f$ in $2Q$, where Q is a cube of \mathbb{R}^n and $f \in L_{comp}^\infty(\mathbb{R}^n)$. Then

$$(2.3) \quad \int_Q |Lu|^2 + V|u|^2 \leq C \left\{ \int_{2Q} |f||u| + \frac{1}{R^2} \int_{2Q} |u|^2 \right\}.$$

We also give some important tools:

Proposition 2.5. Diamagnetic inequality[LL]

For all $u \in W_a^{1,2}(\mathbb{R}^n)$, with

$$W_a^{1,2}(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n), L_k u \in L^2(\mathbb{R}^n), k = 1 \dots, n\},$$

we have

$$(2.4) \quad |\nabla(|u|)| \leq |L(u)|.$$

Proposition 2.6. Kato-Simon inequality:

$$(2.5) \quad |(H(\mathbf{a}, V) + \lambda)^{-1} f| \leq (-\Delta + \lambda)^{-1} |f|; \quad \forall f \in L^2(\mathbb{R}^n), \forall \lambda > 0.$$

We also have the following domination inequality [Si4]:

$$(2.6) \quad |e^{-tH(\mathbf{a}, V)} f| \leq e^{\Delta t} |f|; \quad \forall t \geq 0, f \in L^2(\mathbb{R}^n).$$

Fefferman-Phong inequalities

The usual Fefferman-Phong inequalities are of the form:

$$(2.7) \quad \int_Q |u|^p \min\left\{\int_Q \omega, \frac{1}{R^p}\right\} \leq C \left\{\int_Q |Lu|^p + \omega |u|^p\right\}.$$

In [AB] we established an improved version for these inequalities in absence of the magnetic potential. We can extend this improvement to the magnetic Schrödinger operators:

Lemma 2.7. An improved Fefferman-Phong inequality :

Let $\omega \in A_\infty$ and $1 \leq p < \infty$. Then there are constants $C > 0$ and $\beta \in (0, 1)$ depending only on p, n and the A_∞ constant of ω such that for all cubes Q (with sidelength R) and $u \in C^1(\mathbb{R}^n)$, one has

$$(2.8) \quad \int_Q |Lu|^p + \omega |u|^p \geq \frac{C m_\beta(R^p \int_Q \omega)}{R^p} \int_Q |u|^p$$

where $m_\beta(x) = x$ for $x \leq 1$ and $m_\beta(x) = x^\beta$ for $x \geq 1$.

The proof is the same as that of Lemma 2.1 in [AB], combined with the diamagnetic inequality.

Iwatsuka Gauge transform

Lemma 2.8. Let $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and Q a cube of \mathbb{R}^n . Suppose $B \in C^1(\mathbb{R}^n, M_n(\mathbb{R}))$. Then there exist $\mathbf{h} \in C^1(Q, \mathbb{R}^n)$ and a real function $\phi \in C^2(Q)$, such that $\text{curl} \mathbf{h} = B$ in Q and

$$(2.9) \quad \mathbf{h} = \mathbf{a} - \nabla \phi, \quad \text{in } Q,$$

with

$$(2.10) \quad \sup_Q |\mathbf{h}| \leq C R \sup_Q |B|,$$

and

$$(2.11) \quad \sup_Q |\nabla \mathbf{h}| \leq C \left(\sup_Q |B| + R \sup_Q |\nabla B| \right).$$

See the proof of Lemma 2.4 in [Sh5] which uses the construction of Iwatsuka [I].

Lemma 2.9. Let Q a cube in \mathbb{R}^n . Suppose $B \in L^\infty(Q, M_n(\mathbb{R}))$. Then there exist $\mathbf{h} \in L^\infty(Q, \mathbb{R}^n)$ and a real function $\phi \in W^{1, \infty}(Q)$, such $\text{curl} \mathbf{h} = B$ and

$$(2.12) \quad \mathbf{h} = \mathbf{a} - \nabla \phi \quad \text{a.e in } Q,$$

and

$$(2.13) \quad \sup_Q |\mathbf{h}| \leq C R \sup_Q |B|,$$

and

$$(2.14) \quad \sup_Q |\nabla \mathbf{h}| \leq C \left(\sup_Q |B| + R \sup_Q |\nabla B| \right).$$

Proof. Let $(\mathbf{a}_m)_{m \geq 0}$ be the sequence of C^1 functions obtained by convolution with \mathbf{a} and converge in L^2_{loc} to \mathbf{a} . Set $(B_m)_{m \geq 0}$, $(\phi_m)_{m \geq 0}$ and $(\mathbf{h}_m)_{m \geq 0}$ as the corresponding sequences of the Lemma 2.8. Note that $(\mathbf{h}_m)_{m \geq 0}$ converges in $L^n(Q, \mathbb{R}^n)$. Let \mathbf{h} be this limit, it satisfies (2.12). Note also that $(B_m)_{m \geq 0}$ converges to B in $L^{n/2}_{loc}(Q, M_n(\mathbb{R}))$ and $\text{curl} \mathbf{h} = B$ holds almost everywhere in Q , where curl is defined in the sens of distribution.

We know that for all $m \geq 0$,

$$\left(\int_Q |\mathbf{h}_m|^n \right)^{1/n} \leq c R \left(\int_Q |B_m|^{n/2} \right)^{2/n},$$

uniformly in m . Then applying the limit, we obtain

$$\left(\int_Q |\mathbf{h}|^n \right)^{1/n} \leq c R \left(\int_Q |B|^{n/2} \right)^{2/n}.$$

Hence inequality (2.13) follows easily. By a similar argument, inequality (2.14) holds. \square

3. REVERSE ESTIMATES

The present section talks about certain tools that are handy in the proof of Theorem 1.4. Note that this theorem can be obtained as a consequence of Theorem 1.6 in [Be] if we also assume that $|B|$ is in $RH_{n/2}$. However, condition (1.11) is sufficient to obtain estimate (1.12).

By duality, the L^p boundedness of Riesz transforms for $1 < p \leq 2$ (proved by [Sik] and [DOY]) implies the estimate (1.12) for any $p \geq 2$. For $p < 2$, we follow step by step the proof of the Theorem 1.2 of [AB] once the appropriate Calderón-Zygmund decomposition 3.1 is established. We also use the fact that the time derivatives of the kernel of semigroup e^{-tH} satisfy Gaussian estimates (see [CD], [Da], [G] and [Ou] or, theorem 6.17).

Let us introduce the main technical lemma of this work, which in itself is an interesting result:

Lemma 3.1. *Let $1 \leq p < 2$ and $\alpha > 0$. Under the assumptions of Theorem 1.4, we have: for any function $f \in C^\infty_0(\mathbb{R}^n)$ such that*

$$\|Lf\|_p + \|V^{\frac{1}{2}}f\|_p < \infty.$$

Then, one can find a collection of cubes (Q_k) and functions g and b_k such that

$$(3.1) \quad f = g + \sum_k b_k$$

and the following properties hold:

$$(3.2) \quad \|Lg\|_2 + \|V^{\frac{1}{2}}g\|_2 \leq C \alpha^{1-\frac{p}{2}} (\|Lf\|_p + \|V^{\frac{1}{2}}f\|_p)^{p/2}$$

$$(3.3) \quad \int_{Q_k} |Lb_k|^p + R_k^{-p} |b_k|^p \leq C\alpha^p |Q_k|$$

$$(3.4) \quad \sum_k |Q_k| \leq C\alpha^{-p} \left(\int_{\mathbb{R}^n} |Lf|^p + |V^{\frac{1}{2}}f|^p \right)$$

$$(3.5) \quad \sum_k \mathbf{1}_{Q_k} \leq N,$$

where N depends only on the dimension and C on the dimension, p and the $RH_{n/2}$ constant of $|B|$. Here, R_k denotes the sidelength of Q_k and gradients are taken in the sense of distributions in \mathbb{R}^n .

Remark 3.2. We establish an improved version of estimate (3.2):

$$(3.6) \quad \|Lg\|_\infty \leq C\alpha.$$

Proof. Let Ω be the open set $\{x \in \mathbb{R}^n; M(|Lf|^p + |V^{\frac{1}{2}}f|^p)(x) > \alpha^p\}$, where M is the uncentered maximal operator over the cubes of \mathbb{R}^n . If Ω is empty, then set $g = f$ and $b_i = 0$. Otherwise, our argument is subdivided into six steps.

a) Construction of the cubes:

The maximal theorem gives us

$$|\Omega| \leq C\alpha^{-p} \int_{\mathbb{R}^n} |Lf|^p + |V^{\frac{1}{2}}f|^p < \infty.$$

Set $F = \mathbb{R}^n \setminus \Omega$. Using the Lebesgue derivation Theorem, we have

$$(3.7) \quad |Lf|^p + |V^{\frac{1}{2}}f|^p \leq \alpha^p, \quad \text{a.e in } F.$$

Let (Q_k) be a Whitney decomposition of Ω by dyadic cubes so to say Ω is the disjoint union of the Q_k 's, the cubes $2Q_i$ are contained in Ω and have the bounded overlap property, but the cubes $4Q_k$ intersect F .¹ Hence

$$\sum_k |2Q_k| \leq C|\Omega| \leq C\alpha^{-p} \int_{\mathbb{R}^n} |Lf|^p + |V^{\frac{1}{2}}f|^p.$$

Thus, (3.4) and (3.5) are satisfied by $2Q_k$.

b) Construction of b_k :

Let (χ_k) be a partition of unity on Ω associated to the covering (Q_k) so that for each k , χ_k is a C^1 function supported in $2Q_k$ with

$$(3.8) \quad \|\chi_k\|_\infty + R_k \|\nabla \chi_k\|_\infty \leq c(n),$$

where R_k is the sidelength of Q_k and $\sum \chi_k = 1$ on Ω . We say that a cube Q is of type 1 if $R^2 \int_Q V > 1$, and is of type 2 if $R^2 \int_Q V > 1$.

¹In fact, the factor 2 should be some $c = c(n) > 1$ explicitly given in [[St],Chapter 6]. We use this convention to avoid too many irrelevant constants.

We apply the gauge transformation on the cubes $2Q_k$ such that Q_k is of type 2, hence there exist $\mathbf{h}_k \in L^n(2Q_k, \mathbb{R}^n)$ and a real function $\phi_k \in H^1(2Q_k)$ such that

$$(3.9) \quad \mathbf{h}_k = \mathbf{a} - \nabla \phi \quad \text{in } 2Q_k,$$

and

$$(3.10) \quad \sup_{2Q_k} |\mathbf{h}| \leq C R_k \sup_{2Q_k} |B| \leq C R_k \int_{2Q_k} V.$$

We denote

$$m_{2Q_k}(e^{i\phi_k} f) = \int_{2Q_k} (e^{i\phi_k} f).$$

Let

$$(3.11) \quad b_k = \begin{cases} f\chi_k, & \text{if } 2Q_k \text{ is of type 1,} \\ (f - e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f))\chi_k, & \text{if } 2Q_k \text{ is of type 2.} \end{cases}$$

c) Proof of the estimate (3.3):

Suppose $2Q_k$ is of type 1, $(2R_k)^2 \int_{2Q_k} V > 1$. Then

$$(2R_k)^{-p} \leq \left(\int_{2Q_k} V \right)^{p/2} \leq C \int_{2Q_k} V^{p/2},$$

here we used $V^{p/2} \in RH_{2/p}$ since $p < 2$.

Now we will control Lb_k :

$$Lb_k = L(f\chi_k) = (Lf)\chi_k + \frac{1}{i} f \nabla \chi_k,$$

then

$$\begin{aligned} \int_{2Q_k} |Lb_k|^p + R_k^{-p} |b_k|^p &\leq C \|\chi_k\|_\infty^p \int_{2Q_k} |Lf|^p + \|\nabla \chi_k\|_\infty^p \int_{2Q_k} |f|^p + R_k^{-p} \|\chi_k\|_\infty^p \int_{2Q_k} |f|^p \\ &\leq C \left\{ \int_{2Q_k} |Lf|^p + R_k^{-p} \int_{2Q_k} |f|^p \right\} \leq C \left\{ \int_{\tilde{Q}_k} |Lf|^p + |V^{1/2} f|^p \right\} \leq C \alpha^p |Q_k|, \end{aligned}$$

here we used the L^p version of the Fefferman-Phong inequality (2.7) and the intersection of $4Q_k$ with F . Hence estimation (3.3) holds for the cubes of type 1.

If Q_k is of type 2, $R_k^2 \int_{Q_k} V \leq 1$. $V(x)dx$ is a doubling measure, then there exists $C > 0$, such that $R_k^2 \int_{Q_k} |B| \leq C$.

$$b_k = (f - e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f))\chi_k.$$

Let us estimate Lb_k . By the Gauge invariance, all we require is the estimation of $\tilde{L}(e^{i\phi_k} b_k)$, where

$$\tilde{L} = \frac{1}{i} \nabla - \mathbf{h}_k.$$

We have

$$\tilde{L}(e^{i\phi_k} b_k) = \chi_k(\tilde{L}f_k) + \frac{1}{i}(f_k - m_{2Q_k}f_k)\nabla \chi_k - \left(\int_{2Q_k} f_k \right) \chi_k \mathbf{h}_k,$$

where $f_k = e^{i\phi_k} f$. Thus,

$$\left(\int_{2Q_k} |Lb_k|^p \right)^{1/p} \leq C \left\{ \left(\int_{2Q_k} |\tilde{L}f_k|^p \right)^{1/p} \|\chi_k\|_\infty + \left(\int_{2Q_k} |f_k - m_{2Q_k}f_k|^p \right)^{1/p} \|\nabla \chi_k\|_\infty \right\}$$

$$+ \left(\int_{2Q_k} |\mathbf{h}_k|^p \int_{2Q_k} |f_k|^p \right)^{1/p} \|\chi_k\|_\infty \}.$$

using Poincaré inequality and condition (3.8), we obtain

$$\begin{aligned} \left(\int_{2Q_k} |Lb_k|^p \right)^{1/p} &\leq C \left\{ \left(\int_{2Q_k} |\tilde{L}f_k|^p \right)^{1/p} + \left(\int_{2Q_k} |\nabla f_k|^p \right)^{1/p} + \left(\int_{2Q_k} |\mathbf{h}_k|^p \int_{2Q_k} |f_k|^p \right)^{1/p} \right\} \\ &\leq C \left\{ \left(\int_{2Q_k} |\tilde{L}f_k|^p \right)^{1/p} + \left(\int_{2Q_k} \left| \frac{1}{i} \nabla f_k - \mathbf{h}_k f_k \right|^p \right)^{1/p} \right. \\ &\quad \left. + \left(\int_{2Q_k} |\mathbf{h}_k|^p \int_{2Q_k} |f_k|^p \right)^{1/p} + \left(\int_{2Q_k} |\mathbf{h}_k f_k|^p \right)^{1/p} \right\} \\ &\leq C \left\{ \left(\int_{2Q_k} |\tilde{L}f_k|^p \right)^{1/p} + CR_k \int_{2Q_k} V \left(\int_{2Q_k} |f_k|^p \right)^{1/p} \right\} \\ &\leq C \left\{ \left(\int_{2Q_k} |\tilde{L}f_k|^p \right)^{1/p} + \left(\int_{2Q_k} V \right)^{\frac{1}{2}} \left(\int_{2Q_k} |f_k|^p \right)^{1/p} \right\}. \end{aligned}$$

Fefferman-Phong inequality (2.8) and $\left(\int_{2Q_k} V \right)^{\frac{1}{2}} \leq C \left(\int_{2Q_k} V^{p/2} \right)^{1/p}$ imply

$$\left(\int_{2Q_k} |Lb_k|^p \right)^{1/p} \leq C \left(\int_{2Q_k} |\tilde{L}f_k|^p + |V^{\frac{1}{2}} f_k|^p \right)^{1/p}.$$

Using gauge invariance, it follows $|L(f)| = |\tilde{L}(f_k)|$ and we deduce

$$\int_{2Q_k} |Lb_k|^p \leq C \left\{ \int_{2Q_k} |Lf|^p + |V^{\frac{1}{2}} f|^p \right\} \leq C\alpha^p.$$

Moreover,

$$R_k^{-p} \int_{2Q_k} |b_k|^p = R_k^{-p} \int_{2Q_k} |(f_k - m_{2Q_k} f_k) \chi_k|^p \leq C\alpha^p,$$

here we used the previous argument. Hence (3.3) holds for $2Q_k$ of type 2.

d) Definition and properties of $|B|^{\frac{1}{2}}g$:

Set $g = f - \sum b_k$. Note that, by (3.5), this sum is locally finite. It is clear that $g = f$ on F and $g = \sum_{k \in J} e^{-i\phi_k} m_{2Q_k} (e^{i\phi_k} f) \chi_k$ on Ω , where J is the set of indices k such that Q_k is of type 2.

$$\int |V^{\frac{1}{2}} g|^2 = \int_F |V^{\frac{1}{2}} g|^2 + \int_\Omega |V^{\frac{1}{2}} g|^2 = I + II.$$

By construction,

$$I = \int_F |V^{\frac{1}{2}} g|^2 = \int_F |V^{\frac{1}{2}} f|^2 \leq c\alpha^{2-p} (\|Lf\|_p + \|V^{\frac{1}{2}} f\|_p)^p.$$

Using the L^1 version Fefferman-Phong inequality (2.7) we obtain

$$\begin{aligned} II = \int_\Omega |V^{\frac{1}{2}} g|^2 &\leq c \sum_{k \in J} |Q_k| \left[\int_{2Q_k} V^{\frac{1}{2}} \int_{2Q_k} |f|^2 \right] \leq C \sum_{k \in J} |Q_k| \alpha^2 \\ &\leq c\alpha^2 \cdot \alpha^{-p} \int_{\mathbb{R}^n} |Lf|^p + |V^{\frac{1}{2}} f|^p. \end{aligned}$$

Then

$$(3.12) \quad \left(\int |V^{\frac{1}{2}}g|^2 \right)^{\frac{1}{2}} \leq c\alpha^{1-\frac{p}{2}} (\|Lf\|_p + \|V^{\frac{1}{2}}f\|_p)^{p/2}.$$

e) Calculation of Lg :

Let K the set of indices k . Let $\xi \in C_0^\infty(\mathbb{R}^n)$, a test function. We know that, for all $k \in K$ such that $x \in 2Q_k$, there exists $C > 0$ such that $d(x, F) > CR_k$. Therefore,

$$\int \sum_{k \in K} |b_k| |\xi| \leq C \left(\int \sum_{k \in K} \frac{|b_k|}{R_k} \right) \sup_{x \in \mathbb{R}^n} (d(x, F) |\xi(x)|).$$

The estimate (3.3) gives us

$$\int |b_k|^p \leq CR_k^p \alpha^p |Q_k|.$$

Hence

$$\int \sum_{k \in K} |b_k| |\xi| \leq C\alpha |\Omega| \sup_{x \in \Omega} (d(x, F) |\xi(x)|).$$

We conclude that $\sum_{k \in K} b_k$ converges in the sense of distributions in \mathbb{R}^n . Then

$$\nabla g = \nabla f - \sum_{k \in K} \nabla b_k, \text{ in the sense of distributions in } \mathbb{R}^n.$$

Since the sum is locally finite in Ω and vanishes on F , then $\mathbf{a}g = \mathbf{a}f - \sum_{k \in K} \mathbf{a}b_k$ holds almost everywhere in \mathbb{R}^n . Hence

$$Lg = Lf - \sum_{k \in K} Lb_k, \text{ a.e in } \mathbb{R}^n.$$

f) Proof of estimate (3.2):

To prove this inequality, we have to estimate $\|Lg\|_2$. It suffices to prove that $\|Lg\|_\infty \leq C\alpha$ since $\|Lg\|_p \leq C(\|Lf\|_p + \|V^{\frac{1}{2}}f\|_p)$. We know that $\sum_{k \in K} \nabla \chi_k(x) = 0$ for all $x \in \Omega$, then

$$Lg = (Lf)\mathbf{1}_F + \sum_{k \in J} L(e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f)\chi_k) \quad \text{a.e.}$$

$$L(u) = e^{-i\phi_k} \tilde{L}(e^{i\phi_k} u) \quad \text{avec} \quad \tilde{L} = \frac{1}{i} \nabla - \mathbf{h}_k.$$

Hence

$$\begin{aligned} \sum_{k \in J} L(e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f)\chi_k) &= \frac{1}{i} \sum_{k \in J} e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f) \nabla \chi_k - \sum_{k \in J} e^{-i\phi_k} m_{2Q_k}(e^{i\phi_k} f) \chi_k \mathbf{h}_k \\ &= G_1 + G_2. \end{aligned}$$

Now we will control $\|G_2\|_\infty$, we use (3.10), the L^1 version of Fefferman-Phong inequality (2.7), the fact that $2Q_k$ is of type 2 and $V^{\frac{1}{2}} \in RH_2$:

$$|G_2(x)| = \left| \sum_{k \in J} m_{2Q_k}(e^{i\phi_k} f) \chi_k(x) \mathbf{h}_k(x) \right| \leq C \sum_{k \in J, x \in 2Q_k} R_k \int_{2Q_k} V |m_{2Q_k}(e^{i\phi_k} f)|$$

$$\leq C \sum_{k \in J, x \in 2Q_k} \int_{2Q_k} V^{\frac{1}{2}} \int_{2Q_k} |f| \leq CN\alpha.$$

Thus,

$$(3.13) \quad \|G_2\|_\infty \leq C\alpha.$$

Next, we estimate $\|G_1\|_\infty$. Remember that $G_1(x) = \sum_{k \in J} e^{-i\phi_k(x)} m_{2Q_k}(e^{i\phi_k} f) \nabla \chi_k(x)$. For all $m \in K$, consider $K_m = \{l \in K, 2Q_l \cap 2Q_m \neq \emptyset\}$. By construction of Whitney cubes, there exists a constant $c > 0$ (we can take $c = 18$) such that for any $m \in K$, $2Q_l \subset cQ_m$, for all $l \in K_m$. We denote $\tilde{Q}_m = cQ_m$. Let $\tilde{\phi}_m$ and \tilde{h}_m the functions given by then gauge transform of Lemma 2.9 on \tilde{Q}_m . For a fixed x , we have

$$\begin{aligned} G_1(x) &= \sum_{k \in J} e^{-i\phi_k(x)} m_{2Q_k}(e^{i\phi_k} f) \nabla \chi_k(x) \\ &= \sum_{k \in J} (e^{-i\phi_k(x)} m_{2Q_k}(e^{i\phi_k} f) - e^{-i\tilde{\phi}_m(x)} m_{2Q_k}(e^{i\tilde{\phi}_m} f)) \nabla \chi_k(x) \\ &\quad + \sum_{k \in J} e^{-i\tilde{\phi}_m(x)} (m_{2Q_k}(e^{i\tilde{\phi}_m} f) - m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f)) \nabla \chi_k(x) \\ &\quad + \sum_{k \in J} e^{-i\tilde{\phi}_m(x)} m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f) \nabla \chi_k(x) \\ &= I + II + III. \end{aligned}$$

First,

$$III = \sum_{k \in K_m} \chi_m(x) e^{-i\tilde{\phi}_m(x)} m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f) \nabla \chi_k(x) - \sum_{k \in K_m \setminus J} \chi_m(x) e^{-i\tilde{\phi}_m(x)} m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f) \nabla \chi_k(x).$$

The first term in III vanishes since $\sum_{k \in K_m} \nabla \chi_k(x) = \sum_{k \in K} \nabla \chi_k(x) = 0$, for all $x \in 2Q_m$. Since all cubes $2Q_k$ with $k \in K_m \setminus J$ are of type 1, we obtain

$$\begin{aligned} \left| \sum_{k \in K_m \setminus J} \chi_m(x) e^{-i\tilde{\phi}_m(x)} m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f) \nabla \chi_k(x) \right| &\leq C \sum_{k \in K_m \setminus J} R_k^{-1} \int_{\tilde{Q}_m} |f| \\ &\leq CN\tilde{R}_m^{-1} \int_{\tilde{Q}_m} |f| \leq C \int_{\tilde{Q}_m} |Lf| + V|f| \leq C\alpha, \end{aligned}$$

here we used $|Q_k| \sim |Q_m|$, (3.5), the Fefferman-Phong inequality and $4Q_m \cap F \neq \emptyset$. Hence $\|III\|_\infty \leq C\alpha$.

Secondly,

$$\begin{aligned} |II| &= \left| \sum_{k \in J, x \in 2Q_k} e^{-i\tilde{\phi}_m(x)} (m_{2Q_k}(e^{i\tilde{\phi}_m} f) - m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f)) \nabla \chi_k(x) \right| \\ &\leq \sum_{k \in J, x \in 2Q_k} |m_{2Q_k}(e^{i\tilde{\phi}_m} f) - m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f)| \|\nabla \chi_k\|_\infty \\ &\leq C \sum_{k \in J, x \in 2Q_k} |m_{2Q_k}(e^{i\tilde{\phi}_m} f) - m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f)| R_k^{-1}, \end{aligned}$$

since

$$(3.14) \quad |m_{2Q_k}(e^{i\tilde{\phi}_m} f) - m_{\tilde{Q}_m}(e^{i\tilde{\phi}_m} f)| \leq C\tilde{R}_m\alpha,$$

then

$$\|II\|_\infty \leq CN\alpha.$$

The proof of (3.14) is detailed in [Aus].

Finally, we will estimate I :

$$\begin{aligned} e^{-i\phi_k(x)}m_{2Q_k}(e^{i\phi_k}f) - e^{-i\tilde{\phi}_m(x)}m_{2Q_k}(e^{i\tilde{\phi}_m}f) &= e^{-i\phi_k(x)} \int_{2Q_k} e^{i\phi_k(y)} f(y) dy - e^{-i\tilde{\phi}_m(x)} \int_{2Q_k} e^{i\tilde{\phi}_m(y)} f(y) dy \\ &= \int_{2Q_k} (e^{i(\phi_k(y)-\phi_k(x))} - e^{i(\tilde{\phi}_m(y)-\tilde{\phi}_m(x))}) f(y) dy. \end{aligned}$$

Using inequality

$$|e^{i(\phi_k(y)-\phi_k(x))} - e^{i(\tilde{\phi}_m(y)-\tilde{\phi}_m(x))}| \leq |(\phi_k(y) - \phi_k(x)) - (\tilde{\phi}_m(y) - \tilde{\phi}_m(x))|,$$

we obtain

$$|I| \leq \sum_{k \in J, x \in 2Q_k} R_k^{-1} \left| \int_{2Q_k} |f(y)| |(\phi_k(y) - \phi_k(x)) - (\tilde{\phi}_m(y) - \tilde{\phi}_m(x))| dy \right|.$$

By construction, we have

$$\nabla(\phi_k - \tilde{\phi}_m) = \tilde{\mathbf{h}}_m - \mathbf{h}_k$$

. We also have, for all x and $y \in 2Q_k$

$$\begin{aligned} |(\phi_k - \tilde{\phi}_m)(y) - (\phi_k - \tilde{\phi}_m)(x)| &\leq \int_0^1 |x - y| |(\tilde{\mathbf{h}}_m - \mathbf{h}_k)(x + t(y - x))| dt \\ &\leq C|x - y|R_k \int_{2Q_k} V \leq CR_k^2 \int_{2Q_k} V \leq CR_k \left(\int_{2Q_k} V^{\frac{1}{2}} \right), \end{aligned}$$

here we make use of (3.10), the fact that $2Q_k$ is of type 2 and $V^{\frac{1}{2}} \in RH_2$.

Hence

$$|I| \leq \sum_{k \in J, x \in 2Q_k} \int_{2Q_k} V^{\frac{1}{2}} \int_{2Q_k} |f| \leq CN\alpha \text{ a.e.}$$

It follows

$$(3.15) \quad \|G_1\|_\infty \leq C\alpha.$$

We have $Lg = (Lf)\mathbf{1}_F + G_1 + G_2$ almost everywhere.

Since $|Lf| \leq C\alpha$ on F , then using estimates (3.15) and (3.13), we obtain

$$(3.16) \quad \|Lg\|_\infty \leq C\alpha.$$

Hence

$$\|Lg\|_2 + \|V^{\frac{1}{2}}g\|_2 \leq C\alpha^{1-\frac{p}{2}} (\|Lf\|_p + \|V^{\frac{1}{2}}f\|_p)^{p/2},$$

then estimate (3.2) holds. \square

Remark 3.3. Note that we did not use the fact that V is the electrical potential of H : V can be replaced by any weight function ω in A_∞ :

Suppose $\mathbf{a} \in L_{loc}^2(\mathbb{R}^n)^n$ and we assume the following condition for any cube Q in \mathbb{R}^n :

$$(3.17) \quad \sup_Q |B| \leq C \int_Q \omega.$$

Then for any $1 \leq p < \infty$, there exists a constant $C_p > 0$, which depends on (3.17), such that

$$(3.18) \quad \|H(\mathbf{a}, 0)^{\frac{1}{2}}(f)\|_p \leq C_p \|Lf\|_p + \|\omega^{\frac{1}{2}}f\|_p,$$

for all $f \in C_0^\infty(\mathbb{R}^n)$, if $p > 1$, and

$$(3.19) \quad |\{x \in \mathbb{R}^n; |H(\mathbf{a}, 0)^{\frac{1}{2}}f(x)| > \alpha\}| \leq \frac{C_1}{\alpha} \int |Lf| + \omega^{\frac{1}{2}}|f|,$$

for any $\alpha > 0$ and $f \in C_0^\infty(\mathbb{R}^n)$, if $p = 1$.

4. ESTIMATES FOR WEAK SOLUTION

Fix an open set Ω . A subharmonic function on Ω is a function $v \in L_{loc}^1(\Omega)$ such that $\Delta v \geq 0$ in $D'(\Omega)$.

Lemma 4.1. *Suppose $\mathbf{a} \in L_{loc}^2(\mathbb{R}^n)^n$ and $0 \leq V \in L_{loc}^1(\mathbb{R}^n)$. If u is a weak solution of $H(\mathbf{a}, V)u = 0$ in Ω , then $|u|^2$ is a subharmonic function and*

$$(4.1) \quad \Delta|u|^2 = 2|Lu|^2 + 2V|u|^2.$$

Proof. Since

$$\Delta|u|^2 = \Delta(u\bar{u}) = 2\text{Re}((\Delta u)\bar{u}) + 2|\nabla u|^2,$$

and $H(\mathbf{a}, V)u = 0$, then

$$\Delta u = \sum_{k=1}^n (ia_k \frac{\partial u}{\partial x_k} + i \frac{\partial}{\partial x_k}(a_k u)) + |\mathbf{a}|^2 u + Vu.$$

It follows that

$$\begin{aligned} \Delta|u|^2 &= 2\text{Re} \left(\sum_{k=1}^n (ia_k \frac{\partial u}{\partial x_k} + i \frac{\partial}{\partial x_k}(a_k u)) \bar{u} + |\mathbf{a}|^2 u \bar{u} + Vu \bar{u} \right) + 2|\nabla u|^2 \\ &= 2\text{Re} \left(\sum_{k=1}^n (ia_k \frac{\partial u}{\partial x_k} \bar{u} + i \frac{\partial}{\partial x_k}(a_k u) \bar{u}) \right) + 2|\mathbf{a}|^2 |u|^2 + 2V|u|^2 + 2|\nabla u|^2 \\ &= 2\text{Re} \left(\sum_{k=1}^n (ia_k \frac{\partial u}{\partial x_k} \bar{u} + i \frac{\partial}{\partial x_k}(a_k |u|^2) - ia_k u \frac{\partial \bar{u}}{\partial x_k}) \right) + 2|\mathbf{a}|^2 |u|^2 + 2V|u|^2 + 2|\nabla u|^2 \\ &= 4\text{Im}(\mathbf{a} \nabla u \bar{u}) + 2|\mathbf{a}|^2 |u|^2 + 2|\nabla u|^2 + 2|Vu|^2 = 2|Lu|^2 + 2V|u|^2. \end{aligned}$$

□

Corollary 4.2. *Let Q a cube in \mathbb{R}^n and u a weak solution of $H(\mathbf{a}, V)u = 0$ in a neighbourhood of $2\bar{Q}$. For $V \geq 0$, we have the following inequality*

$$(4.2) \quad \sup_Q |u| \leq C(r, n, \mu) \left(\int_{\mu Q} |u|^r \right)^{1/r},$$

for any $0 < r < \infty$ and $1 < \mu \leq 2$.

The following technical lemma is interesting in its own right. For a detailed proof see [Buc] and [AB]. It states that a form of the mean value inequality for subharmonic functions still holds if the Lebesgue measure is replaced by a weighted measure of Muckenhoupt type. More precisely,

Lemma 4.3. *Let $\omega \in RH_q$ for some $1 < q \leq \infty$ and let $0 < s < \infty$ and $r > q$ (if $q = \infty$, $r = \infty$) such that $\omega \in RH_r$. Then there exists a constant $C \geq 0$ depending only on ω, r, p, s and n , such that for any cube Q and any nonnegative subharmonic function f in a neighbourhood of $\overline{2Q}$ we have for all $1 < \mu \leq 2$,*

$$\left(\int_Q (\omega f^s)^r \right)^{1/r} \leq C \int_{\mu Q} \omega f^s, \text{ for } r < +\infty.$$

And

$$\sup_Q f \leq \frac{C}{\int_Q \omega} \int_{\mu Q} \omega f^s, \text{ for } r = +\infty.$$

Throughout this section we will assume $V \in RH_q$ with $1 < q \leq +\infty$ and B satisfies the assumption (1.8) and u is a weak solution of $H(\mathbf{a}, V)u = 0$ in $4Q$. We will establish some local estimates on $|u|$ and $|Lu|$. Using the gauge transform on $4Q$, we can replace \mathbf{a} and L by \mathbf{h} and $\frac{1}{i}\nabla - \mathbf{h}$ as defined previously in Lemma 2.9.

All the constants are independant of Q and u but they may depend on V and q .

First we give three important results that are the main tools for the proof of Theorem 1.2

Proposition 4.4. *There exists a constant $C > 0$ such that*

$$(4.3) \quad \left(\int_Q |V^{\frac{1}{2}}u|^{2q} \right)^{\frac{1}{2q}} \leq C \left(\int_{3Q} |V^{\frac{1}{2}}u|^2 \right)^{\frac{1}{2}}.$$

Proof. It is a consequence of Lemma 4.3 and Lemma 4.1. □

Proposition 4.5. *For all $k > 0$, there exists a constant C such that*

$$(4.4) \quad \left(\int_Q |Lu|^{q^*} \right)^{1/q^*} \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{3Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}}.$$

Proposition 4.6. *Let $1 < \mu \leq 4$, if $n/2 \leq q < n$ then there exists a constant C such that*

$$(4.5) \quad \left(\int_Q |Lu|^{q^*} \right)^{1/q^*} \leq C \left(\int_{\mu Q} |Lu|^2 \right)^{\frac{1}{2}},$$

If $q \geq n$ then there exists a constant C such that

$$(4.6) \quad \sup_Q |Lu| \leq C \left(\int_{\mu Q} |Lu|^2 \right)^{\frac{1}{2}}.$$

Remark 4.7. *Using Theorem 2 of [IN], we can replace 2 by $\delta \in]0, 2]$ in (4.5) and (4.6).*

We need the following results to prove propositions 4.5 and 4.6:

Lemma 4.8. *Let $1 \leq \mu < \mu' \leq 4$ and $k > 0$, then there exists a constant C such that*

$$\int_{\mu Q} |u|^2 \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{\mu' Q} |u|^2 \right),$$

and

$$\int_{\mu Q} |Lu|^2 + V|u|^2 \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{\mu' Q} |Lu|^2 + V|u|^2 \right).$$

The proof is analogous to that of Lemma 8.1 in [AB], it is based on Caccioppoli (2.3) type inequality and the improved Fefferman-Phong inequality (2.8).

Lemma 4.9. *For any $1 < \mu \leq 4$ and $k > 0$, there exists a constant C such that*

$$(R \int_Q V)^2 \int_Q |u|^2 \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{\mu Q} V |u|^2 \right).$$

Proof. Using Lemma 4.8 with $k > 1$ and $1 < \mu' < \mu$ and subsequently Lemma 4.3, we have:

$$(R \int_Q V)^2 \int_Q |u|^2 \leq \frac{C \int_Q V \int_{\mu' Q} |u|^2}{(1 + R^2 \int_Q V)^{k-1}} \leq \frac{C \int_{\mu' Q} V \sup_{\mu' Q} |u|^2}{(1 + R^2 \int_Q V)^{k-1}} \leq \frac{C \int_{\mu Q} (V |u|^2)}{(1 + R^2 \int_Q V)^{k-1}}.$$

□

Lemma 4.10. *For any $1 < \mu \leq 2$, $k > 0$ and $n < p < \infty$, there exists a constant C such that*

$$(R \int_Q V)^2 \int_Q |u|^2 \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{\mu Q} |Lu|^p \right)^{2/p}.$$

Proof. If $\int_{\mu Q} |Lu|^p = \infty$, there is nothing to prove. Assume, therefore, that $\int_{\mu Q} |Lu|^p < \infty$. Let $1 < \nu < \mu$ and η be a smooth non-negative function, bounded by 1, equal to 1 on νQ with support on μQ and whose gradient is bounded by C/R and Laplacian by C/R^2 .

Integrating the equation $H(\mathbf{a}, 0)u + Vu = 0$ against $\bar{u}\eta^2$.

Since

$$H(\mathbf{a}, V)u = \sum_{j=1}^n L_j^* L_j u + Vu,$$

$$\int H(\mathbf{a}, V)u \bar{u}\eta^2 = \sum_{j=1}^n \int L_j u \overline{L_j(u\eta^2)} + \int V |u|^2 \eta^2,$$

then

$$\int |Lu|^2 \eta^2 + V |u|^2 \eta^2 = 2 \int Lu \cdot \nabla \eta \bar{u}\eta,$$

hence

$$\int V |u|^2 \eta^2 \leq \frac{C}{R} \left(\int_{\mu Q} |Lu|^2 \right)^{1/2} \left(\int |u|^2 \eta^2 \right)^{1/2},$$

$$(4.7) \quad X \leq C (R^2 \int_Q V)^{1/2} |\mu Q|^{1/2} Y^{1/2} Z^{1/2}$$

where we set $X = (R^2 \int_Q V) \int V |u|^2 \eta^2$, $Y = (\int_{\mu Q} |Lu|^p)^{2/p}$ and $Z = \int_Q V \int |u|^2 \eta^2$. By Morrey's embedding theorem and diamagnetic inequality (2.4), u is Hölder continuous with exponent $\alpha = 1 - n/p$. Hence for all $x, y \in \mu Q$, we have

$$\left| |u(x)| - |u(y)| \right| \leq C \left(\frac{|x-y|}{R} \right)^\alpha R \left(\int_{\mu Q} |\nabla |u||^p \right)^{1/p} \leq C \left(\frac{|x-y|}{R} \right)^\alpha R Y^{1/2}.$$

We pick $y \in \overline{Q}$ such that $|u(y)| = \inf_Q |u|$. Then

$$\begin{aligned}
Z &= \int_Q V \int |u|^2 \eta^2 \leq 2 \left(\int_Q V \right) \inf_Q |u|^2 \int \eta^2 + 2 \left(\int_Q V \right) \int ||u(x)| - |u(y)||^2 \eta^2(x) dx \\
&\leq 2 \left(\int_Q (V|u|^2) \right) \int \eta^2 + C \left(\int_Q V \right) R^2 Y \int \left(\frac{|x-y|}{R} \right)^{2\alpha} \eta^2(x) dx \\
&\leq C \left(\int_Q (V|u|^2) \right) |Q| + C \left(\int_Q V \right) R^2 Y |\mu Q| \\
&\leq C \int V |u|^2 \eta^2 + C \left(\int_Q V \right) R^2 Y |\mu Q|.
\end{aligned}$$

where, in the penultimate inequality, we used the support condition on η and $0 \leq \eta \leq 1$, and in the last, $\eta = 1$ on Q . Using the previous inequalities, we obtain

$$X \leq C |\mu Q|^{1/2} Y^{1/2} (CX + C(R^2 \int_Q V)^2 |\mu Q| Y)^{1/2},$$

which, as $2ab \leq \epsilon^{-1}a^2 + \epsilon b^2$ for all $a, b \geq 0$ and $\epsilon > 0$, implies

$$X \leq C(1 + R^2 \int_Q V)^2 |\mu Q| Y.$$

Next, let $1 < \nu' < \nu$. Using $\eta = 1$ on νQ Lemma 4.3 and Lemma 4.8

$$\int V |u|^2 \eta^2 \geq \int_{\nu Q} V |u|^2 \geq C \int_{\nu' Q} V \int_{\nu' Q} |u|^2 \geq C \left(\int_Q V \right) (1 + R^2 \int_Q V)^k \int_Q |u|^2,$$

hence

$$X \geq C(R \int_Q V)^2 (1 + R^2 \int_Q V)^k \int_Q |u|^2.$$

The upper and lower bounds for X yield the lemma. □

Now we will give the proof of 4.5 and 4.6.

Proof of Proposition 4.5: We will assume that $q > \frac{2n}{n+2}$.

a) Preparation:

We remind that modulo a gauge transformation, u is a weak solution of $H(\mathbf{h}, V)u = 0$ on $4Q$, where \mathbf{h} is the potential function defined in Lemma 2.9. We call $L = \frac{1}{i}\nabla - \mathbf{h}$.

Let v a weak solution of $\Delta v = 0$ in $2Q$ with $v = u$ on $\partial(2Q)$. Set $w = u - v$ in $2Q$. From elliptic theory we know that

$$\int_{2Q} |\nabla w|^2 \leq \int_{2Q} |\nabla u|^2,$$

thus,

$$\int_{2Q} |\nabla v|^2 \leq 4 \int_{2Q} |\nabla u|^2.$$

It suffices to establish the following inequality for $1 \leq \mu < 2$,

$$(4.8) \quad \sup_{\mu Q} |Lv| \leq C \left(\int_{\frac{3}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}},$$

and

$$(4.9) \quad \left(\int_Q |Lw|^{q^*} \right)^{1/q^*} \leq \phi(R^2 \int_Q V) \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}},$$

where ϕ is a real function which increases polynomially. The conclusion follows easily through Lemma 4.8.

b) Estimate of v :

Since ∇v is harmonic on $2Q$ then for $1 \leq \mu < 2$,

$$\sup_{\mu Q} |\nabla v| \leq C \int_{2Q} |\nabla v|^2 \leq 4C \int_{2Q} |\nabla u|^2.$$

Using the harmonicity of v and the fact that $v = u$ on $\partial(2Q)$, we obtain

$$\sup_{2Q} |v|^2 \leq \sup_{\partial(2Q)} |v| = \sup_{\partial(2Q)} |u|,$$

and (4.2) gives

$$(4.10) \quad \sup_{2Q} |v| \leq C \left(\int_{\frac{9}{4}Q} |u|^2 \right)^{\frac{1}{2}}.$$

(1.8), (2.13), Lemma 4.9 and the fact that $\int_{2Q} V \sim \int_{\frac{9}{4}Q} V$, yield

$$(4.11) \quad \sup_{2Q} |\mathbf{h}v| \leq C \left[(R \int_{2Q} V)^2 \int_{\frac{9}{4}Q} |u|^2 \right]^{\frac{1}{2}} \leq C \left[\int_{\frac{5}{2}Q} V |u|^2 \right]^{\frac{1}{2}}.$$

Hence, inequality (4.8) holds.

c) Estimate of w :

Using $w = u - v$, (4.10) and (4.2), we obtain

$$(4.12) \quad \sup_{2Q} |w| \leq C \left(\int_{\frac{9}{4}Q} |u|^2 \right)^{\frac{1}{2}},$$

and

$$(4.13) \quad \sup_{2Q} |\mathbf{h}w| \leq C \left(\int_{\frac{5}{2}Q} V |u|^2 \right)^{\frac{1}{2}}.$$

It suffices to estimate ∇w . We have

$$\int_{2Q} |\nabla w|^2 \leq \int_{2Q} |\nabla u|^2 \leq 4 \int_{2Q} |Lu|^2 + \int_{2Q} |\mathbf{h}u|^2,$$

by combining (4.11) and (4.13), it follows

$$(4.14) \quad \int_{2Q} |\nabla w|^2 \leq C \int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2.$$

Now we will control $\int_Q |\nabla w|^{q^*}$. Let $1 < \mu < \mu' < 2$ and η be a smooth non-negative function, bounded by 1, equal to 1 on μQ with support on $\mu' Q$ and whose gradient is bounded by C/R and Laplacian by C/R^2 . We know that

$$0 = H(\mathbf{h}, V)(u) = -\Delta u - \frac{1}{i} \nabla \cdot (\mathbf{h}u) - \frac{1}{i} \mathbf{h} \cdot \nabla u + (|\mathbf{h}|^2 + V)u, \text{ on } 4Q.$$

Then

$$\begin{aligned} \Delta u &= -\frac{1}{i} (\nabla \cdot \mathbf{h})u - \frac{2}{i} \mathbf{h} \cdot \nabla u + (|\mathbf{h}|^2 + V)u \\ &= -\frac{1}{i} (\nabla \cdot \mathbf{h})u - 2\mathbf{h} \cdot Lu + (-|\mathbf{h}|^2 + V)u. \end{aligned}$$

Since $\Delta w = \Delta u$ on $2Q$, then

$$\begin{aligned} \Delta(w\eta) &= (\Delta w)\eta + 2\nabla w \cdot \nabla \eta + w\Delta \eta \\ &= -\frac{1}{i} (\nabla \cdot \mathbf{h})w\eta - 2(\mathbf{h} \cdot Lu)\eta + (-|\mathbf{h}|^2 + V)w\eta + 2\nabla w \cdot \nabla \eta + w\Delta \eta. \end{aligned}$$

Let Γ_0 be the fundamental solution of Δ . We know that

$$|\nabla \Gamma_0(x)| \leq C|x|^{1-n}.$$

Hence, for $x \in \mu' Q$,

$$\begin{aligned} |\nabla w(x)| &\leq C \left(\int_{\mathbb{R}^n} \frac{|\mathbf{h}(y)Lu(y)|\eta(y)}{|x-y|^{n-1}} dy + \int_{\mathbb{R}^n} \frac{(|\nabla \mathbf{h}(y)| + |\mathbf{h}(y)|^2)|u(y)|\eta(y)}{|x-y|^{n-1}} dy \right. \\ &\quad \left. + \int_{\mathbb{R}^n} \frac{V(y)|u(y)|\eta(y)}{|x-y|^{n-1}} dy + \int_{\mu' Q \setminus \mu Q} \left(\frac{|\nabla w(y)|}{R|x-y|^{n-1}} + \frac{|w(y)|}{R^2|x-y|^{n-1}} \right) dy \right). \end{aligned}$$

Using inequalities (2.13), (2.14) and (1.8), we obtain

$$\begin{aligned} |\nabla w(x)| &\leq C \left((R \int_Q V) \int_{\mathbb{R}^n} \frac{|Lu(y)|\eta(y)}{|x-y|^{n-1}} dy + \left(\int_Q V + R \left(\int_Q V \right)^{\frac{3}{2}} + (R \int_Q V)^2 \right) \int_{\mathbb{R}^n} \frac{|u(y)|\eta(y)}{|x-y|^{n-1}} dy \right. \\ (4.15) \quad &\quad \left. + \int_{\mathbb{R}^n} \frac{V(y)|u(y)|\eta(y)}{|x-y|^{n-1}} dy + \left\{ \int_{2Q} |\nabla w| + \frac{1}{R}|w| \right\} \right) = I + II + III + IV. \end{aligned}$$

Poincaré inequality and (4.14) imply

$$\int_{2Q} |\nabla w| + \frac{1}{R}|w| \leq C \left(\int_{2Q} |\nabla w|^2 \right)^{\frac{1}{2}} \leq C \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}}.$$

Now we have to estimate *III*. We use the Hardy-Littlewood-Sobolev theorem, the fact that $V \in RH_q$, estimate (4.2) and Lemma 4.9:

$$\begin{aligned} \left(\int_{\mu' Q} III^{q^*} \right)^{\frac{1}{q^*}} &\leq CR \left(\int_{\mu Q} |Vu|^q \right)^{\frac{1}{q}} \leq CR \int_{2Q} \sup_{2Q} |u| \\ &\leq C \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Since

$$\int_{\mathbb{R}^n} \frac{|u(y)|\eta(y)}{|x-y|^{n-1}} dy \leq \sup_{2Q} |u| \int_{2Q} \frac{dy}{|x-y|^{n-1}} \leq CR \sup_{2Q} |u|$$

Then, (4.2) and Lemma 4.9 imply :

$$\begin{aligned} \sup_{\mu'Q} |II| &\leq C(1 + (R \int_Q V)^{\frac{1}{2}} + R \int_Q V) R \int_Q V \sup_{2Q} |u| \\ &\leq C(1 + R \int_Q V) \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Finally, we apply Hardy-Littlewood-Sobolev theorem and we obtain

$$\left(\int_{\mu'Q} I^{q^*} \right)^{\frac{1}{q^*}} \leq CR^2 \int_Q V \left(\int_{\mu Q} |Lu|^q \right)^{\frac{1}{q}}.$$

Hence

$$\left(\int_{\mu'Q} |\nabla w|^{q^*} \right)^{1/q^*} \leq C(1 + R^2 \int_Q V) \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}} + CR^2 \int_Q V \left(\int_{\mu Q} |Lu|^q \right)^{\frac{1}{q}}.$$

Since $Lw = Lu - Lv$, we combine the above inequality with (4.8) :

$$\left(\int_{\mu'Q} |Lw|^{q^*} \right)^{\frac{1}{q^*}} \leq C(1 + R^2 \int_Q V) \left(\int_{\frac{5}{2}Q} |Lu|^2 + V|u|^2 \right)^{\frac{1}{2}} + C(R^2 \int_Q V) \left(\int_{\mu Q} |Lw|^q \right)^{\frac{1}{q}}.$$

By iterating the previous inequality and using (4.14) we finish the proof of (4.9). \square

Proof of Proposition 4.6: The proof is the same as that of Proposition 4.5, we use Lemma 4.9 instead of Lemma 4.10.

Let us quote two additional results which ensue at once from what precedes but which we shall not use.

Corollary 4.11. *Let $1 < \mu \leq 2$ and $k > 0$. If $q < n$, then there exists a constant C such that*

$$(4.16) \quad \left(\int_Q |Lu|^{q^*} \right)^{1/q^*} \leq \frac{C}{R(1 + R^2 \int_Q V)^k} \left(\sup_{\mu Q} |u| \right).$$

If $q \geq n$, then there exists a constant C such that

$$(4.17) \quad \sup_Q |Lu| \leq \frac{C}{R(1 + R^2 \int_Q V)^k} \left(\sup_{\mu Q} |u| \right).$$

Proof. It remains to use Caccioppoli type inequality (2.3) and Proposition 4.6. \square

Corollary 4.12. *Let $q \geq n/2$. For any $1 < \mu \leq 2$ and $k > 0$ there exists a constant C such that*

$$(R \int_Q V)^2 \int_Q |u|^2 \leq \frac{C}{(1 + R^2 \int_Q V)^k} \left(\int_{\mu Q} |Lu|^2 \right).$$

Proof. The proof is a consequence of Lemma 4.11 and Proposition 4.6. \square

5. MAXIMAL INEQUALITIES

We give some important maximal inequalities that we will use to prove our results around the Riesz transforms. These estimates are essentially a consequence of some L^p estimates proved in [AB] and based on the L^1 inequality deduced from the work of Gallouët and Morel [GM] in the semi-linear setting or from Kato's paper [K2].

Theorem 5.1. *Let $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in RH_q$, $1 < q \leq +\infty$. Then, there exists $\epsilon > 0$, depending only on the RH_q constant of V , such that $VH(\mathbf{a}, V)^{-1}$ and $H(\mathbf{a}, 0)H(\mathbf{a}, V)^{-1}$ are L^p bounded for all $1 \leq p < q + \epsilon$.*

Corollary 5.2. *Let $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in RH_q$, $1 < q \leq +\infty$. Then, there exists an $\epsilon > 0$, depending only on the RH_q constant of V , such that, the operators $V^{1/2}H(\mathbf{a}, V)^{-1/2}$ and $H(\mathbf{a}, 0)^{1/2}H(\mathbf{a}, V)^{-1/2}$ are L^p bounded for all $1 < p < 2q + \epsilon$.*

To prove this result, we shall use complex interpolation relying on the fact that for all $y \in \mathbb{R}$, the imaginary power of Schrödinger operator H^{iy} has a bounded extension on \mathbb{R}^n , $1 < p < \infty$. This result due to Hebisch [H] follows from the Gaussian estimates on the heat kernel e^{-tH} proved by [DR]. Here, H^{iy} is defined as a bounded operator on $L^2(\mathbb{R}^n)$ by functional calculus (see [AB] for more details).

Proof of Theorem 5.1:

The proof of this theorem is identical to that of Theorem 1.1 in [AB]. First we prove an L^1 inequality, then we establish some reverse Hölder type estimates, then finally we apply Theorem 2.3.

Lemma 5.3. *Let $f \in L^\infty_{comp}(\mathbb{R}^n)$ and $u = H(\mathbf{a}, V)^{-1}f$. Then,*

$$(5.1) \quad \int_{\mathbb{R}^n} V|u| \leq \int_{\mathbb{R}^n} |f|,$$

and

$$(5.2) \quad \int_{\mathbb{R}^n} |H(\mathbf{a}, 0)u| \leq 2 \int_{\mathbb{R}^n} |f|.$$

Proof. $V \geq 0$, by Kato-Simon inequality (2.5), we have

$$|H(\mathbf{a}, V)^{-1}f| \leq H(0, V)^{-1}|f|.$$

We know, by [AB] that

$$\int_{\mathbb{R}^n} VH(0, V)^{-1}|f| \leq \int_{\mathbb{R}^n} |f|.$$

Thus, inequality (5.1) holds, and inequality (5.2) follows by difference. \square

Proof of the L^p maximal inequality:

Assume $V \in RH_q$ with $q > 1$. $VH(\mathbf{a}, V)^{-1}$. We know that this operator is bounded on $L^1(\mathbb{R}^n)$, so we apply Theorem 2.3 through the reverse Hölder inequality verified by any weak solution. Set Q a fixed cube and $f \in L^\infty(\mathbb{R}^n)$ a function with compact support in $\mathbb{R}^n \setminus 4Q$. Then $u = H(\mathbf{a}, V)^{-1}f$ is well defined in $\dot{\mathcal{V}}$ and it is a weak solution of $H(\mathbf{a}, 0)u + Vu = 0$ in $4Q$.

Since $|u|^2$ is subharmonic, by Lemma 4.3 with $w = V$, $f = |u|^2$ and $s = 1/2$, we obtain

$$\left(\int_Q |Vu|^q \right)^{1/q} \leq C \int_{2Q} |Vu|.$$

Thus (2.1) holds with $T = VH(\mathbf{a}, V)^{-1}$, $p_0 = 1$, $q_0 = q$, $S = 0$, $\alpha_1 = 2$ and $\alpha_2 = 4$. Hence $VH(\mathbf{a}, V)^{-1}$ is bounded on $L^p(\mathbb{R}^n)$ for all $1 < p < q$ by Theorem 2.3. Due to the properties of RH_q weights, we can replace q by $q + \epsilon$. Taking the difference, we

obtain the same result for $H(\mathbf{a}, 0)H(\mathbf{a}, V)^{-1}$. This completes the proof of Theorem 5.1. \square

6. PROOF OF THEOREM 1.2

Now we will focus on the proof of Theorem 1.2 we first need to establish some important inequalities. First of all we reduce the problem thanks to the following Lemma:

Lemma 6.1. *Under the assumptions of Theorem 1.2. For any $p > 2$, the L^p boundedness of $LH(\mathbf{a}, V)^{-\frac{1}{2}}$ is equivalent to that of $LH(\mathbf{a}, V)^{-1}L^*$ and $LH(\mathbf{a}, V)^{-1}V^{\frac{1}{2}}$.*

Proof. If $LH(\mathbf{a}, V)^{-\frac{1}{2}}$ is L^p bounded, then by duality and using L^p boundedness of Riesz transforms for $1 < p \leq 2$, we have $H(\mathbf{a}, V)^{-\frac{1}{2}}L^*$ is L^q bounded for any $q \geq 2$. Hence $LH(\mathbf{a}, V)^{-1}L^*$ is L^p bounded. By the same argument, $H(\mathbf{a}, V)^{-\frac{1}{2}}V^{\frac{1}{2}}$ is L^p bounded, and hence $LH(\mathbf{a}, V)^{-1}V^{\frac{1}{2}}$ is L^p bounded.

Reciprocally, if $LH(\mathbf{a}, V)^{-1}L^*$ and $LH(\mathbf{a}, V)^{-1}V^{\frac{1}{2}}$ are L^p bounded, then their adjoints $LH(\mathbf{a}, V)^{-1}L^*$ and $V^{\frac{1}{2}}H(\mathbf{a}, V)^{-1}L^*$ are $L^{p'}$ bounded.

Let $\mathbf{F} \in C_0^\infty(\mathbb{R}^n, \mathbb{C}^n)$, then

$$\|H(\mathbf{a}, V)^{-\frac{1}{2}}L^*\mathbf{F}\|_{p'} = \|H(\mathbf{a}, V)^{\frac{1}{2}}H(\mathbf{a}, V)^{-1}L^*\mathbf{F}\|_{p'}.$$

Using assumption (1.8) and inequality (1.12), we obtain

$$\|H(\mathbf{a}, V)^{-\frac{1}{2}}L^*\mathbf{F}\|_{p'} \leq C(\|LH(\mathbf{a}, V)^{-1}L^*\mathbf{F}\|_{p'} + \|V^{\frac{1}{2}}H(\mathbf{a}, V)^{-1}L^*\mathbf{F}\|_{p'}) \leq C\|\mathbf{F}\|_{p'}.$$

Thus $LH(\mathbf{a}, V)^{-\frac{1}{2}}$ is L^p bounded. \square

Next we look at some useful related estimates

Proposition 6.2. *Let $V \in RH_q$, $1 < q \leq +\infty$. Then, for any $2 < p < 2(q + \epsilon)$, and $\epsilon > 0$ depending only on V , $f \in C_0^\infty(\mathbb{R}^n, \mathbb{C})$ and $\mathbf{F} \in C_0^\infty(\mathbb{R}^n, \mathbb{C}^n)$,*

$$\|V^{\frac{1}{2}}H(\mathbf{a}, V)^{-1}V^{\frac{1}{2}}f\|_p \leq C_p\|f\|_p, \quad \|V^{\frac{1}{2}}H(\mathbf{a}, V)^{-1}L^*\mathbf{F}\|_p \leq C_p\|\mathbf{F}\|_p.$$

Proof. To effectuate the proof, we apply Theorem 2.3 and eventually use equation (4.3). \square

To prove Theorem 1.2, it suffices to prove the following Proposition:

Proposition 6.3. *Let $V \in RH_q$, $1 < q \leq +\infty$. Then, for any $2 < p < q^* + \epsilon$ and $\epsilon > 0$ depending only on V , $f \in C_0^\infty(\mathbb{R}^n, \mathbb{C})$ and $\mathbf{F} \in C_0^\infty(\mathbb{R}^n, \mathbb{C}^n)$,*

$$\|LH(\mathbf{a}, V)^{-1}V^{\frac{1}{2}}f\|_p \leq C_p\|f\|_p, \quad \|LH(\mathbf{a}, V)^{-1}L^*\mathbf{F}\|_p \leq C_p\|\mathbf{F}\|_p.$$

Proof. Suppose $q < n/2$. Let Q a cube in \mathbb{R}^n and $\mathbf{F} \in C_0^\infty(\mathbb{R}^n)$ supported away from $4Q$. Set $H = H(\mathbf{a}, V)$, $u = H^{-1}L^*\mathbf{F}$ is well defined on \mathbb{R}^n . In particular, the support condition on \mathbf{F} , implies that u is a weak solution of $Hu = 0$ in $4Q$. Hence, using Proposition 4.5, we have

$$(6.1) \quad \left(\int_Q |LH^{-1}L^*\mathbf{F}|^{q^*} dx \right)^{1/q^*} \leq C \left(\int_{3Q} |LH^{-1}L^*\mathbf{F}|^2 + |V^{\frac{1}{2}}H^{-1}L^*\mathbf{F}|^2 \right)^{\frac{1}{2}}.$$

Then (2.1) holds with $T = LH^{-1}L^*$, $q_0 = q^*$, $p_0 = 2$ and $S = M_2(V^{\frac{1}{2}}H^{-1}L^*)$, where $M_2f = (M|f|^2)^{\frac{1}{2}}$ and M is the maximal Hardy-Littlewood operator. Since S is L^p

bounded for any $1 < p \leq 2q$, then, using Proposition 6.2 and the fact that $q^* \leq 2q$, T is bounded on $L^p(\mathbb{R}^n, \mathbb{C}^n)$, for $2 < p < q^*$.

Now, let $n/2 \leq q < n$. By the same method and using Proposition 4.6, we obtain

$$(6.2) \quad \left(\int_Q |LH^{-1}L^*\mathbf{F}|^{q^*} dx \right)^{1/q^*} \leq C \left(\int_{3Q} |LH^{-1}L^*\mathbf{F}|^2 \right)^{\frac{1}{2}}.$$

Hence, inequality (2.1) holds with $T = LH^{-1}L^*$, $q_0 = q^*$ and $p_0 = 2$. Thus, T is bounded on L^p , $2 < p < q^*$. Finally, if $q \geq n$, then $LH^{-1}L^*$ is L^p bounded for $2 < p < \infty$.

Using the same argument with $u = H^{-1}V^{\frac{1}{2}}f$, it follows that $LH^{-1}V^{\frac{1}{2}}$ is L^p bounded for $2 < p < q^* + \epsilon$.

□

7. SECOND ORDER RIESZ TRANSFORMS

This section is devoted to the study of some operators using results previously established. We are interested in the behaviour of $L_j L_k H(\mathbf{a}, V)^{-1}$ and $V^{\frac{1}{2}} LH(\mathbf{a}, V)^{-1}$. We also need some properties of the kernel of $H(\mathbf{a}, V)^{-1}$. Note that, for $V \in L^1_{loc}(\mathbb{R}^n)$, $V \geq 0$ and $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$, $H(\mathbf{a}, V)^{-1}$ is continuous from $L^1(\mathbb{R}^n)$ to $L^1_{loc}(\mathbb{R}^n)$. Hence there exists $\Gamma(x, y)$ a Schwartz kernel associated to this operator. The following proposition gives some results about this kernel

Proposition 7.1. Γ coincides with a measurable function defined on $\mathbb{R}^n \times \mathbb{R}^n$ in \mathbb{C} , and

(1) We have the following inequality

$$(7.1) \quad |\Gamma(x, y)| \leq C_n |x - y|^{2-n} \text{ a.e.}$$

(2) For almost every $y \in \mathbb{R}^n$, $u : x \mapsto \Gamma(x, y)$ is a weak solution of $H(\mathbf{a}, V)(u) = 0$ on $\mathbb{R}^n \setminus \{y\}$.

Proof. Let $H = H(\mathbf{a}, V)$. Inequality (2.6) and $H^{-1} = \int_0^\infty e^{-tH} dt$ imply that Γ is dominated by the Green function of the Laplacien. Hence, inequality (7.1) follows. Now we will prove (2). Let $f \in C_0^\infty(\mathbb{R}^n)$, using (1) we have the following integral representation

$$H^{-1}(f)(x) = \int_{\mathbb{R}^n} \Gamma(x, y) f(y) dy$$

for almost every x .

Fix a function $\rho \in C_0^\infty(\mathbb{R}^n)$, even, supported in $[-1, 1]^n$ with $\int \rho = 1$. Set $\rho_j^y = 2^{nj} \rho(2^j(z - y))$ for any $y, z \in \mathbb{R}^n$ and $j \in \mathbb{N}$. Using Fubini theorem, we obtain

$$H^{-1}(\rho_j^0 * f)(x) = \int_{\mathbb{R}^n} \Gamma_j(x, y) f(y) dy, \text{ a.e.}$$

where

$$\Gamma_j(x, y) = \int_{\mathbb{R}^n} \Gamma(x, z) 2^{nj} \rho(2^j(z - y)) dz = H^{-1}(\rho_j^y)(x).$$

Using Lebesgue's dominated convergence theorem, we obtain:

$$\lim_{j \rightarrow \infty} H^{-1}(\rho_j^0 * f) = H^{-1}(f) \text{ a.e.}$$

and

$$\lim_{j \rightarrow \infty} \Gamma_j = \Gamma \text{ a.e in } L^1_{loc}.$$

Since $\rho_j^y \in C_0^\infty(\mathbb{R}^n)$, $H^{-1}(\rho_j^y) \in \dot{\mathcal{V}}$ and $H(\Gamma_j(\cdot, y)) = \rho_j^y \in \mathcal{D}'(\mathbb{R}^n)$. Hence $\Gamma_j(\cdot, y)$ is a weak solution of $Hu = 0$ away from the support of ρ_j^y , i.e $\mathbb{R}^n \setminus Q(y, 2^j)$. Here $Q(c, R)$ is the cube centred in c with sidelength R . By (7.1) and a similar argument to that used for the proof of Caccioppoli type inequality, we obtain for $R > 2^{-j+2}$,

$$\int_{\mathbb{R}^n \setminus Q(y, 2R)} |L_x \Gamma_j(x, y)|^2 + V(x) |\Gamma_j(x, y)|^2 dx \leq \frac{C}{R^2} \int_{Q(y, 2R) \setminus Q(y, R)} |\Gamma_j(x, y)|^2 \leq CR^{2-n}.$$

Hence, for any y , $L_x \Gamma_j(\cdot, y)$ and $V^{\frac{1}{2}} \Gamma_j(\cdot, y)$ admit subsequences that weakly converge in $L^2_{loc}(\mathbb{R}^n \setminus \{y\})$. It is easy to prove that their limits are $L_x \Gamma(\cdot, y)$ and $V^{\frac{1}{2}} \Gamma(\cdot, y)$ in $\mathcal{D}'(\mathbb{R}^n)$ for almost every $y \in \mathbb{R}^n$. We deduce that for almost every $y \in \mathbb{R}^n$, $\Gamma(\cdot, y)$ is a weak solution of $Hu = 0$ on $\mathbb{R}^n \setminus \{y\}$ (by taking limits on the equation). \square

Proposition 7.1 and the following technical lemma are two important tools to prove the main results of this section.

Lemma 7.2. *Let $V \in RH_{n/2}$. Then*

$$\sum_{l \in \mathbb{Z}} \frac{(4^l \int_{Q(y, 2^l)} V)^{\frac{1}{2}}}{1 + 4^l \int_{Q(y, 2^l)} V} \leq C(V)$$

uniformly on $y \in \mathbb{R}^n$.

Proof. Shen proved in [Sh1], Lemma 1.2, that there exists an $\alpha > 0$ such that for any r, R with $r < R$ and $y \in \mathbb{R}^n$,

$$r^2 \int_{Q(y, r)} V \leq C \left(\frac{r}{R}\right)^\alpha R^2 \int_{Q(y, R)} V.$$

Let j be the biggest integer such that $4^j \int_{Q(y, 2^j)} V < 1$. We have

$$\sum_{l \in \mathbb{Z}} \frac{(4^l \int_{Q(y, 2^l)} V)^{\frac{1}{2}}}{1 + 4^l \int_{Q(y, 2^l)} V} \leq C(\alpha) \left((4^j \int_{Q(y, 2^j)} V)^{\frac{1}{2}} + (4^{j+1} \int_{Q(y, 2^{j+1})} V)^{-\frac{1}{2}} \right) \leq 2C(\alpha).$$

\square

Remark 7.3. *This lemma does not hold for $V(x) = \frac{1}{|x|^2} \in \cup_{q < \frac{n}{2}} RH_q$.*

Now we will study the L^p boundedness of $V^{\frac{1}{2}} LH(\mathbf{a}, V)^{-1}$:

Proposition 7.4. *Suppose $\mathbf{a} \in L^2_{loc}(\mathbb{R}^n)^n$ and $V \in RH_{n/2}$. Also assume that there exists a constant $C > 0$ such that for any cube Q in \mathbb{R}^n :*

$$(7.2) \quad \begin{cases} \sup_Q |B| \leq C \int_Q V, \\ \sup_Q |\nabla B| \leq C (\int_Q V)^{3/2}. \end{cases}$$

Then, $V^{\frac{1}{2}} LH(\mathbf{a}, V)^{-1}$ is L^1 bounded.

Proof. It suffices to prove

$$\operatorname{esssup}_{y \in \mathbb{R}^n} \int_{\mathbb{R}^n} V^{\frac{1}{2}}(x) |L_x \Gamma(x, y)| dx < +\infty.$$

To simplify the proof we will take $y = 0$. Set $\Gamma(\cdot, 0)$ the weak solution of $Hu = 0$ on $\mathbb{R}^n \setminus \{0\}$. Let $(Q_{l,k})$ be a Whitney decomposition of $\mathbb{R}^n \setminus \{0\}$: $d(Q_{l,k}, 0) \sim 2^l$ the sidelength of $Q_{l,k}$ and k belongs the finite set of indices of cardinality $2^n - 1$. Hence, using the Cauchy-Schwarz inequality and (7.1).

$$\begin{aligned} \int_{\mathbb{R}^n} V^{\frac{1}{2}}(x) |L_x \Gamma(x, 0)| dx &= \sum_{l,k} \int_{Q_{l,k}} V^{\frac{1}{2}}(x) |L_x \Gamma(x, 0)| dx \\ &\leq C \sum_{l,k} |Q_{l,k}| \left(\int_{Q_{l,k}} V \right)^{\frac{1}{2}} \left(\int_{Q_{l,k}} |L_x \Gamma(x, 0)|^2 dx \right)^{\frac{1}{2}} \\ &\leq C \sum_{l,k} |Q_{l,k}| \left(\int_{Q_{l,k}} V \right)^{\frac{1}{2}} (1 + 4^l \int_{Q_{l,k}} V)^{-1} (4^{-l} \int_{Q_{l,k}} |\Gamma(x, 0)|^2 dx)^{\frac{1}{2}} \\ &\leq \sum_{l,k} \frac{(4^l \int_{Q_{l,k}} V)^{\frac{1}{2}}}{(1 + 4^l \int_{Q_{l,k}} V)}. \end{aligned}$$

Since V is in A_∞ , then $\int_{Q_{l,k}} V \sim \int_{Q(0,2^l)} V$ uniformly on l and k . Hence by Lemma 7.2 as $V \in RH_{n/2}$. \square

Proof of Theorem 1.6: Since $T = V^{\frac{1}{2}} LH(\mathbf{a}, V)^{-1}$ is L^1 bounded, then it suffices to apply Theorem 2.3 to the operator T with $p_0 = 1$ and $q_0 = \frac{2qn}{3n-2q}$ if $q < n$ and $q_0 = 2q$ if $q \geq n$.

Let Q a cube of \mathbb{R}^n and $f \in L^\infty_{\text{comp}}(\mathbb{R}^n)$ with support away from $4Q$. We know that $u = H(\mathbf{a}, V)^{-1} f$ is a weak solution of $H(\mathbf{a}, V)u = 0$ in the neighbourhood of $4Q$. There exists $s < q^*$ such that $\frac{1}{q_0} = \frac{1}{2q} + \frac{1}{s}$. Then,

$$\left(\int_Q (V^{\frac{1}{2}} |Lu|)^{q_0} \right)^{1/q_0} \leq \left(\int_Q V^q \right)^{\frac{q}{2}} \left(\int_Q |Lu|^s \right)^{1/s}.$$

Using the remark 4.7 and $V^{\frac{1}{2}} \in RH_{2r}$ (we used Proposition 2.1) we obtain

$$\left(\int_Q (V^{\frac{1}{2}} |Lu|)^{q_0} \right)^{1/q_0} \leq C \left(\int_Q V^{\frac{1}{2}} \right) \left(\int_{2Q} |Lu|^\delta \right)^{1/\delta},$$

for any $\delta > 0$.

Let $\delta > 0$ such that $V^{\frac{1}{2}} \in A_{1/\delta}$, the Muckenhoupt class. We know that ([St], chap V)

$$\left(\int_{2Q} |g|^\delta \right)^{1/\delta} \leq C \left(\frac{1}{\int_{2Q} V^{\frac{1}{2}}} \right) \int_{2Q} V^{\frac{1}{2}} |g|, \text{ for any measurable and non-negative function } g.$$

Hence

$$\left(\int_Q (V^{\frac{1}{2}} |Lu|)^{q_0} \right)^{1/q_0} \leq C \int_{2Q} (V^{\frac{1}{2}} |Lu|),$$

and (2.1) holds with $S = 0$. Thus T is L^p bounded for any $1 < p < q_0$.

We finish the proof using the self-improvement of the reverse Hölder classes.

Now we will use the previous results to get the proof of Theorem 1.5.

Proof of Theorem 1.5: Set $H(\mathbf{a}, V) = H$ and $H(\mathbf{a}, 0) = H_0$.

$$L_s L_k H^{-1} = L_s H^{-1} L_k + L_s [L_k, H^{-1}].$$

For every $j \geq 1$, $L_j H^{-\frac{1}{2}}$ is L^p bounded for any $1 < p < \infty$ then $L_s H^{-1} L_k$ is L^p bounded for any $1 < p < \infty$. Since

$$[L_k, H^{-1}] = -H^{-1} [L_k, H] H^{-1},$$

$$[L_k, H] = [L_k, H(\mathbf{a}, 0)] + [L_k, V].$$

Then

$$L_s H^{-1} [L_k, V] H^{-1} = (L_s H^{-1} L_k)(V H^{-1}) - (L_s H^{-1} V^{\frac{1}{2}})(V^{\frac{1}{2}} L_k H^{-1}).$$

Proposition 6.3, Theorem 5.1 and Theorem 1.6 imply the L^p boundedness of $L_s H^{-1} [L_k, V] H^{-1}$ for any $1 < p < q$.

We would now like to study the behaviour of $L_s H^{-1} [L_k, V] H^{-1}$. We know that

$$[L_k, H_0] = \sum_j b_{kj} L_j - \partial_j b_{kj}.$$

Here b_{kj} are $\partial_j b_{kj}$ the operators of multiplication by b_{kj} and $\partial_j b_{kj}$.

It follows through (1.8) $|B(x)| \leq CV(x)$ and $|\nabla B(x)| \leq CV^{3/2}(x)$, for almost every $x \in \mathbb{R}^n$. Hence, $L_s H^{-1} b_{kj} L_j H^{-1}$ is L^p bounded if $L_s H^{-1} V L_j H^{-1}$ is L^p bounded. Moreover

$$L_s H^{-1} V L_j H^{-1} = (L_s H^{-1} V^{\frac{1}{2}}) (V^{\frac{1}{2}} L_j H^{-1}),$$

is L^p bounded for $1 < p < q_0$, where $q_0 = \frac{2q+n}{3n-2q}$ if $q < n$ and $2q$ if $q \geq n$. Here we have used Proposition 6.3 and Theorem 1.6.

$L_s H^{-1} \partial_j b_{kj} H^{-1}$ is L^p bounded if $L_s H^{-1} V^{3/2} H^{-1}$ is L^p bounded. We could also say

$$L_s H^{-1} V^{3/2} H^{-1} = (L_s H^{-1} V^{\frac{1}{2}}) (V H^{-1}),$$

which is bounded for $1 < p < q$ using Proposition 6.3 and Theorem 5.1. Hence, $L_s [L_k, H^{-1}]$ is L^p bounded for $1 < p < q + \epsilon$. \square

REFERENCES

- [AB] P. Auscher, B. Ben Ali, Maximal inequalities and Riesz transform estimates on L^p spaces for Schrödinger operators with non negative potentials. Annales de l'Institut Fourier. Vol 57 no 6. 2007, p 1975-2013.
- [Aus] P. Auscher, On L^p estimates for square roots of second order elliptic operators on \mathbb{R}^n . Publ. Mat. 48 (2004), no. 1, 159-186.
- [ACDH] P. Auscher, T. Coulhon, X.T. Duong & S. Hofmann, Riesz transform on manifolds and heat kernel regularity. Ann. Sci. Scuola Norm. Sup. (4) 37 (2004), no. 6, 911-957.
- [ADMc] P. Auscher, X. T. Duong, A. McIntosh, Boundedness of Banach space valued singular integral operators and applications to Hardy spaces, non publié.
- [AHS] J. Avron, I. Herbst, B. Simon, Schrödinger operator with magnetic fields; I. General interaction. Duke Math. J. 45(4)(1978), 847-883.
- [AM1] P. Auscher & J.M. Martell, Weighted norm inequalities, off-diagonal estimates and elliptic operators. Part I: General operator theory and weights. Adv. Math. 212 (2007), no. 1, 225-276.
- [BB] N. Badr, B. Ben Ali, L^p boundedness of Riesz transform related to Schrödinger operators (to appear in The Annali della Scuola Normale Superiore di Pisa, arXiv:0812.1265).
- [BBC] P. Bénéilan, H. Brézis & M. Crandall, A semilinear equation in $L^1(\mathbb{R}^N)$. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 2 (1975), no. 4.

- [Be] B. Ben Ali, Maximal inequalities and Riesz transform estimates on L^p spaces for magnetic Schrödinger operators I, submitted.
- [BS] H. Brézis & W. Strauss, Semi-linear second-order elliptic equations in L^1 . J. Math. Soc. Japan 25 (1973), 565-590.
- [Buc] S. Buckley, Pointwise multipliers for reverse Hölder spaces. II. Proc. Roy. Irish Acad. Sect. A 95 (1995), no. 2, 193-204.
- [CD] T. Coulhon & X.T. Duong, Riesz transforms for $1 \leq p \leq 2$. Trans. Amer. Math. Soc. 351 (1999), no. 3, 1151-1169.
- [Da] E.B. Davies, Non-Gaussian aspects of heat kernel behaviour, J. London math. Soc., 55 (1997), 105-125.
- [DOY] X.T. Duong, E. Ouhabaz & L. Yan, Endpoint estimates for Riesz transforms of magnetic Schrödinger operators. Ark Mat, 44 (2006), no2, 261-275.
- [DR] X.T. Duong & D. Robinson, Semigroup kernels, Poisson bounds, and holomorphic functional calculus. (English. English summary) J. Funct. Anal. 142 (1996), no. 1, 89-128.
- [G] A. Grigor'yan, Upper bounds of derivatives of the heat kernel on an arbitrary complete manifolds, J. Funct. Anal., 127 (1995), 363-389.
- [GM] T. Gallouët & J.-M. Morel, Resolution of a semilinear equation in L^1 . Proc. Roy. Soc. Edinburgh Sect. A 96 (1984), no. 3-4, 275-288. Corrigenda: Proc. Roy. Soc. Edinburgh Sect. A 99 (1985), no. 3-4, 399.
- [Gra] L. Grafakos, *Classical and Modern Fourier Analysis*, Pearson Education, New Jersey, 2004.
- [Gui1] D. Guibourg, Inégalités maximales pour l'opérateur de Schrödinger. (French. English, French summary) [Maximal inequalities for Schrödinger operators] C. R. Acad. Sci. Paris Sr. I Math. 316 (1993), no. 3, 249-252.
- [Gui2] D. Guibourg, Inégalités maximales pour l'opérateur de Schrödinger. Ths de doctorant, Université de Rennes 1, 1992.
- [H] W. Hebisch, A multiplier theorem for Schrödinger operators. Colloq. Math. 60/61 (1990), no. 2, 659-664.
- [He] B. Helffer, Semi-classical Analysis for the Schrödinger operator and applications, Lectures notes in Math, 1336(1988), Springer verlag.
- [HM1] B. Helffer, A. Mohamed: Sur le spectre essentiel des opérateurs de Schrödinger avec champ magnétique. Ann. Institut Fourier 38, 951-13 (1988)
- [HM2] B. Helffer, A. Mohamed: Semiclassical analysis for the ground state energy of a Schrödinger operator with magnetic wells. J. Funct. Anal. 138 no. 1, 40-81 (1996)
- [HNW] B. Helffer, J. Nourrigat and X. P. Wang: Sur le spectre de l'équation de Dirac (dans R^2 ou R^3) avec champ magnétique. Ann. scient. Ec. Norm. Sup. 4e serie t. 22, 515-533 (1989).
- [I] A. Iwatsuka, Magnetic Schrödinger operators with compact Resolvent, J. Math. Kyoto Univ. 26-3 (1986), 357-374.
- [N] J. Nourrigat, une inequality L^2 , non published.
- [IN] T. Iwaniec & C. Nolder, A Hardy-Littlewood inequality for quasiregular mappings in certain domains in R^n . Ann. Acad. Sci. Fenn. Ser. A I Math. 10 (1985), 267-282.
- [K1] T. Kato, Schrödinger operators with singular potentials. Proceedings of the International Symposium on Partial Differential Equations and the Geometry of Normed Linear Spaces (Jerusalem, 1972). Israel J. Math. 13 (1972), 135-148 (1973).
- [K2] T. Kato, Remarks on Schrödinger operators with vector potentials. Integral Equations and Operator Theory Vol. 1/1 1978.
- [K3] T. Kato, L^p -theory of Schrödinger operators with a singular potential. Aspects of positivity in functional analysis (Tbingen, 1985), 63-78, North-Holland Math. Stud., 122, North-Holland, Amsterdam, 1986.
- [LL] E.H. Lieb, M. Loss, M. Analysis. Second edition. AMS, Providence, RI, 2001.
- [LS] H. Leinfelder, C. G. Simadar, Schrödinger operators with singular magnetic vector potentials, Math Z. 176 (1981), 1-19.
- [MR] M. Melgaard and G. Rozenblum, Schrödinger operators with singular potentials. Stationary partial differential equations VolIII 407-517.

- [Mra] A.Mohammed , G.Raikov, On the spectral theory of the Schrödinger operator with electromagnetic potential, in: Pseudodifferential Calculus and Mathematical Physics, 298-390. Math. Topics. 5. Akademie-Verlag, Berlin, 1994.
- [O] N. Okazawa, An L^p theory for Schrödinger operators with nonnegative potentials. J. Math. Soc. Japan 36 (1984), no. 4, 675–688.
- [Ou] E. Ouhabaz, *Analysis of heat equations on domains*. London Mathematical Society Monographs Series, 31. Princeton University Press, Princeton, NJ, 2005. xiv+284 pp.
- [Sh1] Z. Shen, L^p estimates for Schrödinger operators with certain potentials, Ann. Inst. Fourier (Grenoble), 45(1995), 513-546.
- [Sh2] Z. Shen, On the eigenvalue asymptotics of Schrödinger operators, unpublished (1995).
- [Sh3] Z. Shen, Eigenvalue asymptotics and exponential decay of eigenfunctions for Schrödinger operators with magnetic fields, Trans. Amer. Math. Soc. 348(11) (1996).
- [Sh4] Z. Shen, Estimates in L^p for magnetic Schrödinger operators. Indiana Univ. Math. J 45 (1996) no3. 817-841.
- [Sh5] Z. Shen, On bounds of $N(\lambda)$ for a magnetic Schrödinger operator, Duke Math. J. 94(3) (1998), 479-507. MR 99e:35171
- [Sh6] Z. Shen, Bounds of Riesz transforms on L^p spaces for second order elliptic operators, Ann. Inst. Fourier (Grenoble) 55(2005), 173-197.
- [Sik] A. Sikora, Riesz transform, Gaussian bounds and the method of wave equation. Math. Z. 247 (2004), no. 3, 643-662.
- [Si1] B. Simon, Schrödinger semigroups. Bull. Amer. Math. Soc. (N.S.) 7 (1982), no. 3, 447-526.
- [Si2] B. Simon, Hardy and Rellich inequalities in nonintegral dimension. J. Operator Theory 9 (1983), no. 1, 143-146.
- [Si3] B. Simon: Kato's inequality and the comparison semigroups. J. Funct. Anal. 32, 97-101 (1979)
- [Si4] B.Simon, Maximal and minimal Schrödinger forms. J. Operator Theory 1 (1979), no. 1, 37-47.
- [Si5] B. Simon: Phase space analysis of simple scattering systems: extension of some work of Enss. Duke Math. J. 46, 119-168 (1979)
- [Si6] B. Simon: Schrödinger operators in the twentieth century. J. Math. Phys. 41, 3523-3555 (2000)
- [St] E. M. Stein, *Harmonic analysis: Real-variable methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.
- [Z] J. Zhong, The Sobolev estimates for some Schrödinger type operators. Math. Sci. Res. Hot-Line 3 (1999), no. 8, 1–48 (and Harmonic Analysis of some Schrödinger type Operators, PhD thesis, Princeton University, 1993.)

B. BEN ALI, UNIVERSITÉ DE PARIS-SUD, UMR DU CNRS 8628, 91405 ORSAY CEDEX, FRANCE

E-mail address: besma.ben-ali@math.u-psud.fr