

NONPROPER PRODUCTS

ANDRZEJ ROSLANOWSKI, SAHARON SHELAH, AND OTMAR SPINAS

ABSTRACT. We show that there exist two proper creature forcings having a simple (Borel) definition, whose product is not proper. We also give a new condition ensuring properness of some forcings with norms.

1. INTRODUCTION

In Roslanowski and Shelah [5] a theory of forcings built with the use of norms was developed and a number of conditions to ensure the properness of the resulting forcings was given. However it is not clear how sharp those results really are and this problem was posed in Shelah [7, Question 4.1]. In particular, he asked about the properness of the forcing notion

$$\mathbb{Q} = \{ \langle w_n : n < \omega \rangle : w_n \subseteq 2^n, w_n \neq \emptyset \text{ and } \lim_{n \rightarrow \omega} |w_n| = \infty \}$$

ordered by $\bar{w} \leq \bar{w}' \Leftrightarrow (\forall n \in \omega)(w'_n \subseteq w_n)$. In the second section we give a general criterion for collapsing the continuum to \aleph_0 and then in Corollary 2.8 we apply it to the forcing \mathbb{Q} , just showing that it is not proper.

That the property of properness is not productive, i.e. is not preserved under taking products, has been observed by Shelah long ago (see [6, XVII, 2.12]). However, his examples are somewhat artificial and certainly it would be desirable to know of some rich enough subclass of proper forcings that is productively closed. It was a natural conjecture put forth by Zapletal, that the class of definable, say analytic or Borel, proper forcings would have this property. Actually, it was only proved recently by Spinas [8] that finite powers of the Miller rational perfect set forcing and finite powers of the Laver forcing notion are proper. These are two of the most frequently used forcings in the set theory of the reals. However, in this paper we shall show that this phenomenon does not extend to all forcing notions defined in the setting of norms on possibilities. In the fourth section of the paper we give an example of a forcing notion with norms which, by the theory developed in the second section, is not proper and yet it can be decomposed as a product of two proper forcing notions of a very similar type, and both of which have a Borel definition. The properness of the factors is a consequence of a quite general theorem presented in the third section (Theorem 3.3). It occurs that a strong version of halving from [5, Section 2.2] implies the properness of forcing notions of the type $\mathbb{Q}_\infty^*(K, \Sigma)$. More on applications of halving can be found in Kellner and Shelah [3, 2] and Roslanowski and Shelah [4].

1991 *Mathematics Subject Classification.* 03E40.

The first two authors acknowledge support from the United States-Israel Binational Science Foundation (Grant no. 2006108). This is publication 941 of the second author.

The research of the third author was partially supported by DFG grant SP 683/1-2 and the Landau foundation.

Notation Most of our notation is standard and compatible with that of classical textbooks on Set Theory (like Bartoszyński and Judah [1]). However in forcing we keep the convention that *a stronger condition is the larger one*.

In this paper \mathbf{H} will stand for a function with domain ω and such that $(\forall m \in \omega)(2 \leq |\mathbf{H}(m)| < \omega)$. We also assume that $0 \in \mathbf{H}(m)$ (for all $m \in \omega$); if it is not the case then we fix an element of $\mathbf{H}(m)$ and we use it whenever appropriate notions refer to 0.

Creature background: Since our results are stated for creating pairs with several special properties, below we present a somewhat restricted context of the creature forcing, introducing *good creating pairs*.

Definition 1.1. (1) A *creature for \mathbf{H}* is a triple

$$t = (\mathbf{nor}, \mathbf{val}, \mathbf{dis}) = (\mathbf{nor}[t], \mathbf{val}[t], \mathbf{dis}[t])$$

such that $\mathbf{nor} \in \mathbb{R}^{\geq 0}$, $\mathbf{dis} \in \mathcal{H}(\omega_1)$, and for some integers $m_{\text{dn}}^t < m_{\text{up}}^t < \omega$

$$\emptyset \neq \mathbf{val} \subseteq \{ \langle u, v \rangle \in \prod_{i < m_{\text{dn}}^t} \mathbf{H}(i) \times \prod_{i < m_{\text{up}}^t} \mathbf{H}(i) : u \triangleleft v \}.$$

The family of all creatures for \mathbf{H} is denoted by $\text{CR}[\mathbf{H}]$.

(2) Let $K \subseteq \text{CR}[\mathbf{H}]$ and $\Sigma : K \rightarrow \mathcal{P}(K)$. We say that (K, Σ) is a *good creating pair* for \mathbf{H} whenever the following conditions are satisfied for each $t \in K$.

(a) [Fullness] $\text{dom}(\mathbf{val}[t]) = \prod_{i < m_{\text{dn}}^t} \mathbf{H}(i)$.

(b) $t \in \Sigma(t)$ and if $s \in \Sigma(t)$, then $\mathbf{val}[s] \subseteq \mathbf{val}[t]$ and so also $m_{\text{dn}}^s = m_{\text{dn}}^t$ and $m_{\text{up}}^s = m_{\text{up}}^t$.

(c) [Transitivity] If $s \in \Sigma(t)$, then $\Sigma(s) \subseteq \Sigma(t)$.

(3) A good creating pair (K, Σ) is

- *local* if $m_{\text{up}}^t = m_{\text{dn}}^t + 1$ for all $t \in K$,

- *forgetful* if for every $t \in K$, $v \in \prod_{i < m_{\text{up}}^t} \mathbf{H}(i)$, and $u \in \prod_{i < m_{\text{dn}}^t} \mathbf{H}(i)$ we

have

$$\langle v \upharpoonright m_{\text{dn}}^t, v \rangle \in \mathbf{val}[t] \quad \Rightarrow \quad \langle u, u \widehat{\vee} v \upharpoonright [m_{\text{dn}}^t, m_{\text{up}}^t] \rangle \in \mathbf{val}[t],$$

- *strongly finitary* if for each $i < \omega$ we have

$$|\mathbf{H}(i)| < \omega \quad \text{and} \quad |\{t \in K : m_{\text{dn}}^t = i\}| < \omega.$$

(4) If $t_0, \dots, t_n \in K$ are such that $m_{\text{up}}^{t_i} = m_{\text{dn}}^{t_{i+1}}$ (for $i < n$) and $w \in \prod_{i < m_{\text{dn}}^{t_0}} \mathbf{H}(i)$,

then we let

$$\text{pos}(w, t_0, \dots, t_n) \stackrel{\text{def}}{=} \{v \in \prod_{j < m_{\text{up}}^{t_n}} \mathbf{H}(j) : w \triangleleft v \ \& \ (\forall i \leq n) (\langle v \upharpoonright m_{\text{dn}}^{t_i}, v \upharpoonright m_{\text{up}}^{t_i} \rangle \in \mathbf{val}[t_i])\}.$$

If K is forgetful and $t \in K$, then we also define

$$\text{pos}(t) = \{v \upharpoonright [m_{\text{dn}}^t, m_{\text{up}}^t] : \langle v \upharpoonright m_{\text{dn}}^t, v \rangle \in \mathbf{val}[t]\}.$$

Note that if K is forgetful, then to describe a creature in K it is enough to give $\text{pos}(t)$, $\mathbf{nor}[t]$ and $\mathbf{dis}[t]$. This is how our examples will be presented (as they all will be forgetful). Also, if K is additionally local, then we may write $\text{pos}(t) = A$ for some $A \subseteq \mathbf{H}(m_{\text{dn}}^t)$ with a natural interpretation of this abuse of notation.

If w, t_0, \dots, t_n are as in 1.1(4) and $s_i \in \Sigma(t_i)$ for $i \leq n$, and $u \in \text{pos}(w, s_0, \dots, s_k)$, $k < n$, then $\text{pos}(u, s_k, \dots, s_n) \subseteq \text{pos}(w, t_0, \dots, t_n)$ (remember 1.1(2b)).

Definition 1.2. Let (K, Σ) be a good creating pair for \mathbf{H} . We define a forcing notion $\mathbb{Q}_\infty^*(K, \Sigma)$ as follows.

A condition in $\mathbb{Q}_\infty^*(K, \Sigma)$ is a sequence $p = (w^p, t_0^p, t_1^p, t_2^p, \dots)$ such that

- (a) $t_i^p \in K$ and $m_{\text{up}}^{t_i^p} = m_{\text{dn}}^{t_{i+1}^p}$ (for $i < \omega$), and
- (b) $w \in \prod_{i < m_{\text{dn}}^{t_0^p}} \mathbf{H}(i)$ and $\lim_{n \rightarrow \infty} \text{nor}[t_n^p] = \infty$.

The relation \leq on $\mathbb{Q}_\infty^*(K, \Sigma)$ is given by: $p \leq q$ if and only if for some $i < \omega$ we have $w^q \in \text{pos}(w^p, t_0^p, \dots, t_{i-1}^p)$ (if $i = 0$ this means $w^q = w^p$) and $t_n^q \in \Sigma(t_{n+i}^p)$ for all $n < \omega$.

For a condition $p \in \mathbb{Q}_\infty^*(K, \Sigma)$ we let $i(p) = \text{lh}(w^p)$.

2. COLLAPSING CREATURES

We will show here that very natural forcing notions of type $\mathbb{Q}_\infty^*(K, \Sigma)$ (for a big local and finitary creating pair (K, Σ)) collapse \mathfrak{c} to \aleph_0 , in particular answering [7, Question 4.1]. The main ingredient of the proof is similar to the “negative theory” presented in [5, Section 1.4], and Definition 2.1 below should be compared with [5, Definition 1.4.4] (but the two properties are somewhat incomparable).

Definition 2.1. Let $h : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ be a non-decreasing unbounded function and let (K, Σ) be a good creating pair for \mathbf{H} . We say that (K, Σ) is *sufficiently h -bad* if there are sequences $\bar{m} = \langle m_i : i < \omega \rangle$, $\bar{A} = \langle A_i : i < \omega \rangle$ and $\bar{F} = \langle F_i : i < \omega \rangle$ such that

- (α) \bar{m} is a strictly increasing sequence of integers, $m_0 = 0$, and

$$(\forall t \in K)(\exists i < \omega)(m_{\text{dn}}^t = m_i \ \& \ m_{\text{up}}^t = m_{i+1}),$$

- (β) A_i are finite non-empty sets,

- (γ) $F_i = (F_i^0, F_i^1) : A_i \times \prod_{m < m_{i+1}} \mathbf{H}(m) \rightarrow A_{i+1} \times 2$,

- (δ) if $i < \omega$, $t \in K$, $m_{\text{dn}}^t = m_i$ and $\text{nor}[t] > 4$, then there is $a \in A_i$ such that for every $x \in A_{i+1} \times 2$, for some $s_x \in \Sigma(t)$ we have

$$\begin{aligned} \text{nor}[s_x] &\geq \min\{h(\text{nor}[t]), h(i)\} \text{ and} \\ (\forall u \in \prod_{m < m_i} \mathbf{H}(m))(\forall v \in \text{pos}(u, s_x))(F_i(a, v) = x). \end{aligned}$$

Proposition 2.2. *Suppose that $h : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ is a non-decreasing unbounded function, and (K, Σ) is a strongly finitary good creating pair for \mathbf{H} . Assume also that (K, Σ) is sufficiently h -bad. Then the forcing notion $\mathbb{Q}_\infty^*(K, \Sigma)$ collapses \mathfrak{c} onto \aleph_0 .*

Proof. The proof is similar to that of [5, Proposition 1.4.5], but for reader’s convenience we present it fully.

Let \bar{m}, \bar{A} and \bar{F} witness that (K, Σ) is sufficiently h -bad. For $i < \omega$ and $a \in A_i$ we define $\mathbb{Q}_\infty^*(K, \Sigma)$ -names $\dot{\rho}_{i,a}$ (for a real in 2^ω) and $\dot{\eta}_{i,a}$ (for an element of $\prod_{j \geq i} A_j$)

as follows:

$$\Vdash_{\mathbb{Q}_\infty^*(K, \Sigma)} \text{ “ } \dot{\eta}_{i,a}(i) = a \text{ and } \dot{\eta}_{i,a}(j) = F_{j-1}^0(\dot{\eta}_{i,a}(j-1), \dot{W} \upharpoonright m_j) \text{ for } j > i \text{ ” ,}$$

and

$$\Vdash_{\mathbb{Q}_\infty^*(K, \Sigma)} \text{“ } \dot{\rho}_{i,a} \upharpoonright i \equiv 0 \text{ and } \dot{\rho}_{i,a}(j) = F_j^1(\dot{\eta}_{i,a}(j), \dot{W} \upharpoonright m_{j+1}) \text{ for } j \geq i \text{”}.$$

Above, \dot{W} is the canonical name for the generic function in $\prod_{i < \omega} \mathbf{H}(i)$, i.e., $p \Vdash_{\mathbb{Q}_\infty^*(K, \Sigma)}$

“ $w^p \triangleleft \dot{W} \in \prod_{i < \omega} \mathbf{H}(i)$ ”. We are going to show that

$$\Vdash_{\mathbb{Q}_\infty^*(K, \Sigma)} \text{“ } (\forall r \in 2^\omega \cap \mathbf{V})(\exists i < \omega)(\exists a \in A_i)(\forall j \geq i)(\dot{\rho}_{i,a}(j) = r(j)) \text{”}.$$

To this end suppose that $p \in \mathbb{Q}_\infty^*(K, \Sigma)$ and $r \in 2^\omega$. Passing to a stronger condition if needed, we may assume that $(\forall j < \omega)(\mathbf{nor}[t_j^p] > 4)$. Let $i < \omega$ be such that $\text{lh}(w^p) = m_i$; then also $m_{\text{dn}}^{t_j^p} = m_{i+j}$ for $j < \omega$ (remember 2.1(α)).

Fix $k < \omega$ for a moment. By downward induction on $j \leq k$ choose $s_j^k \in \Sigma(t_j^p)$ and $a_j^k \in A_{i+j}$ such that

- (a) $\mathbf{nor}[s_j^k] \geq \min\{h(\mathbf{nor}[t_j^p]), h(i+j)\}$ for all $j \leq k$,
- (b) $(\forall u \in \prod_{m < m_{i+k}} \mathbf{H}(m))(\forall v \in \text{pos}(u, s_k^k))(F_{i+k}^1(a_k^k, v) = r(i+k))$,
- (c) for $j < k$:

$$(\forall u \in \prod_{m < m_{i+j}} \mathbf{H}(m))(\forall v \in \text{pos}(u, s_j^k))(F_{i+j}^1(a_j^k, v) = r(i+j) \ \& \ F_{i+j}^0(a_j^k, v) = a_{j+1}^k).$$

(Plainly it is possible by 2.1(δ)).

Since for each $j < \omega$ both $\Sigma(t_j^p)$ and A_{i+j} are finite, we may use König's Lemma to pick an increasing sequence $\bar{k} = \langle k(\ell) : \ell < \omega \rangle$ such that

$$a_j^{k(\ell+1)} = a_j^{k(\ell')} \quad \text{and} \quad s_j^{k(\ell+1)} = s_j^{k(\ell')}$$

for $\ell < \ell' < \omega$ and $j \leq k(\ell)$. Put $w^q = w^p$ and $t_j^q = s_j^{k(j+1)}$, $b_j = a_j^{k(j+1)}$ for $j < \omega$. Easily, $q = (w^q, t_0^q, t_1^q, t_2^q, \dots)$ is a condition in $\mathbb{Q}_\infty^*(K, \Sigma)$ stronger than p . Also, by clause (c) of the choice of s_j^k , we clearly have

$$(\forall j < \omega)(\forall v \in \text{pos}(w^q, t_0^q, \dots, t_j^q))(F_{i+j}^0(b_j, v) = b_{j+1} \ \& \ F_{i+j}^1(b_j, v) = r(i+j)).$$

Hence

$$q \Vdash_{\mathbb{Q}_\infty^*(K, \Sigma)} \text{“ } (\forall j < \omega)(\dot{\eta}_{i,b_0}(i+j) = b_j \ \& \ \dot{\rho}_{i,b_0}(i+j) = r(i+j)) \text{”},$$

finishing the proof. \square

Lemma 2.3. *Suppose that positive integers N, M, d satisfy $(N-2) \cdot 2^M < d$. Let A, B be finite sets such that $|A| \geq 2^M$ and $|B| \leq N$. Then there is a mapping $\hat{F} : A \times {}^d M \rightarrow B$ with the property that:*

- (\otimes) if $2 \leq \ell \leq M$, $\langle c_i : i < d \rangle \in \prod_{i < d} [M]^\ell$, then there is $a \in A$ such that for every $b \in B$, for some $c_i^b \in [c_i]^{[\ell/2]}$ (for $i < d$) we have

$$(\forall u \in \prod_{i < d} c_i^b)(\hat{F}(a, u) = b).$$

Proof. Plainly we may assume that $|A| = 2^M$ and $|B| = N \geq 2$, and then we may pretend that $A = {}^M 2$ and $B = N$.

For $h \in A = M2$ and $u \in {}^dM$ we let $\hat{F}(h, u) < N$ be such that

$$\hat{F}(h, u) \equiv \sum_{i < d} h(u(i)) \pmod{N}.$$

This defines the function $\hat{F} : A \times {}^dM \rightarrow B = N$, and we are going to show that it has the property stated in (\otimes) . To this end suppose that $2 \leq \ell \leq M$ and $\langle c_i : i < d \rangle \in \prod_{i < d} [M]^\ell$. For each $i < d$ we may choose $h_i \in A$ so that

$$|(h_i)^{-1}[\{0\}] \cap c_i| \geq \lfloor \ell/2 \rfloor \quad \text{and} \quad |(h_i)^{-1}[\{1\}] \cap c_i| \geq \lfloor \ell/2 \rfloor$$

Then, for some $h \in A$ and $I \subseteq d$ we have $|I| \geq d/2^M$ and $h_i = h$ for $i \in I$. For $i \in d \setminus I$ we may pick $c_i^* \in [c_i]^{\lfloor \ell/2 \rfloor}$ and $j_i < 2$ such that $h \upharpoonright c_i^* \equiv j_i$.

Now, suppose $b \in B$. Take a set $J \subseteq I$ such that

$$|J| + \sum_{i \in d \setminus I} j_i \equiv b \pmod{N}$$

(possible as $|I| \geq d/2^M > N - 2$, so $|I| \geq N - 1$). By our choices, we may pick $c_i^b \in [c_i]^{\lfloor \ell/2 \rfloor}$ (for $i \in I$) such that

$$\begin{aligned} &\text{if } i \in J, \text{ then } h \upharpoonright c_i^b \equiv 1, \\ &\text{if } i \in I \setminus J, \text{ then } h \upharpoonright c_i^b \equiv 0. \end{aligned}$$

For $i \in d \setminus I$ we let $c_i^b = c_i^*$ (selected earlier). It should be clear that then

$$(\forall u \in \prod_{i < d} c_i^b)(\hat{F}(h, u) = b),$$

as needed. \square

Example 2.4. Let $\bar{m} = \langle m_i : i < \omega \rangle$ be an increasing sequence of integers such that $m_0 = 0$ and $m_{i+1} - m_i > 4^{i+3}$. Let $h(\ell) = \lfloor \ell/2 \rfloor$ for $\ell < \omega$.

For $j < \omega$ we let $\mathbf{H}_{\bar{m}}^0(j) = i + 2$, where i is such that $m_i \leq j < m_{i+1}$. Let $K_{\bar{m}}^0$ consist of all (forgetful) creatures $t \in \text{CR}[\mathbf{H}_{\bar{m}}^0]$ such that

- $\text{dis}[t] = \langle i^t, \langle Z_j^t : m_{i^t} \leq j < m_{i^t+1} \rangle \rangle$ for some $i^t < \omega$ and $\emptyset \neq Z_j^t \subseteq \mathbf{H}_{\bar{m}}^0(j)$ (for $m_{i^t} \leq j < m_{i^t+1}$),
- $\text{nor}[t] = \min\{|Z_j^t| : m_{i^t} \leq j < m_{i^t+1}\}$,
- $\text{pos}(t) = \prod_{j \in [m_{i^t}, m_{i^t+1})} Z_j^t$.

Finally, for $t \in K_{\bar{m}}^0$ we let

$$\Sigma_{\bar{m}}^0(t) = \{s \in K_{\bar{m}}^0 : i^t = i^s \ \& \ (\forall j \in [m_{i^t}, m_{i^t+1})) (Z_j^s \subseteq Z_j^t)\}.$$

Then $(K_{\bar{m}}^0, \Sigma_{\bar{m}}^0)$ is a strongly finitary and sufficiently h -bad good creating pair for $\mathbf{H}_{\bar{m}}^0$. Consequently, the forcing notion $\mathbb{Q}_\infty^*(K_{\bar{m}}^0, \Sigma_{\bar{m}}^0)$ collapses \mathfrak{c} onto \aleph_0 .

Proof. It should be clear that $(K_{\bar{m}}^0, \Sigma_{\bar{m}}^0)$ is a strongly finitary good creating pair for $\mathbf{H}_{\bar{m}}^0$. To show that it is sufficiently h -bad let $A_i = i+2 \cdot 2$, $B_i = A_{i+1} \times 2 = i+3 \cdot 2 \times 2$ and $M_i = i+2$. Since $|B_i| \cdot 2^{M_i} = 2^{i+4+i+2} < m_{i+1} - m_i \stackrel{\text{def}}{=} d_i$, we may apply Lemma 2.3 for $A = A_i$, $B = B_i$, $M = M_i$ and $d = d_i$ to get functions $\hat{F}_i : A_i \times {}^{d_i}M_i \rightarrow B_i$ with the property (\otimes) (for those parameters). For $a \in A_i$ and $v \in \prod_{j < m_{i+1}} \mathbf{H}_{\bar{m}}^0(j)$

we interpret $F_i(a, v)$ as $\hat{F}_i(a, u)$ where $u \in {}^{d_i}(i+1)$ is given by $u(j) = v(m_i + j)$ for

$j < d_i$. It is straightforward to show that \bar{m} , $\bar{A} = \langle A_i : i < \omega \rangle$ and $\bar{F} = \langle F_i : i < \omega \rangle$ witness that $(K_{\bar{m}}^0, \Sigma_{\bar{m}}^0)$ is h -bad. \square

The above example (together with Proposition 2.2) easily gives the answer to [7, Question 4.1]. To show how our problem reduces to this example, let us recall the following.

Definition 2.5 (See [5, Definition 4.2.1]). Suppose $0 < m < \omega$ and for $i < m$ we have $t_i \in \text{CR}[\mathbf{H}]$ such that $m_{\text{up}}^{t_i} \leq m_{\text{dn}}^{t_{i+1}}$. Then we define *the sum of the creatures* t_i as a creature $t = \Sigma^{\text{sum}}(t_i : i < m)$ such that (if well defined then):

- (a) $m_{\text{dn}}^t = m_{\text{dn}}^{t_0}$, $m_{\text{up}}^t = m_{\text{up}}^{t_{m-1}}$,
- (b) $\mathbf{val}[t]$ is the set of all pairs $\langle h_1, h_2 \rangle$ such that:
 - $\text{lh}(h_1) = m_{\text{dn}}^t$, $\text{lh}(h_2) = m_{\text{up}}^t$, $h_1 \triangleleft h_2$,
 - and $\langle h_2 \upharpoonright m_{\text{dn}}^{t_i}, h_2 \upharpoonright m_{\text{up}}^{t_i} \rangle \in \mathbf{val}[t_i]$ for $i < m$,
 - and $h_2 \upharpoonright [m_{\text{up}}^{t_i}, m_{\text{dn}}^{t_{i+1}}]$ is identically zero for $i < m - 1$,
- (c) $\mathbf{nor}[t] = \min\{\mathbf{nor}[t_i] : i < m\}$,
- (d) $\mathbf{dis}[t] = \langle t_i : i < m \rangle$.

If for all $i < m - 1$ we have $m_{\text{up}}^{t_i} = m_{\text{dn}}^{t_{i+1}}$, then we call the sum *tight*.

Definition 2.6. Let (K, Σ) be a local good creating pair for \mathbf{H} , and let $\bar{m} = \langle m_i : i < \omega \rangle$ be a strictly increasing sequence with $m_0 = 0$. We define the \bar{m} -*summarization* $(K^{\bar{m}}, \Sigma^{\bar{m}}, \mathbf{H}^{\bar{m}})$ of (K, Σ, \mathbf{H}) as follows:

- $\mathbf{H}^{\bar{m}}(i) = \prod_{m=m_i}^{m_{i+1}-1} \mathbf{H}(m)$,
- $K^{\bar{m}}$ consists of all tight sums $\Sigma^{\text{sum}}(t_\ell : m_i \leq \ell < m_{i+1})$ such that $i < \omega$, $t_\ell \in K$, $m_{\text{dn}}^{t_\ell} = \ell$,
- if $t = \Sigma^{\text{sum}}(t_\ell : m_i \leq \ell < m_{i+1}) \in K^{\bar{m}}$, then $\Sigma^{\bar{m}}(t)$ consists of all creatures $s \in K^{\bar{m}}$ such that $s = \Sigma^{\text{sum}}(s_\ell : m_i \leq \ell < m_{i+1})$ for some $s_\ell \in \Sigma(t_\ell)$ (for $\ell = m_i, \dots, m_{i+1} - 1$).

Proposition 2.7. Assume that (K, Σ) is a local good creating pair for \mathbf{H} , $\bar{m} = \langle m_i : i < \omega \rangle$ is a strictly increasing sequence with $m_0 = 0$. Then:

- (1) $(K^{\bar{m}}, \Sigma^{\bar{m}})$ is a good creating pair for $\mathbf{H}^{\bar{m}}$;
- (2) the forcing notion $\mathbb{Q}_\infty^*(K^{\bar{m}}, \Sigma^{\bar{m}})$ can be embedded as a dense subset of the forcing notion $\mathbb{Q}_\infty^*(K, \Sigma)$ (so the two forcing notions are equivalent).

Corollary 2.8. Let $\mathbf{H} : \omega \rightarrow \omega$ be increasing, $\mathbf{H}(0) \geq 2$, and let $g : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ be an unbounded non-decreasing function. We define $(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ as follows: $K_g^{\mathbf{H}}$ consists of all creatures $t \in \text{CR}[\mathbf{H}]$ such that

- $\mathbf{dis}[t] = \langle i^t, A^t \rangle$ for some $i^t < \omega$ and $\emptyset \neq A^t \subseteq \mathbf{H}(i^t)$,
- $\mathbf{nor}[t] = g(|A^t|)$, $m_{\text{dn}}^t = i^t$, $m_{\text{up}}^t = i^t + 1$ and $\text{pos}(t) = A^t$.

For $t \in K_g^{\mathbf{H}}$ we let

$$\Sigma_g^{\mathbf{H}}(t) = \{s \in K_g^{\mathbf{H}} : i^t = i^s \ \& \ A^s \subseteq A^t\}.$$

Then $(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ is a local strongly finitary good creating pair for \mathbf{H} . The forcing notion $\mathbb{Q}_\infty^*(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ collapses \mathfrak{c} onto \aleph_0 . In particular, the forcing notion \mathbb{Q} defined in the Introduction is not proper.

Proof. Let $p \in \mathbb{Q}_\infty^*(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$. Plainly, $\lim_{i \rightarrow \infty} |A^{t_i^p}| = \infty$, so we may find a condition $q \geq p$ and an increasing sequence $\bar{m} = \langle m_i : i < \omega \rangle$ such that

- $m_0 = 0$, $m_1 = \text{lh}(w^q)$, $m_{i+1} - m_i > 4^{i+3}$,
- if $m_i \leq m_{\text{dn}}^{t_k^q} < m_{i+1}$, then $|A^{t_k^q}| = i + 2$.

Now we define a condition q^* in $\mathbb{Q}_\infty^*((K_g^{\mathbf{H}})^{\bar{m}}, (\Sigma_g^{\mathbf{H}})^{\bar{m}})$ by

$$w^{q^*} = w^q, \quad t_i^{q^*} = \Sigma^{\text{sum}}(t_k^q : m_{i+1} \leq k < m_{i+2}) \quad (\text{for } i < \omega).$$

The forcing notion $\mathbb{Q}_\infty^*(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ above the condition q is equivalent to the forcing notion $\mathbb{Q}_\infty^*((K_g^{\mathbf{H}})^{\bar{m}}, (\Sigma_g^{\mathbf{H}})^{\bar{m}})$ above q^* . Plainly, $\mathbb{Q}_\infty^*((K_g^{\mathbf{H}})^{\bar{m}}, (\Sigma_g^{\mathbf{H}})^{\bar{m}})$ above q^* is isomorphic to $\mathbb{Q}_\infty^*(K_{\bar{m}}^0, \Sigma_{\bar{m}}^0)$ of Example 2.4 above the minimal condition r with $w^r = w^{q^*}$. The assertion follows now by the last sentence of 2.4. \square

Remark 2.9. (1) If, e.g., $g(x) = \log_2(x)$ then the creating pair $(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ is big (see [5, Definition 2.2.1]), and we may even get “a lot of bigness”. Thus the bigness itself is not enough to guarantee properness of the resulting forcing notion.

- (2) Forcing notions of the form $\mathbb{Q}_\infty^*(K, \Sigma)$ are special cases of $\mathbb{Q}_f^*(K, \Sigma)$ (see [5, Definition 1.1.10 and Section 2.2]). However if the function f is growing very fast (much faster than \mathbf{H}) then our method does not apply. Let us recall that if (K, Σ) is simple, finitary, big and has the Halving Property, and $f : \omega \times \omega \rightarrow \omega$ is \mathbf{H} -fast (see [5, Definition 1.1.12]), then $\mathbb{Q}_f^*(K, \Sigma)$ is proper. Thus one may wonder if we may omit halving - can the forcing notion $\mathbb{Q}_f^*(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ be proper for \mathbf{H} and f suitably “fast”?

3. PROPERNESS FROM HALVING

It was shown in [5, Theorem 2.2.11] that halving and bigness (see [5, Definitions 2.2.1, 2.2.7]) imply properness of the forcings $\mathbb{Q}_f^*(K, \Sigma)$ (for fast f). It occurs that if we have a stronger version of halving, then we may get the properness of $\mathbb{Q}_\infty^*(K, \Sigma)$ even without any bigness assumptions.

Definition 3.1. Let (K, Σ) be a forgetful good creating pair

- (1) Let $t \in K$ and $\varepsilon > 0$. We say that a creature $t^* \in \Sigma(t)$ is an ε -**half** of t if the following hold:
- (i) $\mathbf{nor}[t^*] \geq \mathbf{nor}[t] - \varepsilon$, and
 - (ii) if $s \in \Sigma(t^*)$ and $\mathbf{nor}[s] > 1$, then we can find $t_0 \in \Sigma(t)$ such that

$$\mathbf{nor}[t_0] \geq \mathbf{nor}[t] - \varepsilon \quad \text{and} \quad \text{pos}(t_0) \subseteq \text{pos}(s).$$

- (2) Let $\bar{\varepsilon} = \langle \varepsilon_i : i < \omega \rangle$ be a sequence of positive real numbers and $\bar{m} = \langle m_i : i < \omega \rangle$ be a strictly increasing sequence of integers with $m_0 = 0$. We say that the pair (K, Σ) has the $(\bar{\varepsilon}, \bar{m})$ -*halving property* if for every $t \in K$ with $m_i \leq m_{\text{dn}}^t$ and $\mathbf{nor}[t] \geq 2$ there exists an ε_i -**half** of t in $\Sigma(t)$.

Definition 3.2. Let (K, Σ) be a good creating pair. Suppose that $p \in \mathbb{Q}_\infty^*(K, \Sigma)$ and $I \subseteq \mathbb{Q}_\infty^*(K, \Sigma)$ is open dense. We say that p *essentially belongs to* I , written $p \in^* I$, if there exists $i_* < \omega$ such that for every $v \in \text{pos}(w^p, t_0^p, \dots, t_{i_*-1}^p)$ we have $(v, t_{i_*}^p, t_{i_*+1}^p, t_{i_*+2}^p, \dots) \in I$

Note that if $I \subseteq \mathbb{Q}_\infty^*(K, \Sigma)$ is open dense, $p \in^* I$ and $p \leq q$, then also $q \in^* I$.

Theorem 3.3. *Let $\bar{\varepsilon} = \langle \varepsilon_i : i < \omega \rangle$ be a decreasing sequence of positive numbers and $\bar{m} = \langle m_i : i < \omega \rangle$ be a strictly increasing sequence of integers with $m_0 = 0$. Assume that for each $i < \omega$*

$$|\prod_{n < m_i} \mathbf{H}(n)| \leq 1/\varepsilon_i.$$

Let (K, Σ) be a good creating pair for \mathbf{H} and suppose that (K, Σ) is local, forgetful and has the $(\bar{\varepsilon}, \bar{m})$ -halving property. Then the forcing notion $\mathbb{Q}_\infty^(K, \Sigma)$ is proper.*

Proof. We start with two technical claims.

Claim 3.3.1. *Let $a \geq 2$ and $I \subseteq \mathbb{Q}_\infty^*(K, \Sigma)$ be open dense. Furthermore suppose that $p \in \mathbb{Q}_\infty^*(K, \Sigma)$ and $i < \omega$ is such that $i(p) \leq m_i$ and $\mathbf{nor}[t_n^p] > a$ for every $n \geq m_i - i(p)$. Finally let $v \in \prod_{n < m_i} \mathbf{H}(n)$. Then there exists $q \in \mathbb{Q}_\infty^*(K, \Sigma)$ such that*

- (a) $p \leq q$, $w^p = w^q$ and $t_n^p = t_n^q$ for every $n < m_i - i(p)$;
- (b) $\mathbf{nor}[t_n^q] \geq a - \varepsilon_i$ for every $n \geq m_i - i(p)$;
- (c) either, letting $q^{[v]} = (v, t_{m_i - i(p)}^q, t_{m_i - i(p) + 1}^q, t_{m_i - i(p) + 2}^q, \dots)$, $q^{[v]} \in^* I$ or else there is no $r \geq q^{[v]}$ such that $r \in I$, $w^r = v$ and $\mathbf{nor}[t_n^r] > 1$ for every n .

Proof of the Claim. We know that (K, Σ) has the $(\bar{\varepsilon}, \bar{m})$ -halving property and therefore for each $n \geq m_i - i(p)$ we may choose an ε_i -half $t_n^{q_0} \in \Sigma(t_n^p)$ of t_n^p . For $n < m_i - i(p)$ put $t_n^{q_0} = t_n^p$ and let $w^{q_0} = w^p$. This defines a condition $q_0 = (w^{q_0}, t_0^{q_0}, t_1^{q_0}, t_2^{q_0}, \dots) \in \mathbb{Q}_\infty^*(K, \Sigma)$. Plainly, (a) and (b) hold for q_0 instead of q . Now if there is no $r \geq q_0^{[v]}$ with $r \in I$, $w^r = v$ and $\mathbf{nor}[t_n^r] > 1$ for every $n < \omega$, we can let $q = q_0$. Hence we may assume that such $r = (w^r, t_0^r, t_1^r, t_2^r, \dots)$ does exist.

Pick $j < \omega$ large enough such that $\mathbf{nor}[t_n^r] \geq a - \varepsilon_i$ for every $n \geq j$. Now we define $q \in \mathbb{Q}_\infty^*(K, \Sigma)$:

- $w^q = w^p$, $t_n^q = t_n^p$ for $n < m_i - i(p)$,
- $t_n^q = t_{n - m_i + i(p)}^r$ for $n \geq m_i - i(p) + j$,
- for $m_i - i(p) \leq n < m_i - i(p) + j$ let $t_n^q \in \Sigma(t_n^p)$ be such that

$$\mathbf{nor}[t_n^q] \geq \mathbf{nor}[t_n^p] - \varepsilon_i \geq a - \varepsilon_i \quad \text{and} \quad \text{pos}(t_n^q) \subseteq \text{pos}(t_{n - m_i + i(p)}^r)$$

(exists by the halving property).

Clearly $p \leq q$ and (a), (b) hold. Also, for every $u \in \text{pos}(v, t_{m_i - i(p)}^q, \dots, t_{m_i - i(p) + j}^q)$ we have $q^{[u]} \geq r$, and hence $q^{[u]} \in I$, as I is open. Consequently, $q^{[v]} \in^* I$. \square

Claim 3.3.2. *Let $a \geq 3$ and $I \subseteq \mathbb{Q}_\infty^*(K, \Sigma)$ be open dense. Suppose that $p \in \mathbb{Q}_\infty^*(K, \Sigma)$ and $i < \omega$ is such that $i(p) \leq m_i$ and $\mathbf{nor}[t_n^p] > a$ for every $n \geq m_i - i(p)$. Then there exists $q \in \mathbb{Q}_\infty^*(K, \Sigma)$ such that*

- (a) $p \leq q$, $w^p = w^q$, and $t_n^p = t_n^q$ for $n < m_i - i(p)$;
- (b) $\mathbf{nor}[t_n^q] \geq a - 1$ for every $n \geq m_i - i(p)$;
- (c) for every $v \in \prod_{n < m_i} \mathbf{H}(n)$, either $q^{[v]} \in^* I$, or else there is no $r \in I$ such that $r \geq q^{[v]}$, $w^r = v$ and $\mathbf{nor}[t_n^r] > 1$ for all n .

Proof of the Claim. Let $\langle v_l : l < k \rangle$ enumerate $\prod_{n < m_i} \mathbf{H}(n)$, thus $k \leq 1/\varepsilon_i$. Applying Claim 3.3.1 k times, it is straightforward to construct a sequence $\langle q_l : l \leq k \rangle \subseteq \mathbb{Q}_\infty^*(K, \Sigma)$ such that

- $q_0 = p$, $q_l \leq q_{l+1}$, $w^{q_l} = w^p$, and $t_n^{q_l} = t_n^p$ for every $n < m_i - i(p)$,
- $\mathbf{nor}[t_n^{q_l}] \geq a - l \cdot \varepsilon_i$ for every $n \geq m_i - i(p)$, and
- $\langle q_l, q_{l+1}, v_l, a - l \cdot \varepsilon_i \rangle$ are like $\langle p, q, v, a \rangle$ in Claim 3.3.1.

Then clearly $q = q_k$ is as desired. \square

We will argue now that the forcing notion $\mathbb{Q}_\infty^*(K, \Sigma)$ is proper. So suppose that N is a countable elementary submodel of $(\mathcal{H}(\chi), \in)$ (for some sufficiently large regular cardinal χ), $K, \Sigma, \dots \in N$. Let $p \in N \cap \mathbb{Q}_\infty^*(K, \Sigma)$ and let $\langle I_\ell : \ell < \omega \rangle$ list with ω -repetitions all open dense subsets of $\mathbb{Q}_\infty^*(K, \Sigma)$ from N .

By induction on $\ell < \omega$ we choose integers i_ℓ and conditions $p_\ell \in N \cap \mathbb{Q}_\infty^*(K, \Sigma)$ as follows. We set $p_0 = p$ and $i_0 > i(p)$ is such that $\mathbf{nor}[t_n^{p_0}] > 3$ for all $n \geq m_{i_0} - i(p)$. Now, assume we have defined $p_\ell \in N \cap \mathbb{Q}_\infty^*(K, \Sigma)$ and $i_\ell < \omega$ so that $w^p = w^{p_\ell}$ and $\mathbf{nor}[t_n^{p_\ell}] > 3 + \ell$ for every $n \geq m_{i_\ell} - i(p)$. Applying Claim 3.3.2 (inside N) to $3 + \ell, I_\ell, p_\ell, i_\ell$ here standing for a, I, p, i there we may find a condition $p_{\ell+1} \in N \cap \mathbb{Q}_\infty^*(K, \Sigma)$ such that

- (a) $_\ell$ $p_\ell \leq p_{\ell+1}$, $w^p = w^{p_\ell} = w^{p_{\ell+1}}$, and $t_n^{p_\ell} = t_n^{p_{\ell+1}}$ for all $n < m_{i_\ell} - i(p)$;
- (b) $_\ell$ $\mathbf{nor}[t_n^{p_{\ell+1}}] \geq 2 + \ell$ for every $n \geq m_{i_\ell} - i(p)$;
- (c) $_\ell$ for every sequence $v \in \prod_{n < m_{i_\ell}} \mathbf{H}(n)$, if there exists $r \in I_\ell$ such that $r \geq p_{\ell+1}^{[v]}$,

$$w^r = v \text{ and } \mathbf{nor}[t_n^r] > 1 \text{ for every } n, \text{ then } p_{\ell+1}^{[v]} \in {}^* I_\ell.$$

Then we choose $i_{\ell+1} > i_\ell$ so that $\mathbf{nor}[t_n^{p_{\ell+1}}] > 3 + \ell + 1$ for all $n \geq m_{i_{\ell+1}} - i(p)$.

After the inductive construction is carried out we let q be the natural fusion determined by the p_ℓ (so $w^q = w^p$ and $t_n^q = t_n^{p_\ell}$ whenever $n < m_{i_\ell} - i(q)$). Plainly, $q \in \mathbb{Q}_\infty^*(K, \Sigma)$ (remember (a) $_{\ell+1}$ +(b) $_\ell$) and it is stronger than all p_ℓ (for $\ell < \omega$). Let us show that q is $(N, \mathbb{Q}_\infty^*(K, \Sigma))$ -generic. To this end suppose $I \in N$ is a dense open subset of $\mathbb{Q}_\infty^*(K, \Sigma)$ and $r \in \mathbb{Q}_\infty^*(K, \Sigma)$ is stronger than q . Pick a condition $r_0 = (v, t_0^{r_0}, t_1^{r_0}, t_2^{r_0}, \dots) \geq r$ and $\ell < \omega$ such that

- (*) $r_0 \in I$, $I = I_\ell$ and $\text{lh}(v) = m_{i_\ell}$, and
- (**) $\mathbf{nor}[t_n^{r_0}] > 1$ for every $n < \omega$.

Then $r_0 \geq q^{[v]} \geq p_{\ell+1}^{[v]}$. Therefore, by (c) $_\ell$, we see that $p_{\ell+1}^{[v]} \in {}^* I_\ell$ and hence we may find $u \in \text{pos}(v, t_0^{r_0}, \dots, t_k^{r_0})$ (for some $k < \omega$) such that $p_{\ell+1}^{[u]} \in I_\ell$. Then $p_{\ell+1}^{[u]} \in N \cap I$ is compatible with r .

Note that the above argument shows also that for every open dense subset $I \in N$ of $\mathbb{Q}_\infty^*(K, \Sigma)$, the set $\{q^{[v]} : v \in \prod_{n < m_i} \mathbf{H}(n) \ \& \ i < \omega\} \cap I$ is predense above q . \square

4. A NONPROPER PRODUCT

Here, we will give an example of two proper forcing notions $\mathbb{Q}_\infty^*(K^1, \Sigma^1)$ and $\mathbb{Q}_\infty^*(K^2, \Sigma^2)$ such that their product $\mathbb{Q}_\infty^*(K, \Sigma)$ collapses \mathfrak{c} onto \aleph_0 .

Throughout this section we write \log instead of \log_2 .

Definition 4.1. Let $x, i \in \mathbb{R}$, $x > 0$, $i \geq 0$, and $k \in \omega \setminus \{0\}$. We let

$$f_k(x, i) = \frac{\log(\log(\log(x)) - i)}{k}$$

in the case that all three logarithms are well defined and attain a value ≥ 1 . In all other cases we define $f_k(x, i) = 1$.

- Lemma 4.2.**
- (1) $f_k(\frac{x}{2}, i) \geq f_k(x, i) - \frac{1}{k}$;
 - (2) letting $j = \frac{\log(\log(x)) + i}{2}$, if $f_k(x, i) \geq 2$ then $f_k(x, j) = f_k(x, i) - \frac{1}{k}$;
 - (3) letting j as in (2), if $\min\{f_k(x, i), f_k(y, j)\} > 1$ then $f_k(y, i) \geq f_k(x, i) - \frac{1}{k}$;
 - (4) if $x \geq 2^{2^{4+i}}$ and z such that $\log(\log(z)) = \frac{\log(\log(x)) + i}{2}$, then $f_k(z, i) = f_k(x, i) - \frac{1}{k}$.

Proof. (1) Note that for $x \geq 2$ we have

$$(*) \quad \log(x-1) \geq \log(x) - 1.$$

Indeed, $x \geq 2$ implies $x-1 \geq \frac{x}{2}$. Applying log to both sides we get $\log(x-1) \geq \log(\frac{x}{2}) = \log(x) - 1$.

If $x < 2^{2^{2+i}}$, then $\log(\log(\log(x)) - i) < 1$ (if at all defined), and $f_k(x, i) = 1 = f_k(\frac{x}{2}, i)$. So assume $x \geq 2^{2^{2+i}}$. Then $\log(x) \geq 2^{2+i} \geq 2$ and $\log(\log(x)) - i \geq 2$, and hence we may apply (*) with $\log(x)$ and $\log(\log(x)) - i$ and obtain

$$\begin{aligned} \log(\log(\log(\frac{x}{2})) - i) &= \log(\log(\log(x) - 1) - i) \geq \log(\log(\log(x)) - 1 - i) \\ &\geq \log(\log(\log(x)) - i) - 1. \end{aligned}$$

By dividing both sides by k we arrive at (1).

(2) Note that $f_k(x, i) \geq 2$ implies $\log(\log(x)) - i \geq 4$ and hence $\log(\log(x)) - j = \frac{\log(\log(x)) - i}{2} \geq 2$. Consequently

$$\begin{aligned} f_k(x, j) &= \frac{\log(\log(\log(x)) - j)}{k} = \frac{\log((\log(\log(x)) - i)/2)}{k} \\ &= \frac{\log(\log(\log(x)) - i) - 1}{k} = f_k(x, i) - \frac{1}{k}. \end{aligned}$$

(3) By assumption we have $\log(\log(y)) - j \geq 0$. By plugging in the definition of j and adding $j - i$ to both sides we obtain $\log(\log(y)) - i \geq \frac{1}{2}(\log(\log(x)) - i)$ and hence $\log(\log(\log(y)) - i) \geq \log(\log(\log(x)) - i) - 1$. After dividing by k we reach at (3).

(4) Note that

$$\begin{aligned} f_k(z, i) &= \frac{1}{k} \log(\log(\log(z)) - i) = \frac{1}{k} \log((\log(\log(x)) - i)/2) \\ &= \frac{1}{k} [\log(\log(\log(x)) - i) - 1] = f_k(x, i) - \frac{1}{k}. \end{aligned}$$

□

We are going to modify the example in Corollary 2.8 and Example 2.4.

Let $\bar{m} = \langle m_i : i < \omega \rangle$ be an increasing sequence of integers such that $m_0 = 0$ and $m_{i+1} - m_i > 4^{i+3}$. For $j < \omega$ let $\mathbf{H}(j) = i + 2$, where i is such that $m_i \leq j < m_{i+1}$, and let $g(x) = x$. The local good creating pair $(K_g^{\mathbf{H}}, \Sigma_g^{\mathbf{H}})$ introduced in 2.8 will be denoted by (K^1, Σ^1) . By 2.4 we know that $((K^1)^{\bar{m}}, (\Sigma^1)^{\bar{m}})$ (see 2.6) is sufficiently bad and hence (by 2.7) the forcing $\mathbb{Q}_\infty^*(K^1, \Sigma^1)$ collapses \mathfrak{c} into \aleph_0 .

Recall that for a creature $t \in K^1$ we have

- $\mathbf{dis}[t] = \langle i^t, A^t \rangle$ for some $i^t < \omega$ and $\emptyset \neq A^t \subseteq \mathbf{H}(i^t)$,
- $\mathbf{nor}[t] = |A^t|$ and $\mathbf{pos}(t) = A^t$.

Let $l_n = |\mathbf{H}(n)|$ and

$$k_n = \lfloor \sqrt{\max\{k \in \omega \setminus \{0\} : f_k(l_n, 0) > 1\}} \rfloor \quad \text{if } l_n > 2^{2^{16}},$$

and $k_n = 2$ if $l_n \leq 2^{2^{16}}$. Certainly we have $\lim_{n \rightarrow \infty} l_n = \infty$ and therefore $\lim_{n \rightarrow \infty} k_n = \infty$ as well (and the sequence $\langle k_n : n < \omega \rangle$ is non-decreasing). Note also that $\lim_{n \rightarrow \infty} f_{k_n}(l_n, 0) = \infty$.

Definition 4.3. Let K consist of all creatures $t \in \text{CR}[\mathbf{H}]$ such that

- **dis** $[t] = \langle m^t, A^t, i^t \rangle$ for some $m^t < \omega$ and $\emptyset \neq A^t \subseteq \mathbf{H}(m^t)$, and $i^t \in \omega$, $0 \leq i^t \leq \log(\log(l_{m^t}))$,
- **nor** $[t] = f_{k_{m^t}}(|A^t|, i^t)$, $m_{\text{dn}}^t = m^t$, $m_{\text{up}}^t = m^t + 1$ and $\text{pos}(t) = A^t$.

For $t \in K$ we let

$$\Sigma(t) = \{s \in K : m^s = m^t \ \& \ A^s \subseteq A^t \ \& \ i^s \geq i^t\}.$$

Lemma 4.4. (K, Σ) is a local forgetful strongly finitary good creating pair for \mathbf{H} . The forcing notion $\mathbb{Q}_\infty^*(K, \Sigma)$ collapses \mathfrak{c} to \aleph_0 .

Proof. It is straightforward to check that $(K^{\bar{m}}, \Sigma^{\bar{m}})$ inherits the sufficient badness of $((K^1)^{\bar{m}}, (\Sigma^1)^{\bar{m}})$ (remember 4.2(1)). Then use Proposition 2.7. \square

We are now going to define the desired factoring $\mathbb{Q}_\infty^*(K, \Sigma) \simeq \mathbb{P}^0 \times \mathbb{P}^1$ into proper factors $\mathbb{P}^0, \mathbb{P}^1$. For this we recursively define an increasing sequence $\bar{n} = \langle n_i : i < \omega \rangle$ so that $n_0 = 0$ and n_{i+1} is large enough such that

$$k_{n_{i+1}} \geq 2 \cdot \prod_{j < n_i} \mathbf{H}(j).$$

We put $U^0 = \bigcup_{i < \omega} [n_{2i}, n_{2i+1})$ and $U^1 = \bigcup_{i < \omega} [n_{2i+1}, n_{2i})$ and we let $\pi^0 : \omega \rightarrow U^0$ and $\pi^1 : \omega \rightarrow U^1$ be the increasing enumerations.

Definition 4.5. Let $\ell \in \{0, 1\}$. We define $\mathbf{H}^\ell = \mathbf{H} \circ \pi^\ell$ and we introduce K^ℓ, Σ^ℓ as follows.

- (1) K^ℓ consist of all creatures $t \in \text{CR}[\mathbf{H}^\ell]$ such that
 - **dis** $[t] = \langle m^t, A^t, i^t \rangle$ for some $m^t < \omega$ and $\emptyset \neq A^t \subseteq \mathbf{H}^\ell(m^t)$, and $i^t \in \omega$, $0 \leq i^t \leq \log(\log(l_n))$, where $n = \pi^\ell(m^t)$,
 - $m_{\text{dn}}^t = m^t$, $m_{\text{up}}^t = m^t + 1$, $\text{pos}(t) = A^t$ and **nor** $[t] = f_{k_n}(|A^t|, i^t)$ (where again $n = \pi^\ell(m^t)$).
- (2) For $t \in K^\ell$ we let

$$\Sigma^\ell(t) = \{s \in K^\ell : m^s = m^t \ \& \ A^s \subseteq A^t \ \& \ i^s \geq i^t\}.$$

Lemma 4.6. (1) For $\ell \in \{0, 1\}$, (K^ℓ, Σ^ℓ) is a local forgetful good creating pair for \mathbf{H}^ℓ .

- (2) Let $\bar{m}^0 = \langle m_i^0 : i < \omega \rangle$ and $\bar{\varepsilon}^0 = \langle \varepsilon_i^0 : i < \omega \rangle$ be such that $\pi^0(m_i^0) = n_{2i}$ and $\varepsilon_i^0 = 2/k_{n_{2i}}$. Then (K^0, Σ^0) has the $(\bar{\varepsilon}^0, \bar{m}^0)$ -halving property.
- (3) Let $\bar{m}^1 = \langle m_i^1 : i < \omega \rangle$ and $\bar{\varepsilon}^1 = \langle \varepsilon_i^1 : i < \omega \rangle$ be such that $\pi^1(m_i^1) = n_{2i+1}$ and $\varepsilon_i^1 = 2/k_{n_{2i+1}}$. Then (K^1, Σ^1) has the $(\bar{\varepsilon}^1, \bar{m}^1)$ -halving property.

Proof. (1) Should be clear.

(2) Let $t \in K^0$, **nor** $[t] \geq 2$, **dis** $[t] = \langle m, A, i^* \rangle$. Let $n = \pi^0(m) \geq n_{2i}$ (so $m_i^0 \leq m = m_{\text{dn}}^t$). Define $j = \frac{\log(\log(|A|)) + i^*}{2}$ and let z such that $\log(\log(z)) = j$.

Certainly, $k_n \geq 2$ and $f_{k_n}(|A|, i^*) \geq 2$, so $\log(\log(|A|)) - i^* \geq 16$ and hence $i^* < j \leq [j] < \log(\log(|A|)) \leq \log(\log(l_n))$. Let $t^* \in K^0$ be such that $\mathbf{dis}[t^*] = \langle m, A, [j] \rangle$. Clearly $t^* \in \Sigma^0(t)$. We are going to argue that t^* is an ε_i^0 -half of t (in (K^0, Σ^0)).

By (*) of the proof of Lemma 4.2(1) and then by 4.2(2) we have

$$\begin{aligned} \mathbf{nor}[t^*] &= f_{k_n}(|A|, [j]) = \frac{1}{k_n} \log(\log(\log(|A|)) - [j]) \geq \\ &\frac{1}{k_n} \log((\log(\log(|A|)) - j) - 1) \geq \frac{1}{k_n} \left(\log((\log(\log(|A|)) - j)) - 1 \right) = \\ &f_{k_n}(|A|, j) - \frac{1}{k_n} = f_{k_n}(|A|, i^*) - \frac{2}{k_n} \geq \mathbf{nor}[t] - \varepsilon_i^0. \end{aligned}$$

Now let $s \in \Sigma^0(t^*)$ be such that $\mathbf{nor}[s] > 1$. Let $\mathbf{dis}[s] = \langle m, A', i' \rangle$, thus $A' \subseteq A$ and $i' \geq [j] \geq j$. Let $t_0 \in K^0$ be such that $\mathbf{dis}[t_0] = \langle m, A', i^* \rangle$. Then $t_0 \in \Sigma^0(t)$ and $\text{pos}(t_0) = A' = \text{pos}(s)$. Also, $\mathbf{nor}[s] > 1$ implies $\log(\log(|A'|)) > i' \geq j$. By the definition of z we conclude $|A'| > z$. Noticing that $|A| > 2^{2^{4+i^*}}$ we apply Lemma 4.2(4) to obtain

$$\mathbf{nor}[t_0] = f_{k_n}(|A'|, i^*) \geq f_{k_n}(z, i^*) = f_{k_n}(|A|, i^*) - \frac{1}{k_n} \geq \mathbf{nor}[t] - \varepsilon_i^0.$$

(3) Like (2) above. □

Corollary 4.7. (1) *The forcing notions $\mathbb{Q}_\infty^*(K^\ell, \Sigma^\ell)$ (for $\ell = 0, 1$) are proper.*

(2) *Let $\mathbb{Q} = \{p \in \mathbb{Q}_\infty^*(K, \Sigma) : i(p) = n_i, i < \omega\}$. Then \mathbb{Q} is a dense suborder of $\mathbb{Q}_\infty^*(K, \Sigma)$ and it is isomorphic with a dense suborder of the product $\mathbb{Q}_\infty^*(K^0, \Sigma^0) \times \mathbb{Q}_\infty^*(K^1, \Sigma^1)$. Consequently, the latter forcing collapses \mathfrak{c} to \aleph_0 .*

Proof. (1) Let $\bar{m}^0, \bar{\varepsilon}^0$ be as in 4.6(2). By the choice of \bar{n} we have

$$\left| \prod_{n < \bar{m}_i^0} \mathbf{H}^0(n) \right| = \left| \prod_{\ell < i} \{ \mathbf{H}(j) : j \in \bigcup_{\ell < i} [n_{2\ell}, n_{2\ell+1}] \} \right| \leq \prod_{j < n_{2i-1}} \mathbf{H}(i) \leq \frac{1}{2} k_{n_{2i}} = 1/\varepsilon_i^0.$$

Consequently, Theorem 3.3 and Lemma 4.6(1,2) imply that $\mathbb{Q}_\infty^*(K^0, \Sigma^0)$ is proper.

Similarly for $\mathbb{Q}_\infty^*(K^1, \Sigma^1)$

(2) Should be clear. □

REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory: On the Structure of the Real Line*. A K Peters, Wellesley, Massachusetts, 1995.
- [2] Jakob Kellner and Saharon Shelah. Creature forcing and large continuum. The joy of halving. *preprint*. 1003.3425.
- [3] Jakob Kellner and Saharon Shelah. Decisive creatures and large continuum. *Journal of Symbolic Logic*, 74:73–104, 2009. math.LO/0601083.
- [4] Andrzej Roslanowski and Saharon Shelah. Monotone hulls for $\mathcal{N} \cap \mathcal{M}$. *Periodica Mathematica Hungarica*. 1007.5368.
- [5] Andrzej Roslanowski and Saharon Shelah. Norms on possibilities I: forcing with trees and creatures. *Memoirs of the American Mathematical Society*, 141(671):xii + 167, 1999. math.LO/9807172.
- [6] Saharon Shelah. *Proper and Improper Forcing*. Springer, Berlin, 1998.
- [7] Saharon Shelah. On what I do not understand (and have something to say:) Part I. *Fundamenta Mathematicae*, 166:1–82, 2000. math.LO/9906113.
- [8] Otmar Spinas. Proper Products. *Proceedings of the American Mathematical Society*, 137:2767–2772, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA AT OMAHA, OMAHA NE 68182,
USA

E-mail address: `roslanow@member.ams.org`

EINSTEIN INSTITUTE OF MATHEMATICS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM,
JERUSALEM 91904, ISRAEL AND DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW
BRUNSWICK, NJ 08854, USA

E-mail address: `shelah@math.huji.ac.il`

MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL, LUDEWIG-MEYN-STRASSE
4, 24098 KIEL, GERMANY

E-mail address: `spinas@math.uni-kiel.de`