

ON THE NON-RELATIVISTIC LIMIT OF A MODEL IN QUANTUM ELECTRODYNAMICS

EDGARDO STOCKMEYER

ABSTRACT. We consider a (semi-)relativistic spin-1/2 particle interacting with quantized radiation. The Hamiltonian has the form $\hat{H}_c^V := \{c^2[(\mathbf{p} + \mathbf{A})^2 + \sigma \cdot \mathbf{B}] + (mc^2)^2\}^{1/2} - mc^2 + V + H_f$. Assuming that the potential $|V|$ is bounded with respect to the momentum $|\mathbf{p}|$, we show that \hat{H}_c^V converges in norm-resolvent sense to the usual Pauli-Fierz operator when c , the speed of light, tends to ∞ .

1. INTRODUCTION

We consider a (semi-)relativistic electron interacting with quantized radiation in the presence of an external potential V . The corresponding Hamiltonian is given formally by

$$(1.1) \quad \hat{H}_c^V := \sqrt{c^2[(\mathbf{p} + \mathbf{A})^2 + \sigma \cdot \mathbf{B}] + (mc^2)^2} - mc^2 + V + H_f,$$

acting on $L^2(\mathbb{R}^3, \mathbb{C}^2) \otimes \mathcal{F}$, where \mathcal{F} is the bosonic Fock space. Here \mathbf{p} is the momentum of the electron, \mathbf{A} is the magnetic vector potential (in the Coulomb gauge), \mathbf{B} is the magnetic field, H_f is the free field-energy operator, $m > 0$ is the mass of the electron, $c > 0$ is the speed of light, and $\sigma = (\sigma_1, \sigma_2, \sigma_3)$, where σ_j are the usual Pauli-matrices.

Without an external potential V , the Hamiltonian (1.1) admits a fiber decomposition $\hat{H}_c^0(\mathbf{P})$, since the total momentum $\mathbf{P} \in \mathbb{R}^3$ is conserved. In [9] Miyao and Spohn investigated the ground state of $\hat{H}_c^0(\mathbf{P})$ (the polaron). In [8], Matte and the present author studied exponential localization of the low-lying spectral subspaces for Hydrogen-like atoms modelled by \hat{H}_c^V . In this paper we are interested in the non-relativistic limit of \hat{H}_c^V .

In absence of quantized radiation the non-relativistic limit is well understood (see [12]). However, in the presence of a field, only few results have been presented. Let $\mathcal{D}_c^V := D_{\mathbf{A}} - mc^2 + V + H_f$ be the Dirac operator coupled to the quantized field acting on $L^2(\mathbb{R}^3, \mathbb{C}^4) \otimes \mathcal{F}$, where $D_{\mathbf{A}}$ is the usual Dirac

Date: 07.05.09.

1991 Mathematics Subject Classification. 81Q10.

Key words and phrases. Non-relativistic limit, pseudo-relativistic, quantum electrodynamics.

operator minimally coupled to the vector potential \mathbf{A} . Arai in [2] proved that \mathcal{D}_c^V converges, as c tends to ∞ , in *strong* resolvent sense to the Pauli-Fierz Hamiltonian, denoted by \hat{H}_∞^V acting on $L^2(\mathbb{R}^3, \mathbb{C}^2) \otimes \mathcal{F}$. More precisely,

$$s - \lim_{c \rightarrow \infty} (\mathcal{D}_c^V - i)^{-1} = \begin{pmatrix} (\hat{H}_\infty^V - i)^{-1} & 0_{2 \times 2} \\ 0_{2 \times 2} & 0_{2 \times 2} \end{pmatrix}.$$

The method used cutoffs for V , H_f , and \mathbf{A} in order to include them as perturbations. (We remark that it is unlikely to improve this convergence to norm resolvent sense, since the spectrum of \mathcal{D}_c^0 for $\mathbf{A} = 0$ is equal to the whole real line, i.e., $\sigma(D_0 + H_f) = \mathbb{R}$.) By the same method the non-relativistic limit of the Dirac polaron was investigated by Arai in [3].

In this paper we do not include the field energy H_f as a perturbation but we make strong use of the fact that \hat{H}_c^0 is bounded from below. Assuming that the potential V is (operator) bounded with respect to the (electric) momentum $|\mathbf{p}|$, we show that, as c (the speed of light) tends to infinity, \hat{H}_c^V converges in norm resolvent sense to the usual (non-relativistic) Pauli-Fierz operator \hat{H}_∞^V .

In the next section we introduce the models to be considered and state precisely our main results, Theorems 2.2 and 2.4 below.

2. DEFINITION OF THE MODELS AND MAIN RESULTS

Let $\mathfrak{h} = L^2(\mathbb{R}^3 \times \{1, 2\})$ be the one-photon Hilbert space. We use the convention

$$k = (\mathbf{k}, \lambda) \in \mathbb{R}^3 \times \{1, 2\}; \quad \int dk := \sum_{\lambda \in \mathbb{Z}_2} \int_{\mathbb{R}^3} d^3 \mathbf{k}.$$

The space of the quantized photon field is the bosonic Fock space

$$\mathcal{F} \equiv \mathcal{F}[\mathfrak{h}] := \bigoplus_{n=0}^{\infty} \mathfrak{h}^{\otimes_s n} \ni \psi = (\psi^{(0)}, \psi^{(1)}, \psi^{(2)}, \dots),$$

where $\mathfrak{h}^{\otimes_s n}$ is the n -fold symmetric tensor product of \mathfrak{h} and $\mathfrak{h}^{\otimes_s 0} = \mathbb{C}$. As usual we denote the vacuum vector by $\Omega := (1, 0, 0, \dots) \in \mathcal{F}[\mathfrak{h}]$. Many calculations will be performed on the following dense subspace of $\mathcal{F}[\mathfrak{h}]$,

$$\mathcal{C}_0 := \mathbb{C} \oplus \bigoplus_{n \in \mathbb{N}} C_0((\mathbb{R}^3 \times \{1, 2\})^n) \cap \mathfrak{h}^{\otimes_s n}.$$

The free field energy of the photons is the self-adjoint operator given by

$$\mathcal{D}(H_f) := \left\{ (\psi^{(n)})_{n=0}^\infty \in \mathcal{F}[\mathfrak{h}] : \sum_{n=1}^{\infty} \int \left| \sum_{j=1}^n \omega(k_j) \psi^{(n)}(k_1, \dots, k_n) \right|^2 dk_1 \dots dk_n < \infty \right\},$$

and, for $\psi \in \mathcal{D}(H_f)$,

$$(H_f\psi)^{(0)} = 0, \quad (H_f\psi)^{(n)}(k_1, \dots, k_n) = \sum_{j=1}^n \omega(k_j) \psi^{(n)}(k_1, \dots, k_n), \quad n \in \mathbb{N}.$$

Here the dispersion relation ω is an almost everywhere nonzero multiplication operator on \mathfrak{h} that depends only on \mathbf{k} and not on $\lambda \in \{1, 2\}$.

The annihilation operator $a(f)$ of a photon state $f \in \mathfrak{h}$ is, for any $\varphi \in \mathcal{C}_0$, given by

$$(2.1) \quad a(f)\varphi = \int \bar{f}(k) a(k) \varphi dk,$$

where $a(k)$ annihilates a photon with wave vector/polarization k ,

$$(a(k)\psi)^{(n)}(k_1, \dots, k_n) = (n+1)^{1/2} \psi^{(n+1)}(k, k_1, \dots, k_n), \quad n \in \mathbb{N}_0,$$

almost everywhere, and $a(k)\Omega = 0$. For $f \in \mathfrak{h}$ the creation operator $a^\dagger(f)$ satisfies $\langle a(f)\phi | \varphi \rangle = \langle \phi | a^\dagger(f)\varphi \rangle$, for any $\phi, \varphi \in \mathcal{C}_0$. We define $a^\dagger(f)$ and $a(f)$ on their maximal domains. The following canonical commutation relations hold true on \mathcal{C}_0 , for any $f, g \in \mathfrak{h}$,

$$(2.2) \quad [a(f), a(g)] = [a^\dagger(f), a^\dagger(g)] = 0, \quad [a(f), a^\dagger(g)] = \langle f | g \rangle \mathbb{1}.$$

The full Hilbert space containing electron and photon degrees of freedom is

$$\mathcal{H} := L^2(\mathbb{R}^3, \mathbb{C}^4) \otimes \mathcal{F}.$$

It contains the dense subspace,

$$\mathcal{D}_0 := C_0^\infty(\mathbb{R}^3, \mathbb{C}^4) \otimes \mathcal{C}_0. \quad (\text{Algebraic tensor product.})$$

We consider general form factors fulfilling the following condition:

Hypothesis 1. For every $k \in (\mathbb{R}^3 \setminus \{0\}) \times \mathbb{Z}_2$ and $j \in \{1, 2, 3\}$, $G^{(j)}(k)$ is a bounded continuously differentiable function, $\mathbb{R}^3 \ni \mathbf{x} \mapsto G_{\mathbf{x}}^{(j)}(k)$, satisfying

$$(2.3) \quad 2 \int \omega(k)^\ell \| \mathbf{G}(k) \|_\infty^2 dk \leq d_\ell^2, \quad \ell \in \{-1, 0, 1, 2\},$$

and

$$(2.4) \quad 2 \int \omega(k)^{-1} \| \nabla_{\mathbf{x}} \times \mathbf{G}(k) \|_\infty^2 dk \leq d_1^2,$$

for some $d_{-1}, \dots, d_2 \in (0, \infty)$. Here $\mathbf{G}_{\mathbf{x}}(k) = (G_{\mathbf{x}}^{(1)}(k), G_{\mathbf{x}}^{(2)}(k), G_{\mathbf{x}}^{(3)}(k))$ and $\| \mathbf{G}(k) \|_\infty := \sup_{\mathbf{x}} | \mathbf{G}_{\mathbf{x}}(k) |$.

Remark 2.1. In the specific physical situation the dispersion relation is $\omega^{\text{ph}}(k) = \hbar c|\mathbf{k}|$, where $\hbar > 0$ is Plank's constant divided by 2π . The form factor is given by

$$(2.5) \quad G_{\mathbf{x}}^{\text{ph}}(k) := \frac{\hbar^{1/2}}{2\pi(c|\mathbf{k}|)^{1/2}} \mathbb{1}_{\{|\mathbf{k}| \leq \Lambda\}} e^{-i\mathbf{k} \cdot \mathbf{x}} \boldsymbol{\varepsilon}(k),$$

for $(\mathbf{x}, k) \in \mathbb{R}^3 \times (\mathbb{R}^3 \times \{1, 2\})$ with $\mathbf{k} \neq 0$. Here $\Lambda > 0$ is an ultraviolet cut-off parameter. The polarization vectors, $\boldsymbol{\varepsilon}(\mathbf{k}, \lambda)$, $\lambda \in \mathbb{Z}_2$, are homogeneous of degree zero in \mathbf{k} such that $\{\hat{\mathbf{k}}, \boldsymbol{\varepsilon}(\hat{\mathbf{k}}, 0), \boldsymbol{\varepsilon}(\hat{\mathbf{k}}, 1)\}$ is an orthonormal basis of \mathbb{R}^3 , for every $\hat{\mathbf{k}} \in \mathbb{S}^2$. This corresponds to the Coulomb gauge.

We further remark that the dependence on c of (2.5) and the definition of ω^{ph} is not relevant to us, since we want to study the electron and not the photon non-relativistic limit.

We introduce the self-adjoint Dirac matrices $\alpha_1, \alpha_2, \alpha_3$, and β that act on the four spinor components of an element from \mathcal{H} . They are given by

$$\alpha_j := \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix}, \quad j \in \{1, 2, 3\}, \quad \beta := \alpha_0 := \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix},$$

where $\sigma_1, \sigma_2, \sigma_3$ denote the standard Pauli matrices, and fulfill the relations

$$(2.6) \quad \alpha_i \alpha_j + \alpha_j \alpha_i = 2\delta_{ij} \mathbb{1}, \quad i, j \in \{0, 1, 2, 3\},$$

$$(2.7) \quad \alpha_i \alpha_j = \delta_{ij} + i\epsilon_{ijm} \Sigma_m, \quad i, j \in \{1, 2, 3\},$$

where δ_{ij} is the Kronecker delta, ϵ_{ijm} is the Levi-Civita antisymmetric symbol, and

$$(2.8) \quad \Sigma_j := \begin{pmatrix} \sigma_j & 0 \\ 0 & \sigma_j \end{pmatrix}.$$

The interaction between the electron and the photons is given by

$$(2.9) \quad \boldsymbol{\alpha} \cdot \mathbf{A} \equiv \boldsymbol{\alpha} \cdot \mathbf{A}(\mathbf{x}) := a(\boldsymbol{\alpha} \cdot \mathbf{G}_{\mathbf{x}}) + a^\dagger(\boldsymbol{\alpha} \cdot \mathbf{G}_{\mathbf{x}}).$$

In order to define the semi-relativistic Pauli-Fierz operator we recall that the free Dirac operator, minimally coupled to \mathbf{A} , is given as

$$(2.10) \quad D_{\mathbf{A}} := c\boldsymbol{\alpha} \cdot (-i\hbar\nabla + \mathbf{A}) + \beta mc^2,$$

where $m > 0$ is the mass of the electron. (Note that usually there is a factor in front of \mathbf{A} ; in physical units e/c . We have absorbed this factor in the definition of \mathbf{A} .) An application of Nelson's commutator theorem shows that $D_{\mathbf{A}}$ is essentially self-adjoint on \mathcal{D}_0 [1, 7]. We denote its closure starting from \mathcal{D}_0 again by the same symbol. Henceforth, we set

$$(2.11) \quad \mathbf{p}_{\mathbf{A}} := -i\hbar\nabla + \mathbf{A}, \quad \text{and} \quad \mathbf{p} := -i\hbar\nabla.$$

We have the following relation on \mathcal{D}_0 :

$$(2.12) \quad D_{\mathbf{A}}^2 = c^2(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^2 + (mc^2)^2 = c^2(\mathbf{p}_{\mathbf{A}}^2 + \boldsymbol{\Sigma} \cdot \mathbf{B}) + (mc^2)^2,$$

where $\boldsymbol{\Sigma}$ is defined through (2.8) and the magnetic field \mathbf{B} is given as

$$(2.13) \quad \mathbf{B} \equiv \mathbf{B}(\mathbf{x}) = \nabla \times \mathbf{A}(\mathbf{x}) = a(\nabla \times \mathbf{G}_{\mathbf{x}}) + a^\dagger(\nabla \times \mathbf{G}_{\mathbf{x}}).$$

Equation (2.12) shows, in particular, that $\sigma(D_{\mathbf{A}}) \subset (-\infty, -1] \cup [1, \infty)$.

Next we state the conditions on the potential V .

Hypothesis 2. *Let V be a symmetric (matrix-valued) multiplication operator acting on \mathcal{H} , such that for almost all $\mathbf{x} \in \mathbb{R}^3$,*

$$V(\mathbf{x}) = \begin{pmatrix} V_1(\mathbf{x}) & 0_{2 \times 2} \\ 0_{2 \times 2} & V_2(\mathbf{x}) \end{pmatrix}.$$

We assume that there exist $a, b \in \mathbb{R}^+$, such that for all $\varphi \in \mathcal{D}_0$,

$$(2.14) \quad \langle \varphi | V^2 \varphi \rangle \leq a^2 \langle \varphi | \mathbf{p}^2 \varphi \rangle + b^2 \|\varphi\|^2$$

holds.

We define now the semi-relativistic Pauli-Fierz operator (minus the electron rest energy) through the quadratic form on \mathcal{D}_0 ,

$$(2.15) \quad \langle \varphi | H_c^V \varphi \rangle := \langle \varphi | (|D_{\mathbf{A}}| - mc^2 + V + H_f) \varphi \rangle.$$

Note, that on \mathcal{D}_0 ,

$$(2.16) \quad H_c^0 \equiv H_c = \begin{pmatrix} \hat{d}_{\mathbf{A}} + H_f & 0_{2 \times 2} \\ 0_{2 \times 2} & \hat{d}_{\mathbf{A}} + H_f \end{pmatrix},$$

where

$$(2.17) \quad \hat{d}_{\mathbf{A}} := \sqrt{c^2[(\mathbf{p} + \mathbf{A})^2 + \boldsymbol{\sigma} \cdot \mathbf{B}] + (mc^2)^2} - mc^2$$

acts on $L^2(\mathbb{R}^3, \mathbb{C}^2) \otimes \mathcal{F}$.

Since the quadratic form $\langle \varphi | H_c \varphi \rangle \equiv \langle \varphi | H_c^0 \varphi \rangle$ is positive, there is a unique (positive) self-adjoint extension of $H_c \upharpoonright \mathcal{D}_0$, which we denote again by H_c , whose domain is contained in the form domain of the closure of the quadratic form defined in (2.15) for $V = 0$.

We prove the following result at the end of Section 3.

Theorem 2.2. *Assume that \mathbf{G} fulfills Hypothesis 1 and V fulfills Hypothesis 2. Then, for c (the speed of light) large enough, the operator $H_c^V \upharpoonright \mathcal{D}_0$ has a unique self-adjoint extension, denoted by H_c^V , with form domain $\mathcal{Q}(H_c^V)$ satisfying*

$$(2.18) \quad \mathcal{D}(H_c^V) \subset \mathcal{Q}(H_c^V) = \mathcal{Q}(H_c) = \mathcal{Q}(|\mathbf{p}| + H_f).$$

Moreover, H_c^V is bounded below by a constant independent of c .

Remark 2.3. We note that a similar statement (without the specification of the form domain) was proved in [8] by means of a diamagnetic inequality. Our proof is based on an explicit bound for $|\mathbf{p}|$ in terms of H_c (see Lemma 3.4). This bound turns out to give the key ingredient to show (2.18).

Next we define the (non-relativistic) Pauli-Fierz operator on \mathcal{D}_0 as

$$(2.19) \quad H_\infty^V := \frac{\mathbf{p}_\mathbf{A}^2}{2m} + \frac{1}{2m} \boldsymbol{\Sigma} \cdot \mathbf{B} + V + H_f, \quad H_\infty \equiv H_\infty^0.$$

Actually, the operator defined above is a two-fold copy of the usual Pauli-Fierz operator given by $\hat{H}_\infty := \mathbf{p}_\mathbf{A}^2/(2m) + \sigma \cdot \mathbf{B}/(2m) + H_f$ acting on $L^2(\mathbb{R}^3, \mathbb{C}^2) \otimes \mathcal{F}$:

$$H_\infty^V = \begin{pmatrix} \hat{H}_\infty + V_1 & 0_{2 \times 2} \\ 0_{2 \times 2} & \hat{H}_\infty + V_2 \end{pmatrix}.$$

A potential V satisfying Hypothesis 2 is relatively \mathbf{p}^2 -bounded with bound zero, hence H_∞^V is essentially self-adjoint on any core for $\mathbf{p}^2 + H_f$ and its self-adjoint extension has domain $\mathcal{D}(H_\infty^V) = \mathcal{D}(\mathbf{p}^2 + H_f)$ (see [4, 5]). Henceforth, we set

$$(2.20) \quad T_\mathbf{A} = \frac{\mathbf{p}_\mathbf{A}^2}{2m} + \frac{1}{2m} \boldsymbol{\Sigma} \cdot \mathbf{B}.$$

We can now state the main result of this article whose proof can be found at the end of Section 4.

Theorem 2.4. *Assume that \mathbf{G} and V satisfy Hypotheses 1 and 2 respectively. Then, for any $z \in \mathbb{C} \setminus \mathbb{R}$,*

$$\lim_{c \rightarrow \infty} \|(H_c^V - z)^{-1} - (H_\infty^V - z)^{-1}\| = 0.$$

The rest of the paper is organized as follows: The main technical ingredients and the proof of Theorem 2.2 are presented in Section 3. In Section 4 we prove Theorem 2.4 about the non-relativistic limit. Finally the main text is followed by an Appendix where we prove some known inequalities. Throughout this paper we shall assume that Hypotheses 1 and 2 are fulfilled.

3. MAIN ESTIMATES

The following estimates are well known. A proof of the lemma can be found in Appendix A.

Lemma 3.1 (Field-Bounds). *Assume that $f, f\omega^{-1/2} \in \mathfrak{h}$. Then, for all $\varphi \in \mathcal{D}(H_f^{1/2})$,*

$$(3.1) \quad \|a(f)\varphi\|^2 \leq \|f/\omega^{1/2}\|_{\mathfrak{h}}^2 \|H_f^{1/2}\varphi\|^2,$$

$$(3.2) \quad \|a^\dagger(f)\varphi\|^2 \leq \|f/\omega^{1/2}\|_{\mathfrak{h}}^2 \|H_f^{1/2}\varphi\|^2 + \|f\|_{\mathfrak{h}}^2 \|\varphi\|^2.$$

In particular, for all $\varphi \in \mathcal{D}(H_f^{1/2})$,

$$(3.3) \quad \langle \varphi | \boldsymbol{\Sigma} \cdot \mathbf{B} \varphi \rangle \leq 2d_1 \|H_f^{1/2} \varphi\| \|\varphi\|,$$

$$(3.4) \quad \langle \mathbf{A} \varphi | \mathbf{A} \varphi \rangle \leq 2d_{-1}^2 \|H_f^{1/2} \varphi\|^2 + d_0^2 \|\varphi\|^2.$$

For notational convenience we set

$$(3.5) \quad d_{\mathbf{A}} := |D_{\mathbf{A}}| - mc^2 = \{c^2(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^2 + (mc^2)^2\}^{1/2} - mc^2,$$

acting on $L^2(\mathbb{R}^3, \mathbb{C}^4) \otimes \mathcal{F}$. (Note that $d_{\mathbf{A}} = \hat{d}_{\mathbf{A}} \oplus \hat{d}_{\mathbf{A}}$, where $\hat{d}_{\mathbf{A}}$ is defined in (2.17).) Furthermore,

$$(3.6) \quad R_c^V(z) := (H_c^V + z)^{-1}, \quad R_c(z) := R_c^0(z),$$

$$(3.7) \quad R_{\infty}^V(z) := (H_{\infty}^V + z)^{-1}, \quad R_{\infty}(z) := R_{\infty}^0(z),$$

for some $z \in \mathbb{C}$ with $-z \in \varrho(H_c^V) \cap \varrho(H_{\infty}^V)$, where $\varrho(\cdot)$ denotes the resolvent set.

Lemma 3.2. *For any $\varphi \in \mathcal{D}(T_{\mathbf{A}})$, we have*

$$(3.8) \quad T_{\mathbf{A}} \varphi = \left(d_{\mathbf{A}} + \frac{d_{\mathbf{A}}^2}{2mc^2} \right) \varphi.$$

Moreover, for $\nu \in (0, 2]$, we have the relation $\mathcal{D}(H_{\infty}^V) = \mathcal{D}(\mathbf{p}^2 + H_f) \subset \mathcal{D}(d_{\mathbf{A}}^{\nu})$ and the estimate, for any $-z \in \varrho(H_{\infty}^V)$,

$$(3.9) \quad \|d_{\mathbf{A}}^{\nu} R_{\infty}^V(z)\| = \mathcal{O}(c^{\nu}).$$

Proof. Let $f : \mathbb{R} \rightarrow [0, \infty)$ with $f(t) = \sqrt{(ct)^2 + (mc^2)^2} - mc^2$. Obviously, $t^2/2m = f(t) + f(t)^2/(2mc^2)$. Since $d_{\mathbf{A}} = f(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})$, we find (3.8) by the functional calculus.

Clearly, we have, for all $\nu \in (0, 2]$, the inequality $f(t)^{2\nu} \leq (c|t|)^{2\nu}$, which implies, for $\varphi \in \mathcal{D}_0$, that $\langle \varphi | d_{\mathbf{A}}^{2\nu} \varphi \rangle \leq c^{2\nu} \langle \varphi | |\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}}|^{2\nu} \varphi \rangle$. In view of Young's inequality [$ab \leq a^p/p + b^q/q$, for $p, q \in (1, \infty)$ with $1/p + 1/q = 1$], we obtain, for $t \in \mathbb{R}$ and $\nu \in (0, 2)$, that $|t|^{2\nu} \leq \nu t^4/2 + (2-\nu)/2$. Hence, for any $\nu \in (0, 2]$, we get

$$(3.10) \quad \langle \varphi | d_{\mathbf{A}}^{2\nu} \varphi \rangle \leq c^{2\nu} \nu/2 \langle \varphi | [(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^4 + (2-\nu)/\nu] \varphi \rangle.$$

Recall that $(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^4 = (2m)^2 T_{\mathbf{A}}^2$. Due to (3.10) and the fact that

$$(3.11) \quad \mathcal{D}(H_{\infty}^V) = \mathcal{D}(\mathbf{p}^2 + H_f),$$

we see that, in order to show (3.9), it suffices to prove that there exists a constant $k > 0$, such that for any $\varphi \in \mathcal{D}_0$, $\|T_{\mathbf{A}} \varphi\|^2 \leq k(\|(\mathbf{p}^2 + H_f)\varphi\|^2 + \|\varphi\|^2)$. That this is indeed the case follows from the following consideration: By (3.11),

there exists a constant $k' > 1$, such that $\|H_\infty\varphi\|^2 \leq k'(\|(\mathbf{p}^2 + H_f)\varphi\|^2 + \|\varphi\|^2)$, thus,

$$\begin{aligned}
(3.12) \quad \|T_{\mathbf{A}}\varphi\|^2 &\leq 2(\|(T_{\mathbf{A}} + H_f)\varphi\|^2 + \|H_f\varphi\|^2) \\
&\leq 2k'(\|(\mathbf{p}^2 + H_f)\varphi\|^2 + \|H_f\varphi\|^2 + \|\varphi\|^2) \\
&\leq 2\sqrt{2}k'(\|(\mathbf{p}^2 + H_f)\varphi\|^2 + \|\varphi\|^2).
\end{aligned}$$

This finishes the proof. \square

The next Lemma is proved in [8]. We sketch its proof in Appendix A. We define, for some $E > 0$, the operator

$$(3.13) \quad \tilde{H}_f := H_f + E.$$

Lemma 3.3 ([8]). *Assume that $E \geq (10\pi d_1)^2/(mc)^2$. Then, for all $\varphi \in \mathcal{D}_0$,*

$$(3.14) \quad \operatorname{Re} \langle |D_{\mathbf{A}}|\varphi | \tilde{H}_f \varphi \rangle \geq \left(1 - \frac{10\pi d_1}{mcE^{1/2}}\right) \| |D_{\mathbf{A}}|^{1/2} \tilde{H}_f^{1/2} \varphi \|^2.$$

In the following Lemma we establish that the momentum $|\mathbf{p}|$ is dominated, in the sense of quadratic forms, by the operator H_c (see (2.16)).

Lemma 3.4. *For all $\varepsilon \in (0, 1]$ there exist a constant $k_\varepsilon > 0$, such that for all $c \geq 1$ and $\varphi \in \mathcal{D}_0$,*

$$(3.15) \quad \langle \varphi | |\mathbf{p}|\varphi \rangle \leq \sqrt{(1+\varepsilon)(\varepsilon + 1/c^2)} \langle \varphi | (d_{\mathbf{A}} + H_f)\varphi \rangle + k_\varepsilon \|\varphi\|^2.$$

Proof. We start by observing that, as quadratic form on \mathcal{D}_0 ,

$$(3.16) \quad \mathbf{p}^2 = \mathbf{p}_{\mathbf{A}}^2 - \mathbf{p}_{\mathbf{A}} \cdot \mathbf{A} - \mathbf{A} \cdot \mathbf{p}_{\mathbf{A}} + \mathbf{A}^2.$$

Estimating the cross terms and using (3.4) we get, for $\varepsilon \in (0, 1]$,

$$(3.17) \quad \mathbf{p}^2 \leq (1+\varepsilon)\mathbf{p}_{\mathbf{A}}^2 + \frac{2}{\varepsilon}\mathbf{A}^2 \leq (1+\varepsilon)\mathbf{p}_{\mathbf{A}}^2 + \frac{4d_{-1}^2}{\varepsilon}H_f + \frac{2d_0^2}{\varepsilon}.$$

We use (3.3) to include the spin-term. For some $\delta > 0$ and $\varphi \in \mathcal{D}_0$, we have

$$\begin{aligned}
(3.18) \quad (1+\varepsilon)|\langle \varphi | \boldsymbol{\Sigma} \cdot \mathbf{B}\varphi \rangle| &\leq 4d_1\|H_f^{1/2}\varphi\|\|\varphi\| \\
&\leq \frac{2d_1}{\delta}\langle \varphi | H_f\varphi \rangle + 2d_1\delta\|\varphi\|^2 \\
&\leq \langle \varphi | H_f\varphi \rangle + 4d_1^2\|\varphi\|^2,
\end{aligned}$$

where we choose $\delta = 2d_1$. Inserting (3.18) in (3.17) and using the identity $(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^2 = \mathbf{p}_{\mathbf{A}}^2 + \boldsymbol{\Sigma} \cdot \mathbf{B}$ we get that there exists a constant $k > 0$ such that, for all $\varepsilon \in (0, 1]$,

$$(3.19) \quad \frac{\mathbf{p}^2}{1+\varepsilon} \leq (\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^2 + 1 + \frac{k}{\varepsilon}(H_f + 1) \equiv \tilde{D}_{\mathbf{A}}^2 + \frac{k}{\varepsilon}(H_f + 1),$$

where we define $\tilde{D}_{\mathbf{A}} := \boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}} + \beta = D_{\mathbf{A}}|_{m=c=1}$. In view of Lemma 3.3 (with $m = c = 1$) we have, for $\tilde{H}_f = H_f + E$, $E > \max\{1, (10\pi d_1)^2\}$, and some $\delta > 0$,

$$(3.20) \quad \begin{aligned} \frac{\mathbf{p}^2}{1 + \varepsilon} &\leq \tilde{D}_{\mathbf{A}}^2 + \delta(|\tilde{D}_{\mathbf{A}}|\tilde{H}_f + \tilde{H}_f|\tilde{D}_{\mathbf{A}}|) + \frac{k}{\varepsilon}\tilde{H}_f \\ &\leq (|\tilde{D}_{\mathbf{A}}| + \delta\tilde{H}_f)^2 + \left(\frac{k}{2\varepsilon\delta}\right)^2. \end{aligned}$$

Observing that from $(\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}})^2 = 2md_{\mathbf{A}} + d_{\mathbf{A}}^2/c^2$ follows $|\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}}| \leq \sqrt{1/c^2 + \varepsilon}d_{\mathbf{A}} + m/\varepsilon^{1/2}$ and that the square root is operator monotone, we get that

$$(3.21) \quad \begin{aligned} \frac{|\mathbf{p}|}{\sqrt{1 + \varepsilon}} &\leq |\tilde{D}_{\mathbf{A}}| + \delta\tilde{H}_f + \frac{k}{2\varepsilon\delta} \leq |\boldsymbol{\alpha} \cdot \mathbf{p}_{\mathbf{A}}| + \delta\tilde{H}_f + \frac{k}{2\varepsilon\delta} + 1 \\ &\leq \sqrt{1/c^2 + \varepsilon}d_{\mathbf{A}} + \delta\tilde{H}_f + \frac{k}{2\varepsilon\delta} + 1 + \frac{m}{\varepsilon^{1/2}} \\ &\leq \sqrt{1/c^2 + \varepsilon}(d_{\mathbf{A}} + H_f) + \sqrt{2}E + \frac{k}{2\varepsilon^{3/2}} + 1 + \frac{m}{\varepsilon^{1/2}}, \end{aligned}$$

where in the last inequality we choose $\delta = \sqrt{1/c^2 + \varepsilon} > \varepsilon^{1/2}$ and we used that $\varepsilon \leq 1 \leq c$. Inequality (3.21) proves the lemma. \square

Corollary 3.5. *For any $\lambda \in (0, 1]$ there exists $R_\lambda > 0$, such that for all $\rho > R_\lambda$ and $c > \max\{2/\lambda, 2a/\lambda\}$,*

$$(3.22) \quad \||V|^{1/2}(H_c + \rho)^{-1/2}\|^2 \leq \lambda,$$

where a is the constant appearing in Hypothesis 2.

Proof. We use (3.15) and note that (2.14) implies that

$$(3.23) \quad |V| \leq a|\mathbf{p}| + b \leq \sqrt{2}a\sqrt{\varepsilon + 1/c^2}H_c + b + ak_\varepsilon < \lambda(H_c + \frac{1}{\lambda}(b + ak_\varepsilon)),$$

where we set $\varepsilon = \min\{\lambda^2/4, \lambda^2/(4a^2)\}$ and we used that $1/c^2 < \min\{\lambda^2/4, \lambda/(4a^2)\}$. These facts prove the claim with $R_\lambda := (b + ak_\varepsilon)/\lambda$. \square

Proof of Theorem 2.2. For c large enough we have, by (3.23), that V is H_c -form bounded with form bound less than one. Therefore, by the KLMN theorem (see [10, Theorem X.17]), the operator H_c^V has a unique self-adjoint extension with form domain $\mathcal{Q}(H_c)$ which is bounded below by $-R_1$, where R_1 is defined after Equation (3.23).

In order to prove that $\mathcal{Q}(H_c) = \mathcal{Q}(|\mathbf{p}| + H_f)$, it suffices to show that the quadratic form norms associated to H_c and $|\mathbf{p}| + H_f$, defined on \mathcal{D}_0 , are equivalent. Using the inequality

$$\frac{d_{\mathbf{A}}^2}{c^2} \leq \mathbf{p}_{\mathbf{A}}^2 + \boldsymbol{\Sigma} \cdot \mathbf{B} \leq 2\mathbf{p}^2 + 2\mathbf{A}^2 + \boldsymbol{\Sigma} \cdot \mathbf{B},$$

and the estimates (3.4) and (3.18), we find some constant $K' > 0$, such that

$$d_{\mathbf{A}}^2 \leq (K')^2[(|\mathbf{p}| + H_f)^2 + 1].$$

Therefore, by the (operator) monotonicity of the square root, we get

$$d_{\mathbf{A}} + H_f \leq K'(|\mathbf{p}| + H_f) + H_f + K' \leq (1 + K')(|\mathbf{p}| + H_f + 1).$$

To conclude the proof we note that, by Lemma 3.4, we know that there is a constant K , such that $|\mathbf{p}| \leq K(d_{\mathbf{A}} + H_f + 1)$. Thus, for $\delta \in (0, 1)$, we have, as quadratic forms on \mathcal{D}_0 ,

$$\delta(H_f + |\mathbf{p}|) \leq Kd_{\mathbf{A}} + (K + \delta)H_f + K \leq (K + \delta)(d_{\mathbf{A}} + H_f + 1).$$

This finishes the proof of the assertion. \square

4. THE NON-RELATIVISTIC LIMIT

Lemma 4.1. *Assume that $c \geq 1$ is so large that H_c^V is self-adjoint. Then, for any $z \in \mathbb{C}$ with $-z \in \varrho(H_c^V) \cap \varrho(H_\infty^V)$, the identity*

$$(4.1) \quad R_c^V(z) = \left[1 + R_c^V(z) \frac{d_{\mathbf{A}}^2}{2mc^2}\right] R_\infty^V(z)$$

holds on \mathcal{H} .

Proof. We pick some $\varphi \in \mathcal{D}_0$ and use (3.8) to compute

$$\begin{aligned} R_c^V(z)(H_\infty^V + z)\varphi &= R_c^V(z)(T_{\mathbf{A}} + H_f + V + z)\varphi \\ &= R_c^V(z)\left(d_{\mathbf{A}} + \frac{d_{\mathbf{A}}^2}{2mc^2} + V + H_f + z\right)\varphi = \left(1 + R_c^V(z) \frac{d_{\mathbf{A}}^2}{2mc^2}\right)\varphi \\ &= \left\{\left(1 + R_c^V(z) \frac{d_{\mathbf{A}}^2}{2mc^2}\right) R_\infty^V(z)\right\}(H_\infty^V + z)\varphi. \end{aligned}$$

Due to the fact that H_∞^V is essentially self-adjoint on \mathcal{D}_0 we find (4.1) on some dense subset of \mathcal{H} . Moreover, thanks to (3.9), the term in $\{\dots\}$ above is bounded. Therefore, by a simple limit argument, we obtain (4.1). \square

Proof of Theorem 2.4. We first note that both operators, H_∞^V and H_c^V , are bounded below by a constant independent of the speed of light c . In fact, by (3.8), we have that $H_\infty^V \geq H_c^V$ the last operator being bounded below (independent of c) due to Theorem 2.2. We define $e_V := \inf(\sigma(H_c^V))$ and choose $\rho > |e_V|$.

Next, we observe that it is enough to prove Theorem 2.4 for ρ instead of z . Indeed, define $\mathcal{S}_z := (H_z^V + \rho)/(H_z^V + z)$, for $z \in \mathbb{C} \setminus \mathbb{R}$ and $z = c$ or ∞ . Clearly, $\|\mathcal{S}_z\| \leq 1 + (\rho + |z|)/|\operatorname{Im} z|$. Further, for $z \in \mathbb{C} \setminus \mathbb{R}$, we have the following well-known operator identity

$$(4.2) \quad R_c^V(z) - R_\infty^V(z) = \mathcal{S}_c(R_c^V(\rho) - R_\infty^V(\rho)) \mathcal{S}_z.$$

Therefore, on account of Lemma 4.1 and (4.2), it suffices to show that

$$(4.3) \quad \|R_c^V(\rho) \frac{d_{\mathbf{A}}^2}{2mc^2} R_\infty^V(\rho)\| \rightarrow 0 \quad \text{as} \quad c \rightarrow \infty.$$

We observe that by Tiktopolus' Formula (see [11]) and (3.22), we have, for $c \geq \max\{4, 4a\}$, that

$$(4.4) \quad R_c^V(\rho) = R_c(\rho)^{1/2} (1 + R_c(\rho)^{1/2} V R_c(\rho)^{1/2})^{-1} R_c(\rho)^{1/2}.$$

Therefore, using the obvious fact that $\|d_{\mathbf{A}}^{1/2} R_c(\rho)^{1/2}\| \leq 1$, we find

$$(4.5) \quad \begin{aligned} \|R_c^V(\rho) \frac{d_{\mathbf{A}}^2}{2mc^2} R_\infty^V(\rho)\| &\leq \frac{\|R_c(\rho)^{1/2}\| \|d_{\mathbf{A}}^{1/2} R_c(\rho)^{1/2}\| \|d_{\mathbf{A}}^{3/2} R_\infty^V(\rho)\|}{2mc^2(1 - \| |V|^{1/2} R_c(\rho)^{1/2} \|^2)} \\ &\leq \frac{\|d_{\mathbf{A}}^{3/2} R_\infty^V(\rho)\|}{mc^2 \sqrt{\rho - |eV|}} = \mathcal{O}(c^{-1/2}), \end{aligned}$$

where in the last estimate we use (3.9). This proves (4.3) and concludes the proof of the theorem. \square

Acknowledgement: I am grateful to Oliver Matte and Thomas Østergaard Sørensen for proofreading the manuscript and helpful discussions. This work has been partially supported by the DFG (SFB/TR12).

APPENDIX A.

Proof of Lemma 3.1. Let $\varphi \in \mathcal{C}_0$ and $f, f\omega^{-1/2} \in \mathfrak{h}$. Using (2.1) and the Cauchy-Schwarz inequality, we get that

$$(A.1) \quad \begin{aligned} \|a(f)\varphi\| &= \left\| \int \bar{f}(k) a(k) \varphi dk \right\| \leq \int \frac{|f(k)|}{\omega(k)^{1/2}} \|\omega(k)^{1/2} a(k) \varphi\| dk \\ &\leq \|f\omega^{-1/2}\|_{\mathfrak{h}} \left(\int \|\omega(k)^{1/2} a(k) \varphi\|^2 dk \right)^{1/2}, \end{aligned}$$

from which (3.1) follows, after using that

$$(A.2) \quad \langle H_f^{1/2} \phi | H_f^{1/2} \psi \rangle = \int \omega(k) \langle a(k) \phi | a(k) \psi \rangle dk, \quad \phi, \psi \in \mathcal{D}(H_f^{1/2}),$$

which is a consequence of Fubini's theorem. Equation (3.2) follows from the fact that

$$(A.3) \quad \|a^\dagger(f)\varphi\|^2 = \langle \varphi | a(f) a^\dagger(f) \varphi \rangle = \|a(f)\varphi\|^2 + \|f\|_{\mathfrak{h}}^2 \|\varphi\|^2,$$

where we used the commutation relations (2.2).

Finally equations (3.3) and (3.4) follow from (3.1) and (3.2), the definitions (2.9) and (2.13), and the Hypothesis 1. \square

We shortly present the proof Lemma 3.3 which can be found in [8, Lemma 4.1]. We will keep the notation of [8].

Proof of Lemma 3.3. To begin with we observe that by means of the pull-through formula we have (see [8, Lemma 3.1]), for $\tilde{H}_f = H_f + E$ and $E > 0$,

$$(A.4) \quad \|T_{1/2}\| := \|[\boldsymbol{\alpha} \cdot \mathbf{A}, \tilde{H}_f^{-1/2}] \tilde{H}_f^{1/2}\| \leq 2d_1/E^{1/2}.$$

We note that a simple computation shows (see [8, Corollary 3.2]), for $R_\eta := 1/(D_{\mathbf{A}} - i\eta)$, $\eta \in \mathbb{R}$, and $\psi \in \mathcal{H}$, that

$$(A.5) \quad R_\eta \tilde{H}_f^{-1/2} \psi = (1 + cR_\eta T_{1/2}) \tilde{H}_f^{-1/2} R_\eta \psi,$$

which implies, using that $\|R_\eta\| = 1/\sqrt{(mc^2)^2 + \eta^2}$, that

$$(A.6) \quad \|\tilde{H}_f^{-1/2} R_\eta \tilde{H}_f^{1/2}\| \leq \frac{1}{\sqrt{(mc^2)^2 + \eta^2}} \frac{1}{1 - 2d_1/(mcE^{1/2})},$$

provided $E > (2d_1/(mc))^2$.

Now, we use (A.6) to obtain a bound for $S_{1/2} := [\text{sgn}(D_{\mathbf{A}}), \tilde{H}_f^{-1/2}] \tilde{H}_f^{1/2}$. Applying the formula (see [6, Lemma VI.5.6]) $\text{sgn}(D_{\mathbf{A}})\varphi = \lim_{\tau \rightarrow \infty} \int_{-\tau}^{\tau} R_\eta \varphi \frac{d\eta}{\pi}$, for $\psi \in \mathcal{H}$, we find, for $\varphi \in \mathcal{D}_0$,

$$(A.7) \quad S_{1/2}\varphi = c \int_{-\infty}^{\infty} [R_\eta, \tilde{H}_f^{-1/2}] \tilde{H}_f^{1/2} \varphi \frac{d\eta}{\pi} = c \int_{-\infty}^{\infty} R_\eta T_{1/2} \tilde{H}_f^{-1/2} R_\eta \tilde{H}_f^{1/2} \varphi \frac{d\eta}{\pi}.$$

From this follows, using (A.6), (A.4), and $\int \|R_\eta\|^2 d\eta/\pi = 1/mc^2$, that

$$(A.8) \quad \|S_{1/2}\| \leq \frac{2d_1}{mcE^{1/2}} \frac{1}{1 - 2d_1/(mcE^{1/2})}, \quad \text{for } E > (2d_1/(mc))^2.$$

An analogous calculation and the fact that $\int \| |D_{\mathbf{A}}|^{1/2} R_\eta \| \|R_\eta\| d\eta/\pi \leq 2\pi/\sqrt{mc^2}$, show that

$$(A.9) \quad \| |D_{\mathbf{A}}|^{1/2} S_{1/2} \| \leq \frac{4\pi d_1}{\sqrt{m}E^{1/2}} \frac{1}{1 - 2d_1/(mcE^{1/2})}, \quad \text{for } E > (2d_1/(mc))^2.$$

Finally, we argue as in [8, Lemma 4.1]: Let $\phi \in \mathcal{D}_0$ and set $\psi := \tilde{H}_f^{1/2} \phi$. We have

$$\begin{aligned} & \text{Re} \langle D_{\mathbf{A}} \tilde{H}_f^{-1/2} \psi \mid \text{sgn}(D_{\mathbf{A}}) \tilde{H}_f^{1/2} \psi \rangle \\ &= \text{Re} \langle (D_{\mathbf{A}} - cT_{1/2}^*) \psi \mid \tilde{H}_f^{-1/2} \text{sgn}(D_{\mathbf{A}}) \tilde{H}_f^{1/2} \psi \rangle \\ &= \text{Re} \langle (D_{\mathbf{A}} - cT_{1/2}^*) \psi \mid (\text{sgn}(D_{\mathbf{A}}) - S_{1/2}) \psi \rangle \\ &\geq \langle |D_{\mathbf{A}}| \psi \mid \psi \rangle - \| |D_{\mathbf{A}}|^{1/2} S_{1/2} |D_{\mathbf{A}}|^{-1/2} \| \| |D_{\mathbf{A}}|^{1/2} \psi \|^2 \\ &\quad - c \|T_{1/2}\| \| |D_{\mathbf{A}}|^{-1/2} \|^2 (1 + \|S_{1/2}\|) \| |D_{\mathbf{A}}|^{1/2} \psi \|^2. \end{aligned}$$

A rough estimate shows, choosing $E > 4(2d_1)^2/(mc)^2$ and using (A.4), (A.8), and (A.9), that

$$(A.10) \quad \operatorname{Re} \langle |D_{\mathbf{A}}|\phi | \tilde{H}_f \phi \rangle \geq \left(1 - \frac{10\pi d_1}{mcE^{1/2}}\right) \| |D_{\mathbf{A}}|^{1/2} \tilde{H}_f^{1/2} \phi \|^2.$$

This finishes the proof of Lemma 3.3. \square

REFERENCES

- [1] Asao Arai. A particle-field Hamiltonian in relativistic quantum electrodynamics. *J. Math. Phys.*, 41(7):4271–4283, 2000.
- [2] Asao Arai. Non-relativistic limit of a Dirac-Maxwell operator in relativistic quantum electrodynamics. *Rev. Math. Phys.*, 15(3):245–270, 2003.
- [3] Asao Arai. Non-relativistic limit of a Dirac polaron in relativistic quantum electrodynamics. *Lett. Math. Phys.*, 77(3):283–290, 2006.
- [4] D. Hasler and I. Herbst. On the self-adjointness and domain of Pauli-Fierz type Hamiltonians. *Rev. Math. Phys.*, 20(7):787–800, 2008.
- [5] F. Hiroshima. Self-adjointness of the Pauli-Fierz Hamiltonian for arbitrary values of coupling constants. *Ann. Henri Poincaré*, 3(1):171–201, 2002.
- [6] Tosio Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [7] Elliott H. Lieb and Michael Loss. Stability of a model of relativistic quantum electrodynamics. *Comm. Math. Phys.*, 228(3):561–588, 2002.
- [8] Oliver Matte and Edgardo Stockmeyer. Exponential localization for a Hydrogen-like atom in relativistic quantum electrodynamics. [arXiv:0904.3083v1 \[math-ph\]](https://arxiv.org/abs/0904.3083v1), 2009.
- [9] Tadahiro Miyao and Herbert Spohn. Spectral analysis of the semi-relativistic Pauli-Fierz hamiltonian. *J. Funct. Anal.*, 256(7):2123 – 2156, 2008.
- [10] Michael Reed and Barry Simon. *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975.
- [11] Barry Simon. *Quantum mechanics for Hamiltonians defined as quadratic forms*. Princeton University Press, Princeton, N. J., 1971. Princeton Series in Physics.
- [12] Bernd Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.

EDGARDO STOCKMEYER, MATHEMATISCHES INSTITUT, LUDWIG-MAXIMILIANS-UNIVERSITÄT, THERESIENSTRASSE 39, D-80333 MÜNCHEN, GERMANY.

E-mail address: stock@math.lmu.de