

# Gradient estimate of an eigenfunction on a compact Riemannian manifold without boundary

Yiqian Shi <sup>\*†</sup> and Bin Xu <sup>\*\*</sup>

**Abstract.** Let  $e_\lambda(x)$  be an eigenfunction with respect to the Laplace-Beltrami operator  $\Delta_M$  on a compact Riemannian manifold  $M$  without boundary:  $\Delta_M e_\lambda = \lambda^2 e_\lambda$ . We show the following gradient estimate of  $e_\lambda$ : for every  $\lambda \geq 1$ , there holds  $\lambda \|e_\lambda\|_\infty / C \leq \|\nabla e_\lambda\|_\infty \leq C\lambda \|e_\lambda\|_\infty$ , where  $C$  is a positive constant depending only on  $M$ .

**Mathematics Subject Classification (2000):** Primary 35P20; Secondary 35J05

**Key Words:** Laplace-Beltrami operator, eigenfunction, gradient estimate

## 1 Introduction

Let  $(M, g)$  be an  $n$ -dimensional compact smooth Riemannian manifold without boundary and  $\Delta_M$  the positive Laplace-Beltrami operator on  $M$ . Let  $L^2(M)$  be the space of square integrable functions on  $M$  with respect to the Riemannian density  $dv(M) = \sqrt{g(x)} dx := \sqrt{\det(g_{ij})} dx$ . Let  $e_1(x), e_2(x), \dots$  be a complete orthonormal basis in  $L^2(M)$  for the eigenfunctions of  $\Delta_M$  such that  $0 = \lambda_0^2 < \lambda_1^2 \leq \lambda_2^2 \leq \dots$  for the corresponding eigenvalues, where  $e_j(x)$  ( $j = 1, 2, \dots$ ) are real valued smooth function on  $M$  and  $\lambda_j$  are nonnegative real numbers. Also, let  $e_j$  denote the projection onto the 1-dimensional space  $\mathbb{C}e_j$ . Thus, an  $L^2$  function  $f$  can be written as  $f = \sum_{j=0}^{\infty} e_j(f)$ , where the partial sum converges in the  $L^2$  norm. Let  $\lambda$  be a positive real number  $\geq 1$ . We define the spectral function  $e(x, y, \lambda)$  and the unit band spectral projection operator  $\chi_\lambda$  as follows:

$$e(x, y, \lambda) := \sum_{\lambda_j \leq \lambda} e_j(x) e_j(y),$$

$$\chi_\lambda f := \sum_{\lambda_j \in (\lambda, \lambda+1]} e_j(f).$$

In 1968, Hörmander [6] obtained a one-term asymptotic expansion of the spectral function of a positive definite elliptic linear operator, whose Laplacian case, also called by the

<sup>\*</sup> Department of Mathematics, University of Science and Technology of China, Hefei 230026 China.

<sup>†</sup> Supported in part by the National Natural Science Foundation of China (No. 10671096). e-mail: yqshi@ustc.edu.cn

<sup>\*\*</sup> The correspondent author is supported in part by the National Natural Science Foundation of China (No. 10601053 and No. 10871184). e-mail: bxu@ustc.edu.cn

local Weyl law, has the expression as follows:

$$e(x, x, \lambda) = \lambda^n / ((4\pi)^{n/2} \Gamma(1 + n/2)) + O(\lambda^{n-1}), \lambda \rightarrow \infty.$$

As a consequence Hörmander proved the uniform estimate of eigenfunctions for all  $x \in M$ :

$$\sum_{\lambda_j \in (\lambda, \lambda+1]} |e_j(x)|^2 \leq C\lambda^{n-1}.$$

We note here that in the whole of this paper  $C$  denotes a positive constant which depends only on  $M$  and may take different values at different places. C. D. Sogge noted in (1.7) of [12] the equivalence of the above estimate with the following  $L^\infty$  estimate of  $\chi_\lambda$ ,

$$\|\chi_\lambda f\|_\infty \leq C\lambda^{(n-1)/2} \|f\|_2. \quad (1)$$

Here  $\|f\|_r$  ( $2 \leq r \leq \infty$ ) means the  $L^r$  norm of the function  $f$  on  $M$ . As noted in Lemma 2.7 by the second author [14], there also holds the similar equivalent relationship between the following two gradient estimates for eigenfunctions and  $\chi_\lambda$ :

$$\sum_{\lambda_j \in (\lambda, \lambda+1]} |\nabla e_j(x)|^2 \leq C\lambda^{n+1} \quad \text{for all } x \in M, \quad (2)$$

and

$$\|\nabla \chi_\lambda f\|_\infty \leq C\lambda^{(n+1)/2} \|f\|_2. \quad (3)$$

Here  $\nabla$  is the Levi-Civita connection on  $M$ . In particular,  $\nabla f = \sum_j g^{ij} \partial f / \partial x_j$  is the gradient vector field of a  $C^1$  function  $f$ , the square of whose length equals  $\sum_{i,j} g^{ij} (\partial f / \partial x_i) (\partial f / \partial x_j)$ , where  $(g^{ij})$  is the inverse of the metric matrix  $g_{ij}$ . By using the wave group, Yu Safarov and D. Vassiliev proved a very general theorem (Theorems 1.8.5 and 1.8.5 in [9]) on the spectral function of a positive definite elliptic linear operator so that the gradient estimate (2) is its immediate corollary. By using the parametrix of the wave operator  $\partial^2 / \partial t^2 + \Delta_M$ , the second author [14] also proved in a slightly different way the Laplacian case of the aforementioned theorems of Yu Safarov and D. Vassiliev, which is also sufficient to deduce the estimate (2). X. Xu [17] applied the maximum principle argument to proving (3), and used this estimate to give a new proof for the Hörmander multiplier theorem. A. Seeger and C. D. Sogge [11] firstly proved the the Hörmander multiplier theorem by using the parametrix of the wave kernel. In this paper we stick at estimating the gradient of a single eigenfunction.

**Theorem** *Let  $e_\lambda(x)$  be an eigenfunction with respect to the positive Laplace-Beltrami operator  $\Delta_M$  on an  $n$ -dimensional compact smooth Riemannian manifold  $M$  without boundary:  $\Delta_M e_\lambda = \lambda^2 e_\lambda$ . Then, for every  $\lambda \geq 1$ , there holds*

$$\lambda \|e_\lambda\|_\infty / C \leq \|\nabla e_\lambda\|_\infty \leq C\lambda \|e_\lambda\|_\infty, \quad (4)$$

where  $C$  is a positive constant depending only on  $M$ .

We can also obtain the above estimates (2) and (3) by Hörmander's local Weyl law and the same argument of Theorem. The details will be given in Section 4. The second author announced a more general estimate for  $\|\nabla^k e_\lambda\|_\infty$  with  $k$  a positive integers in Theorem 5.1 of [15]. However, the method we use in this paper can not be applied to deducing higher derivative estimates. We plan to discuss this question in a future paper.

We conclude the introduction by explaining the organization of this paper. Sections 2-4 contain the proof of Theorem. In Section 2, we show the lower bound of the gradient by the equidistributional property of the nodal set of eigenfunctions. In Section 3 we reduce by rescaling the gradient estimate from above to that of an almost harmonic function. In Section 4, we prove the gradient estimate from above for the almost harmonic function by Yau's gradient estimate and the Green function for compact manifolds with boundary. We also in this section sketch a proof of the above estimates (2) and (3) by the same argument with Theorem, which is different from those in [17] and [14]. In the last section we propose a conjecture that a similar gradient estimate as Theorem holds for a compact Riemannian manifold with boundary.

## 2 Nodal sets of eigenfunctions

The nodal set of an eigenfunction  $e_\lambda$  is the zero set

$$Z_{e_\lambda} := \{x \in M : e_\lambda(x) = 0\}.$$

A connected component of the open set  $M \setminus Z_{e_\lambda}$  is called a nodal domain of the eigenfunction  $e_\lambda$ . The following fact, due to J. Brüning [2], is critical in deducing the lower bound estimate. The reader can also find a proof in English in Theorem 4.1 of S. Zelditch [19].

**Fact 1** *Under the assumption of Theorem, there exists a constant  $C$  such that each geodesic ball of radius  $C/\lambda$  in  $M$  must intersect the nodal set  $Z_{e_\lambda}$  of  $e_\lambda$ .*

**PROOF OF THE LOWER BOUND PART OF THEOREM** Take a point  $x$  in  $M$  such that  $|e_\lambda(x)| = \|e_\lambda\|_\infty$ . Then by Fact 1, there exists a point  $y$  in the ball  $B(x, C/\lambda)$  with center  $x$  and radius  $C/\lambda$  such that  $e_\lambda(y) = 0$ . We may assume  $\lambda$  so large that there exists a geodesic normal chart  $(r, \theta) \in [0, C/\lambda] \times \mathbb{S}^{n-1}(1)$  in the ball  $B(x, C/\lambda)$ . By the mean value theorem, there exists a point  $z$  on the geodesic segment connecting  $x$  and  $y$  such that

$$\left| \frac{\partial e_\lambda}{\partial r}(z) \right| \geq \frac{\lambda}{C} |e_\lambda(x)| = \frac{\lambda}{C} \|e_\lambda\|_\infty.$$

### 3 Eigenfunctions on the wavelength scale

In this section we review quickly the following principle:

*On a small scale comparable to the wavelength  $1/\lambda$ , the eigenfunction  $e_\lambda$  behaves like a harmonic function.*

This principle was developed in H. Donnelly and C. Fefferman [3] [4] and N. S. Nadirashvili [8] and was used extensively there. Our setting of the principle in the following is borrowed from D. Mangoubi [7], where he applied it to studying the geometry of nodal domains of eigenfunctions.

Fix an arbitrary point  $p$  in  $M$  and choose a geodesic normal chart  $(B(p, \delta), x = (x_1, \dots, x_n))$  with center  $p$  and radius  $\delta > 0$  depending only on  $M$ . In this chart, we may identify the ball  $B(p, \delta)$  with the  $n$ -dimensional Euclidean ball  $\mathbb{B}(\delta)$  centered at the origin  $0$ , and think of the eigenfunction  $e_\lambda$  as a function in  $\mathbb{B}(\delta)$ . Our aim is to show the inequality  $|\nabla e_\lambda(p)| \leq C\lambda \|e_\lambda\|_\infty$ . Actually we plan to prove a slightly stronger one

$$\sum_j \left| \frac{\partial e_\lambda}{\partial x_j}(0) \right| \leq C\lambda \|e_\lambda(x)\|_{L^\infty(\mathbb{B}(1/\lambda))}, \quad (5)$$

which we remember that  $g_{ij}(0) = \delta_{ij}$  in the normal chart. The rough idea is to consider the rescaled function  $\phi_\lambda(y) = e_\lambda(y/\lambda)$  in the ball  $\mathbb{B}(1)$  instead of the restriction of the eigenfunction  $e_\lambda(x)$  to the ball  $\mathbb{B}(1/\lambda)$ , where we assume  $\lambda$  so large that  $1/\lambda < \delta/2$ . The above estimate is equivalent to its rescaled version

$$\sum_j \left| \frac{\partial \phi_\lambda}{\partial y_j}(0) \right| \leq C \|\phi_\lambda(y)\|_{L^\infty(\mathbb{B}(1))}. \quad (6)$$

On the other hand, rescaling the eigenfunction equation  $\Delta_M e_\lambda = \lambda^2 e_\lambda$  tells us that the function  $\phi_\lambda$  behaves like a harmonic function. The details are given in what follows.

The eigenfunction equation  $\Delta_M e_\lambda = \lambda^2 e_\lambda$  in  $\mathbb{B}(1/\lambda)$  can be written as

$$-\frac{1}{\sqrt{g}} \sum_{i,j} \partial_{x_i} \left( g^{ij} \sqrt{g} \partial_{x_j} e_\lambda \right) = \lambda^2 e_\lambda.$$

Hence the function  $\phi_\lambda$  satisfies the rescaled equation in  $\mathbb{B}(1)$ ,

$$-\frac{1}{\sqrt{g_\lambda}} \sum_{i,j} \partial_{y_i} \left( g_\lambda^{ij} \sqrt{g_\lambda} \partial_{y_j} \phi_\lambda \right) = \phi_\lambda, \quad (7)$$

where  $g_{ij,\lambda}(y) = g_{ij}(y/\lambda)$ ,  $g_\lambda^{ij}(y) = g^{ij}(y/\lambda)$  and  $\sqrt{g_\lambda}(y) = (\sqrt{g})(y/\lambda)$ . We note that the above equation coincides with

$$\Delta_\lambda \phi_\lambda = \phi_\lambda,$$

where  $\Delta_\lambda$  is the positive Laplace-Beltrami operator with respect to the metric  $g_{ij,\lambda}$  on  $\mathbb{B}(1)$ . We observe that the scaling  $x \mapsto y = \lambda x$  gives an isometry from  $(\mathbb{B}(1/\lambda), \lambda^2 g_{ij})$  onto

$(\mathbb{B}(1), g_{ij,\lambda})$ . Therefore, the open Riemannian manifold  $(\mathbb{B}(1), g_{ij,\lambda})$  has uniformly bounded sectional curvature for all  $\lambda \geq 1$ . This fact will be crucial in next section.

## 4 Estimates from above

We prove (6) in this section by using the notations in Section 3.

*Step 1* Recall that  $\Delta_\lambda \phi_\lambda = \phi_\lambda$  in  $\mathbb{B}(1)$ . We can write the function  $\phi_\lambda$  as the sum  $\phi_\lambda = u_\lambda + v_\lambda$  such that the functions  $u_\lambda$  and  $v_\lambda$  satisfy the following boundary-value problems:

$$\begin{cases} \Delta_\lambda u_\lambda = 0 & \text{in } \mathbb{B}(1), \\ u_\lambda = \phi_\lambda & \text{on } \partial\mathbb{B}(1), \end{cases} \quad \begin{cases} \Delta_\lambda v_\lambda = \phi_\lambda & \text{in } \mathbb{B}(1), \\ v_\lambda = 0 & \text{on } \partial\mathbb{B}(1), \end{cases} \quad (8)$$

*Step 2* By the gradient estimate for harmonic functions in page 21 of R. Schoen and S.-T. Yau [10] and the maximum principle, there holds

$$\sum_j \left| \frac{\partial u_\lambda}{\partial y_j}(0) \right| \leq C \|u_\lambda(y)\|_{L^\infty(\mathbb{B}(1))} \leq C \|\phi_\lambda(y)\|_{L^\infty(\mathbb{B}(1))}.$$

Here we also use the fact that the Ricci curvature tensor of the metrics  $g_{ij,\lambda}$  are uniformly bounded for all  $\lambda \geq 1$ .

*Step 3* Let  $\mathbb{G}(y, z)$  be the Green function on the compact manifold  $(\overline{\mathbb{B}(1)}, g_{ij,\lambda})$  with smooth boundary such that in the sense of distribution  $\Delta_{\lambda,z} \mathbb{G}(y, z)$  equals the Dirac delta function  $\delta_y(z)$  at  $y$  and  $\mathbb{G}(y, z)$  vanishes whenever  $y$  or  $z$  belongs to the boundary  $\partial\mathbb{B}(1)$ . Remember that the sectional curvatures of  $g_{ij,\lambda}$  are uniformly bounded for all  $\lambda \geq 1$ . Carefully checking the explicit construction of the Green kernel in pages 106-113 in T. Aubin [1], we find that

$$|(\nabla_y \mathbb{G})(0, z)| \leq C d(0, z)^{1-n}.$$

Taking gradient at 0 in the Green formula for  $v_\lambda$

$$v_\lambda(y) = \int_{\mathbb{B}(1)} \mathbb{G}(y, z) \phi_\lambda(z) dV(z)$$

gives the estimate  $|(\nabla v_\lambda)(0)| \leq C \|\phi_\lambda\|_{L^\infty(\mathbb{B}(1))}$ . Remember that  $g_{ij,\lambda}(0) = \delta_{ij}$  so that  $|(\nabla v_\lambda)(0)|$  is comparable to  $\sum_j |(\partial v_\lambda / \partial y_j)(0)|$ . The proof is completed.

Following the above arguments, we conclude this section by sketching an alternative proof of the estimates (2) and (3), which were proved by X. Xu [17] and the second author [14] by using the maximum principle and the parametrix of the wave kernel, respectively. We also use the notations above. Take a square integrable function  $f$  on  $M$  and consider the smooth function  $u := \chi_\lambda f$ , and the Poisson equation  $v := \Delta_M u$  in the geodesic normal chart  $\mathbb{B}(1/\lambda)$  centered at a given point  $p$  in  $M$  and with radius  $1/\lambda$ . Rescaling the ball  $\mathbb{B}(1/\lambda)$  with metric  $g_{ij}$  to  $\mathbb{B}(1)$  with metric  $g_{ij,\lambda}$ , and using the  $L^\infty$  estimate

$$\|\chi_\lambda f\|_{L^\infty} \leq C \lambda^{(n-1)/2} \|f\|_2$$

implied by the local Weyl law and the Cauchy-Schwarz inequality, we reduce the question to showing

$$\sum_j \left| \frac{\partial u_\lambda}{\partial y_j}(0) \right| \leq C \|u\|_{L^\infty(\mathbb{B}(1/\lambda))} + C\lambda^{-2} \|v_\lambda\|_{L^\infty(\mathbb{B}(1))},$$

where  $u_\lambda(y) = u(y/\lambda)$  and  $v_\lambda(y) = v(y/\lambda)$ . But the last inequality follows from the arguments in Steps 1-3 above. The proof is completed.

## 5 Compact manifold with boundary

D. Grieser [5] and C. D. Sogge [13] obtained a similar sup norm estimate as (1) for  $\chi_\lambda$  associated with either the Dirichlet or the Neumann Laplacian on a compact Riemannian manifold  $N$  with boundary. X. Xu [16] [18] obtained similar gradient estimates as (3) for both the Dirichlet and the Neumann boundary value problems on  $N$  by using a clever maximum principle and the results of Grieser and Sogge. However, we could not use the maximum principle argument in their papers to deduce more refined gradient estimate for a single eigenfunction on  $N$ . We think that new ideas should be introduced to resolve the following conjecture.

**Conjecture** *Let  $N$  be a compact Riemannian manifold with smooth boundary and  $e_\lambda(x)$  be an eigenfunction of either the Dirichlet or the Neumann Laplacian  $\Delta_N$  on  $N$ :  $\Delta_N e_\lambda = \lambda^2 e_\lambda$ . Then there exists a positive constant  $C$  depending only on  $N$  such that for all  $\lambda \geq 1$  there holds*

$$\lambda \|e_\lambda\|_\infty / C \leq \|\nabla e_\lambda\|_\infty \leq C\lambda \|e_\lambda\|_\infty.$$

We should point out that the lower bound estimate for the Dirichlet eigenfunction  $e_\lambda$ ,

$$\|\nabla e_\lambda\|_\infty \geq C\lambda \|e_\lambda\|_\infty,$$

follows easily from the similar argument in Section 2. We also observe that the upper bound estimate  $|\nabla e_\lambda(x)| \leq C\lambda \|e_\lambda\|_\infty$  for every  $x$  outside the boundary layer  $\{x \in N : d(x, \partial N) \leq C/\lambda\}$  follows from the argument in Sections 3 and 4. The authors will discuss this question in a future paper.

### ACKNOWLEDGEMENTS

The authors would like to thank Professor Qing Chen, Professor Yu Safarov, Professor C. D. Sogge and Professor John Toth for their valuable conversations and comments. The second author would like to express his sincere gratitude to Professor Qing Chen and Professor Xiuxiong Chen for their constant encouragement and forceful prodding.

## References

- [1] T. Aubin, *Some Nonlinear Problems in Riemannian Geometry*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [2] J. Brüning, Über knoten von eigenfunktionen des Laplace-Beltrami operators, *Math. Z.* **158** (1975), 15-21.
- [3] H. Donnelly and C. Fefferman, Nodal sets of eigenfunctions on Riemannian manifolds, *Invent. Math.* **93** (1988) no. 1, 161-183.
- [4] H. Donnelly and C. Fefferman, Growth and geometry of eigenfunctions of the Laplacian, *Analysis and partial differential equations*, Lecture Notes in Pure and Appl. Math., vol. 122, Dekker, NY, 1990, pp. 635-655.
- [5] D. Grieser, Uniform bounds for eigenfunctions of the laplacian on manifolds with boundary, *Comm. Partial Differential Equations*, **27** (2002) no. 7-8, 1283-1299.
- [6] L. Hörmander, The Spectral Function of an Elliptic Operator. *Acta Math.* **88** (1968), 341-370.
- [7] D. Mangoubi, Local asymmetry and the inner radius of nodal domain, *Comm. Partial Differential Equations*, **33** (2008), no. 7-9, 1611–1621.
- [8] N. S. Nadirashvili, Metric properties of eigenfunctions of the Laplace operator on manifolds, *Ann. Inst. Fourier (Grenoble)* **41** (1991), no. 1, 259-265.
- [9] Yu Safarov and D. Vassiliev, *The Asymptotic Distribution of Eigenvalues of Partial Differential Operators*, Translations of Mathematical Monograph, **155**, American Mathematical Society, Providence, RI, 1997.
- [10] Y. Schoen, Y. and S.-T. Yau, *Lectures on Differential Geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994.
- [11] A. Seeger and C. D. Sogge, On the boundedness of functions of pseudo-differential operators on a compact manifolds, *Duke Math. J.* **59** (1989), 709-736.
- [12] C. D. Sogge, On the convergence of Riesz means of compact manifolds, *Ann. Math.* **126** (1987), 439-447.
- [13] C. D. Sogge, Eigenfunction and Bochner-Riesz estimates on manifolds with boundary, *Mathematical Research Letters*, **9** (2002), 205-216.
- [14] B. Xu, Derivatives of the spectral function and Sobolev norms of eigenfunctions on a closed Riemannian manifold, *Ann. Glob. Anal. Geom.* **26** (2004), no. 3, 231-252.
- [15] B. Xu, Asymptotic behavior of  $L^2$ -normalized eigenfunctions of the Laplace-Beltrami operator on a closed Riemannian manifold, *Harmonic Analysis and its Applications*, 99–117, Yokohama Publ., Yokohama, 2006.
- [16] X. Xu, Eigenfunction estimates on compact manifolds with boundary and Hörmander multiplier theorem, Ph. D. Thesis, Johns Hopkins University, May 2004.

- [17] X. Xu, New proof of the Hörmander multiplier theorem on compact manifolds without boundary, *Proc. Amer. Math. Soc.* **135** (2007), no. 5, 1585-1595.
- [18] X. Xu, Gradient estimates for eigenfunctions of compact manifolds with boundary and the Hörmander multiplier theorem. *Forum Mathematicum* **21** (2009), no. 3, 455-476.
- [19] S. Zelditch, Local and global analysis of eigenfunctions on Riemannian manifolds, *Handbook of geometric analysis*, no. 1, 545-658, Adv. Lect. Math. **7**, Int. Press, Somerville, MA, 2008.