Dynamics of the Universal Area-Preserving Map Associated with Period Doubling: Hyperbolic Sets

Denis Gaidashev, Tomas Johnson

Department of Mathematics, Uppsala University, Box 480, 751 06 Uppsala,

E-mail: gaidash@math.uu.se, tomas.johnson@math.uu.se

Abstract. It is known that the famous Feigenbaum-Coullet-Tresser period doubling universality has a counterpart for area-preserving maps of \mathbb{R}^2 . A renormalization approach has been used in (Eckmann et al 1982) and (Eckmann et al 1984) in a computer-assisted proof of existence of a "universal" area-preserving map F_* — a map with orbits of all binary periods $2^k, k \in \mathbb{N}$. In this paper, we consider maps in some neighbourhood of F_* and study their dynamics.

We first demonstrate that the map F_* admits a "bi-infinite heteroclinic tangle": a sequence of periodic points $\{z_k\}$, $k \in \mathbb{Z}$, $|z_k| \stackrel{k \to \infty}{\longrightarrow} 0$, $|z_k| \stackrel{k \to -\infty}{\longrightarrow} \infty$, (1) whose stable and unstable manifolds intersect transversally; and, for any $N \in \mathbb{N}$,

$$|z_k| \stackrel{k \to \infty}{\longrightarrow} 0, \quad |z_k| \stackrel{k \to -\infty}{\longrightarrow} \infty,$$
 (1)

a compact invariant set on which F_* is homeomorphic to a topological Markov chain on the space of all two-sided sequences composed of N symbols. A corollary of these results is the existence of unbounded and oscillating orbits.

We also show that the third iterate for all maps close to F_* admits a horseshoe. We use distortion tools to provide rigorous bounds on the Hausdorff dimension of the associated locally maximal invariant hyperbolic set:

$$0.7673 \ge \dim_H(\mathcal{C}_F) \ge \varepsilon \approx 0.00013 \, e^{-7499}$$
.

AMS classification scheme numbers: 37E20, 37F25, 37D05, 37D20, 37C29, 37A05, 37G15, 37M99

Submitted to: Nonlinearity

1. Introduction

Following the pioneering discovery of the Feigenbaum-Coullet-Tresser period doubling universality in unimodal maps (Feigenbaum 1978), (Feigenbaum 1979), (Tresser and Coullet 1978) universality has been demonstrated to be a rather generic phenomenon in dynamics.

To prove universality one usually introduces a *renormalization* operator on a functional space, and demonstrates that this operator has a hyperbolic fixed point.

Such renormalization approach to universality has been very successful in onedimensional dynamics, and has led to explanation of universality in unimodal maps (Epstein 1986), (Epstein 1989), (Lyubich 1999), critical circle maps (de Faria 1992, de Faria 1999), (Yampolsky 2002), (Yampolsky 2003) and holomorphic maps with a Siegel disk (McMullen 1998), (Yampolsky 2007), (?).

Universality has been abundantly observed in higher dimensions, in particular, in two and more dimensional dissipative systems (cf. (Collet et al 1980), (Sparrow 1982)), in area-preserving maps, both as the period-doubling universality (Derrida and Pomeau 1980), (Helleman 1980), (Benettin et al 1980), (Collet et al 1981), (Eckmann et al 1982), (Eckmann et al 1984), (Gaidashev and Koch 2008) and as the universality associated with the break-up of invariant surfaces (Shenker and Kadanoff 1982), (MacKay 1983), (MacKay 1982), (Mehr and Escande 1984), and in Hamiltonian flows (Escande and Doveil 1981), (Abad et al 2000), (Abad et al 1998), (Koch 2002), (Koch 2004), (Koch 2004), (Gaidashev and Koch 2004), (Gaidashev 2005), (Kocić 2005).

It has been established that the universal behaviour in dissipative and conservative higher dimensional systems is fundamentally different. The case of the dissipative systems is often reducible to the one-dimensional Feigenbaum-Coullet-Tresser universality ((Collet et al 1980), (de Carvalho et al 2005), (Lyubich and Martens 2008)). Universality for highly dissipative Hénon like maps

$$F(x, u) = (f(x) - \epsilon(x, u), x),$$

where f is a unimodal map, ϵ - sufficiently small together with its derivatives, has been demonstrated in (de Carvalho *et al* 2005). Specifically, it has been shown that these maps are in the Feigenbaum-Coullet-Tresser universality class, and that the degenerate map

$$F^*(x, u) = (f^*(x), x),$$

where f^* is the Feigenbaum-Coullet-Tresser fixed point, is a renormalization fixed point in that class. The authors of (de Carvalho *et al* 2005) have also constructed a Cantor set (an attractor) for infinitely renormalizable maps, and demonstrated that for such Hénon-like maps universality coexists with non-rigidity: the Hölder exponent of the conjugacy between the actions of any two infinitely renormalizable maps with nonequal average Jacobians on their Cantor sets has an upper bound less than 1.

The case of area-preserving maps seems to be very different, and at present there is no deep understanding of universality in conservative systems, other than in the "trivial" case of the universality for systems "near integrability" (Koch 2002), (Koch 2004), (Gaidashev 2005), (Kocić 2005), (Khanin *et al* 2007).

An infinite period-doubling cascade in families of area-preserving maps was observed by several authors in early 80's (Derrida and Pomeau 1980), (Helleman 1980), (Benettin *et al* 1980), (Bountis 1981), (Collet *et al* 1981). The period-doubling

phenomenon can be illustrated with the area-preserving Hénon family (cf. (Bountis 1981)):

$$H_a(x, u) = (-u + 1 - ax^2, x).$$

Maps H_a have a fixed point $((-1+\sqrt{1+a})/a, (-1+\sqrt{1+a})/a)$ which is stable for -1 < a < 3. When $a_1 = 3$ this fixed point becomes unstable, at the same time an orbit of period two is born with $H_a(x_{\pm}, x_{\mp}) = (x_{\mp}, x_{\pm}), x_{\pm} = (1 \pm \sqrt{a-3})/a$. This orbit, in turn, becomes unstable at $a_2 = 4$, giving birth to a period 4 stable orbit. Generally, there exists a sequence of parameter values a_k , at which the orbit of period 2^{k-1} turns unstable, while at the same time a stable orbit of period 2^k is born. The parameter values a_k accumulate on some a_{∞} . The crucial observation is that the accumulation rate

$$\lim_{k \to \infty} \frac{a_k - a_{k-1}}{a_{k+1} - a_k} = 8.721... \tag{2}$$

is universal for a large class of families, not necessarily Hénon.

Furthermore, the 2^k periodic orbits scale asymptotically with two scaling parameters

$$\lambda = -0.249..., \quad \mu = 0.061...$$
 (3)

To explain how orbits scale with λ and μ we will follow (Bountis 1981). Consider an interval (a_k, a_{k+1}) of parameter values in a "typical" family F_a . For any value $\alpha \in (a_k, a_{k+1})$ the map F_α possesses a stable periodic orbit of period 2^k . We fix some α_k within the interval (a_k, a_{k+1}) in some consistent way; for instance, by requiring that the restriction of $F_{\alpha_k}^{2^k}$ to a neighbourhood of a stable periodic point in the 2^k -periodic orbit is conjugate, via a diffeomorphism H_k , to a rotation with some fixed rotation number r. Let p'_k be some unstable periodic point in the 2^{k-1} -periodic orbit, and let p_k be the further of the two stable 2^k -periodic points that bifurcated from p'_k . Denote with $d_k = |p'_k - p_k|$, the distance between p_k and p'_k . The new elliptic point p_k is surrounded by invariant ellipses; let c_k be the distance between p_k and p'_k in the direction of the minor semi-axis of an invariant ellipse surrounding p_k , see Figure 1. Then,

$$\frac{1}{\lambda} = -\lim_{k \to \infty} \frac{d_k}{d_{k+1}}, \quad \frac{\lambda}{\mu} = -\lim_{k \to \infty} \frac{\rho_k}{\rho_{k+1}}, \quad \frac{1}{\lambda^2} = \lim_{k \to \infty} \frac{c_k}{c_{k+1}},$$

where ρ_k is the ratio of the smaller and larger eigenvalues of $DH_k(p_k)$.




Figure 1. The geometry of the period doubling. p_k is the further elliptic point that has bifurcated from the hyperbolic point p'_k .

This universality can be explained rigorously if one shows that the *renormalization* operator

$$R[F] = \Lambda_F^{-1} \circ F \circ F \circ \Lambda_F, \tag{4}$$

where Λ_F is some F-dependent coordinate transformation, has a fixed point, and the derivative of this operator is hyperbolic at this fixed point.

It has been argued in (Collet *et al* 1981) that Λ_F is a diagonal linear transformation. Furthermore, such Λ_F has been used in (Eckmann *et al* 1982) and (Eckmann *et al* 1984) in a computer assisted proof of existence of a reversible renormalization fixed point F_* and hyperbolicity of the operator R.

An exploration of a possible analytic machinery has been undertaken in (Gaidashev and Koch 2008) where it has been demonstrated that the fixed point F_* is very close, in some appropriate sense, to an area-preserving Hénon-like map

$$H^*(x,u) = (\phi(x) - u, x - \phi(\phi(x) - u)), \tag{5}$$

where ϕ solves the following one-dimensional problem of non-Feigenbaum type:

$$\phi(y) = \frac{2}{\lambda}\phi(\phi(\lambda y)) - y. \tag{6}$$

In this paper we will study the dynamics of the renormalization fixed point F_* and maps in some neighbourhood of F_* . We would like to emphasize that the preservation of area for such maps leads to one important difference between the case at hand and the highly dissipative case of Hénon-like maps: the former maps have no obvious invariant subsets in their domain. In particular, this means, that the construction of invariant Cantor sets carried out in (de Carvalho *et al* 2005), which is using existence of invariant subsets in a crucial way, is not readily applicable to the case of area-preserving maps.

To construct hyperbolic sets, we will use the idea of covering relations (see, e.g. (Zgliczyński and Gidea 2004, Zgliczyński 2009)) in rigorous computations. The Hausdorff dimension of the hyperbolic sets will be estimated with the help of the Duarte Distortion Theorem (see, e.g. (Duarte 2000)) which enables one to use the distortion of a Cantor set to ultimately find bounds on the dimension.

The structure of the paper is as follows: we begin by recalling the basic properties of area-preserving reversible maps in Section 2. In Section 3 we introduce some notation and recall some standard definitions from hyperbolic dynamics. In Section 4 we study the domain of analyticity of maps in a neighbourhood of the renormalization fixed point, and analytic continuation of the renormalization fixed point. Section 5 consists of the statements of our main theorems. In Section 6, we recall the definition and the main properties of the covering relations. In Section 7 we prove that any area-preserving reversible map in a neighbourhood of the renormalization fixed point has a transversal homoclinic orbit in its domain of analyticity. In Section 8 we construct a heteroclinic tangle for the renormalization fixed point. From its existence the existence of unbounded, and oscillating trajectories follow. In Section 9 we recall the Duarte Distortion Theorem. In Sections 10 and 11 we pr ove that the third iterate of any area-preserving reversible map in a neighbourhood of the renormalization fixed point has a horseshoe, and compute bounds on its Hausdorff dimension using the Duarte distortion theorem.

In a satellite paper (Gaidashev and Johnson 2009), we prove that *infinitely* renormalizable maps in the neighbourhood of existence of the hyperbolic set for the third iterate also admit a "stable" set. This set is a bounded invariant set, such that

the maximal Lyapunov exponent for the third iterate is zero. In (Gaidashev and Johnson 2009) we provide an upper bound on the Hausdorff dimension of the stable set, and prove that the Hausdorff dimension is constant for all maps in some subset of infinitely renormalizable maps.

2. Renormalization for area-preserving reversible maps

An "area-preserving map" will mean an exact symplectic diffeomorphism of a subset of \mathbb{R}^2 onto its image.

Recall, that an area-preserving map can be uniquely specified by its generating function S:

$$\begin{pmatrix} x \\ -S_1(x,y) \end{pmatrix} \stackrel{F}{\mapsto} \begin{pmatrix} y \\ S_2(x,y) \end{pmatrix}, \quad S_i \equiv \partial_i S.$$
 (7)

Furthermore, we will assume that F is reversible, that is

$$T \circ F \circ T = F^{-1}$$
, where $T(x, u) = (x, -u)$. (8)

For such maps it follows from (7) that

$$S_1(y,x) = S_2(x,y) \equiv s(x,y), \tag{9}$$

and

$$\begin{pmatrix} x \\ -s(y,x) \end{pmatrix} \stackrel{F}{\hookrightarrow} \begin{pmatrix} y \\ s(x,y) \end{pmatrix}. \tag{10}$$

It is this "little" s that will be referred to below as "the generating function". It follows from (9) that s_1 is symmetric. If the equation -s(y,x) = u has a unique differentiable solution y = y(x,u), then the derivative of such a map F is given by the following formula:

$$DF(x,u) = \begin{bmatrix} -\frac{s_2(y(x,u),x)}{s_1(y(x,u),x)} & -\frac{1}{s_1(y(x,u),x)} \\ s_1(x,y(x,u)) - s_2(x,y(x,u)) \frac{s_2(y(x,u),x)}{s_1(y(x,u),x)} & -\frac{s_2(x,y(x,u))}{s_1(y(x,u),x)} \end{bmatrix}. \quad (11)$$

We will now derive an equation for the generating function of the renormalized map $\Lambda_F \circ F \circ F \circ \Lambda_F^{-1}$.

Applying a reversible F twice we get

$$\begin{pmatrix} x' \\ -s(z',x') \end{pmatrix} \overset{F}{\mapsto} \begin{pmatrix} z' \\ s(x',z') \end{pmatrix} = \begin{pmatrix} z' \\ -s(y',z') \end{pmatrix} \overset{F}{\mapsto} \begin{pmatrix} y' \\ s(z',y') \end{pmatrix}.$$

It has been argued in (Collet et al 1981) that

$$\Lambda_F(x, u) = (\lambda x, \mu u).$$

We therefore set $(x', y') = (\lambda x, \lambda y), z'(\lambda x, \lambda y) = z(x, y)$ to obtain:

$$\begin{pmatrix} x \\ -\frac{1}{\mu}s(z,\lambda x) \end{pmatrix} \stackrel{\Lambda_F}{\mapsto} \begin{pmatrix} \lambda x \\ -s(z,\lambda x) \end{pmatrix} \stackrel{F \circ F}{\mapsto} \begin{pmatrix} \lambda y \\ s(z,\lambda y) \end{pmatrix} \stackrel{\Lambda_F^{-1}}{\mapsto} \begin{pmatrix} y \\ \frac{1}{\mu}s(z,\lambda y) \end{pmatrix}, \quad (12)$$

where z(x, y) solves

$$s(\lambda x, z(x,y)) + s(\lambda y, z(x,y)) = 0. \tag{13}$$

If the solution of (13) is unique, then z(x,y) = z(y,x), and it follows from (12) that the generating function of the renormalized F is given by

$$\tilde{s}(x,y) = \mu^{-1} s(z(x,y), \lambda y). \tag{14}$$

One can fix a set of normalization conditions for \tilde{s} and z which serve to determine scalings λ and μ as functions of s. For example, the normalization

$$s(1,0) = 0$$

is reproduced for \tilde{s} as long as

$$z(1,0) = z(0,1) = 1.$$

In particular, this implies that

$$s(\lambda, 1) + s(0, 1) = 0.$$

Furthermore, the condition

$$\partial_1 s(1,0) = 1 \tag{15}$$

is reproduced as long as

$$\mu = \partial_1 z(1,0).$$

We will now summarize the above discussion in the following definition of the renormalization operator acting on generating functions originally due to the authors of (Eckmann $et\ al\ 1982$) and (Eckmann $et\ al\ 1984$):

Definition 2.1

$$\mathcal{R}_{EKW}[s](x,y) = \mu^{-1}s(z(x,y),\lambda y), \tag{16}$$

where

$$0 = s(\lambda x, z(x, y)) + s(\lambda y, z(x, y)), \tag{17}$$

$$0 = s(\lambda, 1) + s(0, 1)$$
 and $\mu = \partial_1 z(1, 0)$. (18)

Definition 2.2 The Banach space of functions $s(x,y) = \sum_{i,j=0}^{\infty} c_{ij} x^i y^j$, analytic on a bi-disk

$$|x - 0.5| < \rho, |y - 0.5| < \rho,$$

for which the norm

$$||s||_{\rho} = \sum_{i,j=0}^{\infty} |c_{ij}| \rho^{i+j}$$

is finite, will be referred to as $A(\rho)$. $A_s(\rho)$ will denote its symmetric subspace $\{s \in A(\rho) : s_1(x,y) = s_1(y,x)\}.$

As we have already mentioned, the following has been proved with the help of a computer in (Eckmann *et al* 1982) and (Eckmann *et al* 1984):

Theorem 2.3 There exist a polynomial $s_a \in \mathcal{A}_s(\rho)$ and a ball $\mathcal{B}_r(s_a) \subset \mathcal{A}_s(\rho)$, $r = 6.0 \times 10^{-7}$, $\rho = 1.6$, such that the operator \mathcal{R}_{EKW} is well-defined, analytic and compact on \mathcal{B}_r .

Furthermore, its derivative $D\mathcal{R}_{EKW}|_{\mathcal{B}_r}$ has exactly two eigenvalues δ_1 and δ_2 of modulus larger than 1, while

$$\operatorname{spec}(D\mathcal{R}_{EKW}|_{\mathcal{B}_r}) \setminus \{\delta_1, \delta_2\} \subset \{z \in \mathbb{C} : |z| \le \nu < 1\}.$$

Finally, there is an $s^* \in \mathcal{B}_r$ such that

$$\mathcal{R}_{EKW}[s^*] = s^*.$$

The scalings λ_* and μ_* corresponding to the fixed point s^* satisfy

$$\lambda_* \in [-0.24887681, -0.24887376],\tag{19}$$

$$\mu_* \in [0.061107811, 0.061112465].$$
 (20)

Remark 2.4 The radius of the contracting part of the spectrum spec($D\mathcal{R}_{EKW}(s_*) \setminus \{\delta_1, \delta_2\}$ has been estimated in (Eckmann et al 1984) to be $\nu = 0.8$.

It follows from the above theorem that there exist a codimension 2 local stable manifold $W^s_{loc}(s^*) \subset \mathcal{B}_r$.

Definition 2.5 A reversible map F of the form (10) such that $s \in W^s_{loc}(s^*)$ is called infinitely renormalizable. The set of all reversible infinitely renormalizable maps is denoted by \mathbf{W} .

3. Some notation and definitions

We will use the following notation for the sup norm of a function h and a transformation H defined on some set $S \subset \mathbb{R}^2$ or \mathbb{C}^2 :

$$|h|_{\mathcal{S}} \equiv \sup_{(x,u)\in\mathcal{S}} \{|h|\},$$
 (21)

$$|H|_{\mathcal{S}} \equiv \max\{ \sup_{(x,u)\in\mathcal{S}} \{|\mathcal{P}_x H|\}, \sup_{(x,u)\in\mathcal{S}} \{|\mathcal{P}_u H|\}\},$$
(22)

where \mathcal{P}_x and \mathcal{P}_u are projections on the corresponding components.

We will also use the notation $|\cdot|$ for the l_2 norm for vectors in \mathbb{R}^2 .

The interval enclosures of λ_* and μ_* will be denoted

$$\lambda_* \equiv [\lambda_-, \lambda_+]; \quad \lambda_- = -0.24887681, \quad \lambda_+ = -0.24887376,$$
 (23)

$$\mu_* \equiv [\mu_-, \mu_+]; \quad \mu_- = 0.061107811, \quad \mu_+ = 0.061112465.$$
 (24)

The corresponding interval enclosure for the linear map Λ_* will be denoted Λ_* ; if $(x, u) \in \mathbb{C}^2$, then

$$\mathbf{\Lambda}_*(x, u) \equiv \left\{ (\lambda x, \mu u) \in \mathbb{C}^2 : \lambda \in \mathbf{\lambda}_*, \mu \in \boldsymbol{\mu}_* \right\}. \tag{25}$$

The bound on the fixed point generating function s^* will be called s^* :

$$\mathbf{s}^* \equiv \left\{ s \in \mathcal{A}_s(\rho) : \|s - s_{\mathbf{a}}\|_{\rho} \le r = 6.0 \times 10^{-7} \right\},\tag{26}$$

while the bound on the renormalization fixed point F_* will be referred to as F_* :

$$\mathbf{F}_* \equiv \{F : (x, -s(y, x)) \mapsto (y, s(x, y)) : s \in \mathbf{s}^*\},\tag{27}$$

where s_a is as in Theorem 2.3; the third iterate of this bound will be referred to as \mathbf{G}_* . With $D\mathbf{G}_*: p \mapsto [D\mathbf{G}_*(p)]$ we denote an interval matrix valued function such that

$$[DG(p)]_{ij} \in [D\mathbf{G}_*(p)]_{ij}, \text{ for all } G \in \mathbf{G}_*, p \in \mathcal{D}_3,$$

where \mathcal{D}_3 is the domain of \mathbf{G}_* , and the bound on the operator norm of DG for $G = F \circ F \circ F$, $F \in \mathbf{F}_*$ on a set \mathcal{S} will be denoted

$$||D\mathbf{G}_*||_{\mathcal{S}} \equiv \sup_{F \in \mathbf{F}_*} \{||D(F \circ F \circ F)||_{\mathcal{S}}\}.$$

Given a non-empty open set $\mathcal{D} \subset \mathbb{C}^n$ we will denote by $\mathcal{O}_R(\mathcal{D})$ the set of reversible area-preserving maps $F: \mathcal{D} \mapsto \mathbb{C}^n$, analytic on \mathcal{D} .

We will proceed with a collection of classical notations (see, eg, (Katok and Hasselblatt 1995)) relevant to our following discussion.

Definition 3.1 (Hyperbolic set) Let \mathcal{M} be a smooth manifold, and let F be a diffeomorphism of an open subset $\mathcal{U} \subset \mathcal{M}$ onto its image.

A set C is called hyperbolic for the map F if there is a Riemannian metric on a neighbourhood U of C, and $\beta < 1 < \delta$, such that for any $p \in C$ and $n \in \mathbb{N}$ the tangent space $T_{F^n(p)}U$ admits a decomposition in two invariant subspaces:

$$T_{F^n(p)}\mathcal{M} = E_n^+ \oplus E_n^-, \quad DF(F^n(p))E_n^{\pm} = E_{n+1}^{\pm},$$

on which the sequence of differentials is hyperbolic:

$$\|DF(F^n(p))|_{E_n^-}\|<\beta,\quad \|DF^{-1}(F^n(p))|_{E_{n+1}^+}\|<\delta^{-1}.$$

Definition 3.2 (Locally maximal hyperbolic set) Let C be a hyperbolic set for $F : \mathcal{U} \to \mathcal{M}$. If there is a neighbourhood V of C such that $C = \cap_{n \in \mathbb{Z}} F^n(\bar{V})$, then C is called locally maximal.

Definition 3.3 (<u>Bernoulli shift</u>) Let $\{0, 1, ..., N-1\}^{\mathbb{Z}}$ be the space of all two-sided sequences of N symbols:

$$\{0, 1, \dots, N-1\}^{\mathbb{Z}} = \{\omega = (\dots, \omega_{-1}, \omega_0, \omega_1, \dots) : \omega_i \in \{0, 1, \dots, N-1\}, i \in \mathbb{Z}\},\$$

Define the Bernoulli shift on $\{0, 1, \dots, N-1\}^{\mathbb{Z}}$ as

$$\sigma_N(\omega) = \omega', \quad \omega'_n = \omega_{n+1}.$$

Definition 3.4 (<u>Topological Markov chain</u>) Let $A = (a_{i,j})_{i,j=0}^{N-1}$ be an $N \times N$ matrix whose entries are either 0 or 1. Let

$$\{0,1,\ldots,N-1\}_A^{\mathbb{Z}} = \{\omega \in \{0,1,\ldots,N-1\}^{\mathbb{Z}} : a_{\omega_n,\omega_{n+1}} = 1, n \in \mathbb{Z}\}.$$

The restriction

$$\sigma_N|_{\{0,1,\dots,N-1\}_A^{\mathbb{Z}}} \equiv \sigma_A$$

is called a topological Markov chain determined by A.

Definition 3.5 (Homoclinic and heteroclinic points) Let $F: \mathcal{X} \mapsto \mathcal{X}$ be a homeomorphism on a metric space (\mathcal{X}, d) . A point $x \in \mathcal{X}$ is said to be homoclinic to the point $y \in \mathcal{X}$ if

$$\lim_{|n| \to \infty} d(F^n(x), F^n(y)) = 0.$$

A point x is said to be heteroclinic to points y_1 and y_2 if

$$\lim_{n \to \infty} d(F^n(x), F^n(y_1)) = \lim_{n \to -\infty} d(F^n(x), F^n(y_2)) = 0.$$

In this case we will also say that there exists a heteroclinic orbit between points y_1 and y_2 .

If \mathcal{M} is a differentiable manifold, $F \in \text{Diff}^1(\mathcal{M})$ and $x \in \mathcal{M}$ is a hyperbolic fixed point of F, then we say that $q \in \mathcal{M}$ is a transversal homoclinic point to x if it is a point of transversal intersection of the stable and unstable manifolds of x.

Definition 3.6 Let \mathcal{X} be a metric space. If $\mathcal{S} \subset \mathcal{X}$, and $d \in [0, \infty)$, the d-dimensional Hausdorff content of \mathcal{S} is defined as

$$C_d^H(\mathcal{S}) = \inf \left\{ \sum_i \operatorname{diam}(\mathcal{S}_i)^d : \{\mathcal{S}_i\} \text{ is a cover of } \mathcal{S} \right\}.$$
 (28)

The Hausdorff dimension of S is defined as

$$\dim_H(S) = \inf \{ d \ge 0 : C_d^H(S) = 0 \}.$$
 (29)

4. Domain of analyticity of F_{*} and G_{*}

F is defined implicitly by the generating function s, its domain is given as in (Eckmann et al 1984):

$$\mathcal{D}_s = \{(x, y) \in \mathbb{C}^2 : |x - 0.5| < 1.6, |y - 0.5| < 1.6\}. \tag{30}$$

To find the domain of $F \in \mathbf{F}_*$, we note that its second argument is equal to -s(y, x), for some $s \in \mathbf{s}^*$ (see (10)). Thus, the domain of F, \mathcal{D} , is given by:

$$\mathcal{D} = \{(x, u) \in \mathbb{C}^2 : u = -s(y, x), \ s \in \mathbf{s}^*(y, x), \ |x - 0.5| < 1.6, \ |y - 0.5| < 1.6\}.$$
 (31)

We denote by

$$\tilde{\mathcal{D}} = \{ (x, u) \in \mathcal{D} : \Im x = \Im u = 0 \}, \tag{32}$$

the real slice of \mathcal{D} .

To solve nonlinear equations on the computer we use the interval Newton operator, see e.g. (Neumaier 1990).

Definition 4.1 (Interval Newton Operator) Let $F : \mathcal{D} \subseteq \mathbb{R}^n \to \mathbb{R}^n$, and let $D\mathbf{F}$ be an interval matrix valued function such that $[DF(x)]_{ij} \in [D(\mathbf{F})(x)]_{ij}$, for all $x \in \mathcal{D}$. Let $\mathbf{X} \subset \mathcal{D}$ be a Cartesian product of finite intervals, $\hat{x} \in \mathbf{X}$, and assume that if $A \in D\mathbf{F}(\mathbf{X})$, then A is non-singular. We define the interval Newton operator as:

$$N(F, \mathbf{X}, \hat{x}) = \hat{x} - (D\mathbf{F})^{-1}(\mathbf{X})\mathbf{F}(\hat{x}).$$

The main properties of N is that if $N(F, \mathbf{X}, \hat{x}) \subset \text{int} \mathbf{X}$, then there exists a unique solution to F(x) = 0 in \mathbf{X} , which is contained in $N(F, \mathbf{X}, \hat{x})$, and if $N(F, \mathbf{X}, \hat{x}) \cap \mathbf{X} = \emptyset$, then there is no solution to F(x) = 0 in \mathbf{X} .

Lemma 4.2 There exists a non-empty open set $\bar{\mathcal{D}}$,

$$\mathcal{P}_x \bar{\mathcal{D}} \subset \{x \in \mathbb{R} : |x - 0.5| < 1.6\},\$$

such that for every $(x,u) \in \bar{\mathcal{D}}$ there exists a unique solution of the interval Newton operator of the function $h_{(x,u)}(y) = u + s(y,x)$, $s \in \mathbf{s}^*(y,x)$, that satisfies |y - 0.5| < 1.6.

Proof. Set $\mathbf{Y} = \{y \in \mathbb{R} : |y - 0.5| < 1.6\}$. Given (x, u), let $N(h_{(x,u)}, \mathbf{Y}, \hat{y})$ be the interval Newton operator (for some appropriately chosen $\hat{y} \in \mathbf{Y}$). We have verified that there exists a non-empty set $\bar{\mathcal{D}}$, such that for all $(x, u) \in \bar{\mathcal{D}}$

$$N(h_{(x,u)}, \mathbf{Y}, \hat{y}) \subset \mathbf{Y}.$$

This verification is implemented in the program findDomain of (Progs1 2009) It follows, see e.g. (Neumaier 1990), that there is a unique y, such that (x, y) is in the real slice of \mathcal{D}_s and $h_{(x,u)}(y) = 0$. Thus, y is defined as a function of (x, u), and by (31) $(x, u) \in \tilde{\mathcal{D}}$.

Clearly, $\bar{\mathcal{D}} \subset \tilde{\mathcal{D}}$.

The generating function s is analytic on a bi-disk, preferably F should have a similar property. From the above construction, we can at least show that F is defined on a complex neighbourhood of $\bar{\mathcal{D}}$ from the above Lemma 4.2:

Lemma 4.3 \mathcal{D} contains an open complex neighbourhood of the set $\bar{\mathcal{D}}$.

Proof. $N(h_{(x,u)},y)$ is only well defined if $\mathbf{s}^*_1(y,x) \neq 0$, so that the condition of the implicit function theorem is automatically satisfied on the solution set of $N(h_{(x,u)}, \mathbf{Y}, \hat{y})$. Since \mathbf{s}^*_1 is analytic on the bi-disk $\{|x-0.5|<1.6\}\times\{|y-0.5|<1.6\}$, there is an open neighbourhood in \mathbb{C}^2 of the solution set where $\mathbf{s}^*_1(y,x)\neq 0$. It follows from the implicit function theorem that \mathbf{F}_* is analytic on this neighbourhood.

Generally, we will denote the third iterate of a map $F \in \mathcal{O}_R(\mathcal{D})$ as G. The domain of G, \mathcal{D}_3 , is given by

$$\mathcal{D}_3 = F^{-2}(\mathcal{D}'' \cap \mathcal{D}),\tag{33}$$

where

$$\mathcal{D}'' = F(\mathcal{D}' \cap \mathcal{D}), \quad \mathcal{D}' = F(\mathcal{D}).$$

Using the program findDomain, we have verified that the real slice $\tilde{\mathcal{D}}_3$ of \mathcal{D}_3 is an open non-empty set. Approximations of $\tilde{\mathcal{D}}$ and $\tilde{\mathcal{D}}_3$ are shown in Figure 2. We note that by using the renormalization equation $F_* = R[F_*]$, F_* has, for any k, an analytic continuation to domains

$$\mathcal{D}^k \equiv \Lambda_*^{-1}(F_*^{-1}(F_*(\mathcal{D}^{k-1}) \cap \mathcal{D}^{k-1})), \quad \mathcal{D}^0 \equiv \mathcal{D},$$

while G_* has an analytic continuation to

$$\mathcal{D}_3^k \equiv \Lambda_*^{-1}(G_*^{-1}(G_*(\mathcal{D}_3^{k-1}) \cap \mathcal{D}_3^{k-1})), \quad \mathcal{D}_3^0 \equiv \mathcal{D}_3.$$

The real slices of the domains \mathcal{D}_3^0 and \mathcal{D}_3^1 are given in the Figure 3. We will conclude this Section with the following




Figure 2. The real slices of the domains of F_* (blue) G_* (red).

Lemma 4.4 All maps $F \in \mathbf{F}_*$ posses a hyperbolic fixed point $p_0 = p_0(F)$, such that

- 1) $\mathcal{P}_x p_0 \in (0.57761843, 0.57761989)$, and $\mathcal{P}_u p_0 = 0$;
- 2) $DF(p_0)$ has two real eigenvalues:

$$\begin{split} e_+ &\in (-2.05763559, -2.05759928), \\ e_- &\in (-0.48601715, -0.48598084). \end{split}$$

Proof. The bound on the fixed point has been obtained with the help of the interval Newton method. Hyperbolicity has been demonstrated by computing a bound on DF using formula (11).

5. Statement of the main results

We will now summarize our main findings.




Figure 3. The real slices of the domains \mathcal{D}_3^0 (red) and \mathcal{D}_3^1 (blue).

Our first theorem describes the two-sided heteroclinic tangle for the fixed point map F_* .

Main Theorem 1 The renormalization fixed point F_* has the following properties:

- 1) F_* possesses a point p_{\oplus} which is transversally homoclinic to the fixed point p_0 ;
- 2) there exists a positive integer n such that for any negative integer k the map F_*^n has a heteroclinic orbit \mathcal{O}_k between the periodic points $\Lambda_*^k(p_0)$ and $\Lambda_*^{k+1}(p_0)$, and for any positive integer k the map $F_*^{n\cdot 2^k}$ has a heteroclinic orbit \mathcal{O}_k between the periodic points $\Lambda_*^k(p_0)$ and $\Lambda_*^{k-1}(p_0)$;
- 3) for any $N \in \mathbb{N}$ and $M \in \mathbb{N}$ there exists an integer n and an invariant set $\mathcal{Z}_{F_*^n}$, such that

$$F^n_*|_{\mathcal{Z}^n_*}\underset{homeo}{\approx} \sigma_A|_{\{-N,-N+1,\dots,M-1,M\}^{\mathbb{Z}}},$$

where σ_A is the topological Markov chain defined by a $(N+M+1)\times(N+M+1)$

tridiagonal matrix

4) for any $R > \varepsilon > 0$ there exists a point $p \in \mathcal{D}$ and an $n \in \mathbb{N}$ such that $|F_*^n(p)| > R$ (an unbounded orbit), and a point $q \in \mathcal{D}$ and $n, m \in \mathbb{N}$ such that $|F_*^n(q)| > R$ and $|F_*^n(q)| < \varepsilon$ (an oscillating orbit).

Our second result demonstrates that all locally infinitely renormalizable maps admit a hyperbolic set in their domain of analyticity.

Main Theorem 2 Any $F \in \mathbf{F}_*$ admits a hyperbolic set $\mathcal{C}_G \subset \mathcal{D}_3$ for $G \equiv F \circ F \circ F$;

$$G|_{\mathcal{C}_G} \underset{homeo}{pprox} \sigma_2|_{\{0,1\}^{\mathbb{Z}}},$$

whose Hausdorff dimension satisfies:

$$0.7673 \ge \dim_H(\mathcal{C}_G) \ge \varepsilon$$
,

where $\varepsilon \approx 0.00013 \, e^{-7499}$ is strictly positive.

6. Topological tools

The main tools of our proofs are covering relations (Zgliczyński 1997, Zgliczyński and Gidea 2004) and cone conditions (Kokubu et al 2007, Zgliczyński 2009), see also (Galias and Zgliczyński 2001) for proving the existence of homoclinic and heteroclinic orbits. To make the present paper reasonably self-contained we include a brief introduction to the necessary concepts.

6.1. H-sets and Covering relations

The notion of an h-set and a covering relation first appeared in (Zgliczyński 1997), the most thorough treatment is (Zgliczyński and Gidea 2004). The basic idea is to construct computable conditions for the existence of a semi-conjugacy to symbolic dynamics. This is done by constructing h-sets, i.e. hyperbolic-like sets, that cross each other in a (topologically) non-trivial way. We denote by $B_n(c, r)$, the open ball in \mathbb{R}^n with centre c and radius r, and $S^n(c, r) = \partial B_{n+1}(c, r)$.

Definition 6.1 An h-set is a quadruple consisting of

- a compact subset |N| of \mathbb{R}^n ,
- a pair of numbers $u(N), s(N) \in \{0, 1, 2, \ldots\}$, with u(N) + s(N) = n,
- a homeomorphism $c_N : \mathbb{R}^n \to \mathbb{R}^n = \mathbb{R}^{u(N)} \times \mathbb{R}^{s(N)}$, such that

$$c_N(|N|) = \overline{B_u(N)}(0,1) \times \overline{B_s(N)}(0,1).$$

We denote such a quadruple by N. We usually drop the bars on the support and refer to it as N. Furthermore,

$$\begin{split} N_c &= \overline{B_{u(N)}}(0,1) \times \overline{B_{s(N)}}(0,1), \\ N_c^- &= \partial \overline{B_{u(N)}}(0,1) \times \overline{B_{s(N)}}(0,1), \\ N_c^+ &= \overline{B_{u(N)}}(0,1) \times \partial \overline{B_{s(N)}}(0,1), \\ N^- &= c_N^{-1}(N_c^-), \quad N^+ = c_N^{-1}(N_c^+). \end{split}$$

u(N) and s(N) are the nominally unstable and stable directions, respectively. The idea of a covering relation between two h-sets is that the image of the first should be mapped transversally across N^- (the exit set of N) and inside of N^+ (the entrance set of N). We formally define it.

Definition 6.2 Assume N, M are h-sets, such that u(N) = u(M) = u and s(N) = s(M) = s. Let $f: N \to \mathbb{R}^n$ be a continuous map. Let $f_c = c_M \circ f \circ c_N^{-1} : N_c \to \mathbb{R}^u \times \mathbb{R}^s$. We say that

$$N \stackrel{f}{\Longrightarrow} M$$

 $(N \ f-covers \ M)$ iff the following conditions are satisfied

1. There exists a continuous homotopy $h:[0,1]\times N_c\to\mathbb{R}^u\times\mathbb{R}^s$, such that the following conditions hold true

$$h_0 = f_c,$$

 $h([0, 1], N_c^-) \cap M_c = \emptyset,$
 $h([0, 1], N_c) \cap M_c^+ = \emptyset.$

2. There exists a linear map $A: \mathbb{R}^u \to \mathbb{R}^u$, such that

$$h_1(p,q) = (Ap,0)$$
 for $p \in \overline{B_u}(0,1)$ and $q \in \overline{B_s}(0,1)$,
 $A(\partial B_u(0,1)) \subset \mathbb{R}^u \setminus \overline{B_u}(0,1)$.

 h_1 is called a model map for the relation $N \stackrel{f}{\Longrightarrow} M$.

The maps that we study in the present paper are reversible, which we use to reduce the amount of computation. For such maps the following definition is useful, see also the discussion in (Zgliczyński and Gidea 2004).

Definition 6.3 Let N be an h-set. We define the h-set N^T as follows:

- The compact subset of the quadruple N^T is the compact subset of the quadruple N, also denoted by N,
- $u(N^T) = s(N), s(N^T) = u(N).$
- The homeomorphism $c_{N^T}: \mathbb{R}^n \to \mathbb{R}^n = \mathbb{R}^{u(N^T)} \times \mathbb{R}^{s(N^T)}$ is defined by

$$c_{N^T}(x) = j(c_N(x)),$$

where $j: \mathbb{R}^{u(N)} \times \mathbb{R}^{s(N)} \to \mathbb{R}^{s(N)} \times \mathbb{R}^{u(N)}$ is given by j(p,q) = (q,p).

Definition 6.4 Assume N,M are h-sets, such that u(N) = u(M) = u and s(N) = s(M) = s. Let $g: \Omega \subset \mathbb{R}^n \to \mathbb{R}^n$. Assume that $g^{-1}: M \to \mathbb{R}^n$ is well defined and continuous. We say that $N \stackrel{\mathcal{G}}{\rightleftharpoons} M$ (N g-backcovers M) iff $M^T \stackrel{\mathcal{G}^{-1}}{\Longrightarrow} N^T$.

Definition 6.5 Let N and M be h-sets. We say that N generically f-covers M $(N \stackrel{f}{\Longleftrightarrow} M)$ if $N \stackrel{f}{\Longrightarrow} M$ or $N \stackrel{f}{\Longleftrightarrow} M$.

The main property of covering relations is contained in the following theorem (Zgliczyński and Gidea 2004, Corollary 7).

Theorem 6.6 Assume that we have the following chain of covering relations:

$$N_0 \stackrel{f_1}{\Longleftrightarrow} N_1 \stackrel{f_2}{\Longleftrightarrow} N_2 \Longleftrightarrow \dots \stackrel{f_k}{\Longleftrightarrow} N_k$$

then there exists a point $x \in \text{int } N_0$, such that

$$f_i \circ f_{i-1} \circ \ldots \circ f_1(x) \in \operatorname{int} N_i, \quad i = 1, \ldots, k.$$

Moreover, if $N_k = N_0$, then x can be chosen so that

$$f_k \circ f_{k-1} \circ \ldots \circ f_1(x) = x.$$

In our proofs the hypothesis of Theorem 6.6 is verified using the routine checkCoveringRelations of (Progs1 2009).

6.2. Cone conditions for h-sets

Theorem 6.6 gives a computational tool to prove the existence of orbits with prescribed symbolic dynamics. To prove that such orbits are unique one would ideally require hyperbolicity of the map in a neighbourhood of the orbit. Typically, this is proved by constructing invariant cone fields. An alternative method to prove uniqueness is provided by covering relations with cone conditions, first described in (Kokubu $et\ al\ 2007$), the method is studied in further detail in (Zgliczyński 2009), which we follow below.

Definition 6.7 Let $N \subset \mathbb{R}^n$ be an h-set and $Q : \mathbb{R}^n \to \mathbb{R}^n$ be a quadratic form

$$Q((x,y)) = \alpha(x) - \beta(y), \quad (x,y) \in \mathbb{R}^u \times \mathbb{R}^s, \tag{34}$$

where $\alpha: \mathbb{R}^{u(N)} \to \mathbb{R}$, and $\beta: \mathbb{R}^{s(N)} \to \mathbb{R}$ are positive definite quadratic forms.

The pair (N, Q) is called an h-set with cones.

Definition 6.8 Assume that (N, Q_N) and (M, Q_M) are h-sets with cones, such that u(N) = u(M) = u, and let $f: N \to \mathbb{R}^{\dim(M)}$ be continuous. Assume that $N \stackrel{f}{\Longrightarrow} M$. We say that f satisfies the cone condition (with respect to the pair (N, M)) iff for any $p_1, p_2 \in N_c$, $p_1 \neq p_2$ holds

$$Q_M(f_c(p_1) - f_c(p_2)) > Q_N(p_1 - p_2). (35)$$

Definition 6.9 Assume that (N, Q_N) and (M, Q_M) are h-sets with cones, such that u(N) = u(M) = u and s(N) = s(M) = s, and let $f: N \to \mathbb{R}^{u+s}$ be continuous. Assume that $N \stackrel{f}{\longleftarrow} M$. We say that f satisfies the cone condition (with respect to the pair (N, M)) iff for any $q_1, q_2 \in M_c$, $q_1 \neq q_2$ holds

$$Q_M(q_1 - q_2) > Q_N(f_c^{-1}(q_1) - f_c^{-1}(q_2)).$$
 (36)

Remark 6.10 Note, for details see the discussion in (Zqliczyński 2009), that stable manifolds are propagated backwards through a chain of covering relations with cone conditions. Similarly, unstable manifolds are propagated forwards through a chain of covering relations. In the planar case, used in the present paper, the Lipschitz constant of the stable manifold is given by $\sqrt{\frac{\beta}{\alpha}}$, where alpha and beta are the constant coefficients of the corresponding quadratic forms.

To verify that the cone conditions hold, we use the following Lemma (Zgliczyński 2009, Lemma 8).

Lemma 6.11 Assume that for any $B \in Df_c(N_C)$, the quadratic form

$$V(x) = Q_M(Bx) - Q_N(x) \tag{37}$$

is positive definite, then for any $p_1, p_2 \in N_c$ such that $p_1 \neq p_2$

$$Q_M(f_c(p_1) - f_c(p_2)) > Q_N(p_1 - p_2). (38)$$

In our proofs the hypothesis of Lemma 6.11 is verified using the routine checkConeConditions of (Progs1 2009).

Remark 6.12 The proof of Lemma 6.11 also demonstrates that

$$Q_M(f_c(p_1) - f_c(p_2)) - Q_N(p_1 - p_2) = O(|p_1 - p_2|^2),$$

and yields a uniform lower bound on $O(|p_1 - p_2|^2)$ for each pair of h-sets:

$$Q_M(f_c(p_1) - f_c(p_2)) - Q_N(p_1 - p_2) \ge \epsilon_{MN}|p_1 - p_2|^2$$
.

Theorem 6.13 Let $A = (a_{ij})_{i,j=0}^{k-1}$ be a $k \times k$ matrix whose entries are either 0 or 1. Assume that the following set of covering relations with cone conditions holds

$$N_i \stackrel{f}{\iff} N_i \quad \text{if } a_{ij} = 1.$$

Then, each sequence in $\{0,1,\ldots,k-1\}_A^{\mathbb{Z}}$ (see Definition 3.4) is realised by a unique

Proof. Assume that $p_1 \in N_{\omega_0}$ and $p_2 \in N_{\omega_0}$, are such that $p_1 \neq p_2$, $f^i(p_1) \in N_{\omega_i}$ and $f^{i}(p_{2}) \in N_{\omega_{i}}$, for all $i \in \mathbb{Z}$, where $\omega = (..., \omega_{-1}, \omega_{0}, \omega_{1}, ...) \in \{0, 1, ..., k-1\}_{A}^{\mathbb{Z}}$. Let $\epsilon = \min_{\{(i,j): a_{ij}=1\}} \epsilon_{N_j N_i}$, where $\epsilon_{N_j N_i}$ is as in Remark 6.12. There are three cases. (i) Let $Q_{N_{\omega_0}}(p_1 - p_2) > 0$, then we have that for any i > 0

$$Q_{N_{\omega_i}}(f_c^i(p_1) - f_c^i(p_2)) > Q_{N_{\omega_{i-1}}}(f_c^{i-1}(p_1) - f_c^{i-1}(p_2)) + \epsilon |f_c^{i-1}(p_1) - f_c^{i-1}(p_2)|^2 > \cdots$$

$$> Q_{N_{\omega_0}}(p_1 - p_2) + \epsilon \sum_{k=0}^{i-1} |f_c^k(p_1) - f_c^k(p_2)|^2.$$
 (39)

Since by assumption $f^{i}(p_{1}) - f^{i}(p_{2})$ stays inside a compact neighbourhood of the origin, for all $i \in \mathbb{Z}$, where $Q_{N_{\omega_i}}$ is bounded from above, the left hand side is bounded. Hence $f_c^i(p_1) - f_c^i(p_2) \to 0$, and therefore $Q_{N_{\omega_i}}(f_c^i(p_1) - f_c^i(p_2)) \to 0$, but

- $Q_{N_{\omega_i}}(f_c^i(p_1) f_c^i(p_2)) > Q_{N_{\omega_0}}(p_1 p_2) > 0$, since $p_1 \neq p_2$. This is a contradiction. (ii) Let $Q_{N_{\omega_0}}(p_1 p_2) = 0$, then $Q_{N_{\omega_1}}(f_c(p_1) f_c(p_2)) > 0$, and the above argument holds for the pair $f_c(p_1)$ and $f_c(p_2)$.
- (iii) Let $Q_{N_{\omega_0}}(p_1-p_2)<0$ we get the same contradiction for i<0 and ">" replaced by "<".




Figure 4. The homoclinic tangle at the fixed point p_0 . The stable manifold is given in blue, unstable — in red. The black boxes correspond to the h-sets used in the proof.

7. A transversal homoclinic orbit for F_*

The proofs of the theorems in this and the next sections were implemented in a computer program (Progs1 2009) using the software package CAPD (CAPD 2009) for interval arithmetic and covering relations. All computations were performed on an Intel Xeon 2.3GHz, 64bit processor with 16Gb of RAM. The program was compiled with gcc, version 4.1.2.

Numerical experiments indicate that for all $F \in \mathbf{F}_*$ there exists a transversal homoclinic point p_{\pitchfork} , approximately located at (1.067707,0), to the fixed point p_0 , approximately located at (0.577619,0), see Figure 4. We construct a sequence of covering relations and quadratic forms, and verify that the hypotheses of Theorem 6.6 and Lemma 6.11 are satisfied. Here, and in the rest of the paper, we use formula (11) to provide bounds on the derivative of F.

Theorem 7.1 Any map $F \in \mathbf{F}_*$ possesses a point p_{\pitchfork} transversally homoclinic to p_0 .

The map F^{12} admits a locally maximal invariant set $C_{F^{12}} \ni p_*, p_{\pitchfork}$:

$$F^{12}|_{\mathcal{C}_{F^{12}}} \underset{homeo}{\approx} \sigma_2|_{\{0,1\}^{\mathbb{Z}}}.$$

Proof. To avoid the "flip" of the system we study the second iterate of F, following the stable manifold from the homoclinic point to the fixed point. We construct four h-sets, B_i , i=1,2,3,4, as in Table 1: box B_4 around p_0 , box B_1 around p_0 and two intermediate boxes B_2 , and B_3 . B_4 is spanned by the unstable and stable eigenvectors of $DF(p_0)$. The other boxes are experimentally rotated until two sides are roughly parallel with the stable manifold. We prove (using the routines checkCoveringRelations and checkConeConditions in (Progs1 2009)) that the following covering sequence with cone conditions holds

$$B_1 \Rightarrow B_2 \Rightarrow B_3 \Rightarrow B_4 \Rightarrow B_4$$
.

The map F is reversible, by reversing time and reflecting the h-sets in $\{u = 0\}$ our cover is also a back-cover from p_{\uparrow} to p_0 along the unstable manifold of p_0 . Therefore we also have the following covering relations with cone conditions

$$B_4 \Leftarrow T(B_3) \Leftarrow T(B_2) \Leftarrow B_1.$$

Existence of a homoclinic orbit follows from Theorem 6.13. The transversality of the intersection follows from the fact that we use cones with Lipschitz constant less than one, Lip = 0.95.

Note that both p_0 and p_{\pitchfork} depend analytically on F. Part 1 of the first Main Theorem follows immediately, since $F_* \in \mathbf{F}_*$.

Box #	Centre	"Unstable" Direction, \mathbf{e}^u
$\overline{B_1}$	(1.067744, 0.0)	(0.271460881389316, 0.962449473933741)
B_2	(0.90733000, -0.328353)	(0.653774812595729, 0.691649449891044)
B_3	(0.66401000, -0.0727302)	(-0.736700425093906, -0.676219257095255)
B_4	(0.5776305, 0.0)	(0.788578889012330, 0.614933602760558)

Table 1. Boxes B_i used to prove the existence of a transversal homoclinic orbit. The size of each side of B_4 is $0.035|\mathbf{e}_4|$; the size of B_i , i=1,2,3, is $0.01|\mathbf{e}^u|$. The "stable" spanning vector is defined as $\mathbf{e}^s=T(\mathbf{e}^u)$. The covering sequence is $B_1\Rightarrow B_2\Rightarrow B_3\Rightarrow B_4\Rightarrow B_4$.

Remark 7.2 The numbers describing boxes B_4 in the Table 1 are non-representable on the computer. We have used these numbers as the input for the routine checkConeConditions. During its execution, the numbers are rounded up to the nearest representable number; therefore, strictly speaking, Theorem 7.1 has been proved for boxes different from (although very close to) those reported in the table. Since the specific values of the true parameters are largely irrelevant, we will not go into pains of computing and reporting bounds on these parameters. This will be the situation through out the paper: all specific numerical data appearing in various tables and statements of theorems are (very accurate) approximations of the true data for which the results have been proved on the computer.

8. A bi-infinite heteroclinic tangle for F_*

In this Section we will demonstrate that the fixed point F_* has a sequence of hyperbolic periodic points $\{z_k\}, k \in \mathbb{Z}$,

$$|z_k| \stackrel{k \to \infty}{\longrightarrow} 0$$
 and $|z_k| \stackrel{k \to -\infty}{\longrightarrow} \infty$,

such that their stable and unstable manifolds form a heteroclinic tangle.

We first demonstrate that there exists a heteroclinic orbit between the fixed point $z_0 \equiv p_0$ and the period two point $z_1 \equiv \Lambda_*(p_0)$.

We proceed as in the previous section: we construct a covering sequence of h-sets, as in Table 2, consisting of: one box B_0 around p_0 , one box B_1 around $\Lambda_*(p_0)$ and 11 intermediate boxes on the orbit. In this sequence, B_0 is defined as $\Lambda_*^{-1}(B_1)$.

Box #	Centre	l_i^s	l_i^u	\mathbf{e}_i^s	\mathbf{e}_i^u
B_1	(-0.143816,0)	0.00072	0.00072	(0.982114, 0.188285)	(0.982114, -0.188285)
B_2	$\left(-0.141389, -0.000460854\right)$	0.00054	0.00015	(0.982114, 0.188285)	(0.982114, -0.188285)
B_3	(-0.133448, -0.00190821)	0.00045	0.00015	(0.982114, 0.188285)	(0.982114, -0.188285)
B_4	(-0.098565, -0.00713661)	0.00047	0.0002	(0.987, 0.16072)	(0.997, -0.0774016)
B_5	(0.0582909, -0.00620757)	0.00055	0.00015	(0.994, -0.10938)	(0.995, 0.0998749)
B_6	(0.106679, 0.0110872)	0.0004	0.0001	(0.975, -0.222205)	(0.982114, 0.188285)
B_7	(-0.2842, 0.070894)	0.0004	0.0001	(0.84, 0.542586)	(0.866025, -0.5)
B_8	(0.533255, 0.0331362)	0.0003	0.0002	(0.8, -0.6)	(0.788579, 0.614934)
B_9	(0.575172, 0.00190701)	0.0003	0.0002	(0.788579, -0.614934)	(0.788579, 0.614934)
B_{10}	(0.577516, 0.000132348)	0.0005	0.0002	(0.788579, -0.614934)	(0.788579, 0.614934)
B_{11}	(0.577619, 0)	0.001	0.001	(0.788579, -0.614934)	(0.788579, 0.614934)

Table 2. The data used to prove the existence of a heteroclinic orbit for F_*^2 between z_0 and z_1 . Vectors $\mathbf{e}_i^{u,s}$ are the "stable" and "unstable" spanning vectors of the rectangles. The length of the sides of the rectangles B_i is $2 \cdot l_i^{u,s} \cdot |\mathbf{e}_i^{u,s}|$. The covering sequence is $B_1 \Rightarrow B_1 \Rightarrow B_2 \Rightarrow B_3 \Rightarrow B_4 \Rightarrow B_5 \Rightarrow B_6 \Rightarrow B_7 \Rightarrow B_8 \Rightarrow B_9 \Rightarrow B_{10} \Rightarrow B_{11} \Rightarrow B_0 = \Lambda_*^{-1}(B_1) \Rightarrow B_0 = \Lambda_*^{-1}(B_1)$. The first and the last two covering relations are for F_*^2 , the other ones — for F_*^4 .

Since the orbit is on the tangle between the unstable and stable manifolds, there are parts of it where it is hard to construct expanding and contracting directions directly for the second iterate of the map. The fourth iterate, however, exhibits nice hyperbolic like behaviour along a pseudoorbit. We use the programs checkCoveringRelations and checkConeConditions to verify that the hypotheses of Theorem 6.6 and Lemma 6.11 are satisfied for F_*^4 on the sequence of h-sets. In addition, we prove that they are satisfied for F_*^2 for the covering $B_1 \Rightarrow B_1$ (which implies that the covering relations and cone conditions are also satisfied for $B_0 \Rightarrow B_0$). We use cones with Lipschitz constant 0.9 in the proof.

The reversibility of F_* implies that the existence of a heteroclinic orbit from z_1 to z_0 entails the existence of a heteroclinic orbit from z_0 to z_1 . It also implies existence of homoclinic orbits for both points, that enters a small neighbourhood of the other one. The sizes of these neighbourhoods are given by the sizes of the boxes B_0 and B_1 in Table 2.

To construct the pseudoorbit we non-uniformly distribute points in the unstable eigendirection of z_1 , and locate an initial point that passes as close as possible to z_1 and z_0 . The finest discretization of the approximate unstable vector for the period two point is done on the scale 10^{-14} . The first point on the pseudoorbit is in the

unstable eigendirection of z_1 at a distance of 0.00058120854283, and the last point is at a distance of 0.000167835 from z_0 .

We will now demonstrate the following eight covering relations:

$$B_1 \Longrightarrow B_0, \quad B_1 \Longrightarrow B_1; \quad B_0 \Longleftarrow B_1, \quad B_1 \Longleftarrow B_1;$$

$$B_0 \Longrightarrow B_0, \quad B_0 \Longleftarrow B_0; \quad B_0 \Longrightarrow B_1, \quad B_1 \Longleftarrow B_0.$$

Step 1. Set

$${}_{1}B_{1}^{0} = F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap F_{*}^{-4}($$

$$\dots \cap F_{*}^{-4}(F_{*}^{4}(B_{8}) \cap F_{*}^{-4}(F_{*}^{4}(B_{9}) \cap F_{*}^{-4}(F_{*}^{4}(B_{10}) \cap B_{11})))\dots)), \quad (40)$$

$${}_{1}B_{1}^{1} = F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap F_{*}^{-4}(F_{*}^{4}(B_{10}) \cap B_{11})))\dots)$$

$${}_{1}B_{1}^{1} = F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap F_{*}^{-4}($$

$$\dots \cap F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap F_{*}^{-4}(F_{*}^{4}(B_{1}) \cap B_{1})))\dots))$$
(41)

Clearly, the iterate $\hat{F}_* \equiv F_*^{42}$ is well defined and analytic on ${}_1B_1^0 \cup {}_1B_1^1$, and

$$_1B_1^0 \xrightarrow{\hat{F}_*} B_0, \quad _1B_1^1 \xrightarrow{\hat{F}_*} B_1.$$
 (42)

The set B_1 is a parallelogram. Let π_1^s be a projection on the span of \mathbf{e}_1^s "along" the vector \mathbf{e}_1^u , and π_1^u be a projection on the span of \mathbf{e}_1^u "along" the vector \mathbf{e}_1^s . Then the sets ${}_1B_1^i$, i=0,1, are "vertical" in the sense that $\pi_1^s({}_1B_1^i)=\pi_1^s(B_1)$, while the sets $\hat{F}_*({}_1B_1^i)$, i=0,1, are "horizontal" in the sense that $\pi_0^u\hat{F}_*({}_1B_1^0)=\pi_0^u(B_0)$ and $\pi_1^u\hat{F}_*({}_1B_1^1)=\pi_1^u(B_1)$.

Step 2. Since the set B_1 is symmetric: $T(B_1) = B_1$, we have that

$$T(_1B_1^0) \stackrel{\hat{F}_*^{-1}}{\Longrightarrow} B_0, \quad T(_1B_1^1) \stackrel{\hat{F}_*^{-1}}{\Longrightarrow} B_1.$$
 (43)

Step 3. We get from the period doubling equation:

$$\hat{F}_*(\Lambda_*^{-1}(_1B_1^1)) = \Lambda_*^{-1}(\hat{F}_*^2(_1B_1^1)).$$

Clearly, since \hat{F}_* is well-defined and analytic on $_1B_1^1$, and $_1B_1^1 \stackrel{\hat{F}_*}{\Longrightarrow} _1B_1^1$ there exists a vertical subset $_2B_1^1 \subset _1B_1^1$ on which \hat{F}_*^2 is analytic. Set $_1B_0^0 = \Lambda_*^{-1}(_2B_1^1)$. Then, since $B_0 = \Lambda_*^{-1}(B_1)$, $_1B_0^0$ covers B_0 :

$$_1B_0^0 \xrightarrow{\hat{F}_*} B_0, \quad T(_1B_0^0) \xrightarrow{\hat{F}_*^{-1}} B_0.$$
 (44)

Step 4. Let ${}_1\tilde{B}^0_1 \ni {}_1B^0_1$ be any set such that $\epsilon \leq \operatorname{dist}_{p \in \partial_1 \tilde{B}^0_1}(p, {}_1B^0_1) \leq 2\epsilon$ for some sufficiently small $\epsilon > 0$, chosen so that \hat{F}_* is analytic on ${}_1\tilde{B}^0_1$. The set ${}_1B^1_0 = \hat{F}^{-1}_*\left(T({}_1\tilde{B}^0_1)\right) \cap B_0$ satisfies

$$_{1}B_{0}^{1} \xrightarrow{\hat{F}_{*}} B_{1}, \quad T(_{1}B_{0}^{1}) \xrightarrow{\hat{F}_{*}^{-1}} B_{1}.$$

$$\tag{45}$$

Set

$$\mathcal{C}_{\hat{F}_*}^{\infty} = \lim_{k \to \infty} \mathcal{C}_{\hat{F}_*}^k,$$

where

$$\mathcal{C}^k_{\hat{F}_*} = \hat{F}_*(\mathcal{C}^{k-1}_{\hat{F}_*}) \cap \hat{F}_*^{-1}(\mathcal{C}^{k-1}_{\hat{F}_*}), \quad \mathcal{C}^1_{\hat{F}_*} = \hat{F}_*(\cup_{i,j=0}^1 B^j_i) \cap \hat{F}_*^{-1}(\cup_{i,j=0}^1 T(_1B^j_i)).$$

We have proved the following:

Lemma 8.1 The map $\hat{F}_* = F_*^{42}$ admits a locally maximal invariant set $\mathcal{C}_{\hat{F}_*}^{\infty}$ on which its action is homeomorphic to the full Bernoulli shift on $\{0,1\}^{\mathbb{Z}}$.

Proof of Main Theorem 1 (2-4):

The covering relations (42) imply that, for any m, there exist vertical sets ${}_{m}B_{1}^{0} \subset {}_{1}B_{1}^{0}$ and ${}_{m}B_{1}^{1} \subset {}_{1}B_{1}^{1}$ such that \hat{F}_{*}^{m} is analytic on them, and ${}_{m}B_{1}^{0} \stackrel{\hat{F}_{*}^{m}}{\Longrightarrow} B_{0}$ and ${}_{m}B_{1}^{1} \stackrel{\hat{F}_{*}^{m}}{\Longrightarrow} B_{1}$.

Consider the map $\hat{F}_*^{2^k}$ for some integer k > 1. The fixed point equation

$$\Lambda_*^{-k} \circ \hat{F}_*^{2^k} \circ \Lambda_*^k = \hat{F}_* \tag{46}$$

implies that

$$\Lambda_{*}^{-k}(_{n2^{k}}B_{1}^{0}) & \stackrel{\hat{F}_{*}^{n}}{\Longrightarrow} \Lambda_{*}^{-k}(B_{0}) \equiv \Lambda_{*}^{-k-1}(B_{1}),
\Lambda_{*}^{-k}(T(_{n2^{k}}B_{1}^{0})) & \stackrel{\hat{F}_{*}^{-n}}{\Longrightarrow} \Lambda_{*}^{-k}(T(B_{0})) \equiv \Lambda_{*}^{-k-1}(T(B_{1})),
\Lambda_{*}^{-k}(_{n2^{k}}B_{1}^{1}) & \stackrel{\hat{F}_{*}^{n}}{\Longrightarrow} \Lambda_{*}^{-k}(B_{1}),
\Lambda_{*}^{-k}(T(_{n2^{k}}B_{1}^{1})) & \stackrel{\hat{F}_{*}^{-n}}{\Longrightarrow} \Lambda_{*}^{-k}(B_{1}).$$

Below we will use the notation that $A \stackrel{\hat{F}^n_*}{\Longrightarrow} B$ if there exist some vertical set $\tilde{A} \subset A$ (depending on n and A) on which \hat{F}^n_* is analytic, and $\tilde{A} \stackrel{\hat{F}^n_*}{\Longrightarrow} B$. $A \stackrel{\hat{F}^n_*}{\Longleftrightarrow} B$ and $A \stackrel{\hat{F}^n_*}{\Longleftrightarrow} B$ are used similarly. The covering relations (42) together with the fixed point equation (46) imply that, in this notation, we get for any integer k > 0

$$\Lambda_*^k(B_1) \stackrel{\hat{F}_*^{2^k}}{\Longleftrightarrow} \Lambda_*^{k-1}(B_1) \stackrel{\hat{F}_*^{2^{k-1}}}{\Longleftrightarrow} \dots \stackrel{\hat{F}_*^2}{\Longleftrightarrow} B_1,$$

$$\Lambda_*^k(T(B_1)) \stackrel{\hat{F}_*^{-2^k}}{\Longleftrightarrow} \Lambda_*^{k-1}(T(B_1)) \stackrel{\hat{F}_*^{-2^{k-1}}}{\Longleftrightarrow} \dots \stackrel{\hat{F}_*^{-2}}{\Longleftrightarrow} T(B_1).$$

We have, therefore, shown that for any two natural N and M, there exists an integer $n\equiv 2^N,$ such that

$$\Lambda_*^N(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{N-1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \dots \stackrel{\hat{F}_*^n}{\Longleftrightarrow} B_1 \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{-1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \\
\dots \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{-M+1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{-M}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{-M+1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \\
\dots \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{-1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} B_1 \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \dots \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^{N-1}(B_1) \stackrel{\hat{F}_*^n}{\Longleftrightarrow} \Lambda_*^N(B_1)$$

Set

$$\Delta_{N,M} = \bigcup_{k=-M}^{N} \Lambda_*^k \left(\phi(k) B_1^0 \right) \cup \Lambda_*^k \left(\phi(k) B_1^1 \right),$$

where $\phi(k) = 1$ if $k \ge 0$ and $\phi(k) = 2^{N-k}$ if k < 0, and define recursively:

$$\mathcal{Z}^1_{\hat{F}^n} \equiv \hat{F}^n_*(\Delta_{N,M}) \cap \hat{F}^{-n}_*(T(\Delta_{N,M})) \quad \text{and} \quad \mathcal{Z}^k_{\hat{F}^n} \equiv \hat{F}^n_*(\mathcal{Z}^{k-1}_{\hat{F}^n}) \cap \hat{F}^{-n}_*(\mathcal{Z}^{k-1}_{\hat{F}^n}).$$

Clearly, the set $\mathcal{Z}_{F_*^n} \equiv \mathcal{Z}_{F_*^n}^{\infty}$ is a locally maximal invariant set for F_*^n , appearing in part 3) of the Main Theorem 1.

 $F^n_*|_{\mathcal{Z}_{F^n_*}}$ is homeomorphic to a topological Markov chain, which is mixing. This together with $\operatorname{dist}(\Lambda^{-k}_*(B_1),0) \stackrel{k\to\infty}{\longrightarrow} \infty$ and $\operatorname{dist}(\Lambda^k_*(B_1),0) \stackrel{k\to\infty}{\longrightarrow} 0$ implies the existence of *unbounded* and *oscillating* orbits. This proves part 4) the Main Theorem 1.

Finally, we notice that if \mathcal{O}_0 is a heteroclinic orbit between points z_1 and z_0 for the iterate $\bar{F}_* \equiv F_*^4$ (or vice versa), then the fixed point equation (46) implies that, on one hand $\mathcal{O}_k \equiv \Lambda_*^k(\mathcal{O}_0)$ is a heteroclinic orbit for iterate $\bar{F}_*^{2^k}$ between periodic points z_{k+1} and z_k :

$$z_k \equiv \Lambda_*^k(z_0);$$

on the other hand if $p \in \mathcal{O}_{-k} \equiv \Lambda_*^{-k}(\mathcal{O}_0)$ then $\Lambda_*^k(p) \in \mathcal{O}_0$, and so is $q = \bar{F}_*^{2^k}(\Lambda_*^k(p))$, and

$$\bar{F}_*(p) = \Lambda_*^{-k}(\bar{F}_*^{2^k}(\Lambda_*^k(p))) = \Lambda_*^{-k}(q) \in \mathcal{O}_{-k}.$$

Therefore, \mathcal{O}_{-k} contains a heteroclinic orbit between fixed points z_{1-k} and z_{-k} for iterate \bar{F}_* . Stable and unstable manifolds of points z_k and z_{k+1} intersect in a two-sided tangle, see Figure 5. This proves part 2) of the Main Theorem 1.

9. Distortion tools

We will say that a pair (\mathcal{K}, ψ) is a dynamically defined Cantor set if $\mathcal{K} \subseteq R$ is a Cantor set and $\psi : \mathcal{K} \mapsto \mathcal{K}$ is a locally Lipschitz expanding map, topologically conjugate to the full Bernoulli shift $\sigma : \{0,1\}^{\mathbb{N}} \mapsto \{0,1\}^{\mathbb{N}}$.

A Markov partition $\mathcal{P} = \{\mathcal{K}_0, \mathcal{K}_1\}$ of (\mathcal{K}, ψ) is a partition of \mathcal{K} in two disjoint subsets, $\mathcal{K} = \mathcal{K}_0 \cup \mathcal{K}_1$, $\mathcal{K}_0 \cap \mathcal{K}_1 = \emptyset$, such that $\psi|_{\mathcal{K}_i} : \mathcal{K}_i \mapsto \mathcal{K}$ is a strictly monotone Lipschitz expanding homeomorphism.

Given a sequence $(a_0, \ldots, a_{n-1}) \in \{0, 1\}^n$, denote

$$\mathcal{K}(a_0,\ldots,a_{n-1}) = \bigcap_{i=0}^{n-1} \psi^{-i}(\mathcal{K}_{a_i}).$$

A bounded component of $\mathbb{R} \setminus \mathcal{K}$ is called a gap of \mathcal{K} . Gaps can be ordered as follows. The unique gap of order 0 is the interval $U_0 \equiv \hat{\mathcal{K}} \setminus \left(\hat{\mathcal{K}_0} \cup \hat{\mathcal{K}_1}\right)$, where \hat{A} denotes the convex hull of a set $A \subseteq \mathbb{R}$. A connected component of

$$\hat{\mathcal{K}} \setminus \bigcup_{(a_0,\dots,a_1)\in\{0,1\}^n} \widehat{\mathcal{K}(a_0,\dots,a_{n-1})}$$

is called a gap of order n if it is not a gap of order k < n. Any gap of \mathcal{K} is a gap of some finite order.

Definition 9.1 Given a gap U of K of order n, we denote L_U (respectively, R_U) the unique left (respectively, right) adjacent to U interval of the form $K(a_0, \ldots, a_{n-1})$, $(a_0, \ldots, a_{n-1}) \in \{0, 1\}^n$.

The numbers

$$\tau_L(\mathcal{K}) \equiv \inf \left\{ \frac{|L_U|}{|U|} : \text{U is a gap of } \mathcal{K} \right\},$$
(47)

$$\tau_R(\mathcal{K}) \equiv \inf \left\{ \frac{|R_U|}{|U|} : \text{U is a gap of } \mathcal{K} \right\}$$
(48)




Figure 5. Intersections of a) the stable manifolds of z_1 and $F_*(z_1)$ (blue) with the unstable ones for z_2 , $F_*(z_2)$, $F_*^2(z_2)$ and $F_*^3(z_2)$ (red and magenta); b) the stable manifold of z_0 (blue) with the unstable ones for z_1 and $F_*(z_1)$ (red); c) the stable manifold of z_{-1} (blue) with the unstable one for z_0 (red).

will be called the left and right thicknesses of K. The numbers

$$\tau_L(\mathcal{P}) \equiv \frac{|L_{U_0}|}{|U_0|} \quad \text{and} \quad \tau_R(\mathcal{P}) \equiv \frac{|R_{U_0}|}{|U_0|}$$

are called the left and right thicknesses of the Markov partition \mathcal{P} . Given a Lipschitz expanding map $\psi: I \mapsto \mathbb{R}, I \subset \mathbb{R}$, define the distortion of ψ on I:

$$\operatorname{Dist}(\psi, I) \equiv \sup_{\{x, y, z \in I : z \neq x, y \neq x\}} \log \left\{ \frac{|\psi(y) - \psi(x)||z - x|}{|\psi(z) - \psi(x)||y - x|} \right\} \in [0, \infty]$$
 (49)

(notice, $\psi(z) \neq \psi(x)$, $\psi(y) \neq \psi(x)$).

Definition 9.2 The distortion of the dynamically defined Cantor set (K, ψ) is

$$\operatorname{Dist}_{\psi}(\mathcal{K}) \equiv \sup_{(a_0, \dots, a_{n-1}) \in \{0,1\}^n} \operatorname{Dist}(\psi^n, \mathcal{K}(a_0, \dots, a_{n-1})). \tag{50}$$

Lemma 9.3 (see (Palis and Takens 1993), (Duarte 2000)) Let (K, ψ) dynamically defined Cantor set with a Markov partition \mathcal{P} and distortion $\mathrm{Dist}_{\psi}(\mathcal{K}) = c$. Then

$$e^{-c}\tau_L(\mathcal{P}) \le \tau_L(\mathcal{K}) \le e^c\tau_L(\mathcal{P}), \quad e^{-c}\tau_R(\mathcal{P}) \le \tau_R(\mathcal{K}) \le e^c\tau_R(\mathcal{P}).$$
 (51)

This lemma is important since it allows one to estimate the thicknesses of a Cantor set in terms of thicknesses of its Markov partition.

Definition 9.4 (see (Duarte 2000), (Gorodetsky and Kaloshin 2008)) Define \mathcal{F} to be the set of all maps $F: \Delta_0 \cup \Delta_1 \mapsto \mathbb{R}^2$ such that:

- 1) Δ_0 and Δ_1 are compact sets, diffeomorphic to rectangles, with non-empty interior;
- 2) F is C^2 on a neighbourhood of $\Delta_0 \cup \Delta_1$, that maps $\Delta_0 \cup \Delta_1$ diffeomorphically
- 3) the locally maximal invariant set $C_F = \bigcap_{n \in \mathbb{Z}} F^{-n}(\Delta_0 \cup \Delta_1)$ is a hyperbolic set, and the action of F on C_F is conjugated to the Bernoulli shift $\sigma: \{0,1\}^{\mathbb{Z}} \mapsto \{0,1\}^{\mathbb{Z}}$.
- 4) $\mathcal{P} = \{\Delta_0, \Delta_1\}$ is a Markov partition for $F : \mathcal{C}_F \mapsto \mathcal{C}_F$, in particular, F has two fixed points $p_0 \in \Delta_0$ and $p_1 \in \Delta_1$, whose stable and unstable manifolds contain the boundaries of Δ_0 and Δ_1 .

We will now introduce our main tool for computing the Hausdorff dimension the Duarte Distortion Theorem (see (Duarte 2000), (Gorodetsky and Kaloshin 2008)).

Definition 9.5 Given positive constants C, ϵ and γ define $\mathcal{F}(C, \epsilon, \gamma)$ to be the class of maps $F: \Delta_0 \cup \Delta_1 \mapsto \mathbb{R}^2$, $F \in \mathcal{F}$, such that

- 1) $\operatorname{diam}(\Delta_0 \cup \Delta_1) \le 1$, $\operatorname{diam}(F(\Delta_0) \cup F(\Delta_1)) \le 1$;
- 2) the derivative of F, $DF(x,u) = \begin{bmatrix} a(x,u) & b(x,u) \\ c(x,u) & d(x,u) \end{bmatrix}$, where a, b, c and d are C^1 , satisfies on $\Delta_0 \cup \Delta_1$
 - a) $\det DF(x, u) \equiv 1$,

 - b) $|d| < 1 < |a| \le \frac{C}{\epsilon}$, c) $|b|, |c| \le \epsilon(|a| 1)$;

3) functions $\tilde{a} \equiv a \circ F^{-1}$, $\tilde{b} \equiv b \circ F^{-1}$, $\tilde{c} \equiv c \circ F^{-1}$, $\tilde{c} \equiv c \circ F^{-1}$ on $F(\Delta_0) \cup F(\Delta_1)$, appearing in $DF^{-1}(x,u) \equiv \begin{bmatrix} \tilde{a} & -\tilde{b} \\ -\tilde{c} & \tilde{a} \end{bmatrix}$, satisfy

$$a) \quad \left| \frac{\partial \tilde{b}}{\partial x} \right| = \left| \frac{\partial \tilde{d}}{\partial u} \right|, \quad \left| \frac{\partial \tilde{b}}{\partial u} \right|, \quad \left| \frac{\partial \tilde{c}}{\partial x} \right|, \quad \left| \frac{\partial \tilde{a}}{\partial x} \right| = \left| \frac{\partial \tilde{c}}{\partial y} \right| \le \gamma (|\tilde{a}| - (52))$$

$$b) \quad \left| \frac{\partial a}{\partial u} \right| = \left| \frac{\partial b}{\partial x} \right|, \quad \left| \frac{\partial b}{\partial u} \right|, \quad \left| \frac{\partial c}{\partial x} \right|, \quad \left| \frac{\partial \tilde{c}}{\partial u} \right| = \left| \frac{\partial d}{\partial x} \right| \le \gamma (|\tilde{a}| - 5)$$

c)
$$\left| \frac{\partial \tilde{a}}{\partial u} \right|$$
, $\left| \frac{\partial \tilde{d}}{\partial x} \right| \le \gamma |\tilde{a}|(|\tilde{a}| - 1),$ (54)

$$d) \quad \left| \frac{\partial a}{\partial x} \right|, \quad \left| \frac{\partial d}{\partial u} \right| \le \gamma |\tilde{a}|(|\tilde{a}| - 1); \tag{55}$$

- 4) the variation of $\log |a(x,u)|$ in each rectangle Δ_i is less or equal to $\gamma(1-\alpha_i)$, $\alpha_i = \max_{(x,u) \in \Delta_i} |a(x,u)|$;
- 5) the gap sizes satisfy:

$$\operatorname{dist}(\Delta_0, \Delta_1) \geq \frac{\epsilon}{\gamma}, \quad \operatorname{dist}(F(\Delta_0), F(\Delta_1)) \geq \frac{\epsilon}{\gamma}.$$

Remark 9.6 Unlike in (Duarte 2000) and (Gorodetsky and Kaloshin 2008) we do not require the eigenvalues of the maps $F \in \mathcal{F}$ at the fixed points p_0 and p_1 to be positive. The positivity of the eigenvalues is important only in the construction of the boundaries of the two components Δ_0 and Δ_1 of the Markov partition \mathcal{P} (see Lemma 10.5 for the construction of \mathcal{P} in our case), but does not enter the proof of Theorem 9.7.

We will now continue with the notion of stable and unstable Cantor sets. Denote the stable and unstable foliations of \mathcal{C}_F as \mathcal{F}^s and \mathcal{F}^u :

$$\mathcal{F}^s \equiv \{ \text{connected comp. of } \mathcal{W}^s(\mathcal{C}_F) \cap (\Delta_0' \cup \Delta_1') \}, \tag{56}$$

$$\mathcal{F}^u \equiv \{ \text{connected comp. of } \mathcal{W}^u(\mathcal{C}_F) \cap (F(\Delta_0') \cup F(\Delta_1')) \}.$$
 (57)

Also, define

$$\mathcal{I}^s \equiv \mathcal{W}^s_{loc}(p_0) \cap \Delta'_0$$
 and $\mathcal{I}^u \equiv \mathcal{W}^u_{loc}(p_0) \cap \Delta_0$.

Then, the Cantor sets

$$\mathcal{K}^s \equiv \mathcal{C}_F \cap \mathcal{I}^u \quad \text{and} \quad \mathcal{K}^u \equiv \mathcal{C}_F \cap \mathcal{I}^s$$
 (58)

can be identified with the set of stable leaves of \mathcal{F}^s , respectively unstable leaves of \mathcal{F}^u . Define the projections $\pi_s: \mathcal{C}_F \mapsto \mathcal{K}^s$ and $\pi_u: \mathcal{C}_F \mapsto \mathcal{K}^u$ in the following way: if $p \in \mathcal{C}_F$ then

$$\pi_s(p) = \mathcal{W}_{loc}^s(p) \cap \mathcal{I}^u$$
 and $\pi_u(p) = \mathcal{W}_{loc}^u(p) \cap \mathcal{I}^s$,

and the maps $\psi^s: \mathcal{K}^s \mapsto \mathcal{K}^s$ and $\psi^u: \mathcal{K}^u \mapsto \mathcal{K}^u$ as

$$\psi^s \equiv \pi_s \circ F$$
 and $\psi^u \equiv \pi_u \circ F^{-1}$.

The pairs (K^s, ψ^s) and (K^u, ψ^u) are dynamically defined Cantor sets with Markov partitions

$$\mathcal{P}^s \equiv \{ \mathcal{I}^u \cap \Delta_0, \mathcal{I}^u \cap \Delta_1 \}$$
 and $\mathcal{P}^u \equiv \{ \mathcal{I}^s \cap F(\Delta_0), \mathcal{I}^s \cap F(\Delta_1) \}.$

Theorem 9.7 (Duarte Distortion Theorem, (Duarte 2000)) If $F \in \mathcal{F}(C, \epsilon, \gamma)$, then the distortion of the dynamically defined Cantor sets $(K^{u,s}, \psi_{u,s})$, $\psi_{u,s} = \pi_{u,s} \circ F$, is bounded:

$$Dist_{\psi_{u,s}}(\mathcal{K}^{u,s}) \le D(C,\epsilon,\gamma) \equiv 4(C+3)\gamma + 2\epsilon. \tag{59}$$

In particular,

$$e^{-D(C,\epsilon,\gamma)}\tau_L(\mathcal{P}^{u,s}) \le \tau_L(\mathcal{K}^{u,s}) \le e^{D(C,\epsilon,\gamma)}\tau_L(\mathcal{P}^{u,s}),$$

$$e^{-D(C,\epsilon,\gamma)}\tau_R(\mathcal{P}^{u,s}) \le \tau_R(\mathcal{K}^{u,s}) \le e^{D(C,\epsilon,\gamma)}\tau_R(\mathcal{P}^{u,s}).$$

The Duarte Distortion Theorem has been used as a powerful tool in conservative dynamics on several instances. It has been proved, and applied in (Duarte 2000) to demonstrate that accumulation of a locally maximal invariant set by periodic elliptic points is generic in two parameter families. Furthermore, (Gorodetsky and Kaloshin 2008) uses it to show that generic unfoldings of homoclinic tangencies of two-dimensional area-preserving diffeomorphisms give rise to hyperbolic sets of dimension arbitrarily close to 2. To show this, the authors of (Gorodetsky and Kaloshin 2008) complement the Duarte Distortion theorem with the following results, which will be useful to us:

Lemma 9.8 The Hausdorff dimension of a Cantor set K satisfies

$$\dim_H \mathcal{K} > d$$
,

where d is the solution of

$$\tau_L(\mathcal{K})^d + \tau_R(\mathcal{K})^d = (1 + \tau_L(\mathcal{K}) + \tau_R(\mathcal{K}))^d.$$

Lemma 9.9 Suppose that for given $t_L > 0$, $t_R > 0$ the solution d of the equation

$$t_L^d + t_R^d = (1 + t_L + t_R)^d$$

is in (0,1), then

$$d > \max \left\{ \frac{\log\left(1 + \frac{t_R}{1 + t_L}\right)}{\log\left(1 + \frac{1 + t_R}{t_L}\right)}, \frac{\log\left(1 + \frac{t_L}{1 + t_R}\right)}{\log\left(1 + \frac{1 + t_L}{t_R}\right)} \right\}.$$

$$(60)$$

10. A horseshoe for G_*

In this Section we demonstrate the existence of a horseshoe for the third iterate of $G \in \mathbf{G}_*$. We start with definitions.

Definition 10.1 (Full component) Suppose $\Delta \subset \mathcal{U} \subset \mathbb{R}^n = \mathbb{R}^k \oplus \mathbb{R}^l$ is homeomorphic to a rectangle (a set of the form $D_1 \times D_2 \subset \mathbb{R}^k \oplus \mathbb{R}^l$), and let $F : \mathcal{U} \mapsto \mathbb{R}^n$ be a diffeomorphism. A connected component $\Delta_0 = F(\bar{\Delta}_0)$ of $\Delta \cap F(\Delta)$ is called full, if

- 1) $\tilde{P}_2(\bar{\Delta}_0) = D_2$,
- 2) for any $z \in \bar{\Delta}_0$, $\tilde{\mathcal{P}}_1|_{F(\bar{\Delta}_0 \cap (D_1 \times \tilde{\mathcal{P}}_2(z)))}$ is a bijection onto D_1 .

Here, $\tilde{\mathcal{P}}_{1,2} \equiv \mathcal{P}_{1,2} \circ h$, h the homeomorphism $\Delta_0 \stackrel{\approx}{h} D_1 \times D_2$, \mathcal{P}_1 and \mathcal{P}_2 are the canonical projections on \mathbb{R}^k and \mathbb{R}^l .

Definition 10.2 (<u>Two-component horseshoe</u>) Let $\mathcal{U} \subset \mathbb{R}^n$ be an open set, then the set $\Delta \subset \mathcal{U}$ homeomorphic to a rectangle $D_1 \times D_2$ is called a two-component horseshoe for the diffeomorphism $F : \mathcal{U} \mapsto \mathbb{R}^n$ if $\Delta \cap F(\Delta)$ contains at least two full components Δ_0 and Δ_1 such that,

- 1) $\tilde{\mathcal{P}}_2(\Delta_0 \cup \Delta_1) \subset \operatorname{int} D_2$, $\tilde{\mathcal{P}}_1(F^{-1}(\Delta_0 \cup \Delta_1)) \subset \operatorname{int} D_1$;
- 2) $DF|_{F^{-1}(\Delta_0 \cup \Delta_1)}$ preserves and expands an unstable ("horizontal") cone family on $F^{-1}(\Delta_0 \cup \Delta_1)$;
- 2) $DF^{-1}|_{\Delta_0\cup\Delta_1}$ preserves and expands a stable ("vertical") cone family on $\Delta_0\cup\Delta_1$.

By a standard result of the theory of dynamical systems, if Δ is a two-component horseshoe for F, then the action of F on the locally maximal hyperbolic set $C_F \equiv \bigcap_{i=-\infty}^{\infty} F^i(\Delta)$ is homeomorphic to the Bernoulli shift σ_2 on $\{0,1\}^{\mathbb{Z}}$.

To demonstrate the existence of a horseshoe for the third iterate of $G \in \mathbf{G}_*$, we construct two rectangles: Δ'_0 which contains the fixed point p_0 and Δ'_1 which contains the period three point p_1 . We will prove that these two rectangles constitute two full components of a horseshoe, and are such that $G(\Delta'_0) \cap \mathcal{W}^s_{\text{loc}}(p_0)$ and $G(\Delta'_1) \cap \mathcal{W}^s_{\text{loc}}(p_0)$ contain the components of the *Markov partition* of the locally maximal hyperbolic set \mathcal{K}^u_* .

The centres of the rectangles and sizes of the spanning vectors, are given in Table 3. Note that these boxes are symmetric with respect to the involution $T:(x,u)\mapsto (x,-u)$. Therefore, we only have to prove that the covering relations hold for \mathbf{G}_* .

Theorem 10.3 For any $G \in \mathbf{G}_*$, the inverse map G^{-1} admits a horseshoe whose two full components are the sets Δ_0' and Δ_1' , defined in Table 3. Δ_0' and Δ_1' are independent of G.

Proof. To demonstrate the existence of the horseshoe we first use the routine checkCoveringRelations to show that

$$\Delta_0' \overset{\mathbf{G}_*}{\Longrightarrow} \Delta_0' \overset{\mathbf{G}_*}{\Longrightarrow} \Delta_1' \overset{\mathbf{G}_*}{\Longrightarrow} \Delta_1' \overset{\mathbf{G}_*}{\Longrightarrow} \Delta_0',$$

where the h-sets are as in Table 3. This implies that 1) in the Definition 10.2 holds.

The remainder of the proof of the existence of a horseshoe is done for the map $\tilde{\mathbf{G}}_{*}^{-1} \equiv \mathcal{T} \circ \mathbf{G}_{*}^{-1} \circ \mathcal{T}^{-1}$, where $\mathcal{T}(x,u)$ is some coordinate transformation that approximately diagonalizes the derivative $D\mathbf{G}_{*}$ on $\Delta'_{0} \cup \Delta'_{1}$. (See (Progs2 2009) for

Component	Centre	"Stable" Scale	"Unstable" Scale
Δ'_0	(0.670198, 0.0)	0.083	0.083
Δ_1^{\prime}	(-0.441811, 0.0)	0.0655	0.0655

Table 3. The rectangles that approximate the Markov partition for the horseshoe of \mathbf{G}_* . The spanning vectors are approximate stable and unstable vectors for $D\mathbf{G}_*(\mathbf{p}_{0,1})$. The "stable" vectors for \mathbf{p}_0 and \mathbf{p}_1 , respectively, are $\mathbf{e}_0^s = (0.788578889012330, -0.614933602760558)$ and $\mathbf{e}_1^s = (0.750925931392967773, 0.660386436536671957)$. The "unstable" vectors are $\mathbf{e}_0^u = T(\mathbf{e}_0^s)$ and $\mathbf{e}_1^u = T(\mathbf{e}_1^s)$. The length of the sides of the rectangles Δ_0' and Δ_1' is $2 \cdot \text{stable/unstable}$ scale $\cdot |\mathbf{e}_{0,1}^{u,s}|$. The covering sequence, which holds for \mathbf{G}_* , is $\Delta_0' \Rightarrow \Delta_0' \Rightarrow \Delta_1' \Rightarrow \Delta_1' \Rightarrow \Delta_0'$.

programs used in this part of the proof). We have used the following coordinate change:

$$T(x, u) = 0.55 (x - 0.5, u + 0.0265461632116977382 - 0.658388496175704694(x - 0.577619) - 0.611583723237529069(x - 0.577619)^{2} + 0.102408008658008658(x - 0.577619)^{3}).$$
(61)

Denote $\tilde{\Delta}'_{0,1} = \mathcal{T}(\Delta'_{0,1})$. We show that there exist an invariant unstable cone field $\tilde{\mathcal{C}}_u$ on $\tilde{\Delta}'_0 \cup \tilde{\Delta}'_1$,

$$\tilde{\mathcal{C}}_u(p) \equiv \{ \mathbf{v} = (v_1, v_2) \in \mathbb{R}^2 : -0.15 \le \frac{v_2}{v_1} \le 0.2 \}, \quad p \in \tilde{\Delta}'_0,$$
 (62)

$$\tilde{\mathcal{C}}_u(p) \equiv \{ \mathbf{v} = (v_1, v_2) \in \mathbb{R}^2 : -0.1 \le \frac{v_2}{v_1} \le 0.05 \}, \quad p \in \tilde{\Delta}'_1.$$
 (63)

We have verified that

$$D\tilde{\mathbf{G}}_{*}(p) \cdot \mathbf{v} \in \operatorname{int} \tilde{\mathcal{C}}_{u}(\tilde{\mathbf{G}}_{*}(p)), \text{ for all } \mathbf{v} \in \tilde{\mathcal{C}}_{u}(p), \quad p \in \tilde{\Delta}'_{0} \cup \tilde{\Delta}'_{1},$$
 (64)

and that vectors inside the cone field are expanded:

$$|D\tilde{\mathbf{G}}_*(p)\cdot\mathbf{v}| = A|\mathbf{v}|, \quad \mathbf{v} \in \operatorname{int} \tilde{\mathcal{C}}_u(\tilde{\mathbf{G}}_*(p)),$$

where the expansion rate A is in $\mathbf{A} = [A_-, A_+]$

$$A_{-} = 6.091, \quad A_{+} = 26.892.$$
 (65)

Verification of invariance and expansion of the cone fields is carried out (among other things) in the routine Distortion of (Progs2 2009).

Reversibility and symmetry of the partition imply that there exists a stable cone field

$$\tilde{\mathcal{C}}_s(p) \equiv D\mathcal{T}(\mathcal{T}^{-1}(p))(T(D\mathcal{T}^{-1}(p)\mathcal{C}_u(p)))$$
(66)

on $\tilde{\mathbf{G}}_*(\tilde{\Delta}_0' \cup \tilde{\Delta}_1') \cap (\tilde{\Delta}_0' \cup \tilde{\Delta}_1')$, which is mapped into its interior and expanded with the rate A by $D\tilde{G}^{-1}$.

Lemma 10.4 The cone fields

$$C_u(p) = D\mathcal{T}^{-1}(\mathcal{T}(p))\,\tilde{C}_u(\mathcal{T}(p)), \quad p \in \Delta_0' \cup \Delta_1', \tag{67}$$

$$C_s(q) = T(C_u(q)), \quad q \in \mathbf{G}_*(\Delta_0' \cup \Delta_1') \cap (\Delta_0' \cup \Delta_1'), \tag{68}$$

are transversal in the sense that the angle between any $\mathbf{u} \in \mathcal{C}_u(q)$ and $\mathbf{v} \in \mathcal{C}_s(q)$, $q \in \mathbf{G}_*(\Delta_0' \cup \Delta_1') \cap (\Delta_0' \cup \Delta_1')$ is bounded from below.

Furthermore, the leaves of the foliations \mathcal{F}^s and \mathcal{F}^u , defined in (56) —(57), are graphs over the x-axis.

Proof. We verify on the computer (this is done in the routine Distortion in (Progs2 2009)) that any $\mathbf{v} \in D\mathcal{T}^{-1}(\mathcal{T}(p))\tilde{\mathcal{C}}_u(\mathcal{T}(p)), p \in \Delta'_0$, satisfies:

$$\frac{v_2}{v_1} \ge 0.461,$$

and that any $\mathbf{v} \in D\mathcal{T}^{-1}(\mathcal{T}(p))\tilde{\mathcal{C}}_u(\mathcal{T}(p)), p \in \Delta'_1$, satisfies:

$$\frac{v_2}{v_1} \le -0.675.$$

This implies that the angle between any $\mathbf{u} \in \mathcal{C}_u(q)$ and $\mathbf{v} \in \mathcal{C}_s(q)$, $q \in \mathbf{G}_*(\Delta'_0 \cup \Delta'_1) \cap (\Delta'_0 \cup \Delta'_1)$ is bounded from below.

The horseshoe for G is illustrated in Figure 6, for both original (a) and \mathcal{T} (b) coordinates.

Note that the full components Δ'_0 , and Δ'_1 are chosen uniformly for all $G \in \mathbf{G}_*$. This is not compatible with the definition of full components in Definition 9.4. We can, however, prove that any $G \in \mathbf{G}_*$ has full components, as in Definition 9.4, contained in the components constructed in Theorem 10.3.

Lemma 10.5 For any $G \in \mathbf{G}_*$ there exist full components $\tilde{\Delta}_0 \subset \tilde{\Delta}_0'$ and $\tilde{\Delta}_1 \subset \tilde{\Delta}_1'$ as in Definition 9.4.

Proof. (see (Progs2 2009) for programs) Let $G \in \mathbf{G}_*$, by Theorem 10.3

$$\tilde{p}_1 \equiv \mathcal{T}(p_1) \in \tilde{\Delta}'_1 \cap \tilde{G}(\tilde{\Delta}'),$$

hence there exists $\epsilon > 0$ such that $B_{\epsilon}(\tilde{p}_1) \subset \tilde{\Delta}'$. Therefore, $\tilde{W}^u_{\epsilon} \equiv \tilde{W}^u_{\text{loc}}(\tilde{p}_1) \cap B_{\epsilon}(\tilde{p}_1) \subset \tilde{\Delta}'$. Let $t \mapsto \tilde{W}^u_{\epsilon}(t)$ be some parametrization of \tilde{W}^u_{ϵ} . For sufficiently small ϵ the tangent vector $\tau(t)$ to $\tilde{W}^u_{\epsilon}(t)$ is close to $\mathbf{e}^u_1(\tilde{p}_1)$, that is, there exists $\epsilon > 0$ such that $\tau(t) \in \tilde{\mathcal{C}}_u(\tilde{W}^u_{\epsilon}(t))$.

The cone conditions imply that $\tilde{\mathcal{W}}^u_{\epsilon}(t)$ is expanded in the forward (unstable) cone into $\operatorname{cvx} \tilde{G}(\tilde{\Delta}')$: there exists a positive integer N such that $\tilde{G}^N(\tilde{\mathcal{W}}^u_{\epsilon}) \cap (\tilde{\Delta}'_{0,1})_1^- \neq \emptyset$ and $\tilde{G}^N(\tilde{\mathcal{W}}^u_{\epsilon}) \cap (\tilde{\Delta}'_{1,0})_2^- \neq \emptyset$ where $(\tilde{\Delta}'_{0,1})_i^-$, i=1,2, denotes the two components of the exit set of $\tilde{\Delta}'_{0,1}$ (see Definition 6.1).

Set $b_0 \equiv \tilde{G}^N(\tilde{\mathcal{W}}^u_{\epsilon}) \cap \tilde{\Delta}'_0$ and $b_1 \equiv \tilde{G}^N(\tilde{\mathcal{W}}^u_{\epsilon}) \cap \tilde{\Delta}'_1$. $\tilde{G}(b_0)$, stretches across $\tilde{\Delta}'$, by the set of covering relations in Table 3, that is $\tilde{G}(b_0) \cap (\tilde{\Delta}'_{0,1})_1^- \neq \emptyset$ and $\tilde{G}(b_0) \cap (\tilde{\Delta}'_{1,0})_2^- \neq \emptyset$. We set $t_0 \equiv \tilde{G}(b_0) \cap \tilde{\Delta}'_0$ and $t_1 \equiv \tilde{G}(b_0) \cap \tilde{\Delta}'_1$. Via a similar construction for \tilde{G}^{-1} , we can show that there is a pair of stable

Via a similar construction for G^{-1} , we can show that there is a pair of stable leaves l_1 and r_1 such that $l_1 \cap (\tilde{\Delta}'_1)^+_i \neq \emptyset$, i = 1, 2, and $r_1 \cap (\tilde{\Delta}'_1)^+_i \neq \emptyset$, i = 1, 2. Here $(\tilde{\Delta}'_{0,1})^+_i$, i = 1, 2, denote the two components of the entrance set of $\tilde{\Delta}'_{0,1}$ (see Definition 6.1). Furthermore, there is a pair l_0 and r_0 with similar properties.

Since b_1 and t_1 stretch across $\tilde{\Delta}'_1$ "horizontally", and l_1 and r_1 stretch across $\tilde{\Delta}'_1$ "vertically", b_1 intersects l_1 and r_1 at least once, and t_1 intersects l_1 and r_1 at least once. Clearly, \tilde{p}_1 is one of these intersection points. Therefore, b_1 , t_1 , r_1 and l_1 bound an open set $\mathcal{U} \in \tilde{\Delta}'_1$ with at least one connected component. We define $\tilde{\Delta}_1$ as the closure of the connected component that contains \tilde{p}_1 in its boundary.

 Δ_0 is constructed in a similar way: it is the closure of the connected component bounded by b_0 , t_0 , l_0 and r_0 that contains p_0 in its interior.




Figure 6. a) The two components Δ_0' and Δ_1' of the horseshoe map across $\Delta_0' \cup \Delta_1'$ under G and G^{-1} . Δ_0' is the diamond on the right, Δ_0' — the diamond on the left. The entrance sets are given in red and magenta, the exit sets are in green and blue. The two sets whose longer sides in red and magenta are $G(\Delta_i')$, i=0,1, the sets with longer blue and green sides are $G^{-1}(\Delta_i')$, i=0,1. b) The same horseshoe in the coordinate T. The local stable manifold is given in cyan, the boxes signify the endpoints of the Markov partition for \mathcal{K}_*^u .

11. Hausdorff dimension of the locally maximal invariant set for G_*

According to Lemma 9.8 and the Duarte Distortion Theorem 9.7, the knowledge of the left and right thicknesses of the Markov partition for the unstable Cantor set \mathcal{K}^u

together with the distortion of this Cantor set is sufficient to compute a lower bound on its Hausdorff dimension.

Lemma 11.1 For any $G \in \mathbf{G}_*$ the intersections of the local stable manifold $\tilde{\mathcal{W}}^s_{\mathrm{loc}}(\tilde{p}_0)$ with $\partial \tilde{G}(\tilde{\Delta}_0)$ and $\partial \tilde{G}(\tilde{\Delta}_1)$ are at the points k_0^b, k_0^t and k_1^b, k_1^t , respectively. Where,

$$k_0^b \in [0.041476215, 0.04509142] \times [0.01162711, 0.015242315], \\ k_0^t \in [0.034578335, 0.03698849] \times [0.023247785, 0.02565794],$$

and

$$\begin{aligned} k_1^b &\in [0.099814, 0.1022241] \times [-0.07865165, -0.07624155], \\ k_1^t &\in [0.09487115, 0.09660585] \times [-0.06980765, -0.06710275], \end{aligned}$$

respectively.

Proof. (see (Progs1 2009) for programs) We prove that there exists two points $q_1, q_2 \in \mathcal{W}^u_{loc}(p_1) \cap G(\Delta_1) \cap \mathcal{W}^s(p_0)$, such that

$$G(q_1) \cap \mathcal{W}_{loc}^s(p_0) \neq \emptyset$$

$$G(q_2) \cap \Delta_0 \neq \emptyset$$

$$G^2(q_2) \cap \Delta_1 \neq \emptyset$$

$$G^3(q_2) \cap \mathcal{W}_{loc}^s(p_0) \neq \emptyset$$

We now describe the construction of q_1 . Numerical experiments indicate that there exists a heteroclinic orbit between p_0 , and the period three point p_1 , approximately located at (-0.527156, 0). To prove this we construct a covering sequence of h-sets, as in Table 4, consisting of: one box around p_0 , one box around p_1 and five intermediate boxes on the orbit; two of them close to the unstable vector of p_1 and three of them close to the stable vector of p_0 . We use the programs checkCoveringRelations and checkConeConditions to verify that the hypotheses of Theorem 6.6 and Lemma 6.11 are satisfied on the sequence of h-sets.

We emphasize that the choice of boxes is very delicate. To construct the sequence we first construct relatively large boxes around p_0 and p_1 , spanned by their unstable and stable eigenvectors. Second, we reduce the size of these boxes until we are able to prove that the hypothesis of Lemma 6.11 is satisfied. Third, we non-uniformly distribute points in the unstable eigendirection inside of the box containing the period three point; on the smallest scale the points are only separated by 10^{-11} . Fourth, we iterate, non-rigorously, these points to locate a heteroclinic pseudoorbit, which starts within the box containing p_1 and ends in the box containing p_0 . The first point on the orbit is in the unstable eigendirection of the period three point, at a distance of 0.000158885 from our approximation of the period three point, and the last point is at a distance of 0.00177401 from the fixed point. Finally, we use the points on this pseudoorbit as centres of our covering sequence. The first two intermediate points are on the local unstable manifold of the period three point, and we use its unstable and stable eigenvectors to span the corresponding h-sets. The other three intermediate points are on the local stable manifold of the fixed point, and we use its unstable and stable eigenvectors to span the corresponding h-sets. Note that if we change the location of the original point on the stable vector by as little as 10^{-8} , the proof fails. The existence of \tilde{q}_1 follows from the set of covering relations constructed in Table 4.

To find q_2 we do a similar non-rigorous computation to locate a pseudoorbit with the properties above. The corresponding sequence of h-sets and covering relations is given in Table 5. Finally, we put $k_1^b = \mathcal{T}G(q_1)$, $k_0^t = \mathcal{T}G^2(q_1)$, $k_0^b = \mathcal{T}G^3(q_1)$, and $k_1^t = \mathcal{T}G^3(q_2)$. By the construction of the components of the Markov partition $\tilde{\Delta}_0$ and $\tilde{\Delta}_1$, it is immediate that $\{k_0^t, k_0^b\} = \tilde{\mathcal{W}}_{\text{loc}}^s(\tilde{p}_0) \cap \partial \tilde{G}(\tilde{\Delta}_0)$ and $\{k_1^t, k_1^b\} = \tilde{\mathcal{W}}_{\text{loc}}^s(\tilde{p}_0) \cap \partial \tilde{G}(\tilde{\Delta}_1)$. Evaluating $k_{0/1}^{b/t}$ at $\mathbf{q}_{0/1}$ with \mathbf{G}_* yields the enclosures given in the statement of the lemma.

Consider a $G \in \mathbf{G}_*$, by Lemma 10.5 the two components of the Markov partition for \mathcal{K}^u are contained in the intersections

$$\tilde{G}(\tilde{\Delta}_0) \cap \left(\tilde{\mathcal{W}}^s_{\mathrm{loc}}(\tilde{p}_0) \cap \tilde{\Delta}_0\right), \quad \tilde{G}(\tilde{\Delta}_1) \cap \left(\tilde{\mathcal{W}}^s_{\mathrm{loc}}(\tilde{p}_0) \cap \tilde{\Delta}_0\right).$$

Box #	Centre	Eigenvector	Scale
0	(0.577619,0)	Fixed	0.004
1	(-0.527156, 0)	Period Three	0.0005
2	(-0.524339,-0.00302553)	Period Three	0.001
3	(-0.458451,-0.0695428)	Period Three	0.002
4	(0.683671,-0.0907786)	Fixed	0.002
5	(0.565061, 0.00974195)	Fixed	0.002
6	(0.578698,-0.0014086)	Fixed	0.003

Table 4. The boxes used to prove the existence of a heteroclinic orbit between the period three point and the fixed point, passing through q_1 . Note that the numbers in this table are in the original coordinates. The point q_1 corresponds to box #3. The spanning vectors are the stable and unstable vectors for either the fixed point or the period three point. Note, the stable vector is constructed by reflecting the unstable vector in $\{u=0\}$. The approximate unstable eigenvectors for the fixed and period three points are (0.788578889012330, 0.614933602760558) and (0.682259082166558, -0.731107710380897), respectively. To construct the corresponding parallelogram the spanning vectors are scaled as indicated in the table. The covering sequence is $1 \Rightarrow 1 \Rightarrow 2 \Rightarrow 3 \Rightarrow 4 \Rightarrow 5 \Rightarrow 6 \Rightarrow 0 \Rightarrow 0$.

Box #	Centre	Eigenvector	Scale
0	(0.577619,0)	Fixed	0.002
1	(-0.527156, 0)	Period Three	0.00015
2	(-0.527035, -0.000143356)	Period Three	0.0001(0,9)
3	(-0.523879,-0.00351678)	Period Three	0.0003
4	(-0.447416, -0.0799474)	Period Three	0.0003
5	(0.787595, -0.00888005)	Fixed	0.0005
6	(-0.36687, 0.0109682)	Period Three	0.0003
7	(0.67407, -0.0819091)	Fixed	0.001
8	(0.566215, 0.00882267)	Fixed	0.001

Table 5. The boxes used to prove the existence of a heteroclinic orbit between the period three point and the fixed point, passing through q_2 . Note that the numbers in this table are in the original coordinates. k_1^t corresponds to box #7 and q_2 to box #4. The spanning vectors are the stable and unstable vectors for either the fixed point or the period three point. Note, the stable vector is constructed by reflecting the unstable vector in $\{u=0\}$. The approximate unstable eigenvectors for the fixed and period three points are (0.788578889012330, 0.614933602760558) and (0.682259082166558, -0.731107710380897), respectively. To construct the corresponding parallelogram the spanning vectors are scaled as indicated in the table. The covering sequence is $1 \Rightarrow 1 \Rightarrow 2 \Rightarrow 3 \Rightarrow 4 \Rightarrow 5 \Rightarrow 6 \Rightarrow 7 \Rightarrow 8 \Rightarrow 0 \Rightarrow 0$.

Therefore, our next task will be two find an enclosure of the local stable manifold at the fixed point p_0 .

By the property of h-sets with cone conditions mentioned in Remark 6.10, it is possible to propagate stable manifolds backwards through a sequence of covering relations. We use this property in the program encloseStableManifold to enclose the local stable manifold $\tilde{W}^s_{loc}(\tilde{p}_0) \cap \tilde{\Delta}_0$; estimates of the length of its intersection with $\tilde{G}_*(\tilde{\Delta})$ are needed to estimate the left and right thicknesses of the Markov partition for K^u , and ultimately the Hausdorff dimension.

To enclose the stable manifold of the fixed point, we work in the coordinates (61). First, we choose 200 points q_i distributed uniformly on the interval

$$(p_0, p_0 + 0.00011(0.788578889012330, -0.614933602760558)) \subset \mathbb{R}^2,$$

and iterate each of them 10 times with F_*^{-1} . These 2200 points $z_{i,n} \equiv F_*^{-n}(q_i)$, $1 \le i \le 200$, $0 \le n \le 10$, provide an approximation of the stable manifold.

We next construct rectangles

$$B_{i,n} \equiv \{(x,u) \in \mathbb{R}^2 : |(x - \mathcal{P}_x \mathcal{T}(z_{i,n}), u - \mathcal{P}_u \mathcal{T}(z_{i,n})) \cdot \mathbf{e}^u| < 0.001, \\ |(x - \mathcal{P}_x \mathcal{T}(z_{i,n}), u - \mathcal{P}_u \mathcal{T}(z_{i,n})) \cdot \mathbf{e}^s| < 0.001\},$$

where

$$\mathbf{e}^{u} = (0.570868623900820281, -0.821041420541974730),$$

 $\mathbf{e}^{s} = (0.992704972028756258, 0.120568812341277774)$

are the approximate unstable and stable eigenvectors of $D\tilde{F}_*$ at p_0 , and verify that

$$B_{i,10} \stackrel{\tilde{F}}{\Longrightarrow} B_{i,9} \stackrel{\tilde{F}}{\Longrightarrow} \dots \stackrel{\tilde{F}}{\Longrightarrow} B_{i,0} \stackrel{\tilde{F}}{\Longrightarrow} B_0,$$

for all $F \in \mathbf{F}_*$, where

$$B_0 \equiv \{ (x, u) \in \mathbb{R}^2 : |(x - \mathcal{P}_x \mathcal{T}(p_0), u - \mathcal{P}_u \mathcal{T}(p_0)) \cdot \mathbf{e}^u| < 0.001, |(x - \mathcal{P}_x \mathcal{T}(p_0), u - \mathcal{P}_u \mathcal{T}(p_0)) \cdot \mathbf{e}^s| < 0.001 \}.$$

To prove that the cone conditions are satisfied we verify the hypothesis of Lemma 6.11 using the routine checkConeConditions. This proves that $\left(\bigcup_{i,n}B_{i,n}\right)\cup B_0$ covers the local stable manifold.

The sets $\mathbf{k}_0^b, \mathbf{k}_0^t, \mathbf{k}_1^b$, and \mathbf{k}_1^t , from Lemma 11.1, serve as bounds on the end points of the sets $\mathcal{K}_{0,1}^u$ in the Markov partition $\mathcal{P} = \{\mathcal{K}_0^u, \mathcal{K}_1^u\}$. Therefore,

$$\tau_L(\mathcal{P}) \ge \frac{\operatorname{dist}(\mathbf{k}_1^b, \mathbf{k}_1^t)}{\sqrt{1 + \operatorname{Lip}^2} \sup_{q \in \mathbf{k}_1^t, q' \in \mathbf{k}_0^b} |q - q'|} \ge 0.0650166,\tag{69}$$

$$\tau_R(\mathcal{P}) \ge \frac{\operatorname{dist}(\mathbf{k}_0^b, \mathbf{k}_0^t)}{\sqrt{1 + \operatorname{Lip}^2} \sup_{q \in \mathbf{k}_0^t, q' \in \mathbf{k}_0^b} |q - q'|} \ge 0.0514139 \tag{70}$$

(these quantities are computed in the routine Gaps in (Progs2 2009)).

Finally, we compute (in routine Distortion of (Progs2 2009)) the quantities C, γ and ϵ appearing in the Definition 9.5, and verify parts 3), 4) and 5) of this definition. In particular, we have set

$$\epsilon = \sup_{\Delta_0 \cup \Delta_1} \left\{ \frac{\max\{|b|,|c|\}}{|a|-1} \right\}, \quad C = \sup_{\Delta_0 \cup \Delta_1} \left\{ |a| \frac{\max\{|b|,|c|\}}{|a|-1} \right\},$$

while γ has been chosen as the supremum of all possible values that verify the conditions (52)–(53) over $\Delta_0 \cup \Delta_1$. To verify part 4) we have bound the total variation of $\eta(x, u) = \log |a(x, u)|$ on Δ'_i from above by $\text{Area}(\Delta'_i) \cdot \sup_{(x, u) \in \Delta'_i} |\nabla \eta(x, u)|$.

Lemma 11.2 Any function $G \in \mathbf{G}_* : \Delta_0 \cup \Delta_1 \mapsto \mathbb{R}^2$ is of class $\mathcal{F}(C, \gamma, \epsilon)$ with

$$C = 16.6, \quad \gamma = 47.8, \quad \epsilon = 0.88,$$

A straightforward implementation of the Duarte Distortion Theorem and Lemma 9.8 gives us a bound on the Hausdorff dimension of the unstable Cantor set:

Corollary 11.3 For all $G \in \mathbf{G}_*$, the distortion of the unstable Cantor set \mathcal{K}_G^u is less or equal to

$$D(C, \gamma, \epsilon) = 3749.3,$$

while its Hausdorff dimension satisfies:

$$\dim_{H}(\mathcal{K}_{G}^{u}) \geq \left\{ \frac{\log\left(1 + \frac{e^{-D}\tau_{R}(\mathcal{P})}{1 + e^{D}\tau_{L}(\mathcal{P})}\right)}{\log\left(1 + \frac{1 + e^{D}\tau_{R}(\mathcal{P})}{e^{-D}\tau_{L}(\mathcal{P})}\right)}, \frac{\log\left(1 + \frac{e^{-D}\tau_{L}(\mathcal{P})}{1 + e^{D}\tau_{R}(\mathcal{P})}\right)}{\log\left(1 + \frac{1 + e^{D}\tau_{L}(\mathcal{P})}{e^{-D}\tau_{R}(\mathcal{P})}\right)} \right\} \approx \frac{1}{2D} e^{-2D} \frac{\tau_{R}(\mathcal{P})}{\tau_{L}(\mathcal{P})},$$

where $\tau_L(\mathcal{P})$ and τ_R are as in (69)–(70).

The Hausdorff dimension of the set C_G satisfies

$$\dim_H(\mathcal{C}_G) \ge 2\dim_H(\mathcal{K}_G^u) \approx 0.00013 e^{-7499}.$$

Proof. The fact that the stable and unstable foliations are transversal (see Lemma 10.4) implies that $\mathcal{C}_G = \mathcal{K}_G^u \times \mathcal{K}_G^s$. Reversibility and the symmetry of the Markov partition $\Delta_0 \cap \Delta_1$ implies that $\mathcal{K}_G^s = T(\mathcal{K}_G^u)$, and $\dim_H(\mathcal{K}_G^s) = \dim_H(\mathcal{K}_G^u)$, which in turn implies the last claim.

Corollary 11.3 together with Theorem 10.3 proves the first part of the second Main Theorem.

Let us now construct a convergent sequence of approximations of the Cantor sets \mathcal{C}_G . As before, we denote Δ_0 and Δ_1 the two full components of the Markov partition as in Definition 9.5 and Lemma 10.5, $\Delta \equiv \Delta_0 \cup \Delta_1$. Define recursively:

$$\mathcal{U}_G^1 \equiv G(\Delta) \cap G^{-1}(\Delta)$$
 and $\mathcal{U}_G^k \equiv G(\mathcal{U}_G^{k-1}) \cap G^{-1}(\mathcal{U}_G^{k-1}).$

Each of the sets \mathcal{U}_G^k contains $2 \cdot 4^k$ components $\mathcal{U}_G^{k,n}$, $n = 1...2 \cdot 4^k$.

Recall that $\partial \Delta_j = b_j \cup t_j \cup l_j \cup r_j$, j = 0, 1. By definition of $\mathcal{U}_G^{k,n}$, the boundary $\partial \mathcal{U}_G^{k,n}$ consists of two images, $t^{k,n}$ and $b^{k,n}$, of some subsets of t_j and b_j , j = 0, 1, under G^k , and two images, $t^{k,n}$ and $t^{k,n}$, of subsets of t_j and t_j , $t_j = 0, 1$, under $t_$ as "corners" of $\mathcal{U}_G^{k,n}$.

Lemma 11.4 *Let*

$$\rho_{k,n} = \sup_{B_{\rho} \subset \mathcal{U}_G^{k,n}} (\rho),$$

and set $\rho_k = \min_n \{\rho_{k,n}\}$. There exist constants C > 0 and c > 0 such that

diam
$$\left(\mathcal{U}_{G}^{k,n}\right) \leq C A_{-}^{-k} \equiv C \kappa_{+}^{k},$$

 $\rho_{k} \geq c A_{+}^{-k} \equiv c \kappa_{-}^{k},$

$$\rho_k \geq c A_+^{-k} \equiv c \kappa_-^k,$$

where A_{\pm} are as in (65).

Proof. The length of an edge of $\mathcal{U}_G^{k,n}$ is bounded by cA^{-k} for some $A \in \mathbf{A}$. This follows from the fact that Δ_j , j=0,1, are constructed so that the tangent vectors to t_j and b_j , j=0,1, are contained in \mathcal{C}_u , and those to l_j and r_j , j=0,1, — in \mathcal{C}_s (see (67) and (68)), and from the fact that an edge of $\mathcal{U}_G^{k,n}$ is contained in the image of an edge of $\mathcal{U}_G^{k-1,j}$ for some j.

Consider an edge of \mathcal{U}_G^k , say $t^{k,n}$. Set

$$\alpha_t^+ \equiv \sup_{p \in t^{k,n}} \left\{ \text{maximum } \angle \mathbf{v} : \mathbf{v} \in \mathcal{C}_u(p) \right\}, \quad \alpha_t^- \equiv \inf_{p \in t^{k,n}} \left\{ \text{minimum } \angle \mathbf{v} : \mathbf{v} \in \mathcal{C}_u(p) \right\},$$

where $\angle \mathbf{v}$ signifies the angle between vector \mathbf{v} and the x-axis, measured, say, counterclockwise. Similarly for $b^{k,n}$, $l^{k,n}$ and $r^{k,n}$. Straightforward geometric considerations demonstrate that the angles $\alpha_{t,b,l,r}^{\pm}$ specify two quadrilaterals, containing and contained in $\mathcal{U}_{G}^{k,n}$, respectively.

Since the angle between any $\mathbf{v} \in \mathcal{C}^u(p)$ and any $\mathbf{u} \in \mathcal{C}^s(p)$ is bounded from below (see Lemma 10.4), the angles of the quadrilaterals are bounded away from 0 and π , and their diameters satisfy:

$$c A_+^{-k} \leq \text{diam} (\text{Inner Quadrilateral}),$$

 $C A_-^{-k} \geq \text{diam} (\text{Outer Quadrilateral}).$

The claim follows.

Corollary 11.5 The Hausdorff dimension of the Cantor set C_G satisfies

$$\dim_H(\mathcal{C}_G) \le \frac{\log(4)}{-\log \kappa_+} \le 0.7673.$$

Proof. Clearly $C_G \subset \mathcal{U}_G^k$, for all k. As above, $\mathcal{U}_G^k = \bigcup_{n=1}^{2\cdot 4^k} \mathcal{U}_G^{k,n}$, where

$$\operatorname{diam}(\mathcal{U}_G^{k,n}) \le \operatorname{const} \kappa_+^k, \quad \text{ for all } 1 \le n \le 2 \cdot 4^k.$$

Hence, for any k > 1, $C_d^H(\mathcal{C}_G) \leq \text{const } 4^k \kappa_+^{kd}$, and if $4\kappa_+^d < 1$, then $C_d^H(\mathcal{C}_G) = 0$.

12. Overview of the programs

We will now give a brief overview of the programs used in the proofs.

A subset of the domain of a map $F \in \mathbf{F}_*$ is determined using the program findDomain of (Progs1 2009), used in Lemma 4.2. findDomain discretizes the domain of the generating function in the x-domain, and computes rigorous bounds on the maximum and minimum of the generating function, s(y,x), on this slice of the domain. During the process of computing the maximum and minimum, the program also verifies that the conditions of the implicit function theorem are verified on the slice.

To verify that a pair of h-sets satisfies the hypothesis of Theorem 6.6, we use the routine checkCoveringRelations of (Progs1 2009). It discretizes the boundary of

a h-set, and verifies that each piece of the boundary is mapped so that the covering relations hold.

To verify that a pair of h-sets with cones satisfies the hypothesis of Lemma 6.11, we use the routine <code>checkConeConditions</code> of (Progs1 2009). It computes the derivative of the map on the initial h-set in appropriate coordinates, and checks the positive definite condition of the quadratic form V in the statement of the lemma using the Sylvester criterion.

The sequences of h-sets used to enclose the stable manifold in Section 11, are constructed in the program encloseStableManifold of (Progs1 2009). This program uses the routines checkCoveringRelations and checkConeConditions, to verify that on each sequence, the hypotheses of Theorem 6.6 and Lemma 6.11, respectively, are satisfied.

As we have already mentioned parts of Theorem 10.3, Lemmas 10.4 and 11.2 are proved with the help of the routine Distortion of (Progs2 2009). The input of the routine is a set of parameters that define the components Δ'_0 and Δ'_1 (see Table 3) and the cone fields (see (62)–(63)). The routine verifies that the cone fields are invariant and computes the expansion rates (65). It also computes the quantities C, γ and ϵ appearing in Lemma 11.2.

The thicknesses (69) and (70) of the Markov partition for the horseshoe of G_* are computed in the routine Gaps of (Progs2 2009).

13. Concluding remarks

An obvious question is whether the Hausdorff dimension of the set C_G for all G in some subset of G_* is independent of G.

In the satellite work (Gaidashev and Johnson 2009) we were able to use a renormalization approach to show that the Hausdorff dimension of the "stable" set, that is the set on which the maximal Lyapunov exponent is zero, is indeed invariant for a subset of *infinitely renormalizable* maps. An essential ingredient of that proof is the fact that the renormalizations of all infinitely renormalizable maps in a neighbourhood of the fixed point converge to that fixed point on their domain of analyticity \mathcal{D} .

In a somewhat similar fashion, the invariance of the Hausdorff dimension of the set \mathcal{C}_G could be demonstrated if one can show that renormalizations of the infinitely renormalizable maps converge to G_* on neighbourhoods \mathcal{V}_k of the rescalings $\Lambda_{k,G}^{-1}(\mathcal{C}_G)$ of the hyperbolic sets. However, $\mathcal{D} \cup_{k=1...\infty} \mathcal{V}_k$ is not a connected domain, and convergence on \mathcal{D} does not imply that on $\cup_{k=1...\infty} \mathcal{V}_k$.

Acknowledgments

The authors would like to thank Daniel Wilczak for many interesting discussions and his help with the usage of the software package (CAPD 2009). We are also grateful to Hans Koch for his many useful insights into period-doubling in area-preserving maps, as well as for his great help with understanding the original computer assisted proof (Eckmann *et al* 1984).

References

J. J. Abad, H. Koch, Renormalization and periodic orbits for Hamiltonian flows, Comm. Math. Phys. 212 (2000) # 2 371–394.

- J. J. Abad, H. Koch and P. Wittwer, A renormalization group for Hamiltonians: numerical results, Nonlinearity 11 (1998) 1185–1194.
- G. Benettin et al, Universal properties in conservative dynamical systems, Lettere al Nuovo Cimento 28 (1980) 1–4.
- T. Bountis, Period doubling bifurcations and universality in conservative Systems, Physica 3D (1981) 577–589.
- CAPD-Computer Assisted Proofs in Dynamics group, a C++ package for rigorous numerics, http://capd.wsb-nlu.edu.pl
- A. de Carvalho, M. Lyubich, M. Martens, Renormalization in the Hénon family, I: Universality but non-rigidity, J. Stat. Phys 121 (2005) 611–669.
- P. Collet, J.-P. Eckmann and H. Koch, Period doubling bifurcations for families of maps on \mathbb{R}^n , J. Stat. Phys. **3D** (1980).
- P. Collet, J.-P. Eckmann and H. Koch, On universality for area-preserving maps of the plane , *Physica* 3D (1981) 457–467.
- B. Derrida, Y. Pomeau, Feigenbaum's ratios of two dimensional area preserving maps, Phys. Lett. A80 (1980) 217–219.
- P. Duarte, Persistent homoclinic tangencies for conservative maps near identity, *Ergod. Th. & Dynam. Sys.* **20** (2000) 393–438.
- J.-P. Eckmann, H. Koch and P. Wittwer, Existence of a fixed point of the doubling transformation for area-preserving maps of the plane, Phys. Rev. A 26 (1982) # 1 720–722.
- J.-P. Eckmann, H. Koch and P. Wittwer, A Computer-Assisted Proof of Universality for Area-Preserving Maps, Memoirs of the American Mathematical Society 47 (1984), 1–121.
- H. Epstein, New proofs of the existence of the Feigenbaum functions, Commun. Math. Phys. 106 (1986) 395–426.
- H. Epstein, Fixed points of composition operators II, Nonlinearity 2 (1989) 305-310.
- D. F. Escande, F. Doveil, Renormalization method for computing the threshold of the large scale stochastic instability in two degree of freedom Hamiltonian systems, J. Stat. Phys. 26 (1981) 257–284
- E. de Faria, Proof of universality for critical circle mappings, Thesis, CUNY, 1992.
- E. de Faria, Asymptotic rigidity of scaling ratios for critical circle mappings, Ergodic Theory Dynam. Systems 19(1999), no. 4, 995–1035.
- M. J. Feigenbaum, Quantitative universality for a class of nonlinear transformations, J. Stat. Phys. 19 (1978) 25–52.
- M. J. Feigenbaum, Universal metric properties of non-linear transformations, J. Stat. Phys. 21 (1979) 669–706.
- D. Gaidashev, Renormalization of isoenergetically degenerate Hamiltonian flows and associated bifurcations of invariant tori, Discrete Contin. Dyn. Syst. 13(2005), no. 1, 63–102.
- D. Gaidashev, Cylinder renormalization for Siegel disks and a constructive Measurable Riemann Mapping Theorem, Nonlinearity 20 (2007), no 3, 713–742.
- D. Gaidashev, H. Koch, Renormalization and shearless invariant tori: numerical results, Nonlinearity 17(2004), no. 5, 1713–1722.
- D. Gaidashev, H. Koch, Period doubling in area-preserving maps: an associated one-dimenisonal problem, e-print arXiv:math.DS/0811.2588.
- D. Gaidashev, T. Johnson, Dynamics of the Universal Area-Preserving Map Associated with Period Doubling: Stable Sets, e-print math.DS/0905.4527 at Arxiv.org.
- Z. Galias and P. Zgliczynski, Abundance of homoclinic and heteroclinic orbits and rigorous bounds for the topological entropy for the Hénon map, Nonlinearity 14 no. 5 (2001) 909–932.
- A. Gorodetsky and V. Kaloshin, Conservative homoclinic bifurcations and some applications, preprint (2008), to appear in Steklov Institute Proceedings.
- R. H. G. Helleman, Self-generated chaotic behavior in nonlinear mechanics, in "Fundamental problems in statistical mechanics", Ed. by E. G. D. Cohen, North-Holland, Amsterdam, p.165, (1980).
- A. Katok, B. Hasselblat, Introduction to the Modern Theory of Dynamical Systems, Cambridge University Press, Cambridge (1995).
- K. Khanin, J. Lopes Dias, J. Marklof, Multidimensional continued fractions, dynamic renormalization and KAM theory, Comm. Math. Phys., 270 (2007), no. 1, 197–231.
- H. Koch, On the renormalization of Hamiltonian flows, and critical invariant tori, Discrete Contin. Dyn. Syst. 8 (2002), 633–646.
- H. Koch, A renormalization group fixed point associated with the breakup of golden invariant tori, Discrete Contin. Dyn. Syst. 11 (2004), no. 4, 881–909.
- H. Koch, Existence of critical invariant tori, Ergodic Theory Dynam. Systems, 28 (2008), no. 6, 1879–1894.

- S. Kocić, Renormalization of Hamiltonians for Diophantine frequency vectors and KAM tori, Nonlinearity 18 (2005) 2513–2544.
- H. Kokubu, D. Wilczak and P. Zgliczyński, Rigorous verification of cocoon bifurcations in the Michelson system, Nonlinearity 20 no. 9 (2007), 2147–2174.
- M. Lyubich, Feigenbaum-Coullet-Tresser universality and Milnor's hairness conjecture, Annals of Mathematics 149 (1999) 319–420.
- M. Lyubich, M. Martens, Renormalization in the Hénon family, II: Homoclinic tangle, e-print, arXiv:mathDS/0804.0780.
- M. Lyubich, M. Yampolsky, Dynamics of quadratic polynomials: complex bounds for real maps, Ann. Ins. Fourier 47 4 (1997) 1219-1255.
- R. S. MacKay, Renormalization approach to invariant circles in area-preserving maps, *Physica D7* (1983) 283–300.
- R. S. MacKay, Renormalisation in area preserving maps, Thesis, Princeton (1982). World Scientific, London (1993).
- A. Mehr and D.F. Escande, Destruction of KAM tori in Hamiltonian systems: link with the distabilization of nearby cycles and calculation of residues, *Physica* D13 (1984) 302–338.
- C. McMullen, Self-similarity of Siegel disks and Hausdorff dimension of Julia sets, Acta Math. 180(1998), 247-292.
- A. Neumaier, Interval Methods for Systems of Equations. Encyclopedia of Mathematics and its Applications 37, Cambridge Univ. Press, Cambridge, 1990
- J. Palis, F. Takens, Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations. Fractal dimensions and infinitely many attractors. (Cambridge Studies in Advanced Mathematics, 35.) Cambridge University Press, Cambridge, 1993.
- S. J. Shenker, L. P. Kadanoff, Critical behaviour of KAM surfaces. I Empirical results, J. Stat. Phys. 27 (1982) 631–656.
- C. Sparrow, The Lorenz equations : bifurcations, chaos, and strange attractors, New York ; Berlin : Springer (1982).
- C. Tresser and P. Coullet, Itérations d'endomorphismes et groupe de renormalisation, C. R. Acad. Sci. Paris 287A(1978), 577–580.
- M. Yampolsky, Hyperbolicity of renormalization of critical circle maps, Publ. Math. Inst. Hautes Etudes Sci. 96(2002), 1–41.
- M. Yampolsky, Renormalization horseshoe for critical circle maps, Commun. Math. Physics 240(2003), 75–96.
- M. Yampolsky, Siegel disks and renormalization fixed points, Holomorphic dynamics and renormalization, 377–393, Fields Inst. Commun., 53, Amer. Math. Soc., Providence, RI, 2008.
- P. Zgliczyński, Computer assisted proof of chaos in the Rossler equations and in the Hénon map, Nonlinearity 10, no. 1 (1997), 243–252.
- P. Zgliczyński, Covering relations, cone conditions and the stable manifold theorem, Journal of Differential Equations 246 issue 5 (2009), 1774–1819.
- P. Zgliczyński and M. Gidea 2004 Covering relations for multidimensional dynamical systems. J. Differential Equations 202 (2004), no. 1, 32–58.

Programs available at $http://www.math.uu.se/\sim johnson$

Programs available at http://www.math.uu.se/~gaidash