

A SUM-DIVISION ESTIMATE OF REALS

LIANGPAN LI AND JIAN SHEN

ABSTRACT. Let A be a finite set of positive real numbers. We present a sum-division estimate:

$$|A + A|^2 |A/A| \geq \frac{|A|^4}{4}.$$

1. INTRODUCTION

Let A be a finite set of positive real numbers throughout. The sum-set, product-set and ratio-set of A are defined respectively to be

$$\begin{aligned} A + A &= \{a + b : a, b \in A\}, \\ AA &= \{ab : a, b \in A\}, \\ A/A &= \{a/b : a, b \in A\}. \end{aligned}$$

A famous conjecture of Erdős and Szemerédi [6] asserts that for any $\alpha < 2$, there exists a constant $C_\alpha > 0$ such that

$$\max \{|A + A|, |AA|\} \geq C_\alpha |A|^\alpha.$$

In a series of papers [1, 2, 7, 11, 12, 13], upper bounds on α were found by many authors. One highlight in this direction was a proof by Elekes [2] that α can be taken $\frac{5}{4}$. His argument utilized a clever application of the Szemerédi-Trotter theorem on point-line incidences. Recently, using the concept of multiplicative energy and an ingenious geometric observation, Solymosi [14] obtained if A is not a singleton, then

$$(1) \quad |A + A|^2 |AA| \geq \frac{|A|^4}{4^{\lceil \log_2 |A| \rceil}},$$

which yields

$$(2) \quad \max \{|A + A|, |AA|\} \geq \frac{|A|^{4/3}}{2^{\lceil \log_2 |A| \rceil^{1/3}}}.$$

2000 *Mathematics Subject Classification.* 11B75 .

Key words and phrases. sum-product estimate, sum-division estimate.

One cannot completely drop the logarithmic term in (2) since if we choose $\tilde{A} = \{1, 2, \dots, n\}$, then [4, 5, 8, 15]

$$(3) \quad |\tilde{A}\tilde{A}| = \frac{n^2}{(\ln n)^{\beta+o(1)}}, \quad \beta = 1 - \frac{1 + \ln \ln 2}{\ln 2} = 0.0860713\dots$$

There is a subtle difference between $|\tilde{A}\tilde{A}|$ and $|\tilde{A}/\tilde{A}|$. In fact, Elekes and Ruzsa [3] showed that there exists a universal constant $\gamma > 0$ such that

$$(4) \quad |A + A|^6 |A/A| \geq \gamma |A|^8,$$

which yields

$$|\tilde{A}/\tilde{A}| \geq \frac{\gamma}{64} |\tilde{A}|^2$$

by choosing $A = \tilde{A}$. This leads to a natural question: how to give a joint estimate on $|A + A|$ and $|A/A|$? It is not difficult to use the Szemerédi-Trotter theorem on point-line incidences to show that

$$(5) \quad |A + A| |A/A| \geq C |A|^{5/2}$$

holds for some universal constant $C > 0$. Besides, if we carefully analyze Solymosi's proof of (1), then

$$(6) \quad |A + A|^2 |A/A| \geq \frac{|A|^4}{4 \lceil \log_2 |A| \rceil}.$$

The main purpose of this note is to drop the term $\lceil \log_2 |A| \rceil$ in (6):

Theorem 1. *Let A be a finite set of positive real numbers. Then*

$$|A + A|^2 |A/A| \geq \frac{|A|^4}{4}.$$

This implies a sum-division estimate

$$\max \{|A + A|, |A/A|\} \geq \frac{|A|^{4/3}}{2}.$$

There is an explanation on Theorem 1 in plane geometry. View \mathbb{R}^2 naturally as the complex plane \mathbb{C} . Given a finite set A of positive real numbers, denote by $\text{Rad}(A \times A)$ and $\text{Ang}(A \times A)$ respectively the radius-set and the angle-set of $A \times A$. Applying Theorem 1 with $\hat{A} = \{a^2 : a \in A\}$ yields

$$\max \{|\text{Rad}(A \times A)|, |\text{Ang}(A \times A)|\} \geq \frac{|A|^{4/3}}{2}.$$

This shows the angle-set and the radius-set of $A \times A$ cannot be small simultaneously.

2. PROOF OF THE MAIN RESULT

Suppose $|A/A| = y$ and $A/A = \{z_i\}_{i=1}^y$. Suppose z_i has m_i representations in $A \times A$, that is,

$$m_i = \left| \left\{ (a, b) \in A \times A : \frac{a}{b} = z_i \right\} \right| \quad (i = 1, 2, \dots, y).$$

Without loss of generality we may order all m_i 's as follows:

$$(7) \quad m_1 \leq m_2 \leq \dots \leq m_y.$$

Since $|A|^2 = \sum_{i=1}^y m_i$, there exists a unique integer k , $1 \leq k \leq y$, such that

$$\sum_{i=1}^{k-1} m_i < \frac{|A|^2}{2} \leq \sum_{i=1}^k m_i \leq km_k.$$

Hence

$$(8) \quad |A/A| \geq k \geq \frac{|A|^2}{2m_k}$$

and

$$(9) \quad \sum_{i=k}^y m_i = \left(|A|^2 - \sum_{i=1}^{k-1} m_i \right) \geq \frac{|A|^2}{2}.$$

By (7) and Solymosi's geometric observation [14],

$$(10) \quad |A + A|^2 = |(A \times A) + (A \times A)| \geq m_k \sum_{i=k}^y m_i.$$

Multiplying (8), (9) and (10) yields

$$|A + A|^2 |A/A| \geq \frac{|A|^4}{4}.$$

This proves Theorem 1.

Remark. Let $F_n = \{a/q : 1 \leq a \leq q \leq n, (a, q) = 1\}$ be the set of Farey fractions of order n . It is well-known ([10]) that $|F_n| \sim \frac{3}{\pi^2} n^2$ as $n \rightarrow \infty$. Besides, it is not difficult to deduce from (3) (see also [8, 9]) that

$$\max\{|F_n + F_n|, |F_n - F_n|, |F_n F_n|, |F_n/F_n|\} \leq C \frac{n^4}{(\ln n)^{\beta+o(1)}} \quad (n \rightarrow \infty).$$

This shows generally one can not expect the estimate

$$\max\{|A + A|, |A/A|\} \asymp |A|^2 \quad (|A| \rightarrow \infty).$$

We thank Dimitris Koukoulopoulos for communicating this example to us.

ACKNOWLEDGMENT

Li's research was supported by the Mathematical Tianyuan Foundation of China (No. 10826088). Shen's research was partially supported by NSF (CNS 0835834) and Texas Higher Education Coordinating Board (ARP 003615-0039-2007).

REFERENCES

- [1] Y. G. Chen, *On sums and products of integers*. Proc. Amer. Math. Soc. 127 (1999) 1927–1933.
- [2] Gy. Elekes, *On the number of sums and products*. Acta Arith. 81 (1997) 365–367.
- [3] Gy. Elekes, I. Z. Ruzsa, *Few sums, many products*. Studia Sci. Math. Hungar. 40 (2003) 301–308.
- [4] P. Erdős, *Some remarks on number theory*. Riveon Lematematika 9 (1955) 45–48.
- [5] P. Erdős, *An asymptotic inequality in the theory of numbers*. Vestnik Leningrad. Univ. 15 (1960) 41–49.
- [6] P. Erdős and E. Szemerédi, *On sums and products of integers*. In: Studies in Pure Mathematics (Birkhauser, Basel, 1983) 213–218.
- [7] K. Ford, *Sums and products from a finite set of real numbers*. Ramanujan J. 2 (1998) 59–66.
- [8] K. Ford, *The distribution of integers with a divisor in a given interval*. Ann. of Math. 168 (2008) 367–433.
- [9] A. K. Haynes, K. Homma, *The group ring of \mathbb{Q}/\mathbb{Z} and an application of a divisor problem*. Proc. Amer. Math. Soc. 137 (2009) 1285–1293.
- [10] K. L. Hua, *Introduction to Number Theory*. Springer-Verlag Berlin, Heidelberg, New York, 1982.
- [11] M. B. Nathanson, *On sums and products of integers*. Proc. Amer. Math. Soc. 125 (1997) 9–16.
- [12] J. Solymosi, *On sum-sets and product-sets of complex numbers*. J. Théor. Nombres Bordeaux 17 (2005) 921–924.
- [13] J. Solymosi, *On the number of sums and products*. Bull. London Math. Soc. 37 (2005) 491–494.
- [14] J. Solymosi, *Bounding multiplicative energy by the sumset*. Adv. Math. (2009) doi:10.1016/j.aim.2009.04.006.
- [15] G. Tenenbaum, *Sur la probabilité qu'un entier possède un diviseur dans un intervalle donné*. Compositio Math. 51 (1984) 243–263.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAO TONG UNIVERSITY, SHANGHAI 200240, CHINA & DEPARTMENT OF MATHEMATICS, TEXAS STATE UNIVERSITY, SAN MARCOS, TEXAS 78666, USA

E-mail address: liliangpan@yahoo.com.cn

DEPARTMENT OF MATHEMATICS, TEXAS STATE UNIVERSITY, SAN MARCOS, TEXAS 78666, USA

E-mail address: js48@txstate.edu