

# ASYMPTOTIC ESTIMATES FOR SOME NUMBER THEORETIC POWER SERIES

STEFAN GERHOLD

ABSTRACT. We derive asymptotic bounds for the ordinary generating functions of several classical arithmetic functions, including the Möbius, Liouville, and von Mangoldt functions. The estimates result from the Korobov-Vinogradov zero-free region for the Riemann zeta-function, and are sharper than those obtained by Abelian theorems from bounds for the summatory functions.

*Such functions as  $\sum \mu(n)x^n$ ,  $\sum \phi(n)x^n$ ,  $\sum \Lambda(n)x^n$  are extremely difficult to handle.*

— G.H. HARDY, E.M. WRIGHT [15]

## 1. INTRODUCTION

While Hardy and Wright are of course right in that ordinary generating functions of arithmetic functions do not share the versatility and usefulness of their well-known Dirichlet counterparts, several non-trivial results – both old and new – have been obtained for them. For instance, the analysis of

$$\sum_{n=1}^{\infty} \tau(n)z^n = \sum_{n=1}^{\infty} \frac{z^n}{1-z^n},$$

where  $\tau(n)$  denotes the number of divisors of  $n$ , goes back to Lambert [16], and the expansion

$$(1) \quad \sum_{n=1}^{\infty} \tau(n)e^{-nt} \sim \frac{1}{t} \log \frac{1}{t} + \frac{\gamma}{t} - \sum_{n=0}^{\infty} \frac{B_{n+1}^2}{(n+1)!(n+1)} t^n,$$

where  $t \rightarrow 0$ ,  $|\arg(t)| < \frac{1}{2}\pi - \theta$  for some  $\theta > 0$ ,

involving Euler's constant and Bernoulli numbers, has been known for a long time [13, 18, 22, 24]. Titchmarsh [22] has applied (1) in a result on mean values of the Riemann zeta-function, and Canfield et al. [9] have extended (1) to the case of the arithmetic function that counts only divisors in some fixed residue class. Another generalization has been obtained by Berndt and Evans [6], who also proved the formula

$$\sum_{n=1}^{\infty} p_n z^n \sim \frac{1}{(1-z)^2} \log \frac{1}{1-z}, \quad z \rightarrow 1^- \text{ in } \mathbb{R},$$

where  $p_n$  is the  $n$ -th prime number.

Recently, the transcendence of number theoretic power series has been of interest to several authors. Banks et al. [2] have established the irrationality of  $\sum \mu(n)z^n$ ,

---

*Date:* November 28, 2021.

*2000 Mathematics Subject Classification.* Primary 11N37; Secondary 30B10.

*Key words and phrases.* Arithmetic function, asymptotics, power series, Mellin transform.

This work was financially supported by CDG, BA-CA, AFFA, and the joint centre Microsoft Research-INRIA.

$\sum p_n z^n$ ,  $\sum \tau(n) z^n$ , and several other similar series, over  $\mathbb{Q}(z)$ . Later it was noted [4, 8] that the transcendence of these series follows easily from the fact that they have the unit circle as a natural boundary. This property even shows that they are not  $D$ -finite [5, 12, 20]. General results about the transcendence of  $\sum f(n) z^n$  with  $f$  multiplicative have recently been obtained by Borwein and Coons [8] and by Bell and Coons [4].

The present note is concerned with asymptotic estimates for power series  $\sum a_n z^n$ , where the Dirichlet generating function  $\sum a_n n^{-s}$  has singularities at the zeros of the Riemann zeta-function. For instance,  $a_n = \mu(n)$  falls under this category. Delange [11] has noted that the prime number theorem in the form

$$M(x) := \sum_{n \leq x} \mu(n) = o(x), \quad x \rightarrow \infty,$$

where  $M(x)$  denotes the Mertens function, readily implies

$$\sum_{n=1}^{\infty} \mu(n) z^n = o\left(\frac{1}{1-z}\right), \quad z \rightarrow 1^- \text{ in } \mathbb{R}.$$

A quick way to improve this starts from Walfisz' deep result [23]

$$(2) \quad M(x) = O\left(x \exp\left(-\frac{c(\log x)^{3/5}}{(\log \log x)^{1/5}}\right)\right).$$

Recall the following a basic Abelian theorem [7, 12, 19]:

**Lemma 1.** *Suppose that  $(a_n)$  is an ultimately monotone real sequence with  $a_n \sim n^\alpha \ell(n)$ , where  $\alpha > 0$ , and  $\ell$  is positive and varies slowly at infinity. Then*

$$\sum_{n=1}^{\infty} a_n z^n \sim \frac{\Gamma(\alpha+1)}{(1-z)^{\alpha+1}} \ell\left(\frac{1}{1-z}\right)$$

as  $z \rightarrow 1$  in any sector

$$(3) \quad S_\theta := \{z \in \mathbb{C} : |\arg(1-z)| \leq \frac{1}{2}\pi - \theta\}, \quad \theta > 0.$$

From the lemma (applied here only for real  $z$ ) and (2) we obtain

$$(4) \quad \begin{aligned} \sum_{n=1}^{\infty} \mu(n) z^n &= (1-z) \sum_{n=1}^{\infty} M(n) z^n \\ &= O\left(\frac{1}{t} \exp\left(-\frac{c(\log 1/t)^{3/5}}{(\log \log 1/t)^{1/5}}\right)\right), \quad t = -\log z \sim 1-z \rightarrow 0^+ \text{ in } \mathbb{R}. \end{aligned}$$

There seems to be no Tauberian result available to translate (4) back into an estimate for the Mertens function  $M(x)$ . This typical asymmetry suggests that we might be able to do a little better than (4) by using dedicated methods. Indeed, our main result (Theorem 3 below) improves (4) to

$$(5) \quad \sum_{n=1}^{\infty} \mu(n) z^n = O\left(\frac{1}{t} \exp\left(-\frac{0.0203 \times \log(1/t)}{(\log \log 1/t)^{2/3} (\log \log \log 1/t)^{1/3}}\right)\right), \quad t = -\log z,$$

where  $z \rightarrow 0$  in an arbitrary sector  $S_\theta$ ,  $\theta > 0$ . The proof rests on the contour integral representation [22]

$$(6) \quad \sum_{n=1}^{\infty} \mu(n) e^{-nt} = \frac{1}{2\pi i} \int_{\kappa-i\infty}^{\kappa+i\infty} \frac{\Gamma(s)}{\zeta(s)} t^{-s} ds, \quad \kappa > 1.$$

In a way that is familiar from the prime number theorem or the Selberg-Delange method [21], one can deform the integration contour a little bit into the critical strip  $0 < \Re(s) < 1$ , and then estimate the resulting integral. The exponential

decrease of the Gamma function along vertical lines is a convenient feature of (6), which is not present in the Perron summation formula [21]

$$(7) \quad \sum_{n \leq x} \mu(n) = \frac{1}{2\pi i} \int_{\kappa - i\infty}^{\kappa + i\infty} \frac{1}{\zeta(s)} \frac{x^s}{s} ds, \quad \kappa > 1, x \in \mathbb{R}^+ \setminus \mathbb{Z}.$$

Power series thus tend to be easier to estimate than summatory functions. The fact that  $x$  is real in (7), whereas in (6) it is natural to consider also complex  $t$ , causes no great difficulties. (At least if  $|\arg(t)|$  stays bounded away from  $\frac{1}{2}\pi$ .)

In the following section we put (5) into perspective by relating the growth of  $\sum \mu(n)z^n$  to the Riemann Hypothesis. Section 3 contains our main result, from which (5) follows. A few related power series will be estimated in Section 4. Section 5 collects some open problems.

## 2. CONNECTION TO THE RIEMANN HYPOTHESIS

In conjunction with (2) and (5), the following proposition shows that the gap between the Riemann Hypothesis and what is provable today is slightly smaller in the power series case than in the case of the summatory function  $M(x)$ .

**Proposition 2.** *Let  $\frac{1}{2} \leq \eta < 1$ . Then the following are equivalent:*

- (i)  $\zeta(s)$  has no zeros for  $\Re(s) > \eta$ ,
- (ii)  $M(x) = O(x^{\eta+\varepsilon})$ ,
- (iii)  $\sum_{n \geq 1} \mu(n)z^n = O((1-z)^{-\eta+\varepsilon})$  as  $z \rightarrow 1^-$  in  $\mathbb{R}$ .

*Proof.* The equivalence of (i) and (ii) is classical for  $\eta = \frac{1}{2}$ , see Titchmarsh [22], and the proof of the more general case is an easy modification. (The implication (ii)  $\Rightarrow$  (i), which we actually do not require, is posed as Exercise 13.4 in Apostol's textbook [1].) If (ii) holds, then (iii) follows by Lemma 1. Finally, if we assume that (iii) is true, we have that

$$F(t) := \sum_{n=1}^{\infty} \mu(n)e^{-nt} = O(t^{-(\eta+\varepsilon)}), \quad t \rightarrow 0^+ \text{ in } \mathbb{R}.$$

Hence the Mellin transform [13]

$$\int_0^{\infty} F(t)t^{s-1}dt = \frac{\Gamma(s)}{\zeta(s)}$$

defines an analytic function for  $\Re(s) > \eta$ . □

Under the Riemann Hypothesis, one would expect that we can push the integration contour in (6) across the critical line  $\Re(s) = \frac{1}{2}$  to obtain an expansion of the form

$$(8) \quad \sum_{n=1}^{\infty} \mu(n)e^{-nt} \stackrel{?}{=} t^{-1/2}H(\log(1/t)) - 2 + o(1), \quad t \rightarrow 0,$$

where  $H$ , a bounded oscillating function, is a sum of infinitely many harmonics corresponding to the non-trivial zeros of the zeta-function. The fast decrease of the Gamma function makes the residues of  $\Gamma(s)/\zeta(s)$  at these zeros rather small, so that the term  $-2$  will dominate in (8) unless  $1-z$  is very close to zero. Indeed, the  $\Omega(t^{-1/2})$  term becomes numerically visible only from about  $1-z = 10^{-10}$  onwards [P. Flajolet, private communication]. This “fake asymptotics” property has also been noted by Bateman and Diamond [3]. Without assuming the Riemann Hypothesis, Delange [11] has shown that

$$\sum_{n=1}^{\infty} \mu(n)z^n = \Omega_{\pm} \left( \frac{1}{\sqrt{1-z}} \right), \quad z \rightarrow 1^- \text{ in } \mathbb{R},$$

which is in line with (8), and shows that the left-hand side *does not* converge to  $-2$ .

### 3. MAIN RESULT

We write

$$D(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}, \quad s = \sigma + i\tau,$$

for the Dirichlet generating function of a sequence  $a_n$ . The following theorem gives an estimate for the power series  $\sum a_n z^n$  near  $z = 1$ , assuming analyticity and growth conditions for  $D(s)$ .

**Theorem 3.** *Let  $a_n$  be a sequence of complex numbers such that  $D(s)$  is absolutely convergent for  $\Re(s) > 1$  and has an analytic continuation to a set  $\Omega$  of the form*

$$(9) \quad \sigma \geq g(\tau) := \begin{cases} 1 - b(\log |\tau|)^{-\alpha} (\log \log |\tau|)^{-\beta} & |\tau| \geq w \\ 1 - b(\log w)^{-\alpha} (\log \log w)^{-\beta} & |\tau| \leq w \end{cases}$$

for some positive parameters  $\alpha, \beta, b, w$ . Assume furthermore that

$$D(s) = O(\tau^\nu),$$

uniformly as  $s \rightarrow \infty$  in  $\Omega$ , for some  $\nu > 0$ . Then for any  $\varepsilon > 0$

$$\sum_{n=1}^{\infty} a_n z^n = O\left(\frac{1}{t} \exp\left(-\frac{(b-\varepsilon) \log(1/t)}{(\log \log 1/t)^\alpha (\log \log \log 1/t)^\beta}\right)\right), \quad t = -\log z \sim 1-z.$$

The variable  $z$  may tend to 1 in an arbitrary sector of the form (3).

This result immediately implies the bound (5), by noting that  $D(s) = 1/\zeta(s)$  for  $a_n = \mu(n)$  and putting  $\alpha = \frac{2}{3}$  and  $\beta = \frac{1}{3}$ . The required analyticity and growth of  $1/\zeta(s)$  are the content of Korobov and Vinogradov's famous theorem [22], which describes the largest known zero-free region for the Riemann zeta function. (Recall that it leads to the best known error term in the prime number theorem.) For the constant  $b$  in (9) one may take  $b = 0.05507 \times (4.45)^{-2/3} > 0.0203$  in this case, by a result of Ford [14].

*Proof of Theorem 3.* The convergence assumption on  $D(s)$  clearly implies that the radius of convergence of  $\sum a_n z^n$  is at least one. We assume that  $z$  stays inside  $S_{2\theta}$ ; then  $t = -\log z$  satisfies

$$|\arg(t)| \leq \frac{1}{2}\pi - \theta$$

for small  $|t|$ . For  $\kappa > 1$  we have [22, p. 151]

$$(10) \quad \sum_{n=1}^{\infty} a_n e^{-nt} = \frac{1}{2\pi i} \int_{\kappa-i\infty}^{\kappa+i\infty} D(s) \Gamma(s) t^{-s} ds.$$

Now we deform the integration contour as indicated in Figure 1, where

$$\kappa = 1 - 1/\log |t| > 1,$$

and  $T = T(t) > 0$ , to be fixed later, tends to infinity as  $t \rightarrow 0$ . Between  $\pm iT$ , the contour is defined by the function  $g(\tau)$  from (9).

We will repeatedly apply the following version of Stirling's formula [10]: If  $\Re(s) = \sigma$  is confined to a finite interval, then

$$|\Gamma(s)| \sim \sqrt{2\pi} e^{-\pi|\tau|/2} |\tau|^{\sigma-1/2}, \quad |\tau| \rightarrow \infty,$$

uniformly w.r.t.  $\sigma$ .




FIGURE 1. The deformed integration contour, extending into the strip  $0 < \Re(s) < 1$ .

To bound the integral over the upper vertical line  $[\kappa + iT, \kappa + i\infty[$ , note that there we have

$$\begin{aligned} |\Gamma(s)t^{-s}| &\ll_t |t|^{-\kappa} \tau^{\kappa-1/2} e^{-\tau(\pi/2-\arg(t))} \\ &\leq e|t|^{-1} \tau^{\kappa-1/2} e^{-\theta\tau}. \end{aligned}$$

(Here and in the following, we write  $A \ll_t B$  for  $A = O(B)$  as  $t \rightarrow 0$ , where the estimate holds uniformly in  $\tau$ , if  $\tau = \Im(s)$  is a free variable in the right-hand side  $B$ .) Hence the integral over the upper vertical line satisfies

$$(11) \quad |I_{\text{vert}}| \ll_t \frac{1}{|t|} \int_T^\infty e^{-\theta\tau} \tau^{\kappa+\nu-1/2} d\tau \ll_t |t|^{-1} e^{-\theta T} T^{\kappa+\nu-1/2}.$$

We next estimate the contribution of the horizontal segment  $[g(T) + iT, \kappa + iT[$  to the integral. In this range we have

$$|t^{-s}| \leq |t|^{-\kappa} e^{T(\pi/2-\theta)} = |t|^{-1} e^{T(\pi/2-\theta)+1}$$

and

$$|\Gamma(s)| \ll_t T^{\kappa-1/2} e^{-T\pi/2},$$

hence this portion of the integral is

$$(12) \quad |I_{\text{hor}}| \ll_t |t|^{-1} e^{-\theta T} T^{\kappa+\nu-1/2},$$

so that we obtain the same estimate as in (11).

Finally, we bound the integral over the arc  $\sigma = g(\tau)$ , which we call  $I_{\text{arc}}$ . The integral from  $g(w)$  to  $g(w) + iw$  is plainly  $O(t^{\delta-1})$  for some positive  $\delta$ , hence negligible compared to (12). In the remaining range  $\tau > w$ , we have

$$|t^{-s}| = |t|^{-g(\tau)} e^{\tau \arg(t)} \leq |t|^{-g(T)} e^{\tau(\pi/2-\theta)}$$

and

$$|\Gamma(s)| \ll_t \tau^{g(\tau)-1/2} e^{-\tau\pi/2} \leq \tau^{g(T)-1/2} e^{-\tau\pi/2},$$

so that we have the bound

$$\begin{aligned}
|I_{\text{arc}}| &\ll_t |t|^{-g(T)} T^\nu \int_w^T e^{-\tau\theta} \tau^{g(T)-1/2} d\tau \\
&\ll_t |t|^{-g(T)} T^\nu \Gamma(g(T) + \tfrac{1}{2}) \\
(13) \quad &\ll_t |t|^{-g(T)} T^\nu.
\end{aligned}$$

To complete the proof, we have to pick  $T$  wisely in order to balance the estimates (12) and (13). We would like to have  $T$  as large as possible in (12), whereas (13) calls for a small  $T$ . We therefore choose

$$T = \frac{\log(1/|t|)}{(\log \log 1/|t|)^\alpha},$$

which makes (12) and (13) approximately equal. The former then implies

$$|I_{\text{hor}}| \ll_t \frac{1}{t} \exp\left(-\frac{(\theta - \varepsilon) \log(1/t)}{(\log \log 1/t)^\alpha}\right),$$

whereas (13) yields

$$|I_{\text{arc}}| \ll_t \frac{1}{t} \exp\left(-\frac{(b - \varepsilon) \log(1/t)}{(\log \log 1/t)^\alpha (\log \log \log 1/t)^\beta}\right),$$

both for arbitrarily small  $\varepsilon > 0$ . □

#### 4. FURTHER EXAMPLES

Besides (5), Theorem 3 yields also estimates for other number theoretic power series. In what follows, we let  $\Lambda, \lambda, \omega$ , and  $\tau$  denote, as usual, the von Mangoldt function, the Liouville function, the number-of-distinct-prime-factors function, and the number-of-divisors function. Applying Theorem 3 to the Dirichlet generating functions [11, 21]

$$(14) \quad \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \mu(n)}{n^s} = \frac{1}{\zeta(s)} \frac{2^s + 1}{2^s - 1},$$

$$(15) \quad \sum_{n=1}^{\infty} \frac{\Lambda(n) - 1}{n^s} = -\frac{\zeta'(s)}{\zeta(s)} - \zeta(s),$$

$$(16) \quad \sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s} = \frac{\zeta(2s)}{\zeta(s)},$$

$$(17) \quad \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \lambda(n)}{n^s} = (1 + 2^{1-s}) \frac{\zeta(2s)}{\zeta(s)},$$

$$(18) \quad \sum_{n=1}^{\infty} \frac{2^{\omega(n)} - \tau(n)}{n^s} = \frac{\zeta(s)^2}{\zeta(2s)} - \zeta(s)^2$$

yields the following result.

**Corollary 4.** *Let  $E(z)$  denote the function in the error term in (5). Then we have*

$$\begin{aligned}
 & \sum_{n=1}^{\infty} (-1)^n \mu(n) z^n = O(E(z)), \\
 & \sum_{n=1}^{\infty} \Lambda(n) z^n = \frac{1}{1-z} + O(E(z)), \\
 & \sum_{n=1}^{\infty} \lambda(n) z^n = O(E(z)), \\
 & \sum_{n=1}^{\infty} (-1)^n \lambda(n) z^n = O(E(z)), \\
 (19) \quad & \sum_{n=1}^{\infty} 2^{\omega(n)} z^n = \frac{1}{1-z} \log \frac{1}{1-z} + \frac{1+\gamma}{1-z} + O(E(z)),
 \end{aligned}$$

as  $z$  tends to 1 in an arbitrary sector of the form (3).

*Proof.* The Dirichlet series (14)–(18) satisfy the assumptions of Theorem 3; see, e.g., Titchmarsh [22]. As for the case of  $2^{\omega(n)}$ , formula (1) provides the required expansion of  $\sum \tau(n) z^n$ .  $\square$

Recall that Selberg and Delange [21, II.5] established expansions for summatory functions  $\sum_{n \leq x} a_n$  in the scale

$$(20) \quad x(\log x)^{\rho-k}, \quad k = 1, 2, \dots,$$

assuming that the corresponding Dirichlet series  $\sum a_n n^{-s}$  is sufficiently close to a power  $\zeta(s)^{-\rho}$  of the zeta-function, where  $\rho \in \mathbb{C}$ . This is proved from Perron's summation formula, using a contour akin to Figure 1, but circumventing the possible singularity at  $s = 1$  by a narrow loop. The same programme could be carried out for power series, too, but this seems not worthwhile. Note that Dirichlet series with a pole at  $s = 1$  can be handled by Theorem 3 after subtracting a singular element, as we did in the proof of (19). An algebraic singularity at  $s = 1$  leads to an infinite expansion in the scale (20), which readily translates into an expansion for  $\sum a_n z^n$  at  $z = 1$  by an Abelian theorem (Lemma 1).

## 5. OPEN PROBLEMS

As noted in the introduction, the unit circle is a natural boundary of  $\sum \mu(n) z^n$ . Hence one would expect that, if  $z$  tends to 1 along a path that comes very close to the unit circle, the function picks up too much growth from neighboring singularities to be bounded in any scale involving only  $1/(1-z)$ . So the restriction of  $z$  to sectors in Theorem 3 is presumably essential. More precisely, we pose the following question: If  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is an arbitrary function, does it follow that

$$\sum_{n=1}^{\infty} \mu(n) z^n = \Omega \left( f \left( \frac{1}{1-z} \right) \right)$$

as  $z \rightarrow 1$  in the unit disk?

On another register, a natural continuation of the transcendence results mentioned in the introduction would be to investigate whether the power series  $\sum f(n) z^n$ , with  $f$  any of the classical arithmetic functions, can satisfy an algebraic differential equation [17].

**Acknowledgement.** I thank Philippe Flajolet and Florian Luca for helpful comments.

## REFERENCES

- [1] T. M. APOSTOL, *Introduction to Analytic Number Theory*, Springer-Verlag, 1976.
- [2] W. D. BANKS, F. LUCA, AND I. E. SHPARLINSKI, *Irrationality of power series for various number theoretic functions.*, Manuscr. Math., 117 (2005), pp. 183–197.
- [3] P. T. BATEMAN AND H. G. DIAMOND, *On the oscillation theorems of Pringsheim and Landau.*, in Number Theory, Bambah, R. P. et al., ed., Trends in Mathematics, Birkhäuser, 2000, pp. 43–54.
- [4] J. P. BELL AND M. COONS, *Transcendence of generating functions whose coefficients are multiplicative.* Preprint, Mar. 2009. Available from [arXiv:0903.5240v1](https://arxiv.org/abs/0903.5240v1) [math.NT].
- [5] J. P. BELL, S. GERHOLD, M. KLAZAR, AND F. LUCA, *Non-holonomicity of sequences defined via elementary functions*, Annals of Combinatorics, 12 (2008), pp. 1–16.
- [6] B. C. BERNDT AND R. J. EVANS, *Extensions of asymptotic expansions from Chapter 15 of Ramanujan's second notebook*, Journal für die reine und angewandte Mathematik, 361 (1985), pp. 118–134.
- [7] N. H. BINGHAM, C. M. GOLDIE, AND J. L. TEUGELS, *Regular variation*, vol. 27 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, 1989.
- [8] P. BORWEIN AND M. COONS, *Transcendence of power series for some number theoretic functions*, Proc. Amer. Math. Soc., 137 (2009), pp. 1303–1305.
- [9] E. R. CANFIELD, C. D. SAVAGE, AND H. S. WILF, *Regularly spaced subsums of integer partitions.*, Acta Arith., 115 (2004), pp. 205–216.
- [10] E. T. COPSON, *An introduction to the theory of functions of a complex variable.*, IV + 448 p. 8 fig. Oxford, Clarendon Press, 1935.
- [11] H. DELANGE, *On some special entire series. (Sur certaines séries entières particulières.)*, Acta Arith., 92 (2000), pp. 59–70.
- [12] P. FLAJOLET, S. GERHOLD, AND B. SALVY, *On the non-holonomic character of logarithms, powers, and the  $n$ th prime function*, Electronic Journal of Combinatorics, 11(2) (2005), pp. 1–16.
- [13] P. FLAJOLET, X. GOURDON, AND P. DUMAS, *Mellin transforms and asymptotics: Harmonic sums*, Theoretical Computer Science, 144 (1995), pp. 3–58.
- [14] K. FORD, *Zero-free regions for the Riemann zeta function*. Bennett, M. A. (ed.) et al., Number theory for the millennium II. Proceedings of the millennial conference on number theory, Urbana-Champaign, IL, USA, May 21–26, 2000. Natick, MA: A K Peters. 25–56, 2002.
- [15] G. H. HARDY AND E. M. WRIGHT, *An Introduction to the Theory of Numbers*, Oxford University Press, fifth ed., 1979.
- [16] K. KNOPP, *Über Lambertsche Reihen.*, Journal für die reine und angewandte Mathematik, 142 (1913), pp. 283–315.
- [17] L. A. RUBEL, *A survey of transcendently transcendental functions.*, Am. Math. Mon., 96 (1989), pp. 777–788.
- [18] O. SCHLÖMILCH, *Compendium der höheren Analysis, Band 2*, Verlag Vieweg und Sohn, second ed., 1874.
- [19] A. M. SEDLETSKIJ, *Complex variants of Abelian theorem for Laplace transform*, Integral Transforms Spec. Funct., 3 (1995), pp. 107–112.
- [20] R. P. STANLEY, *Differentiably finite power series*, European J. Combin., 1 (1980), pp. 175–188.
- [21] G. TENENBAUM, *Introduction to analytic and probabilistic number theory*, Cambridge University Press, 1995. Translated from the second French edition (1995) by C. B. Thomas.
- [22] E. C. TITCHMARSH AND D. R. HEATH-BROWN, *The Theory of the Riemann Zeta-function*, Oxford Science Publications, second ed., 1986.
- [23] A. WALFISZ, *Weylsche Exponentialsummen in der neueren Zahlentheorie.*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.
- [24] S. WIGERT, *Sur la série de Lambert et son application à la théorie des nombres.*, Acta Math., 41 (1917), pp. 197–218.

VIENNA UNIVERSITY OF TECHNOLOGY AND MICROSOFT RESEARCH-INRIA, ORSAY  
*E-mail address:* sgerhold at fam.tuwien.ac.at