

A conjecture on square roots of Laplacian and signless Laplacian eigenvalues of graphs

S. Akbari^{a,b}, E. Ghorbani^{a,b}, M. R. Oboudi^a

^a*Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran*

^b*School of Mathematics, Institute for Research in Fundamental Sciences (IPM),*

P.O. Box 19395-5746, Tehran, Iran

Let G be a graph with n vertices, m edges, the incidence matrix X , and the directed incidence matrix D for a given orientation. X and D are n -by- m matrices, X is a $(0, 1)$ -matrix, and D is a $(-1, 0, 1)$ -matrix. Moreover

$$XX^T = \Delta + A, \text{ and } DD^T = \Delta - A,$$

where A is the adjacency matrix and Δ is the diagonal matrix whose diagonal entries are the vertex degrees of G . Note that $\Delta - A$ is the Laplacian matrix and $\Delta + A$ is the signless Laplacian matrix of G . Thus the singular values of X and D are the square roots of the signless Laplacian and the Laplacian eigenvalues of G , respectively. We denote the sum of all singular values of a matrix M by $\mathcal{E}(M)$. We have the following conjecture:

Conjecture. For any graph, $\mathcal{E}(D) \leq \mathcal{E}(X)$.

It is not hard to see that for bipartite graphs, $\mathcal{E}(D) = \mathcal{E}(X)$. A computer aid search showed that the conjecture is true for all graphs up to 9 vertices. For k -regular graphs with adjacency eigenvalues $\lambda_1, \dots, \lambda_n$ the conjecture is equivalent to the following inequality:

$$\sqrt{k - \lambda_1} + \dots + \sqrt{k - \lambda_n} \leq \sqrt{k + \lambda_1} + \dots + \sqrt{k + \lambda_n}.$$