

On the Mazur–Ulam theorem in fuzzy normed spaces

M. Eshaghi Gordji

Department of Mathematics, Semnan University,

P. O. Box 35195-363, Semnan, Iran

Tel:0098-231-4459905

Fax:0098-231-3354082

e-mail: madjid.eshaghi@gmail.com

N. Ghobadipour

Department of Mathematics, Semnan University,

P. O. Box 35195-363, Semnan, Iran

e-mail: ghobadipour.n@gmail.com

Abstract. In this paper, we establish the Mazur–Ulam theorem in the fuzzy strictly convex real normed spaces.

Mathematics Subject Classification. Primary 46S40; Secondary 39B52, 39B82, 26E50, 46S50.

Key words and phrases: Fuzzy normed space; Mazur–Ulam theorem; Fuzzy isometry

1. INTRODUCTION

In 1984, Katsaras [10] defined a fuzzy norm on a linear space and at the same year Wu and Fang [18] also introduced a notion of fuzzy normed space and gave the generalization of the Kolmogoroff normalized theorem for fuzzy topological linear space. In [6], Biswas defined and studied fuzzy inner product spaces in linear space. Since then some mathematicians have defined fuzzy metrics and norms on a linear space from various points of view [5, 9, 12, 15, 17]. In 1994, Cheng and Mordeson introduced a definition of fuzzy norm on a linear space in such a manner that the corresponding induced fuzzy metric is of Kramosil and Michalek type [11]. In 2003, Bag and Samanta [5] modified the definition of Cheng and Mordeson [8] by removing a regular condition. They also established a decomposition theorem of a fuzzy norm into a family of crisp norms and investigated some properties of fuzzy norms (see [3]). Following [1], we give the employing notion of a fuzzy norm.

Let X be a real linear space. A function $N : X \times \mathbb{R} \rightarrow [0, 1]$ (the so-called fuzzy subset) is said to be a fuzzy norm on X if for all $x, y \in X$ and all $a, b \in \mathbb{R}$:

(N_1) $N(x, a) = 0$ for $a \leq 0$;

(N_2) $x = 0$ if and only if $N(x, a) = 1$ for all $a > 0$;

(N_3) $N(ax, b) = N(x, \frac{b}{|a|})$ if $a \neq 0$;

(N_4) $N(x + y, a + b) \geq \min\{N(x, a), N(y, b)\}$;

(N_5) $N(x, \cdot)$ is non-decreasing function on \mathbb{R} and $\lim_{a \rightarrow \infty} N(x, a) = 1$;

(N_6) For $x \neq 0$, $N(x, \cdot)$ is (upper semi) continuous on \mathbb{R} .

The pair (X, N) is called a fuzzy normed linear space. One may regard $N(x, a)$ as the truth value of the statement "the norm of x is less than or equal to the real number a ".

Example 1.1. Let $(X, \|\cdot\|)$ be a normed linear space. Then

$$N(x, a) = \begin{cases} \frac{a}{a+\|x\|}, & a > 0, x \in X, \\ 0, & a \leq 0, x \in X \end{cases}$$

is a fuzzy norm on X .

Let (X, N) be a fuzzy normed linear space. Let $\{x_n\}$ be a sequence in X . Then $\{x_n\}$ is said to be convergent if there exists $x \in X$ such that $\lim_{n \rightarrow \infty} N(x_n - x, a) = 1$ for all $a > 0$. In that case, x is called the limit of the sequence $\{x_n\}$ and we denote it by $N - \lim_{n \rightarrow \infty} x_n = x$. A sequence $\{x_n\}$ in X is called Cauchy if for each $\epsilon > 0$ and each a_0 there exists n_0 such that for all $n \geq n_0$ and all $p > 0$, we have $N(x_{n+p} - x_n, a) > 1 - \epsilon$. It is known that every convergent sequence in fuzzy normed space is Cauchy. If each Cauchy sequence is convergent, then the fuzzy norm is said to be complete and the fuzzy normed space is called a fuzzy Banach space.

Definition 1.2. A fuzzy normed space is called strictly convex if $N(x+y, a+b) = \min\{N(x, a), N(y, b)\}$ and $N(x, a) = N(y, b)$ imply $x = y$ and $a = b$.

Definition 1.3. Let (X, N) and (Y, N) be two fuzzy normed spaces. We call $f : (X, N) \rightarrow (Y, N)$ a fuzzy isometry if $N(x - y, a) = N(f(x) - f(y), a)$ for all $x, y \in X$ and all $a > 0$.

Definition 1.4. Let X be a real linear space and x, y, z mutually disjoint elements of X . Then x, y and z are said to be collinear if $y - z = t(x - z)$ for some real number t .

The theory of isometry had its beginning in the important paper by Mazur and Ulam [13] in 1932.

In [13], Mazur and Ulam proved the following theorem.

Theorem 1.5. (Mazur–Ulam) Every isometry of a real normed linear space onto a real normed linear space is a linear mapping up to translation.

Also, Baker [4] investigated the Mazur–Ulam problem and obtained the following result.

Theorem 1.6. [4] Let X and Y be real normed linear spaces and suppose that Y is strictly convex. If $f : X \rightarrow Y$ is an isometry, then f is affine.

Recently, Väisälä [16] proved that every bijective isometry $f : X \rightarrow Y$ between normed linear spaces is affine.

In this paper, by using ideas of [4, 14], we investigate a Mazur–Ulam type problem in the framework of strictly convex fuzzy normed spaces.

2. MAIL RESULT

In this section we establish the Mazur–Ulam theorem in the fuzzy strictly convex real normed spaces. We need the following lemma to prove the main theorem of our paper.

Lemma 2.1. Let X be a fuzzy normed space which is strictly convex and let $a, b \in X$ and $s > 0$. Then $x = \frac{a+b}{2}$ is the unique member of X satisfying

$$N(a - x, s) = N(a - b, 2s)$$

and

$$N(b - x, s) = N(a - b, 2s).$$

Proof. There is nothing to prove if $a = b$. Let $a \neq b$. Then by (N_3) , we have

$$N(a - x, s) = N(a - \frac{a+b}{2}, s) = N(a - b, 2s)$$

and

$$N(b - x, s) = N(b - \frac{a+b}{2}, s) = N(a - b, 2s).$$

To show the uniqueness, assume that u and v are two elements of X satisfying

$$N(a - u, s) = N(a - v, s) = N(b - u, s) = N(b - v, s) = N(a - b, 2s).$$

Then

$$N(a - \frac{u+v}{2}, s) \geq \min\{N(a - u, s), N(a - v, s)\} = N(a - b, 2s) \quad (2.1)$$

and

$$N(b - \frac{u+v}{2}, s) \geq \min\{N(b - u, s), N(b - v, s)\} = N(a - b, 2s). \quad (2.2)$$

If both of inequalities (2.1) and (2.2) were strict we would have

$$\begin{aligned} N(a - b, 2s) &= N(a - \frac{u+v}{2} + \frac{u+v}{2} - b, 2s) \\ &\geq \min\{N(a - \frac{u+v}{2}, s), N(b - \frac{u+v}{2}, s)\} \\ &> N(a - b, 2s), \end{aligned}$$

a contradiction. So at least one of the equalities holds in (2.1) and (2.2). Without loss of generality assume that equality holds in (2.1). Then

$$N(a - \frac{u+v}{2}, s) = \min\{N(a - u, s), N(a - v, s)\}.$$

By the strict convexity, we obtain $N(a - u, s) = N(a - v, s)$ that is $u = v$. \square

Theorem 2.2. *Let X and Y be fuzzy real normed spaces and let Y be strictly convex. Suppose $f : X \rightarrow Y$ is a fuzzy isometry satisfies $f(a), f(b)$ and $f(c)$ are collinear when a, b and c are collinear. Then f is affine.*

Proof. We prove that $f - f(0)$ is linear. Let $h(a) = f(a) - f(0)$. Then h is a fuzzy isometry and $h(0) = 0$. Also we have

$$N(h(\frac{a+b}{2}) - h(a), s) = N(\frac{a+b}{2} - a, s) = N(a - b, 2s)$$

and similarly

$$N(h(\frac{a+b}{2}) - h(b), s) = N(\frac{a+b}{2} - b, s) = N(a - b, 2s)$$

for all $a, b \in X$ and all $s > 0$. It follows from Lemma 2.1 that

$$h(\frac{a+b}{2}) = \frac{1}{2}h(a) + \frac{1}{2}h(b).$$

Since $h(0) = 0$, we can easily show that h is additive. It follows that h is \mathbb{Q} -linear. Let $r \in \mathbb{R}$ with $r \neq 1$ and $a \in X$. Since $h(0) = 0$ and $h(0)$ is collinear, then there exists a real number r' such that $h(ra) = r'h(a)$. Now, we prove that $r = r'$. Since a and b are collinear, then $a \neq b$. Hence,

$$N(r(a - b), s) = N(h(ra) - h(rb), s) = N(r'(h(a) - h(b)), s) = N(r'(a - b), s).$$

By the strict convexity we obtain $r(a - b) = r'(a - b)$. Thus $h(ra) = rh(a)$ for all $a \in X$ and all $r \in \mathbb{R}$. Hence, f is affine. \square

3. CONCLUSION

We study the notion of fuzzy strictly convex normed spaces, and then we establish the Mazur–Ulam theorem in the fuzzy strictly convex real normed spaces.

REFERENCES

- [1] T. Bag and S. K. Samanta, Finite dimensional fuzzy normed linear spaces, *J. Fuzzy Math.* 11 (3) (2003) 687–705.
- [2] T. Bag and S. K. Samanta, Fuzzy bounded linear operators, *Fuzzy Sets Syst.* 151 (2005) 513–547.
- [3] T. Bag and S. K. Samanta, Fuzzy bounded linear operators, *Fuzzy Sets Syst.* 151 (2005) 513–547.
- [4] J. A. Baker, Isometries in normed spaces, *Amer. Math. Monthly* (1971) 655–658.
- [5] V. Balopoulos, A. G. Hatzimichailidis and Basil K. Papadopoulos, Distance and similarity measures for fuzzy operators, *Inform. Sci.* 177 (2007) 2336–2348.
- [6] R. Biswas, Fuzzy inner product spaces and fuzzy norm functions, *Inform. Sci.* 53 (1991) 185–190.
- [7] D. G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Math. J.* 16 (1949) 385–397.
- [8] S. C. Cheng and J. N. Mordeson, Fuzzy linear operator and fuzzy normed linear spaces, *Bull. Calcutta Math. Soc.* 86 (1994) 429–436.
- [9] C. Felbin, Finite dimensional fuzzy normed linear space, *Fuzzy Sets Syst.* 48 (1992) 239–248.
- [10] A. K. Katsaras, Fuzzy topological vector spaces II, *Fuzzy Sets Syst.* 12 (1984) 143–154.
- [11] I. Kramosil and J. Michalek, Fuzzy metric and statistical metric spaces, *Kybernetika* 11 (1975) 326–334.
- [12] S.V. Krishna and K. K. M. Sarma, Separation of fuzzy normed linear spaces, *Fuzzy Sets Syst.* 63 (1994) 207–217.
- [13] S. Mazur, S. Ulam, Sur les transformation isometriques d’espaces vectoriels normes, *C. R. Acad. Sci. Paris* 194 (1932) 946–948.
- [14] M. S. Moslehian and Gh. Sadeghi, A Mazur-Ulam theorem in non-Archimedean normed spaces, *Nonlinear Anal. (TMA)* 69 (2008), no. 10, 3405–3408.
- [15] B. Shieh, Infinite fuzzy relation equations with continuous t-norms, *Inform. Sci.* 178 (2008) 1961–1967.
- [16] J. Väisälä, A proof of the Mazur-Ulam theorem, *Amer. Math. Monthly* 110 (2003) 633–635.
- [17] J. Z. Xiao and X.-H. Zhu, Fuzzy normed spaces of operators and its completeness, *Fuzzy Sets Syst.* 133 (2003) 389–399.
- [18] C. Wu and J. Fang, Fuzzy generalization of Klomogoroffs theorem, *J. Harbin Inst. Technol.* 1 (1984) 1–7.