

# SEGAL-BARGMANN TRANSFORM AND PALEY-WIENER THEOREMS ON $M(2)$

E. K. NARAYANAN AND SUPARNA SEN

ABSTRACT. We study the Segal-Bargmann transform on  $M(2)$ . The range of this transform is characterized as a weighted Bergman space. In a similar fashion Poisson integrals are investigated. Using a Gutzmer's type formula we characterize the range as a class of functions extending holomorphically to an appropriate domain in the complexification of  $M(2)$ . We also prove a Paley-Wiener theorem for the inverse Fourier transform.

MSC 2000 : Primary 22E30; Secondary 22E45.

Keywords : Segal-Bargmann transform, Poisson integrals, Paley-Wiener theorem.

## 1. INTRODUCTION

Consider the following results from Euclidean Fourier analysis :

(I): A function  $f \in L^2(\mathbb{R}^n)$  admits a factorization  $f(x) = g * p_t(x)$  where  $g \in L^2(\mathbb{R}^n)$  and  $p_t(x) = \frac{1}{(4\pi t)^{\frac{n}{2}}} e^{-\frac{|x|^2}{4t}}$  (the heat kernel on  $\mathbb{R}^n$ ) if and only if  $f$  extends as an entire function to  $\mathbb{C}^n$  and we have  $\frac{1}{(2\pi t)^{n/2}} \int_{\mathbb{C}^n} |f(z)|^2 e^{-\frac{|y|^2}{2t}} dx dy < \infty$  ( $z = x + iy$ ). In this case we also have

$$\|g\|_2^2 = \frac{1}{(2\pi t)^{n/2}} \int_{\mathbb{C}^n} |f(z)|^2 e^{-\frac{|y|^2}{2t}} dx dy.$$

The mapping  $g \rightarrow g * p_t$  is called the Segal-Bargmann transform and the above says that the Segal-Bargmann transform is a unitary map from  $L^2(\mathbb{R}^n)$  onto  $\mathcal{O}(\mathbb{C}^n) \cap L^2(\mathbb{C}^n, \mu)$ , where  $d\mu(z) = \frac{1}{(2\pi t)^{n/2}} e^{-\frac{|y|^2}{2t}} dx dy$  and  $\mathcal{O}(\mathbb{C}^n)$  denotes the space of entire functions on  $\mathbb{C}^n$ .

---

The first author was supported in part by a grant from UGC via DSA-SAP and the second author was supported by Shyama Prasad Mukherjee Fellowship from Council of Scientific and Industrial Research, India.

(II) A function  $f \in L^2(\mathbb{R})$  admits a holomorphic extension to the strip  $\{x + iy : |y| < t\}$  such that

$$\sup_{|y| \leq s} \int_{\mathbb{R}} |f(x + iy)|^2 dx < \infty \quad \forall s < t$$

if and only if

$$e^{s|\xi|} \tilde{f}(\xi) \in L^2(\mathbb{R}) \quad \forall s < t$$

where  $\tilde{f}$  denotes the Fourier transform of  $f$ .

(III) An  $f \in L^2(\mathbb{R}^n)$  admits an entire extension to  $\mathbb{C}^n$  such that

$$|f(z)| \leq C_N (1 + |z|)^{-N} e^{R|Imz|} \quad \forall z \in \mathbb{C}^n$$

if and only if  $\tilde{f} \in C_c^\infty(\mathbb{R}^n)$  and  $\text{supp } \tilde{f} \subseteq \mathbb{B}(0, R)$ , where  $\mathbb{B}(0, R)$  is the ball of radius  $R$  centered around 0 in  $\mathbb{R}^n$ .

In this paper we aim to prove similar results for the non-commutative group  $M(2) = \mathbb{R}^2 \rtimes SO(2)$ . Some remarks are in order.

As noted above the map  $g \rightarrow g * p_t$  in (I) is called the Segal-Bargmann transform. This transform has attracted a lot of attention in the recent years mainly due to the work of Hall [3] where a similar result was established for an arbitrary compact Lie group  $K$ . Let  $q_t$  be the heat kernel on  $K$  and let  $K_{\mathbb{C}}$  be the complexification of  $K$ . Then Hall's result, Theorem 2 in [3], states that the map  $f \rightarrow f * q_t$  is a unitary map from  $L^2(K)$  onto the Hilbert space of  $\nu$ -square integrable holomorphic functions on  $K_{\mathbb{C}}$  for an appropriate positive  $K$ -invariant measure  $\nu$  on  $K_{\mathbb{C}}$ . Soon after Hall's paper a similar result was proved for compact symmetric spaces by Stenzel in [9]. We also refer to [4], [5], [6] and [7] for similar results for other groups and spaces.

The second result above (II) is originally due to Paley and Wiener. Let

$\mathcal{H}_t = \{f \in L^2(\mathbb{R}), f \text{ has a holomorphic extension to } |Imz| < t \text{ and}$

$$\sup_{|y| \leq s} \int_{\mathbb{R}} |f(x + iy)|^2 dx < \infty \quad \forall s < t\}.$$

Then  $\bigcup_{t>0} \mathcal{H}_t$  may be viewed as the space of all analytic vectors for the regular representation of  $\mathbb{R}$  on  $L^2(\mathbb{R})$ . This point of view was further developed by R. Goodman (see [1] and [2]) who studied analytic vectors for representations of Lie groups. The theorem of Paley-Wiener (II) characterizes analytic vectors for the regular representation of  $\mathbb{R}$  via a condition on the Fourier transform.

The third result (III) is the classical Paley-Wiener theorem. For a long time the Paley-Wiener theorem has been looked at as a characterization of the image (under Fourier transform) of  $\mathcal{C}_c^\infty$  functions on the space you are interested in. Recently, using Gutzmer's formula Thangavelu [11] has proved a Paley-Wiener type result for the inverse Fourier transform. See [8] also for a similar result.

The plan of this paper is as follows : In the remaining of this section we recall the representation theory and Plancherel theorem of  $M(2)$  and we prove the unitarity of the Segal-Bargmann transform. In the next section we study generalized Segal-Bargmann transform and prove an analogue of Theorem 8 and Theorem 10 in [3]. Third section is devoted to a study of Poisson integrals on  $M(2)$ . This section is modeled after the work of Goodman [1] and [2]. In the final section we establish a Paley-Wiener type result for the inverse Fourier transform on  $M(2)$ .

The rigid motion group  $M(2)$  is the semi-direct product of  $\mathbb{R}^2$  with  $SO(2)$  (which will be identified with the circle group  $S^1$ ) with the group law

$$(x, e^{i\alpha}) \cdot (y, e^{i\beta}) = (x + e^{i\alpha}y, e^{i(\alpha+\beta)}) \text{ where } x, y \in \mathbb{R}^2; e^{i\alpha}, e^{i\beta} \in S^1.$$

This group may be identified with a matrix subgroup of  $GL(2, \mathbb{C})$  via the map

$$(x, e^{i\alpha}) \rightarrow \begin{pmatrix} e^{i\alpha} & x \\ 0 & 1 \end{pmatrix}.$$

Unitary irreducible representations of  $M(2)$  are completely described by Mackey's theory of induced representations. For any  $\xi \in \mathbb{R}^2$  and  $g \in M(2)$ , we define  $U_g^\xi$  as follows :

$$U_g^\xi F(s) = e^{i\langle x, s\xi \rangle} F(r^{-1}s)$$

for  $g = (x, r)$  and  $F \in L^2(S^1)$ .

It is known that  $U_\xi$  is equivalent to  $U_{\xi'}$  iff  $|\xi| = |\xi'|$ . The above collection gives all the unitary irreducible representations of  $M(2)$  sufficient for the Plancherel theorem to be true. The Plancherel theorem (see Theorem 4.2 of [10]) reads

$$\int_{M(2)} |f(g)|^2 dg = \int_{\mathbb{R}^2} \|\hat{f}(\xi)\|_{HS}^2 d\xi$$

where  $\hat{f}(\xi)$  is the 'group Fourier transform' defined as an operator from  $L^2(S^1)$  to  $L^2(S^1)$  by

$$\hat{f}(\xi) = \int_{M(2)} f(g) U_g^\xi dg.$$

Moreover, the group Fourier Transform  $\hat{f}(\xi)$ , for  $\xi \in \mathbb{R}^2$  of  $f \in L^1(M(2))$  is an integral operator with the kernel  $k_f(\xi, e^{i\alpha}, e^{i\beta})$  where

$$k_f(\xi, e^{i\alpha}, e^{i\beta}) = \tilde{f}(e^{i\beta}\xi, e^{i(\beta-\alpha)}),$$

and  $\tilde{f}$  is the Euclidean Fourier Transform of  $f$  in the  $\mathbb{R}^2$ -variable.

The Lie algebra of  $M(2)$  is given by  $\left\{ \begin{pmatrix} i\alpha & x \\ 0 & 0 \end{pmatrix} : (x, e^{i\alpha}) \in M(2) \right\}$ .

Let

$$X_1 = \begin{pmatrix} i & 0 \\ 0 & 0 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix}$$

Then it is easy to see that  $\{X_1, X_2, X_3\}$  forms a basis for the Lie algebra of  $M(2)$ . The "Laplacian"  $\Delta_{M(2)} = \Delta$  is defined by

$$\Delta = -(X_1^2 + X_2^2 + X_3^2).$$

A simple computation shows that  $\Delta = -\Delta_{\mathbb{R}^2} - \frac{\partial^2}{\partial \alpha^2}$  where  $\Delta_{\mathbb{R}^2}$  is the Laplacian on  $\mathbb{R}^2$  given by  $\Delta_{\mathbb{R}^2} = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ . Since  $\Delta_{\mathbb{R}^2}$  and  $\frac{\partial^2}{\partial \alpha^2}$  commute, it follows that the heat kernel  $\psi_t$  associated to  $\Delta_{M(2)}$  is given by the product of the heat kernels  $p_t$  on  $\mathbb{R}^2$  and  $q_t$  on  $SO(2)$ . In other words,

$$\psi_t(x, e^{i\alpha}) = p_t(x)q_t(e^{i\alpha}) = \frac{1}{4\pi t} e^{-\frac{|x|^2}{4t}} \sum_{n \in \mathbb{Z}} e^{-n^2 t} e^{in\alpha}.$$

Let  $f \in L^2(M(2))$ . Expanding  $f$  in the  $SO(2)$  variable we obtain

$$f(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} f_m(x) e^{im\alpha}$$

where  $f_m(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x, e^{i\alpha}) e^{-im\alpha} d\alpha$  and the convergence is understood in the  $L^2$ -sense. Since  $p_t$  is radial (as a function on  $\mathbb{R}^2$ ) a simple computation shows that

$$f * p_t(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} f_m * p_t(x) e^{-m^2 t} e^{im\alpha}.$$

Let  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$  and  $\mathcal{H}(\mathbb{C}^2 \times \mathbb{C}^*)$  be the Hilbert space of holomorphic functions on  $\mathbb{C}^2 \times \mathbb{C}^*$  which are square integrable with respect to  $\mu \otimes \nu(z, w)$  where

$$d\mu(z) = \frac{1}{2\pi t} e^{-\frac{|z|^2}{2t}} dx dy \text{ on } \mathbb{C}^2$$

and

$$d\nu(w) = \frac{1}{2\pi} \frac{1}{\sqrt{2\pi t}} \frac{e^{-\frac{(\ln|w|)^2}{2t}}}{|w|^2} dw \text{ on } \mathbb{C}^*.$$

Using the Segal-Bargmann result for  $\mathbb{R}^2$  and  $S^1$  we can easily prove the following theorem:

**Theorem 1.1.** *If  $f \in L^2(M(2))$ , then  $f * p_t$  extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$ . Moreover, the map  $f \rightarrow f * p_t$  is a unitary map from  $L^2(M(2))$  onto  $\mathcal{H}(\mathbb{C}^2 \times \mathbb{C}^*)$ .*

## 2. GENERALIZATIONS OF SEGAL-BARGMANN TRANSFORM

In this section we study generalizations of the Segal-Bargmann transform and prove an analogue of Theorem 8 and Theorem 10 in [3].

Let  $\mu$  be any radial real-valued function on  $\mathbb{R}^2$  such that

- $\mu(x) > 0 \forall x \in \mathbb{R}^2$  and is locally bounded away from zero,
- $\forall x \in \mathbb{R}^2, \sigma(x) = \int_{\mathbb{R}^2} e^{2x \cdot y} \mu(y) dy < \infty$ .

Define, for  $z \in \mathbb{C}^2$

$$\psi(z) = \int_{\mathbb{R}^2} \frac{e^{ia(y)}}{\sqrt{\sigma(y)}} e^{-iy \cdot z} dy,$$

where  $a$  is a real valued measurable function on  $\mathbb{R}^2$ . Next, let  $\nu$  be a measure on  $\mathbb{C}^*$  such that

- $\nu$  is  $S^1$ -invariant,
- $\nu$  is given by a positive density which is locally bounded away from zero,
- $\forall n \in \mathbb{Z}, \delta(n) = \int_{\mathbb{C}^*} |w|^{2n} d\nu(w) < \infty$ .

Define  $\chi(w) = \sum_{n \in \mathbb{Z}} \frac{c_n}{\sqrt{\delta(n)}} w^n$  for  $w \in \mathbb{C}^*$  and  $c_n \in \mathbb{C}$  such that  $|c_n| = 1$ .

Also define  $\phi(z, w) = \psi(z)\chi(w)$  for  $z \in \mathbb{C}^2, w \in \mathbb{C}^*$ . It is easy to see that  $\phi(z, w)$  is a holomorphic function on  $\mathbb{C}^2 \times \mathbb{C}^*$ . We have the following Paley-Wiener type theorem.

**Theorem 2.1.** *The mapping*

$$C_\phi f(z, w) = \int_{M(2)} f(\xi, e^{i\alpha}) \phi((\xi, e^{i\alpha})^{-1}(z, w)) d\xi d\alpha$$

*is an isometric isomorphism of  $L^2(M(2))$  onto*

$$\mathcal{O}(\mathbb{C}^2 \times \mathbb{C}^*) \cap L^2(\mathbb{C}^2 \times \mathbb{C}^*, \mu(y) dx dy d\nu(w)).$$

*Proof.* Let  $f \in L^2(M(2))$  and

$$(2.1) \quad f(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} f_m(x) e^{im\alpha}$$

where  $f_m(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x, e^{i\alpha}) e^{-im\alpha} d\alpha$ . Since the function  $\phi(x, e^{i\alpha})$ , for  $(x, e^{i\alpha}) \in M(2)$  is radial in the  $\mathbb{R}^2$  variable  $x$ , a simple computation shows that the Fourier series of  $f * \phi(x, e^{i\alpha})$  is given by

$$f * \phi(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} f_m * \psi(x) \frac{c_m}{\sqrt{\delta(m)}} e^{im\alpha}.$$

Now, notice that  $f_m \in L^2(\mathbb{R}^2)$ ,  $\forall m \in \mathbb{Z}$  and  $f_m * \psi$  is a holomorphic function on  $\mathbb{C}^2$ . Moreover, by Theorem 8 of [3] we have

$$(2.2) \quad \int_{\mathbb{C}^2} |f_m * \psi(z)|^2 \mu(y) dx dy = \int_{\mathbb{R}^2} |f_m(x)|^2 dx.$$

Naturally, the analytic continuation of  $f * \phi(x, e^{i\alpha})$  to  $\mathbb{C}^2 \times \mathbb{C}^*$  is given by

$$(2.3) \quad f * \phi(z, w) = \sum_{m \in \mathbb{Z}} f_m * \psi(z) \frac{c_m}{\sqrt{\delta(m)}} w^m.$$

We show that the series in (2.3) converges uniformly on compact sets in  $\mathbb{C}^2 \times \mathbb{C}^*$  proving the holomorphicity. Let  $K$  be a compact set in  $\mathbb{C}^2 \times \mathbb{C}^*$ . For  $(z, w) \in K$  we have

$$(2.4) \quad \left| \sum_{m \in \mathbb{Z}} f_m * \psi(z) \frac{c_m}{\sqrt{\delta(m)}} w^m \right| \leq \sum_{m \in \mathbb{Z}} |f_m * \psi(z)| \frac{|w|^m}{\sqrt{\delta(m)}}.$$

By Fourier inversion (see also Theorem 8 in [3])

$$f_m * \psi(z) = \int_{\mathbb{R}^2} \widetilde{f}_m(\xi) \frac{e^{ia(\xi)}}{\sqrt{\sigma(\xi)}} e^{-i\xi(x+iy)} d\xi$$

where  $z = x + iy \in \mathbb{C}^2$  and  $\widetilde{f}_m$  is the Fourier transform of  $f_m$ . Hence, if  $z$  varies in a compact subset of  $\mathbb{C}^2$ , we have

$$\begin{aligned} |f_m * \psi(z)| &\leq \|f_m\|_2 \left( \int_{\mathbb{R}^2} \frac{e^{2\xi \cdot y}}{\sigma(\xi)} d\xi \right)^{\frac{1}{2}} \\ &\leq C \|f_m\|_2. \end{aligned}$$

Using the above in (2.4) and assuming  $|w| \leq R$  (as  $w$  varies in a compact set in  $\mathbb{C}^*$ ) we have

$$\left| \sum_{m \in \mathbb{Z}} f_m * \psi(z) \frac{c_m}{\sqrt{\delta(m)}} w^m \right| \leq C \sum_{m \in \mathbb{Z}} \|f_m\|_2 \frac{R^m}{\sqrt{\delta(m)}}.$$

Applying Cauchy-Schwarz inequality to the above, noting that

$$\sum_{m \in \mathbb{Z}} \|f_m\|_2^2 = \|f\|_2^2 \text{ and } \sum_{m \in \mathbb{Z}} \frac{R^{2m}}{\delta(m)} < \infty$$

we prove the above claim. Applying Theorem 10 in [3] for  $S^1$  we obtain

$$\int_{\mathbb{C}^*} |f * \phi(z, w)|^2 d\nu(w) = \sum_{m \in \mathbb{Z}} |f_m * \psi(z)|^2.$$

Integrating the above against  $\mu(y) dx dy$  on  $\mathbb{C}^2$  and using (2.2) we obtain that

$$\int_{\mathbb{C}^2} \int_{\mathbb{C}^*} |f * \phi(z, w)|^2 \mu(y) dx dy d\nu(w) = \|f\|_2^2.$$

To prove that the map  $C_\phi$  is surjective it suffices to prove that the range of  $C_\phi$  is dense in  $\mathcal{O}(\mathbb{C}^2 \times \mathbb{C}^*) \cap L^2(\mathbb{C}^2 \times \mathbb{C}^*, \mu(y) dx dy d\nu(w))$ . For this, consider functions of the form  $f(x, e^{i\alpha}) = g(x) e^{im\alpha} \in L^2(M(2))$  where  $g \in L^2(\mathbb{R}^2)$ . Then a simple computation shows that

$$f * \phi(z, w) = g * \psi(z) w^m \text{ for } (z, w) \in \mathbb{C}^2 \times \mathbb{C}^*.$$

Suppose  $F \in \mathcal{O}(\mathbb{C}^2 \times \mathbb{C}^*) \cap L^2(\mathbb{C}^2 \times \mathbb{C}^*, \mu(y) dx dy d\nu(w))$  be such that

$$(2.5) \quad \int_{\mathbb{C}^2 \times \mathbb{C}^*} F(z, w) \overline{g * \psi(z)} \bar{w}^m \mu(y) dx dy d\nu(w) = 0$$

$\forall g \in L^2(\mathbb{R}^2)$  and  $\forall m \in \mathbb{Z}$ . From (2.5) we have

$$\int_{\mathbb{C}^*} \left( \int_{\mathbb{C}^2} F(z, w) \overline{g * \psi(z)} d\mu(z) \right) \bar{w}^m d\nu(w) = 0,$$

which by Theorem 10 of [3] implies that

$$\int_{\mathbb{C}^2} F(z, w) \overline{g * \psi(z)} d\mu(z) = 0.$$

Finally, an application of Theorem 8 of [3] shows that  $F \equiv 0$ . Hence the proof.  $\square$

### 3. POISSON INTEGRALS AND PALEY-WIENER TYPE THEOREMS

In this section we study the Poisson integrals on  $M(2)$ . We also find conditions on the 'group Fourier transform' of a function so that it extends holomorphically to an appropriate domain in the complexification of the group. We start with the following Gutzmer-type lemma:

**Lemma 3.1.** *Let  $f \in L^2(M(2))$  extend holomorphically to the domain*

$$\{(z, w) \in \mathbb{C}^2 \times \mathbb{C}^* : |Imz| < t, \frac{1}{R} < |w| < R\}$$

and

$$\sup_{\{|y| < s, \frac{1}{N} < |w| < N\}} \int_{M(2)} |f(x + iy, |w|e^{i\theta})|^2 dx d\theta < \infty$$

$\forall s < t$  and  $N < R$ . Then

$$\int_{M(2)} |f(x + iy, |w|e^{i\theta})|^2 dx d\theta = \sum_{n \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\tilde{f}_n(\xi)|^2 e^{-2\xi \cdot y} d\xi \right) |w|^{2n}$$

provided  $|y| < t$  and  $\frac{1}{R} < |w| < R$ . Conversely, if

$$\sup_{\{|y| < s, \frac{1}{N} < |w| < N\}} \sum_{n \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\tilde{f}_n(\xi)|^2 e^{-2\xi \cdot y} d\xi \right) |w|^{2n} < \infty \quad \forall s < t \text{ and } N < R$$

then  $f$  extends holomorphically to the domain

$$\left\{ (z, w) \in \mathbb{C}^2 \times \mathbb{C}^* : |Imz| < t, \frac{1}{R} < |w| < R \right\}$$

and

$$\sup_{\{|y| < s, \frac{1}{N} < |w| < N\}} \int_{M(2)} |f(x + iy, |w|e^{i\theta})|^2 dx d\theta < \infty \quad \forall s < t \text{ and } N < R.$$

*Proof.* Notice that  $f_n(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x, e^{i\alpha}) e^{-in\alpha} d\alpha$ . It follows that  $f_n(x)$  has a holomorphic extension to  $\{z \in \mathbb{C}^2 : |Imz| < t\}$  and

$$\sup_{|y| < s} \int_{\mathbb{R}^2} |f_n(x + iy)|^2 dx < \infty \quad \forall s < t.$$

Consequently,

$$\int_{\mathbb{R}^2} |f_n(x + iy)|^2 dx = \int_{\mathbb{R}^2} |\tilde{f}_n(\xi)|^2 e^{-2\xi \cdot y} d\xi \text{ for } |y| < s \quad \forall s < t.$$

Now, for each fixed  $z \in \mathbb{C}^2$  with  $|Imz| < s$  the function  $w \rightarrow f(z, w)$  is holomorphic in the annulus  $\{w \in \mathbb{C}^* : \frac{1}{N} < |w| < N\}$  for every  $s < t$  and  $N < R$  and so admits a Laurent series expansion

$$f(z, w) = \sum_{m \in \mathbb{Z}} a_m(z) w^m.$$

It follows that  $a_m(z) = f_m(z) \quad \forall m \in \mathbb{Z}$ . First part of the lemma is proved now by appealing to the Plancherel theorem on  $S^1$  and  $\mathbb{R}^2$ . Converse can also be proved similarly.

□

Recall from the introduction that the Laplacian  $\Delta$  on  $M(2)$  is given by  $\Delta = -\Delta_{\mathbb{R}^2} - \frac{\partial^2}{\partial \alpha^2}$ . If  $f \in L^2(M(2))$  it is easy to see that

$$e^{-t\Delta^{\frac{1}{2}}} f(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} \widetilde{f}_m(\xi) e^{-t(|\xi|^2 + m^2)^{\frac{1}{2}}} e^{ix \cdot \xi} d\xi \right) e^{im\alpha}.$$

We have the following (almost) characterization of the Poisson integrals. Let  $\Omega_s$  denote the domain in  $\mathbb{C}^2 \times \mathbb{C}^*$  defined by

$$\Omega_s = \{(z, w) : |Imz| < s, e^{-s} < |w| < e^s\}.$$

**Theorem 3.2.** *Let  $f \in L^2(M(2))$ . Then  $g = e^{-t\Delta^{\frac{1}{2}}} f$  extends to a holomorphic function on the domain  $\Omega_{\frac{t}{\sqrt{2}}}$  and*

$$\sup_{\left\{ |y| < \frac{t}{\sqrt{2}}, e^{-\frac{t}{\sqrt{2}}} < |w| < e^{\frac{t}{\sqrt{2}}} \right\}} \int_{M(2)} |g(x + iy, |w|e^{i\alpha})|^2 dx d\alpha < \infty.$$

Conversely, let  $g$  be a holomorphic function on  $\Omega_t$  and

$$\sup_{\{|y| < s, e^{-s} < |w| < e^s\}} \int_{M(2)} |g(x + iy, |w|e^{i\alpha})|^2 dx d\alpha < \infty \text{ for } s < t.$$

Then  $\forall s < t, \exists f \in L^2(M(2))$  such that  $e^{-s\Delta^{\frac{1}{2}}} f = g$ .

*Proof.* We know that, if  $f \in L^2(M(2))$  then

$$g(x, e^{i\alpha}) = e^{-t\Delta^{\frac{1}{2}}} f(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} \widetilde{f}_m(\xi) e^{-t(|\xi|^2 + m^2)^{\frac{1}{2}}} e^{ix \cdot \xi} d\xi \right) e^{im\alpha}.$$

Also,  $g(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} g_m(x) e^{im\alpha}$  with  $\widetilde{g}_m(\xi) = \widetilde{f}_m(\xi) e^{-t(|\xi|^2 + m^2)^{\frac{1}{2}}}$ . If  $s \leq \frac{t}{\sqrt{2}}$  it is easy to show that

$$\sup_{\{\xi \in \mathbb{R}^2, m \in \mathbb{Z}\}} e^{-2t(|\xi|^2 + m^2)^{\frac{1}{2}}} e^{2|\xi||y|} e^{2|m|s} \leq C < \infty \text{ for } |y| \leq \frac{t}{\sqrt{2}}.$$

It follows that

$$\sup_{\left\{ |y| < \frac{t}{\sqrt{2}}, e^{-\frac{t}{\sqrt{2}}} < |w| < e^{\frac{t}{\sqrt{2}}} \right\}} \sum_{m \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\widetilde{g}_m(\xi)|^2 e^{-2\xi \cdot y} d\xi \right) |w|^{2m} < \infty.$$

By the previous lemma we prove the first part of the theorem.

Conversely, let  $g$  be a holomorphic function on  $\Omega_t$  and

$$\sup_{\{|y| < s, e^{-s} < |w| < e^s\}} \int_{M(2)} |g(x + iy, |w|e^{i\alpha})|^2 dx d\alpha < \infty \text{ for } s < t.$$

By Lemma 3.1 we have

$$\sup_{\{|y| < s, e^{-s} < |w| < e^s\}} \sum_{n \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\tilde{g}_n(\xi)|^2 e^{-2\xi \cdot y} d\xi \right) |w|^{2n} < \infty \text{ for } s < t.$$

Integrating the above over  $|y| = s < t$ , we obtain

$$\sum_{n \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\tilde{g}_n(\xi)|^2 J_0(i2s|\xi|) d\xi \right) |w|^{2n} < \infty,$$

where  $J_0$  is the Bessel function of first kind. Noting that  $J_0(i2s|\xi|) \sim e^{2s|\xi|}$  for large  $|\xi|$  we obtain

$$\sum_{n \in \mathbb{Z}} \int_{\mathbb{R}^2} |\tilde{g}_n(\xi)|^2 e^{2s|\xi|} e^{2|n|s} d\xi < \infty \text{ for } s < t.$$

This surely implies that

$$\sum_{n \in \mathbb{Z}} \int_{\mathbb{R}^2} |\tilde{g}_n(\xi)|^2 e^{2s(|\xi|^2 + m^2)^{\frac{1}{2}}} d\xi < \infty \text{ for } s < t.$$

Defining  $\tilde{f}_m(\xi)$  by  $\tilde{f}_m(\xi) = \tilde{g}_m(\xi) e^{s(|\xi|^2 + m^2)^{\frac{1}{2}}}$  we obtain

$$f(x, e^{i\alpha}) = \sum_{m \in \mathbb{Z}} f_m(x) e^{im\alpha} \in L^2(M(2))$$

and  $g = e^{-s\Delta^{\frac{1}{2}}} f$ .

□

*Remark 3.3.* A similar result may be proved for the operator  $e^{-t\Delta_{\mathbb{R}^2}^{\frac{1}{2}}} e^{-t\Delta_{S^1}^{\frac{1}{2}}}$ .

**Analytic vectors.** Let  $\pi$  be a unitary representation of a Lie group  $G$  on a Hilbert space  $H$ . A vector  $v \in H$  is called an analytic vector for  $\pi$  if the function  $g \rightarrow \pi(g)v$  is analytic.

Recall the representations  $U_g^\xi$  from the introduction. Denote by  $U_g^a$  the representations  $U_g^{(a,0)}$  for  $a > 0$ . If  $e_n(\theta) = e^{in\theta} \in L^2(S^1)$ , it is easy to see that  $e_n$ 's are analytic vectors for these representations. For  $g = (x, e^{i\alpha}) \in M(2)$ , we have

$$(U_g^a e_n)(\theta) = e^{i\langle x, ae^{i\theta} \rangle} e^{in(\theta - \alpha)}.$$

This action of  $U_g^a$  on  $e_n$  can clearly be analytically continued to  $\mathbb{C}^2 \times \mathbb{C}^*$  and we obtain

$$(U_{(z,w)}^a e_n)(\theta) = e^{i\langle x, ae^{i\theta} \rangle} e^{-\langle y, ae^{i\theta} \rangle} w^{-n} e^{in\theta}$$

where  $(z, w) \in \mathbb{C}^2 \times \mathbb{C}^*$  and  $z = x + iy \in \mathbb{C}^2$ .

We also note that the action of  $S^1$  on  $\mathbb{R}^2$  naturally extends to an action of  $\mathbb{C}^*$  on  $\mathbb{C}^2$  given by

$$w(z_1, z_2) = (z_1 \cos \zeta - z_2 \sin \zeta, z_1 \sin \zeta + z_2 \cos \zeta)$$

where  $w = e^{i\zeta} \in \mathbb{C}^*$  and  $(z_1, z_2) \in \mathbb{C}^2$ .

Our next theorem is in the same spirit of Theorem 3.1 from Goodman [2].

**Theorem 3.4.** *Let  $f \in L^2(M(2))$ . Then  $f$  extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$  with*

$$\int_{|y|=r} \int_{M(2)} |f(w^{-1}(x + iy), |w|e^{i\alpha})|^2 dx d\alpha d\sigma_r(y) < \infty$$

(where  $\sigma_r$  is the normalized surface area measure on the sphere  $\{|y| = r\} \subseteq \mathbb{R}^2$ ) iff

$$\int_0^\infty \int_{|y|=r} \|U_{(z,w)}^a \hat{f}(a)\|_{HS}^2 d\sigma_r(y) da < \infty$$

where  $z = x + iy \in \mathbb{C}^2$  and  $w \in \mathbb{C}^*$ . In this case we also have

$$\begin{aligned} & \int_0^\infty \int_{|y|=r} \|U_{(z,w)}^a \hat{f}(a)\|_{HS}^2 d\sigma_r(y) da \\ &= \int_{|y|=r} \int_{M(2)} |f(w^{-1}(x + iy), |w|e^{i\alpha})|^2 dx d\alpha d\sigma_r(y). \end{aligned}$$

*Proof.* First assume that  $f \in L^2(M(2))$  satisfies the transformation property

$$(3.1) \quad f(e^{i\theta}x, e^{i\alpha}) = e^{im\theta} f(x, e^{i\alpha})$$

for some fixed  $m \in \mathbb{Z}$  and  $\forall (x, e^{i\alpha}) \in M(2)$ . As earlier we have

$$\left( \hat{f}(a)e_n \right) (\theta) = \tilde{f}_n(ae^{i\theta})e^{in\theta}.$$

By the Hecke-Bochner identity, we have

$$\tilde{f}_n(ae^{i\theta}) = i^{-|m|} a^{|m|} (\mathcal{F}_{2+2|m|}g)(a)e^{im\theta}$$

where  $\mathcal{F}_{2+2|m|}(g)$  is the  $2+2|m|$ -dimensional Fourier transform of  $g(x) = \frac{f_n(|x|)}{|x|^{|m|}}$ , considered as a radial function on  $\mathbb{R}^{2+2|m|}$ .

Hence,

$$\left( \hat{f}(a)e_n \right) (\theta) = i^{-|m|} a^{|m|} (\mathcal{F}_{2+2|m|}g)(a)e^{i(m+n)\theta}.$$

It follows that  $\hat{f}(a)e_n$  is an analytic vector and we can apply  $U_{(z,w)}^a$  to the above. We obtain

$$\begin{aligned} & \left( U_{(z,w)}^a \hat{f}(a)e_n \right) (\theta) \\ &= e^{i\langle x, ae^{i\theta} \rangle} e^{-\langle y, ae^{i\theta} \rangle} i^{-|m|} a^{|m|} (\mathcal{F}_{2+2|m|} g)(a) w^{-(m+n)} e^{i(m+n)\theta}. \end{aligned}$$

Thus,

$$\begin{aligned} & \int_{S^1} \left| \left[ U_{(z,w)}^a \hat{f}(a)e_n \right] (\theta) \right|^2 d\theta \\ &= |w|^{-2(m+n)} \int_{S^1} a^{2m} |(\mathcal{F}_{2+2m} g)(a)|^2 e^{-2\langle y, ae^{i\theta} \rangle} d\theta \\ &= |w|^{-2(m+n)} \int_{S^1} e^{-2\langle y, ae^{i\theta} \rangle} |\tilde{f}_n(ae^{i\theta})|^2 d\theta. \end{aligned}$$

Hence,

$$(3.2) \quad \begin{aligned} & \int_0^\infty \|U_{(z,w)}^a \hat{f}(a)\|_{HS}^2 ada \\ &= |w|^{-2m} \sum_{n \in \mathbb{Z}} |w|^{-2n} \left( \int_{\mathbb{R}^2} e^{-2\langle y, \xi \rangle} |\tilde{f}_n(\xi)|^2 d\xi \right). \end{aligned}$$

Notice that, if  $f$  extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$  we must have  $f(w^{-1}z, w) = w^{-m} f(z, w) \forall (z, w) \in \mathbb{C}^2 \times \mathbb{C}^*$  because of (3.1).

In view of Lemma 3.1, the above remark and the identity (3.2), the theorem is established for functions with transformation property (3.1) and we obtain

$$(3.3) \quad \begin{aligned} & \int_0^\infty \int_{|y|=r} \|U_{(z,w)}^a \hat{f}(a)\|_{HS}^2 d\sigma_r(y) ada \\ &= \int_{|y|=r} \int_{M(2)} |f(w^{-1}(x+iy), |w|e^{i\alpha})|^2 dx d\alpha d\sigma_r(y). \end{aligned}$$

Next, we deal with the general case. For  $f \in L^2(M(2))$  define

$$f^m(x, e^{i\alpha}) = \int_{S^1} f(e^{i\theta}x, e^{i\alpha}) e^{-im\theta} d\theta.$$

Then  $f^m(e^{i\theta}x, e^{i\alpha}) = e^{im\theta} f^m(x, e^{i\alpha})$  and  $f_m$ 's are orthogonal on  $M(2)$ . Assume that  $f$  extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$ . Then, so does  $f^m \forall m \in \mathbb{Z}$  and we have

$$(3.4) \quad \begin{aligned} & \int_{|y|=r} \int_{M(2)} |f(w^{-1}(x+iy), |w|e^{i\alpha})|^2 dx d\alpha d\sigma_r(y) \\ &= \sum_{m \in \mathbb{Z}} \int_{|y|=r} \int_{M(2)} |f^m(w^{-1}(x+iy), |w|e^{i\alpha})|^2 dx d\alpha d\sigma_r(y). \end{aligned}$$

This follows from the fact that

$$\int_{|y|=r} \int_{\mathbb{R}^2} f^m(w^{-1}(x+iy), w) \overline{f^l(w^{-1}(x+iy), w)} dx d\sigma_r(y) = 0 \text{ if } m \neq l.$$

Applying identity (3.3) we get from (3.4)

$$(3.5) \quad \sum_{m \in \mathbb{Z}} \int_0^\infty \int_{|y|=r} \|U_{(z,w)}^a \widehat{f^m}(a)\|_{HS}^2 adad\sigma_r(y) < \infty.$$

Now, let  $\langle T, S \rangle_{HS} = \sum_{n \in \mathbb{Z}} \langle Te_n, Se_n \rangle$  denote the inner product in the space of Hilbert-Schmidt operators on  $L^2(S^1)$ . Then we notice that

$$(3.6) \quad \begin{aligned} & \int_{|y|=r} \left\langle U_{(z,w)}^a \widehat{f^m}(a), U_{(z,w)}^a \widehat{f^l}(a) \right\rangle_{HS} d\sigma_r(y) \\ &= \delta_{ml} \sum_{n \in \mathbb{Z}} (-1)^{-l} \left(\frac{a}{i}\right)^{m+l} (\mathcal{F}_{2+2|m|} g^m)(a) \\ & \quad \frac{\overline{(\mathcal{F}_{2+2|l|} g^l)(a)}}{w^{-(m+n)} (\overline{w})^{-(l+n)}} J_0(2ira). \end{aligned}$$

Hence (3.5) implies that

$$\int_0^\infty \int_{|y|=r} \|U_{(z,w)}^a \widehat{f}(a)\|_{HS}^2 adad\sigma_r(y) < \infty.$$

To prove the converse, we first show that  $f$  has a holomorphic extension to whole of  $\mathbb{C}^2 \times \mathbb{C}^*$ . Recall that we have

$$\left( \widehat{f}(a) e_n \right) (\theta) = \widetilde{f}_n(ae^{i\theta}) e^{in\theta}.$$

Expanding  $\widetilde{f}_n(ae^{i\theta})$  into Fourier series we have

$$\widetilde{f}_n(ae^{i\theta}) = \sum_{k \in \mathbb{Z}} C_{a,n}(k) e^{ik\theta}.$$

Hence  $\left( U_{(z,w)}^a \widehat{f}(a) e_n \right) (\theta)$  is given by

$$\sum_{k \in \mathbb{Z}} C_{a,n}(k) e^{i\langle x, ae^{i\theta} \rangle} e^{-\langle y, ae^{i\theta} \rangle} w^{-(k+n)} e^{i(k+n)\theta}.$$

Thus

$$\int_{|y|=r} \int_{S^1} \left| \left[ U_{(z,w)}^a \widehat{f}(a) e_n \right] (\theta) \right|^2 d\theta d\sigma_r(y) = J_0(2ira) \sum_{k \in \mathbb{Z}} |C_{a,n}(k)|^2 |w|^{-(k+n)}.$$

Notice that  $J_0(2ira) \sim e^{2ra}$  for large  $a$  and  $\sum_{k \in \mathbb{Z}} |C_{a,n}(k)|^2 = \int_{S^1} |\tilde{f}_n(ae^{i\theta})|^2 d\theta$ .

If  $e^{-r} < |w| < e^r$ , we obtain

$$\sum_{n \in \mathbb{Z}} \left( \int_0^\infty \int_{S^1} |\tilde{f}_n(ae^{i\theta})|^2 e^{2ra} d\theta da \right) |w|^{2n} < \infty,$$

which implies

$$\sum_{n \in \mathbb{Z}} \left( \int_{\mathbb{R}^2} |\tilde{f}_n(\xi)|^2 e^{2r|\xi|} d\xi \right) |w|^{2n} < \infty.$$

Since this is true for all  $r > 0$  and  $w \in \mathbb{C}^*$ , by Lemma 3.1  $f$  extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$ . It follows that  $f^m(x, e^{i\alpha})$  defined by

$$f^m(x, e^{i\alpha}) = \int_{S^1} f(e^{i\theta}x, e^{i\alpha}) e^{-im\theta} d\theta$$

also extends holomorphically to  $\mathbb{C}^2 \times \mathbb{C}^*$  and

$$\int_{M(2)} |f^m((x+iy), |w|e^{i\alpha})|^2 dx d\alpha < \infty.$$

Now the proof can be completed using the identity (3.3), orthogonality of  $U_{(z,w)}^a \widehat{f^m}(a)$  (see (3.6)) and (3.4). □

#### 4. A PALEY-WIENER THEOREM FOR THE INVERSE FOURIER TRANSFORM

Recall from the introduction that the 'group Fourier Transform'  $\hat{f}(\xi)$ , for  $\xi \in \mathbb{R}^2$  of  $f \in L^1(M(2))$  is an integral operator with the kernel  $k_f(\xi, e^{i\alpha}, e^{i\beta})$  where  $k_f(\xi, e^{i\alpha}, e^{i\beta}) = \tilde{f}(e^{i\beta}\xi, e^{i(\beta-\alpha)})$ ,  $\tilde{f}$  being the Euclidean Fourier Transform of  $f$  in the  $\mathbb{R}^2$ -variable. We have the following Paley-Wiener theorem for the inverse Fourier Transform :

**Theorem 4.1.** *Let  $f \in L^1(M(2))$  be such that  $\hat{f}(\xi) \equiv 0 \forall |\xi| > R$  and the kernel  $k_f$  of  $\hat{f}(\xi)$  is smooth on  $\mathbb{R}^2 \times S^1 \times S^1$ . Then  $x \rightarrow f(x, e^{i\alpha})$  extends to an entire function of exponential type  $R$  such that*

$$(4.1) \quad \sup_{e^{i\alpha} \in S^1} |z^m f(z, e^{i\alpha})| \leq c_m e^{R|Imz|}, \quad \forall z \in \mathbb{C}^2, \quad \forall m \in \mathbb{N}^2.$$

*Conversely, if  $f$  extends to an entire function on  $\mathbb{C}^2$  in the first variable and satisfies (4.1) then  $\hat{f}(\xi) = 0$  for  $|\xi| > R$  and  $k_f$  is smooth on  $\mathbb{R}^2$ .*

*Proof.* We have

$$(\hat{f}(\xi)F)(e^{i\alpha}) = \int_{S^1} k_f(\xi, e^{i\alpha}, e^{i\beta})F(e^{i\beta})d\beta, \quad \text{for } F \in L^2(S^1)$$

where  $k_f(\xi, e^{i\alpha}, e^{i\beta}) = \tilde{f}(e^{i\beta}\xi, e^{i(\beta-\alpha)})$ .

Assume that  $\hat{f}(\xi) \equiv 0 \forall |\xi| > R$ . Since  $k_f$  is smooth we have  $\tilde{f}(\cdot, e^{i\alpha}) \in \mathcal{C}_c^\infty(\mathbb{R}^2)$ . By the Paley-Wiener theorem on  $\mathbb{R}^2$  we obtain that  $f(\cdot, e^{i\alpha})$  extends to an entire function on  $\mathbb{C}^2$  of exponential type. Moreover, if  $m \in \mathbb{N}^2$

$$z^m f(z, e^{i\alpha}) = \int_{|\xi| \leq R} \frac{\partial^m \tilde{f}}{\partial \xi^m}(\xi, e^{i\alpha}) e^{i\xi \cdot z} d\xi.$$

It follows that

$$\sup_{e^{i\alpha} \in S^1} |z^m f(z, e^{i\alpha})| \leq \sup_{e^{i\alpha} \in S^1} \left\| \frac{\partial^m \tilde{f}}{\partial \xi^m}(\cdot, e^{i\alpha}) \right\|_1 e^{R|Imz|}.$$

Conversely, if  $f$  is holomorphic on  $\mathbb{C}^2$  and satisfies (4.1), by the Paley-Wiener theorem on  $\mathbb{R}^2$  we get  $\tilde{f}(\cdot, e^{i\alpha}) \in \mathcal{C}_c^\infty(\mathbb{R}^2)$  and  $\tilde{f}(\xi, e^{i\alpha}) = 0 \forall |\xi| > R$ . Hence  $\hat{f}(\xi) \equiv 0 \forall |\xi| > R$ . Moreover,  $k_f$  is smooth on  $\mathbb{R}^2$  since  $\tilde{f}$  is smooth.  $\square$

## REFERENCES

- [1] R. W. Goodman, *Analytic and entire vectors for representations of Lie groups*, Trans. Amer. Math. Soc., 143 (1969), 55–76.
- [2] R. W. Goodman, *Complex Fourier analysis on a nilpotent Lie group*, Trans. Amer. Math. Soc., 160 (1971), 373–391.
- [3] B. C. Hall, *The Segal-Bargmann “coherent state” transform for compact Lie groups*, J. Funct. Anal., 122 (1994) no. 1, 103–151.
- [4] B. C. Hall, W. Lewkeeratiyutkul, *Holomorphic Sobolev spaces and the generalized Segal-Bargmann transform*, J. Funct. Anal., 217 (2004) no. 1, 192–220.
- [5] B. C. Hall, J. J. Mitchell, *The Segal-Bargmann transform for non compact symmetric spaces of the complex type*, J. Funct. Anal., 227 (2005) no. 2, 338–371.
- [6] B. Krötz, G. Ólafsson, R. J. Stanton, *The image of the heat kernel transform on Riemannian symmetric spaces of the non compact type*, Int. Math. Res. Not., (2005) no. 22, 1307–1329.
- [7] B. Krötz, S. Thangavelu, Y. Xu, *The heat kernel transform for the Heisenberg group*, J. Funct. Anal., 225 (2005), no. 2, 301–336.
- [8] A. Pasquale, *A Paley-Wiener theorem for the inverse spherical transform*, Pacific J. Math., 193 (2000), no. 1, 143–176.
- [9] M. B. Stenzel, *The Segal-Bargmann transform on a symmetric space of compact type*, J. Funct. Anal., 165 (1999) no. 1, 44–58.
- [10] M. Sugiura, *Fourier series of smooth functions on compact Lie groups*, Osaka J. Math., 8 (1971), 33–47.

- [11] S. Thangavelu, *A Paley-Wiener theorem for the inverse Fourier transform on some homogeneous spaces*, Hiroshima Math. J., 37 (2007), no. 2, 145–159.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE  
- 560012, INDIA.

*E-mail address:* `naru@math.iisc.ernet.in`, `suparna@math.iisc.ernet.in`.