# THE TOTEM EXPERIMENT AT THE LHC

G. LATINO

(on behalf of the TOTEM Collaboration)

INFN Pisa & University of Siena, Department of Physics, Via Roma, 56 - 53100 Siena, Italy

The TOTEM experiment at the LHC is dedicated to the measurement of the total *pp* cross section and to the study of elastic scattering and of diffractive dissociation processes. TOTEM is here presented with a general overview on the main features of its experimental apparatus and of its physics programme.

### 1 Introduction

In order to fulfil its physics programme the TOTEM experiment<sup>1</sup> has to cope with the challenge of triggering and recording events in the very forward region with a good acceptance for particles produced at very small angles with respect to the beam. The evaluation of the total cross section with an error down to  $1\div 2\%$  will in particular require the simultaneous measurement of the elastic scattering at low values for the squared four-momentum transfer  $(|t| \sim (p\theta)^2)$  as well as of the inelastic interaction rate with losses reduced to few per-cents. This involves the detection of scattered protons at a location very close to the beam, together with efficient forward particle detection in inelastic events. A flexible trigger provided by its detectors will allow to take data under all LHC running scenarios. The combination of the CMS<sup>2</sup> and TOTEM experiments, representing the largest acceptance detector ever built at a hadron collider, will also allow the study of a wide range of diffractive processes with an unprecedented coverage in rapidity. For this purpose the TOTEM trigger and data acquisition (DAQ) systems are designed to be compatible with the CMS ones, in order to allow common data taking periods foreseen at a later stage<sup>3</sup>. In the following, after a general overview of the experimental apparatus, the main features of the TOTEM physics programme will be described.

### 2 Detector Components

Located on both sides of the interaction point IP5 (the same as for CMS) the TOTEM experimental apparatus comprises "Roman Pots" (RPs) detectors and the T1 and T2 inelastic telescopes (Fig. 1). The RPs, placed on the beam-pipe of the outgoing beam in two stations at about 147 m and 220 m from IP5, are special movable beam-pipe insertions designed to detect "leading" protons with a scattering angle down to few  $\mu$ rad. T1 and T2, embedded inside the forward region of CMS, provide charged track reconstruction for  $3.1 < |\eta| < 6.5$  ( $\eta = -ln(tan\frac{\theta}{2})$ ) with a  $2\pi$  coverage and with a very good efficiency. These detectors, providing trigger signals with acceptance grater than 95% for all inelastic events, will allow the measurement of inelastic rates with small losses and will be also used for the reconstruction of the event interaction vertex, so to reject background events<sup>3</sup>. The read-out of all TOTEM sub-detectors is based on the digital VFAT chip<sup>4</sup>, specifically designed for TOTEM and characterized by trigger capabilities.



Figure 1: Right: Roman Pots location along the LHC beam-line. Left: T1/T2 location in the forward region of the CMS detector. All TOTEM detectors are located on both sides of IP5.

The RPs host silicon detectors which are moved very close to the beam when it is in stable conditions. Each RP station is composed of two units in order to have a lever harm for local track reconstruction and trigger selections by track angle. Each unit consists of three pots, two vertical and one horizontal completing the acceptance for diffractively scattered protons. Each pot contains a stack of 10 planes of silicon strip detectors (Fig. 2, left). Each plane has 512 strips (pitch of 66  $\mu$ m) allowing a single hit resolution of about 20  $\mu$ m. As the detection of protons elastically scattered at angles down to few  $\mu$ rads requires a detector active area as close to the beam as ~ 1 mm, a novel "edgeless planar silicon" detector technology has been developed for TOTEM RPs in order to have an edge dead zone minimized to only about 50  $\mu$ m<sup>5</sup>.

Each T1 telescope arm, covering the range  $3.1 < |\eta| < 4.7$  and located at ~ 9 m from IP5, consists of five planes formed by six trapezoidal "Cathode Strip Chambers" (CSC)<sup>1</sup> (Fig. 2, centre). These CSCs, with 10 mm thick gas gap and a gas mixture of Ar/CO<sub>2</sub>/CF<sub>4</sub> (40%/50%/10%), give three measurements of the charged particle coordinates with a spatial resolution of about 1 mm: anode wires (pitch of 3 mm), also giving level-1 trigger information, are parallel to the trapezoid base; cathode strips (pitch of 5 mm) are rotated by  $\pm 60^{\circ}$  with respect to the wires.



Figure 2: Left: silicon detectors hosted in one pot. Centre (right): one half-arm of the T1 (T2) telescope.

The T2 telescope, based on "Gas Electron Multiplier" (GEM) technology<sup>6</sup>, extends charged track reconstruction to the range  $5.3 < |\eta| < 6.5^{1}$ . Ten aligned detectors planes, having an almost semicircular shape, are combined to form one of the half-arms located at ~ 13.5 m from IP5 (Fig. 2, right). This novel gas detector technology is optimal for the T2 telescope thanks to its good spatial resolution, excellent rate capability and good resistance to radiation. The T2 GEMs are characterized by a triple-GEM structure and a gas mixture of Ar/CO<sub>2</sub> (70%/30%)<sup>1</sup>. The read-out board has two separate layers with different patterns: one with 256x2 concentric circular strips (80  $\mu$ m wide, pitch of 400  $\mu$ m), allowing track radial coordinate reconstruction with a resolution of about 100  $\mu$ m; the other with a matrix of 24x65 pads (from 2x2 mm<sup>2</sup> to 7x7 mm<sup>2</sup> in size) providing level-1 trigger information and track azimuthal coordinate reconstruction.

## 3 Physics Programme

The physics goals of the TOTEM experiment are the measurement of the total pp cross section  $(\sigma_{tot})$  with a precision down to  $1 \div 2\%$ , the study of the nuclear elastic pp differential cross section  $(d\sigma_{el}/dt)$  over a wide range of  $|t| (\sim 10^{-3} < |t| < 10 \,\text{GeV}^2)$  and the development of a comprehensive physics programme on diffractive dissociation, partially in cooperation with CMS. These studies will allow to distinguish among different models of soft proton interactions, giving a deeper understanding of proton structure. Furthermore, the study of charged particle flow in the forward region will give precious input to cosmic ray MonteCarlo simulations.



Figure 3: Left: Fits from the COMPETE Coll. to available pp and  $p\bar{p}$  scattering and cosmic ray data. Right:  $d\sigma_{el}/dt$  at LHC as predicted by different models; t-acceptance ranges for different machine optics are also shown.

Fig. 3 (left), summarizing the existing measurements on pp and  $p\bar{p}$  scattering, shows predictions for  $\sigma_{tot}$  from the COMPETE Collaboration based on fits according to different models<sup>7</sup>. The best fit predicts  $\sigma_{tot} = 111.5 \pm 1.2^{+4.1}_{-2.1}$  mb for the LHC energy ( $\sqrt{s} = 14$  TeV). The large uncertainties on available high energy data give a big error (90÷130 mb), depending on the model used for the extrapolation. TOTEM aims to measure  $\sigma_{tot}$  with a precision down to 1÷2%, therefore allowing to discriminate among the different models. The measurement will be based on the "luminosity independent method" which, combining the optical theorem with the total rate, gives  $\sigma_{tot}$  and the machine luminosity ( $\mathcal{L}$ ) as functions of measurable rates:

$$\sigma_{tot} = \frac{16\pi}{1+\rho^2} \cdot \frac{dN_{el}/dt|_{t=0}}{N_{el}+N_{inel}} \qquad \qquad \mathcal{L} = \frac{1+\rho^2}{16\pi} \cdot \frac{(N_{el}+N_{inel})^2}{dN_{el}/dt|_{t=0}} \tag{1}$$

where  $N_{el}$  and  $N_{inel}$  are respectively the elastic and inelastic rate and  $\rho$  is the ratio of real to imaginary part of the forward nuclear elastic scattering amplitude (given by theoretical predictions). Therefore, TOTEM will also provide an absolute measurement of  $\mathcal{L}$  to be used for calibration purposes. The uncertainty on the extrapolation of  $dN_{el}/dt$  to t = 0 (optical point) depends on the acceptance for protons scattered at small |t| values, hence at small angles. This requires a small beam angular divergence at the IP, which can be achieved in special runs with high  $\beta^*$  machine optics and typically low  $\mathcal{L}$ . An approved optics with  $\beta^* = 1540$  m (and  $\mathcal{L} \sim$  $10^{28}$  cm<sup>-2</sup>s<sup>-1</sup>) will give a  $\sigma_{tot}$  ( $\mathcal{L}$ ) measurement at the level of  $1 \div 2\%$  (2%). Another approved optics with  $\beta^* = 90$  m (and  $\mathcal{L} \sim 10^{30}$  cm<sup>-2</sup>s<sup>-1</sup>), achievable without modifying the standard LHC injection optics, is expected to allow a preliminary  $\sigma_{tot}$  ( $\mathcal{L}$ ) measurement at the level of  $\sim 5\%$  (7%) in the first period of the LHC running. The experimental systematic error for the measurement with  $\beta^* = 90$  m will be dominated by the evaluation of  $dN_{el}/dt|_{t=0}$ , while with  $\beta^* = 1540$  m it will be dominated by the uncertainty on the corrections to trigger losses in Single Diffraction events for masses below ~  $10 \,\text{GeV/c}^2$ <sup>4</sup>. Given the high rates involved, the statistical error on  $\sigma_{tot}$  will be negligible after few hours of data taking even at low  $\mathcal{L}$ . The theoretical uncertainty related to the estimate of the  $\rho$  parameter is expected to give a contribution on the relative uncertainty of less than 1.2% (considering for instance the full error band on  $\rho$  extrapolation as derived in ref<sup>7</sup>).

Fig. 3 (right) shows the distributions of  $d\sigma_{el}/dt$  at  $\sqrt{s} = 14$  TeV as predicted by different models in the whole |t|-range accessible by TOTEM according to the different LHC optics settings<sup>4</sup>. Several regions are found at increasing |t|: for  $|t| < 6.5 \times 10^{-4}$  GeV<sup>2</sup> (the Coulomb region)  $d\sigma_{el}/dt \sim 1/|t|^2$  is dominated by photon exchange; for |t| up to  $\sim 10^{-3}$  GeV<sup>2</sup>, the hadronic and Coulomb scattering interfere; for  $10^{-3} < |t| < 0.5$  GeV<sup>2</sup> there is the hadronic region, e.g. described by "single-Pomeron exchange", characterized by an approximately exponential fall  $(d\sigma_{el}/dt \sim e^{-B|t|})$ ; the diffractive structure of the proton is then expected for 0.5 < |t| < 1GeV<sup>2</sup>; for |t| > 1 GeV<sup>2</sup> elastic collisions are described by pQCD, e.g. in terms of triple-gluon exchange with  $d\sigma_{el}/dt \sim |t|^{-8}$ . TOTEM will allow to discriminate among different models with a precise measurement of  $d\sigma_{el}/dt$  over all the accessible t-region, where  $d\sigma_{el}/dt$  spans for over 11 orders of magnitude. In the hadronic region, important for the extrapolation of  $d\sigma_{el}/dt$  to t = 0, a fit on B(|t|) is typically performed in the  $|t|_{min} < |t| < 0.25$  GeV<sup>2</sup> range,  $|t|_{min}$  depending on the acceptance for protons scattered at small angles, hence on the beam optics.

Diffractive (due to colour singlet exchange) and non-diffractive (due to colour exchange) inelastic interactions represent a big fraction (around  $60 \div 70 \%$ ) of  $\sigma_{tot}$ . Nevertheless many details of these processes, with close ties to proton structure and low-energy QCD, are still poorly understood. The majority of diffractive events exhibits intact ("leading") protons characterized by their t and fractional momentum loss  $\xi \equiv \Delta p/p$ . TOTEM will be able to measure  $\xi$ -, t- and mass-distributions with acceptances depending on the beam optics. It will also study the charged particle flow in the forward region, providing a significant contribution to the understanding of cosmic ray physics. The existing models give in fact predictions on energy flow, multiplicity and other quantities related to cosmic ray air showers, with significant inconsistencies in the forward region. Furthermore, the integration of TOTEM with the CMS detector will offer the possibility of more detailed studies on inelastic events, including hard diffraction<sup>3</sup>.

# 4 Summary and Conclusions

The TOTEM experiment will be ready for data taking at the LHC start. Running under all beam conditions, it will perform an important and exciting physics programme involving the measurement of  $\sigma_{tot}$  and  $d\sigma_{el}/dt$  in pp interactions as well as studies on diffractive precesses and on forward charged particle production. Special high  $\beta^*$  runs will be required in order to measure  $\sigma_{tot}$  at the level of ~ 5 % (early measurement with  $\beta^* = 90$  m) and ~ 1 ÷ 2 % ( $\beta^* = 1540$  m).  $d\sigma_{el}/dt$  will be studied in the range ~  $10^{-3} < |t| < 10 \text{ GeV}^2$  allowing to distinguish among several theoretical predictions. A common physics programme with CMS on soft and hard diffraction as well as on forward particle production studies is also foreseen in a later stage.

# References

- 1. TOTEM TDR, CERN-LHCC-2004-002; addendum CERN-LHCC-2004-020 (2004).
- 2. CMS Physics TDR, Volume I, CERN-LHCC-2006-001 (2006).
- 3. The CMS and TOTEM Diffr. and Fwd Phys. WG, CERN-LHCC-2006-039/G-124 (2006).
- 4. G. Anelli et al. (TOTEM Collaboration), JINST 3:S08007 (2008).
- 5. E. Noschis et al., Nucl. Instrum. Methods A 563, 41 (2006).
- 6. F. Sauli, Nucl. Instrum. Methods A **386**, 531 (1997).
- 7. J.R. Cudell et al. (COMPETE Collaboration), Phys. Rev. Lett. 89, 201801 (2002).