

Identification of All Dark Matter as Black Holes

Paul H. Frampton*

School of Physics, Korean Institute for Advanced Study
207-43 Cheongryangri-dong, Dongdaemun-gu, Seoul 130-012, Korea
and

Department of Physics and Astronomy, University of North Carolina at
Chapel Hill
CB3255, Chapel Hill, NC 27599-3255, USA[†]

Abstract

For the universe I use dimensionless entropy $S/k = \ln \Omega$ for which the most convenient unit is the googol (10^{100}) and take all dark matter as black holes whereupon the present entropy is about a thousand googols independently of whether dark energy possesses entropy. While the energy of the universe has been established to be about 0.04 baryons, 0.24 dark matter and 0.72 dark energy, the cosmological entropy is almost entirely, about $(1 - 10^{-15})$, from black holes and only 10^{-15} from everything else. This identification of all dark matter as black holes is natural in statistical mechanics.

*frampton@physics.unc.edu

[†]permanent address

1 Introduction

The present note proposes a solution for the cosmological dark matter problem. The problem is over 75 years old [1] and has been the subject of innumerable papers, conferences and books.

There is about six times more dark matter than baryonic matter in the universe. A well-known book [2] on the connection between particle phenomenology and theoretical cosmology discusses possible candidates – axions, WIMPs, MACHOs – for the constituents of the dark matter. In the present article I shall provide a complete solution to the dark matter problem which is in the MACHO category and surprisingly simple: the dark matter is all black holes!

Let me first consider low mass dark matter candidates suggested by particle phenomenology. There are two especially popular candidates.

The lighter is the invisible axion predicted by [3–6]. Its mass [‡] must be in an allowed window $10^{-15}GeV < M_{axion} < 10^{-12}GeV$. Although there are compelling arguments [7] that the invisible axion requires stronger fine tuning than the strong CP problem it purports to solve, it has staying power and remains the subject of experimental searches. The other very popular low-mass dark matter candidate, at a proposed mass typically a trillion times that of the invisible axion, is the WIMP (Weakly-Interacting Massive Particle) exemplified by the neutralino of supersymmetry. The fact that WIMPs can naturally annihilate to an acceptable present abundance of dark matter is sometimes cited as support for WIMPs as dark matter.

But if my proposal for identification of the constituents of dark matter as black holes with masses above $30M_{\odot}$ is correct, particle-phenomenology-inspired dark matter candidates are, unfortunately for their creators, irrelevant.

Higher mass dark matter constituents are generically called MACHOs (Massive Astrophysical Compact Halo Objects). My proposed constituents fall into this class with masses in the range $30M_{\odot} < M_{MACHO} < 500M_{\odot}$ where M_{\odot} is the solar mass; equivalently $10^{26}kg < M_{MACHO} < 10^{27}kg$ or $10^{55}GeV < M_{MACHO} < 10^{56} GeV$. Even the most die-hard opponent to the Large Hadron Collider (LHC) could not take seriously that the LHC could create a black hole with fifty orders of magnitude times the available LHC energy of 10^4GeV so the present article thus might convince that the LHC is safe.

The MACHOs are truly compact, all being smaller in physical size than the Earth. It is the small size and large mass which have enabled these dark matter constituents to escape detection for 75 years. We discuss two methods for their detection, their formation then what, for me, is the best motivation - the entropy of the universe.

[‡]Throughout we adopt an order-of-magnitude notation where 10^x denotes any integer between 10^{x-1} and 10^{x+1} .

2 MACHO Searches by Microlensing

In the late twentieth century heroic searches were made for microlensing events and many spectacular examples [8, 9] of MACHOs were identified with masses up to $30M_{\odot}$. The length of time of a microlensing event depends on the velocity and mass of the lensing MACHO. Since the velocities do not vary widely it is the MACHO mass which is most crucial in determining longevity.

The longevity is directly proportional to the square root of the lens mass. MACHOs were discovered with microlensing longevities up to about 400 days. Insufficient MACHOs to account for more than a few percent of the dark matter were discovered. To detect MACHOs up to mass $500M_{\odot}$, the current upper limit for halo black holes, necessitates study of microlensing events with duration up to 4.5 years.

3 Wide Binaries

Nature has provided delicate astrophysical clocks whose accuracy is sensitive to the possible proximity of MACHOs. These clocks are binaries with separations up to 1 pc and therefore can be bigger than the Solar System in which the distance from the Sun to the outermost planet is only 10^{-4} pc.

Because these wide binaries are so weakly bound they are the most sensitive to disruption by nearby MACHOs. After formation they retain their original orbital parameters as a reliable clock except when affected by gravitational encounters with MACHOs.

The detailed study of wide binaries is a young and promising technique for MACHO detection. In the first pioneering presentation [10] the range of masses allowed for MACHOs which provide all dark matter was $30M_{\odot} < M_{MACHO} < 43M_{\odot}$. It seemed unlikely, though logically possible, that all dark matter MACHOs could have masses within such a narrow range.

A revisit and reanalysis [11] of wide binaries has, however, recently [§] discovered that one of the key examples used in [10] was spurious. This leads to quite different and much more promising restrictions on MACHOs, with the allowed mass range extending up to $500M_{\odot}$. The new constraint is displayed in Fig. 1. and it is now far more likely that all dark matter is in the form of black hole MACHOs.

4 Formation

Because intergalactic dust shows evidence of metallicity which involves nuclei heavier than lithium, the heaviest element formed during big-bang nucleosynthesis, it has been hypothesized [12] that there must have existed Pop III stars in an era before galaxy formation. These subsequently exploded producing the dust nuclei heavier than lithium, and leaving massive black holes which are MACHO candidates.

[§]I am grateful to Julio Chaname for bringing [11] to my attention.




Figure 1: Fig. 1. The vertical axis is the total intermediate-mass black hole (IMBH) mass as the percent of the halo dark matter; the horizontal axis is the individual IMBH mass in terms of the solar mass M_\odot . Note that for the range of masses between $30M_\odot$ and $500M_\odot$ all halo mass can be IMBHs. Taken from [11].

Numerical simulations of dark matter halos [13] have given insight into the likely general features of the halo profile. The spatial resolution of such simulations is, however, far too coarse to study the formation of MACHOs smaller than the Earth whose radius is $10^{-10}pc$. Nevertheless, dark matter is known to clump at large scales and may well do so at such far smaller scales.

The formation of the IBMHs was therefore in the early universe as primordial black holes (PBHs) as suggested in [14–16] and on which limits have been placed in [17].

5 Irrelevance of Dark Energy

I have assumed that dark energy possesses no entropy. Let me argue that this is a very weak assumption.

Recently the Planck satellite was launched and is on its way to Lagrangian point 2 where it will attempt to measure with unprecedented accuracy the dark energy equation of state $w = p/\rho$ where p, ρ are respectively pressure, density.

The key quantity is $\phi = (1 + w)$. If ϕ vanishes identically the assumption of zero entropy for dark energy is justified because it is fully described by one parameter, the cosmological constant. If non-zero ϕ were observationally established, it would imply a dynamical dark energy with concomitant entropy. However, without black holes, the dark energy dimensionless entropy would be much less than one googol and so the identification of the dark matter as black holes and the rest of the present discussion would be unchanged.

6 Discussion

The identification of the dark matter constituent has been an intellectual pursuit for over 75 years. Particle theorists have naturally suggested light mass solutions which include the invisible axion and WIMPs. According to theoretical cosmology, however, the most likely solution is that all dark matter is black holes. According to the recently updated analysis of wide binaries the permitted mass range is between 30 and 500 solar masses. The present known (SMBH) dimensionless entropy of the universe is 1000 googols. This is useful for model building in cyclic cosmology as an alternative to the big bang [18–22].

Taking a universe comprised of 10^{11} halos each of mass $10^{12}M_{\odot}$, and using a high MACHO mass 10^5M_{\odot} , there are 10^7 MACHOs per halo and 10^{18} in the universe. The dimensionless entropy per MACHO, using the PBH formula [23–25] is 10^{88} and the total is a million googols ¶. For galactic-core supermassive black holes (SMBHs), there are 10^{11} of them so that for a central mass 10^7M_{\odot} and individual entropy 10^{92} their total entropy is only 1000 googols.

¶It was suggested at a conference June 2009 in Sweden that such IBMHs be called Framptons.

||Note that the constraints in [26] apply at the recombination era not directly to the present era because of possible mergers.

Because $S \propto M^2$, the total mass of MACHOs is much larger than that of SMBHs. For the central values it is 10^5 times larger. The total MACHO mass is $10^{23} M_{\odot} = 10^{53} kg = 10^{80} GeV$ so I hereby predict several thousand trillion trillion trillion trillion kilograms of black holes.

The existence of such numerous black holes will hopefully be confirmed by observational astronomers. The two most promising known techniques are microlensing and study of wide binaries. It is possible that X-ray emission from accretion disks of the IMBHs can also be detected.

Acknowledgement

I thank Ki-Myeong Lee for hospitality at the Korean Institute for Advanced Study. This work was supported by U.S. Department of Energy grant number DE-FG02-06ER41418.

References

- [1] F. Zwicky, *Helv. Phys. Acta*, **6**, 110 (1933).
- [2] E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley (1990).
- [3] M. Dine, *et al.* *Phys. Lett.* **104B**, 199 (1981).
- [4] J.E. Kim, *Phys. Rev. Lett.* **43**, 103 (1979).
- [5] M.A. Shifman *et al.*, *Nucl. Phys.* **B166**, 493 (1980).
- [6] A.R. Zhitnitsky, *Sov. J. Nucl. Phys.* **31**, 260 (1980).
- [7] R. Holman, *et al.* *Phys. Lett.* **B282**, 132 (1992).
- [8] C. Alcock *et al.* *Astrophys. J.* **486**, 697 (1997).
- [9] C. Alcock, *et al.* *Astrophys. J.* **542**, 281 (200)
- [10] J. Yoo, J. Chaname, and A. Gould, *Astrophys.J.* **601**, 311 (2004).
- [11] D.P. Quinn, *et al.* [arXiv:0903.1664](https://arxiv.org/abs/0903.1664)[astro-ph]
- [12] P. Madau and M.J. Rees, *Astrophys. J.* **551**, L27 (2001).
- [13] J.F. Navarro, C.S. Frenk, and S.D.M. White, *Astrophys. J.* **462**, 563 (1996).
- [14] J. Yokoyama, *Astron. Astrophys.* **318**, 673 (1997).
- [15] J. Yokoyama, *Phys. Rev. bf D58*, 083510 (1998).

- [16] T. Kawaguchi *et al*, Mon. Not. R. Astron. Soc. **388**, 1426 (2008).
- [17] R. Saito and J. Yokoyama, Phys. Rev.Lett. **102**, 161101 (2009).
- [18] L. Baum and P.H. Frampton, Phys. Rev. Lett. **98**, 071301 (2007).
- [19] L. Baum and P.H. Frampton, Mod. Phys. Lett. **A23**, 33 (2008).
- [20] L. Baum, P.H. Frampton and S. Matsuzaki, JCAP 0804:032 (2008).
- [21] P.H. Frampton, Mod. Phys. Lett. A22, 2587 (2007)
- [22] P.H. Frampton, [arXiv:0903.4309](https://arxiv.org/abs/0903.4309) [hep-ph].
- [23] L. Parker, Phys. Rev. **183**, 1057 (1969).
- [24] J.D. Bekenstein, Phys. Rev. **D7**, 2333 (1973).
- [25] S.W. Hawking, Commun. Math. Phys. **43**, 1974 (1975).
- [26] M. Ricotti, J.P. Ostriker and K.J. Mack. [arXiv:0709.0524](https://arxiv.org/abs/0709.0524) [astro-ph].