

ENTIRE SELF-SIMILAR SOLUTIONS TO LAGRANGIAN MEAN CURVATURE FLOW

ALBERT CHAU, JINGYI CHEN, AND WEIYONG HE

ABSTRACT. We consider self-similar solutions to mean curvature evolution of entire Lagrangian graphs. When the Hessian of the potential function u has eigenvalues strictly uniformly between -1 and 1 , we show that on the potential level all the shrinking solitons are quadratic polynomials while the expanding solitons are in one-to-one correspondence to functions of homogenous of degree 2 with the Hessian bound. We also show that if the initial potential function is cone-like at infinity then the scaled flow converges to an expanding soliton as time goes to infinity.

1. INTRODUCTION

Let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function. Then $L_u = \{(x, Du(x)) : x \in \mathbb{R}^n\}$ defines a graphical Lagrangian submanifold in $\mathbb{R}^{2n} = \mathbb{R}^n \oplus \sqrt{-1}\mathbb{R}^n = \mathbb{C}^n$. On the other hand, any entire graphical Lagrangian submanifold in \mathbb{R}^{2n} can be obtained by such a potential u , up to addition of constants. Let

$$G(D^2u) = \frac{1}{\sqrt{-1}} \log \frac{\det(I_n + \sqrt{-1}D^2u)}{\sqrt{\det(I_n + (D^2u)^2)}},$$

where I_n is the n dimensional identity matrix. The operator $G(D^2u)$ is strictly elliptic and arises in the equation for minimal Lagrangian graphs. Namely, L_u is a minimal submanifold in \mathbb{R}^{2n} exactly when $G(D^2u)$ is equal to some constant Θ . In this note, we will consider the following two equations for $u(x)$:

$$(1) \quad G(D^2u) - u + \frac{1}{2}\nabla u \cdot x = 0,$$

$$(2) \quad G(D^2u) + u - \frac{1}{2}\nabla u \cdot x = 0.$$

Date: May 19, 2009.

2000 Mathematics Subject Classification. Primary 53C44, 53A10.

The first two authors are partially supported by NSERC, and the third author is partially supported by a PIMS postdoctoral fellowship.

Our motivation is that (1) and (2) are the defining equations for self-expanding and self-shrinking solutions to the Lagrangian mean curvature flow of entire graphs in \mathbb{R}^{2n} . We describe this in more detail in the next section. We will refer to an entire solution to (1) as a *self-expanding soliton* and an entire solution to (2) as a *self-shrinking soliton*.

Definition 1.1. We say that u_0 satisfies *Condition A* if $D^2u_0 \in L^\infty(\mathbb{R}^n)$ and

$$-(1 - \delta)I_n \leq D^2u_0 \leq (1 - \delta)I_n$$

for some $\delta \in (0, 1)$ where I_n is the n dimensional identity matrix. We say u_0 satisfies *Condition B* if $D^2u_0 \in L^\infty(\mathbb{R}^n)$ and

$$u_0(x) = \frac{1}{\lambda^2}u_0(\lambda x)$$

for any $\lambda > 0$.

We now state our main result

Theorem 1.1. *If u is a smooth self-shrinking soliton satisfying Condition A, then u is a quadratic. There exists a one-to-one correspondence between smooth self-expanding solitons satisfying Condition A and functions which satisfy Condition A and Condition B. In particular, there are infinitely many nontrivial smooth self-expanding solitons satisfying Condition A.*

We also obtain a convergence result for the following fully nonlinear parabolic equation which arises from the Lagrangian mean curvature flow (see §2):

$$(3) \quad \begin{cases} \frac{du}{dt} = G(D^2u) \\ u(x, 0) = u_0(x). \end{cases}$$

Theorem 1.2. *Let u_0 satisfy Condition A and suppose that*

$$(4) \quad \lim_{\lambda \rightarrow \infty} \lambda^{-2}u_0(\lambda x) \rightarrow U_0(x)$$

for some $U_0(x)$ as $\lambda \rightarrow \infty$. Let $u(x, t)$ and $U(x, t)$ be solutions to (3) with initial data $u_0(x)$ and $U_0(x)$ respectively. Then $t^{-1}u(\sqrt{t}x, t)$ converges to $U(x, 1)$ uniformly and smoothly in compact subsets of \mathbb{R}^n when $t \rightarrow \infty$, where $U(x, 1)$ is a self-expanding soliton.

If we define $v(x, s) = t^{-1}u(\sqrt{t}x, t)$ where $s = \log t$ and $t \geq 1$. Then $v(x, s)$ satisfies the following equation

$$\frac{\partial v}{\partial s} = G(D^2v) - v + \frac{1}{2}\nabla v \cdot x.$$

Theorem 1.2 implies that $v(x, s)$ converges to $U(x, 1)$ when $s \rightarrow \infty$. For mean curvature flow of entire graphic hypersurface, if the initial surface is cone-like at infinity, in the sense that the normal component of the graph decays to zero at infinity at a certain rate, then the scaled mean curvature flow converges to a self-expanding soliton [4], where the monotonicity formula plays an important role. Our method is different from [4] and is based on the existence result and the estimates in [2] and the uniqueness result in [3].

Some interesting examples of non-graphical self-similar solutions to Lagrangian mean curvature flow in \mathbb{C}^n have recently been constructed [5, 6]; also see references therein.

2. PRELIMINARIES

We now describe the connections among the elliptic equations (1) and (2), the parabolic equation (3) and the Lagrangian mean curvature flow.

Let $F_0 : M^n \rightarrow \mathbb{R}^{n+m}$ be an embedding of a manifold M^n into \mathbb{R}^{n+m} . Then the mean curvature flow with initial condition F_0 is the equation

$$(5) \quad \begin{cases} \frac{dF}{dt} = H \\ F(x, 0) = F_0(x) \end{cases}$$

where $H(x, t)$ is the mean curvature vector of the submanifold $F(\cdot, t)(M)$ of \mathbb{R}^{2n} at $F(x, t)$. The solution $F(\cdot, t) : M^n \rightarrow \mathbb{R}^{n+m}$ is called *self-expanding* if it is defined for all $t > 0$ and $F(\cdot, t)$ has the form

$$(6) \quad M_t = \sqrt{t} M_1, \quad \text{for all } t > 0$$

where $M_t = F(\cdot, t)(M)$ and \sqrt{t} represents the homothety $p \mapsto \sqrt{t}p$, $p \in \mathbb{R}^{n+m}$. Similarly, $F(\cdot, t) : M^n \rightarrow \mathbb{R}^{n+m}$ is called *self-shrinking* if it is defined for all $t < 0$ and $F(x, t)$ has the form

$$(7) \quad M_t = \sqrt{-t} M_{-1}, \quad \text{for all } t < 0.$$

Now when $F_0(x) = (x, Du_0(x))$ for $x \in \mathbb{R}^n$ defines an entire Lagrangian graph for some potential $u_0 : \mathbb{R}^n \rightarrow \mathbb{R}$, then (5) is equivalent to the following fully nonlinear parabolic equation (3):

$$\begin{cases} \frac{du}{dt} = G(D^2u) \\ u(x, 0) = u_0(x). \end{cases}$$

In particular, if u is a solution to (3) then there exists a family of diffeomorphisms [7] $r_t : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $F(x, t) = (r_t(x), Du(r_t(x), t))$ solves (5). Now suppose the family $F(x, t)$ satisfies (6). Then we have

$$(8) \quad D \left(u(x, t) - tu \left(\frac{x}{\sqrt{t}}, 1 \right) \right) = 0, \quad \forall t > 0.$$

Thus by letting $t = 1$, we have

$$(9) \quad u(x, t) = tu \left(\frac{x}{\sqrt{t}}, 1 \right), \quad \forall t > 0.$$

Using (3) and (9) we verify directly that $u(x, 1)$ satisfies (1), in other words, $u(x, 1)$ is a self-expanding soliton. Similarly, we can derive that if $F(x, t)$ is a self-shrinking solution to (5) with corresponding solution $u(x, t)$ to (3), then $u(x, t)$ satisfies

$$(10) \quad u(x, t) = -tu \left(\frac{x}{\sqrt{-t}}, -1 \right), \quad \forall t > 0,$$

and $u(x, -1)$ satisfies (2). So $u(x, 1)$ is a self-shrinking soliton.

Conversely, it is not hard to show that if $u(x)$ solves either (1) or (2), then using (9) or (10) respectively we can generate a solution $F(x, t)$ to (5) which is either shrinking or expanding. We illustrate this in detail in the expanding case. The case for shrinking solutions is similar. Suppose $u(x)$ solves (1) and define $u(x, t) := tu(x/\sqrt{t})$. Then as above, the family $M_t := \{(x, Du(x, t)) : x \in \mathbb{R}^n\}$ is easily checked to satisfy (6). On the other hand, we also have

$$(11) \quad \begin{aligned} \frac{du}{dt}(x, t) &= u \left(\frac{x}{\sqrt{t}} \right) - \frac{1}{2} \nabla u(x\sqrt{t}) \cdot \frac{x}{\sqrt{t}} \\ &= G \left(D_y^2 u \left(\frac{x}{\sqrt{t}} \right) \right) \\ &= G(D^2 u(x, t)) \end{aligned}$$

where $y = x/\sqrt{t}$. In other words, $u(x, t)$ solves (3) and by the discussion above there exists a family r_t such that $F(x, t) = (r_t(x), Du(r_t(x), t))$ is solution to (5) which is self-expanding.

3. PROOFS OF THEOREMS

The proof of Theorem 1.1 is to consider the parabolic equation (3) with Lipschitz initial data. Let us recall the main theorem in [2]:

Theorem 3.1. *Let $u_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfy Condition A. Then (3) has a longtime smooth solution $u(x, t)$ for all $t > 0$ with initial condition u_0 such that the following estimates hold:*

- (i) $-(1 - \delta)I_n \leq D^2u \leq (1 - \delta)I_n$ for all $t > 0$,
- (ii) $\sup_{x \in \mathbb{R}^n} |D^l u(x, t)|^2 \leq C(l, \delta)/t^{l-2}$, for $l \geq 3$ and some $C(l, \delta)$.
- (iii) $u(x, t) \in C^\infty(\mathbb{R}^n \times (0, \infty))$ and $u(x, t)$ converges to $u_0(x)$ in the Lipschitz norm when $t \rightarrow 0$.

We will apply Theorem 3.1 and the uniqueness result in [3] for the parabolic equation (3) to prove Theorem 1.1. The proof consists of the following two lemmas.

Lemma 3.1. *Let $u_0 : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfy Condition A and Condition B. Then (3) has a unique longtime smooth solution $u(x, t)$ for all $t > 0$ with initial condition u_0 such that $u(x, t) = tu(x/\sqrt{t}, 1)$. In particular $u(x, 1)$ is a smooth self-expanding soliton satisfying (1). Conversely, if v is smooth a self-expanding soliton satisfying Condition A, then there exists u_0 satisfying Condition A and Condition B such that $u(x, t)$ is the unique solution of (3) with initial condition u_0 and $v = u(x, 1)$.*

Proof. If u_0 satisfies Condition A, then by Theorem 3.1, there exists a smooth solution $u(x, t)$ to (3) for all $t > 0$ with initial data u_0 . It is clear that

$$u_\lambda(x, t) := \lambda^{-2}u(\lambda x, \lambda^2 t)$$

is also a solution to (3) with initial data

$$u_\lambda(x, 0) = \lambda^{-2}u_0(\lambda x) = u_0(x)$$

where we have used that u_0 satisfies Condition B. Since $u_\lambda(x, 0) = u_0$, the uniqueness result in [3] implies

$$u(x, t) = u_\lambda(x, t).$$

for any $\lambda > 0$. Therefore $u(x, t)$ satisfies (9), and hence $u(x, 1)$ solves (1). In other words, $u(x, 1)$ is a smooth self-expanding soliton.

Now suppose that v is a smooth solution to (1) satisfying Condition A (in fact, it suffices to assume D^2v is bounded). Define $u(x, t)$ for $t > 0$ by

$$u(x, t) = tv \left(\frac{x}{\sqrt{t}} \right).$$

It is clear that $u(x, t)$ satisfies the evolution equation in (3) since v satisfies (1). Now we claim that the limit of $u(x, t)$ exists when t goes to zero. To see this, begin by noting $u(0, t) = tv(0)$ for $t > 0$ and so

$$\lim_{t \rightarrow 0} u(0, t) = 0.$$

Moreover, it is clear that

$$Du(0, t) = \sqrt{t}Dv(0)$$

and that for any $t > 0$

$$-(1 - \delta)I_n \leq D^2u(x, t) \leq (1 - \delta)I_n$$

since

$$D^2u(x, t) = D^2 \left(tv \left(\frac{x}{\sqrt{t}} \right) \right) = D^2v \left(\frac{x}{\sqrt{t}} \right).$$

We may then conclude that for any sequence $t_i \rightarrow 0$ there is a subsequence t_{k_i} such that $u(x, t_{k_i})$ converges in $C^{1,\alpha}$ uniformly in compact subsets of \mathbb{R}^n for any $0 < \alpha < 1$. This limit is in fact independent of the sequence t_i . Indeed, let u_1 and u_2 be two such limits along sequences t_i and t'_i respectively. Since $u(x, t)$ is a solution to (3), $\partial u / \partial t$ is also uniformly bounded for any $t > 0$, $x \in \mathbb{R}^n$. Thus for any $x \in \mathbb{R}^n$ we may have

$$|u(x, t_i) - u(x, t'_i)| \leq C|t_i - t'_i|$$

for some C independent of i . Letting $i \rightarrow \infty$, we conclude that $u_1(x) = u_2(x)$. So for different sequences $t_i \rightarrow 0$, the limit is unique. Let

$$u_0(x) = \lim_{t \rightarrow 0} u(x, t).$$

Then $D^2u_0 \in L^\infty$ and we have

$$-(1 - \delta)I_n \leq D^2u_0 \leq (1 - \delta)I_n.$$

Further,

$$\begin{aligned} \frac{1}{\lambda^2}u_0(\lambda x) &= \frac{1}{\lambda^2} \lim_{t \rightarrow 0} tv \left(\frac{\lambda x}{\sqrt{t}} \right) \\ &= \lim_{t \rightarrow 0} \left(\frac{\sqrt{t}}{\lambda} \right)^2 v \left(\frac{\lambda x}{\sqrt{t}} \right) \\ &= u_0(x). \end{aligned}$$

Therefore u_0 satisfies Condition B. □

Remark 3.1. Lemma 3.1 provides many examples of entire graphical Lagrangian self-expanding solitons. For instance, we can choose u_0 such that

$$u_0(x) = \begin{cases} a_1x_1^2 + \sum_{i=2}^n a_ix_i^2, & \text{if } x_1 \geq 0 \\ -a_1x_1^2 + \sum_{i=2}^n a_ix_i^2, & \text{if } x_1 < 0. \end{cases}$$

If $a_i \in (-1 + \delta, 1 - \delta)$ for any fixed $\delta \in (0, 1)$, then u_0 satisfies Condition A and Condition B and $u(x, 1)$ is a self-expanding soliton which satisfies (1). As u_0 is not smooth, $u(x, 1)$ cannot be a quadratic polynomial.

Next, we show that smooth self-shrinking solitons are trivial if Condition A holds.

Lemma 3.2. *If v is a smooth solution of (2) such that $-(1 - \delta)I_n \leq D^2v \leq (1 - \delta)I_n$, then v is a quadratic polynomial.*

Proof. If v is a smooth solution to (2), then

$$u(x, t) = (1 - t)v\left(\frac{x}{\sqrt{1 - t}}\right)$$

is a solution to (3) for $t \in (0, 1)$ and $u(x, 0) = v(x)$, hence Theorem 3.1 applies to $u(x, t)$ since this solution is unique, and in particular $|D^3u(x, t)| \leq C$ for some constant C when $t \geq 1/2$ for any x . But one checks directly that

$$D^3u(x, t) = \frac{1}{\sqrt{1 - t}}D^3v\left(\frac{x}{\sqrt{1 - t}}\right).$$

This implies

$$|D^3v(x)| = \left|D^3v\left(\frac{x\sqrt{1 - t}}{\sqrt{1 - t}}\right)\right| \leq C\sqrt{1 - t}$$

for any x . It follows that $D^3v(x) = 0$, thus v is quadratic. \square

Now we prove Theorem 1.2.

Proof. Let $u(x, t)$ be the solution to (3) with initial data u_0 satisfying Condition A. It is clear that for any λ ,

$$u_\lambda(x, t) = \lambda^{-2}u(\lambda x, \lambda^2 t)$$

is a solution of (3) with initial data $u_\lambda(x, 0) = \lambda^{-2}u_0(\lambda x)$ satisfying Condition A. By the uniqueness result in [3], we may then apply the estimates in Theorem 3.1 to $u_\lambda(x, t)$. For any sequence $\lambda_i \rightarrow \infty$, consider the solutions $u_{\lambda_i}(x, t)$. For $t > 0$, it is clear that

$$D^2u_{\lambda_i}(x, t) = D^2u(\lambda_i x, \lambda_i^2 t)$$

and so by Theorem 3.1,

$$-(1 - \delta)I_n \leq D^2u_{\lambda_i}(x, t) \leq (1 - \delta)I_n$$

for all x and $t \geq 0$, and for $t > 0$ and $l \geq 3$

$$|D^l u_{\lambda_i}(x, t)| \leq C(l, \delta)\sqrt{t}^{2-l}.$$

Thus by (3) we can also get the following estimates that, for any $m \geq 1, l \geq 0$, there is a constant $C(m, l, \delta)$ such that

$$\left| \frac{\partial^m}{\partial t^m} D^l u_{\lambda_i} \right| \leq C(m, l, \delta) \sqrt{t}^{2-l-2m}.$$

In particular, there are constants C and $C(\delta)$ such that

$$\left| \frac{\partial u_{\lambda_i}}{\partial t} \right| \leq C, \quad \left| \frac{\partial D u_{\lambda_i}}{\partial t} \right| \leq \frac{C(\delta)}{\sqrt{t}}$$

for all $t > 0$. We observe that

$$u_{\lambda_i}(0, 0) = \lambda_i^{-2} u_0(0)$$

and

$$D u_{\lambda_i}(0, 0) = \lambda_i^{-1} D u_0(0)$$

are both bounded, thus $u_{\lambda_i}(0, t)$ and $D u_{\lambda_i}(0, t)$ are uniformly bounded in i for any fixed t . By Arzelà-Ascoli theorem, there exists a subsequence λ_{k_i} such that $u_{\lambda_{k_i}}(x, t)$ converges smoothly and uniformly in compact subsets of $\mathbb{R}^n \times (0, \infty)$ to a solution $U(x, t)$ of (3). Moreover, $U(x, t)$ satisfies the estimates in Theorem 3.1. Since $\partial U / \partial t$ is uniformly bounded for any $t > 0$, $U(x, t)$ converges to some function $U(x, 0)$ when $t \rightarrow 0$. In particular we have

$$U(x, 0) = \lim_{\lambda_{k_i} \rightarrow \infty} \lambda_{k_i}^{-2} u_0(\lambda_{k_i} x).$$

It is clear that $U(x, 0)$ satisfies Condition A. Similarly as in the proof of Lemma 2.1, the hypothesis (4) implies that $U(x, 0)$ satisfies Condition B. Thus by Theorem 1.1 we know that $U(x, 1)$ is a self-expanding soliton, and $U(x, t)$ satisfies (9).

Moreover, notice that $U(x, 0) = U_0(x)$ by (4) and $U_0(x)$ does not depend on the sequence λ_i , it then follows from the uniqueness result in [3] that the solution $U(x, t)$ is also independent of the sequence λ_i and we may write

$$u_{\lambda}(x, t) \rightarrow U(x, t).$$

as $\lambda \rightarrow \infty$. In particular, letting $\lambda = \sqrt{t}$ we have $t^{-1} u(\sqrt{t}x, t)$ converges to $U(x, 1)$ smoothly and uniformly in compact subsets of \mathbb{R}^n when $t \rightarrow \infty$. \square

Remark 3.2. The family of Lagrangian graphs $(x + at, D u_0(x) + bt)$ is a translating solution to (5) precisely when u satisfies

$$(12) \quad \sum_i \arctan \lambda_i + \sum_i a_i \frac{\partial u_0}{\partial x_i} - \sum_i b_i x_i = c,$$

for some constant c . In [2] the authors proved that if u is a smooth solution to (12) which satisfies condition A, then u is a quadratic function. We observe here that this result can also be obtained from the uniqueness result in [3] together with Theorem 3.1. Namely, if $u_0(x)$ solves (12) then $v(x, t) = u_0(x - at) + b \cdot x t + ct$ solves (3) with initial condition $v(x, 0) = u_0(x)$. On the other hand, Theorem 3.1 guarantees a longtime solution $u(x, t)$ to (3) with initial condition u_0 for which $\sup_{x \in \mathbb{R}^n} |D^3 u(x, t)| \rightarrow 0$ as $t \rightarrow \infty$. Then as $u(x, t) = v(x, t)$ by the uniqueness result in [3], $\sup_{x \in \mathbb{R}^n} |D^3 u_0(x - at)| = \sup_{x \in \mathbb{R}^n} |D^3 v(x, t)| \rightarrow 0$ as $t \rightarrow \infty$ and we conclude that $\sup_{x \in \mathbb{R}^n} |D^3 u_0(x)| = 0$, in other words u_0 must be quadratic.

REFERENCES

1. G. Barles, S. Biton, O. Ley, *Uniqueness for parabolic equations without growth condition and applications to the mean curvature flow in \mathbf{R}^2* , J. Differential Equations **187** (1987), 456-472.
2. A. Chau, J. Chen, W. He, *Lagrangian Mean Curvature flow for entire Lipschitz graphs*, arXiv:0902.3300.
3. J. Chen, C. Pang, *Uniqueness of viscosity solutions of a geometric fully nonlinear parabolic equation*, arXiv:0905.3868.
4. K. Ecker, G. Huisken, *Mean curvature evolution of entire graphs*, Ann. of Math. (2) **130** (1989), no.3, 453-471.
5. H. Anciaux, *Construction of Lagrangian self-similar solutions to the mean curvature flow in \mathbb{C}^n* , Geom. Dedicata 120 (2006), 37-48.
6. D. Joyce, Y.I. Lee, M.P. Tsui, *Self-similar solutions and translating solitons for Lagrangian mean curvature flow*. arXiv:0801.372.
7. K. Smoczyk, *Longtime existence for the Lagrangian mean curvature flow*, Calc. Var. (2004), **20**, 25-46.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, B.C., V6T 1Z2, CANADA

E-mail address: `chau@math.ubc.ca`

E-mail address: `jychen@math.ubc.ca`

E-mail address: `whe@math.ubc.ca`