

On MHD stability of gravitating plasmas with field aligned flows

H. Tasso* G. N. Throumoulopoulos[†]

* Max-Planck-Institut für Plasmaphysik
Euratom Association
85748 Garching bei München, Germany

[†] University of Ioannina,
Association Euratom-Hellenic Republic,
Department of Physics, GR 451 10 Ioannina, Greece

October 29, 2018

Abstract

A previous stability condition (see Throumoulopoulos and Tasso, Physics of Plasmas 14, 122104 (2007)) for incompressible plasmas with field aligned flows is extended to gravitating plasmas including self-gravitation. It turns out that the stability condition is affected by gravitation through the equilibrium values only.

Key words: Magnetohydrodynamics, stability

*het@ipp.mpg.de

[†]gthroum@cc.uoi.gr

The system under consideration is a constant density plasma with an incompressible flow aligned with the magnetic field, and maintained in a magnetohydrodynamic equilibrium by the Lorenz force and external and self-gravitation. It is governed by the following equations

$$\mathbf{v} = \lambda \mathbf{B} \quad (1)$$

with λ an arbitrary function,

$$\nabla \cdot \mathbf{v} = 0, \quad (2)$$

$$\mathbf{v} \cdot \nabla \mathbf{v} = \mathbf{J} \times \mathbf{B} - \nabla P + \nabla \phi, \quad (3)$$

where \mathbf{v} is the velocity, \mathbf{B} is the magnetic field and $\mathbf{J} = \nabla \times \mathbf{B}$ is the current density. ϕ is the gravitational potential. The equations are written in convenient units and the constant density is set to 1. (In fact, as will be proved below it suffices to assume constant density only at equilibrium.) In addition we assume the existence of well defined equilibrium magnetic surfaces labelled by a smooth function ψ . These equations differ from the corresponding equations of Ref.[1] by the inclusion of the gravitational potential.

Equations (1) and (2) lead to $\lambda = \lambda(\psi)$, and using the identity $\mathbf{v} \cdot \nabla \mathbf{v} = \nabla v^2/2 - \mathbf{v} \times \nabla \times \mathbf{v}$, (1)-(3) lead to

$$(1 - \lambda^2) \mathbf{J} \times \mathbf{B} = \nabla \left(P + \frac{\lambda^2 B^2}{2} - \phi \right) - B^2 \nabla \frac{\lambda^2}{2}, \quad (4)$$

where B and v are the magnetic field and velocity field moduli respectively. Taking the scalar product of Eq.(4) with \mathbf{B} gives

$$P + \frac{\lambda^2 B^2}{2} - \phi = P_{s\phi}(\psi), \quad (5)$$

where $P_{s\phi}(\psi)$ is the arbitrary function in the presence of gravity.

Consequently, Eq.(4) can be put in the form

$$(1 - \lambda^2) \mathbf{J} \times \mathbf{B} = P'_{s\phi} \nabla \psi - (\lambda^2)' \frac{B^2}{2} \nabla \psi, \quad (6)$$

with

$$\mathbf{J} \times \mathbf{B} = g_\phi(\psi, B^2) \nabla \psi, \quad (7)$$

where

$$g_\phi(\psi, B^2) = \frac{P'_{s\phi}}{1 - \lambda^2} - \frac{(\lambda^2)' B^2}{1 - \lambda^2} \frac{1}{2}. \quad (8)$$

The only difference with Ref.[1] is the presence of the gravitational potential ϕ in (4)-(8). This potential obeys Poisson's equation

$$\Delta\phi = -1, \quad (9)$$

whose solution is the superposition of the internal potential ϕ_i given by $\phi_i = \int_V \frac{1}{|\mathbf{x}-\mathbf{x}'|} d^3\mathbf{x}'$, where V is the volume of the plasma considered, and an external potential ϕ_e due to fixed masses outside the plasma.

Let us now come to the stability of our flowing and gravitating equilibria. The general problem without gravity has been already formulated in Ref.[2] in terms of the Lagrangian displacement $\boldsymbol{\xi}$. Its application to equilibria with field aligned incompressible flows and homogeneous density was investigated in Ref.[1] and Ref.[3]. The correct derivation of the final condition was done in Ref.[1]. This condition states that a general steady state of a plasma of uniform density and incompressible flow parallel to \mathbf{B} is stable to small three-dimensional perturbations if the flow is sub-Alfvénic ($\lambda^2 < 1$) and $A \geq 0$, where

$$A = -g^2[(1-\lambda^2)(\mathbf{J} \times \nabla\psi) \cdot (\mathbf{B} \cdot \nabla)\nabla\psi + \frac{(\lambda^2)'}{2} |\nabla\psi|^2 (\nabla\psi \cdot \nabla \frac{B^2}{2} + g|\nabla\psi|^2)]. \quad (10)$$

The quantity g is like the g_ϕ of Eq.(8) but the P_s in it does not contain the gravitational potential ϕ . Now, $A \geq 0$ remains valid if g in (10) is replaced by g_ϕ . The reason is that the perturbed gravity in the linearized dynamical equations vanishes for the $\nabla \cdot \boldsymbol{\xi} = 0$ and homogeneous density at equilibrium. To prove this statement we employ the relations[2]

$$\rho_1 = -\nabla \cdot (\rho\boldsymbol{\xi}), \quad (11)$$

$$\mathbf{v}_1 = \frac{\partial\boldsymbol{\xi}}{\partial t} + (\mathbf{v} \cdot \nabla\boldsymbol{\xi} - \boldsymbol{\xi} \cdot \nabla\mathbf{v}), \quad (12)$$

where the subscript 1 denotes first order quantities. Acting the divergence operator on (12) and using the identity $\nabla \times (\boldsymbol{\xi} \times \mathbf{v}) = \mathbf{v} \cdot \nabla\boldsymbol{\xi} - \boldsymbol{\xi} \cdot \nabla\mathbf{v} + \boldsymbol{\xi}(\nabla \cdot \mathbf{v}) - \mathbf{v}(\nabla \cdot \boldsymbol{\xi})$ with $\nabla \cdot \mathbf{v} = 0$ yields

$$\nabla \cdot \mathbf{v}_1 = \nabla \cdot \frac{\partial\boldsymbol{\xi}}{\partial t} + \mathbf{v} \cdot \nabla(\nabla \cdot \boldsymbol{\xi}) \equiv \frac{D}{Dt}(\nabla \cdot \boldsymbol{\xi}). \quad (13)$$

Therefore, if $\nabla \cdot \boldsymbol{\xi} = 0$ it follows $\nabla \cdot \mathbf{v}_1 = 0$ and, inversely, $\nabla \cdot \mathbf{v}_1 = 0$ implies $\nabla \cdot \boldsymbol{\xi} = 0$ for $t > t_0$ if this relation is satisfied initially at $t = t_0$. Then, (11)

in conjunction with the linearized continuity equation, $\partial\rho_1/\partial t + \mathbf{v} \cdot \nabla\rho_1 = 0$, lead to $\rho_1 = 0$. The fact that for uniform equilibrium density, the density remains uniform in the perturbed state because of incompressibility was not noticed in Refs.[1] and [3].

We conclude by stating that the above mentioned condition is sufficient for stability if we replace g in (10) by g_ϕ [Eq. (8)], which means that this stability condition is affected by gravity through the equilibrium values only.

Acknowledgements

Part of this work was conducted during a visit of the author G.N.T. to the Max-Planck-Institut für Plasmaphysik, Garching. The hospitality of that Institute is greatly appreciated.

This work was performed within the participation of the University of Ioannina in the Association Euratom-Hellenic Republic, which is supported in part by the European Union and by the general Secretariat of Research and Technology of Greece. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

References

- [1] G.N. Throumoulopoulos and H. Tasso, *Phys. Plasmas* 14, 122104 (2007).
- [2] E. Frieman and Manuel Rotenberg, *Rev. Mod. Phys.* 32, 898 (1960).
- [3] V.A. Vladimirov and K.I. Ilin, *Phys. Plasmas* 5, 4199 (1998).