

A CATEGORICAL QUOTIENT IN THE CATEGORY OF DENSE CONSTRUCTIBLE SUBSETS

DEVIRIM CELIK (TÜBINGEN)

ABSTRACT. A. A'Campo-Neuen and J. Hausen gave an example of an algebraic torus action on an open subset of the affine four space that admits no quotient in the category of algebraic varieties. We show that this example admits a quotient in the category of dense constructible subsets and thereby answer a question of A. Białynicki-Birula.

The purpose of this note is to answer a question by A. Białynicki-Birula on categorical quotients in the category of dense constructible subsets (dc-subsets) introduced in [4]. The objects of this category are pairs $X \subseteq X'$, where X' is a complex algebraic variety and X is a dense constructible subset of X' . A morphism from $X \subseteq X'$ to $Y \subseteq Y'$ is a map $X \rightarrow Y$ that extends to a morphism $X'_1 \rightarrow Y'$ for some open neighbourhood $X'_1 \subseteq X'$ of X . Consider the \mathbb{C}^* -variety

$$X := \mathbb{C}^2 \times (\mathbb{C}^*)^2 \cup (\mathbb{C}^*)^2 \times \mathbb{C}^2, \quad tx := (tx_1, tx_2, x_3, t^{-1}x_3).$$

A categorical quotient for this \mathbb{C}^* -variety is a \mathbb{C}^* -invariant morphism $\pi: X \rightarrow Y$, i.e. a morphism being constant along the \mathbb{C}^* -orbits, such that any other \mathbb{C}^* -invariant morphism $\varphi: X \rightarrow Z$ has a unique factorization $\varphi = \psi \circ \pi$ with a morphism $\psi: Y \rightarrow Z$. This concept depends strongly on the category one works with. For example, in the category of complex algebraic varieties, there is no such quotient for X , see [2, Prop. 5.1 (ii)], whereas in the category of toric varieties, X admits a quotient, namely the toric morphism

$$\pi: X \rightarrow \mathbb{C}^3, \quad (x_1, x_2, x_3, x_4) \mapsto (x_1x_4, x_2x_4, x_3),$$

use [1, Thm. 2.3]. Now, set $Y := \pi(X)$. Then $Y \subseteq \mathbb{C}^3$ is a proper dc-subset, and π defines a morphism of dc-subsets from $X \subseteq X$ to $Y \subseteq \mathbb{C}^3$. A. Białynicki-Birula asks [4, p. 53], whether or not this is a categorical quotient in the category of dc-subsets.

Theorem. *The map $\pi: X \rightarrow Y$ is a categorical quotient for the \mathbb{C}^* -action on X in the category of dc-subsets.*

2000 *Mathematics Subject Classification.* 14L30, 14L24.

Key words and phrases. Algebraic torus action, categorical quotient, dense constructible subsets.

$$P(e_1) := e_1, \quad P(e_2) := e_2, \quad P(e_3) := e_3, \quad P(e_4) := e_1 + e_2.$$
$$\begin{aligned} Y &= \mathbb{C}^3 \setminus (\{0\} \times \mathbb{C}^* \times \{0\} \cup \mathbb{C}^* \times \{0\} \times \{0\}) \\ &= \mathbb{T}y_0 \cup \bigcup_{i=1}^3 \mathbb{T}y_{\rho_i} \cup \mathbb{T}y_{\tau_1} \cup \mathbb{T}y_{\delta} \end{aligned}$$

Moreover, let \tilde{Y} denote the toric prevariety obtained by gluing the affine toric varieties $\tilde{Y}_{\tau_i} := \text{Spec}(\mathbb{C}[\tau_i^\vee \cap \mathbb{Z}^3])$, where $i = 1, 2$, along the big torus $\mathbb{T} := (\mathbb{C}^*)^3 \subseteq \tilde{Y}_{\tau_i}$. Then $P: \mathbb{Z}^4 \rightarrow \mathbb{Z}^3$ defines a \mathbb{C}^* -invariant toric morphism $\tilde{\pi}: X \rightarrow \tilde{Y}$. Finally, the lattice homomorphism $\text{id}: \mathbb{Z}^3 \rightarrow \mathbb{Z}^3$ gives us a toric morphism $\kappa: \tilde{Y} \rightarrow \mathbb{C}^3$ with $\kappa(\tilde{Y}) = Y$ and $\pi = \kappa \circ \tilde{\pi}$.

In order to verify the universal property of the morphism $\pi: X \rightarrow Y$, let $Z \subseteq Z'$ be a dc-subset and $\varphi: X \rightarrow Z$ a \mathbb{C}^* -invariant morphism. According to [2, Prop. 5.1 (i)], the toric morphism $\tilde{\pi}: X \rightarrow \tilde{Y}$ is a categorical quotient for the \mathbb{C}^* -action on X in the category of complex prevarieties. Consequently, we have $\varphi = \tilde{\psi} \circ \tilde{\pi}$ with a morphism $\tilde{\psi}: \tilde{Y} \rightarrow Z'$. Our task is to show that there is a morphism $\psi: Y \rightarrow Z$ of dc-subsets making the

following diagram commutative

$$\begin{array}{ccccc}
 X & \xrightarrow{\varphi} & Z & \subseteq & Z' \\
 \tilde{\pi} \downarrow & \searrow \pi & \nearrow \psi & & \nearrow \\
 \tilde{Y} & \xrightarrow{\kappa} & Y & &
 \end{array}$$

$\tilde{\psi}$ (curved arrow from \tilde{Y} to Z')

In a first step, we show that such a ψ exists as a set theoretical map. Since we have $\tilde{\psi}(\tilde{Y}) = \varphi(X) \subseteq Z$, this amounts to verifying that $\tilde{\psi}$ is constant along the fibers of κ . Let $\tilde{y}_0, \tilde{y}_{\rho_1}, \tilde{y}_{\rho_2}, \tilde{y}_{\tau_1} \in \tilde{Y}_{\tau_1}$ and $\tilde{y}_0, \tilde{y}_{\rho_3}, \tilde{y}_{\rho_4}, \tilde{y}_{\tau_2} \in \tilde{Y}_{\tau_2}$ denote the distinguished points corresponding to the cones $0, \rho_i$ etc.. Further denote by $\mathbb{T}_{y_{\tau_i}}$ the isotropy groups of y_{τ_i} . Then, using e.g. [3, Prop. 3.5], we obtain that the κ -fibres are

$$\kappa^{-1}(ty_0) = t\tilde{y}_0, \quad \kappa^{-1}(ty_{\rho_i}) = t\tilde{y}_{\rho_i}, \quad i = 1, 2, 3,$$

$$\kappa^{-1}(ty_{\tau_1}) = t\mathbb{T}_{y_{\tau_1}}\tilde{y}_{\tau_1} \cup t\mathbb{T}_{y_{\tau_1}}\tilde{y}_{\rho_4}, \quad \kappa^{-1}(y_{\delta}) = \mathbb{T}_{\tilde{y}_{\tau_2}},$$

where $t \in \mathbb{T}$. Thus, only for $\kappa^{-1}(ty_{\tau_1})$ and $\kappa^{-1}(y_{\delta})$, there is something to show. Take $v \in \mathbb{Z}^3$ from the relative interior ρ_4 and let $\lambda_v: \mathbb{C}^* \rightarrow \mathbb{T}$ denote the corresponding one-parameter-subgroup. Then, in \tilde{Y} , we have

$$\overline{\lambda_v(\mathbb{C}^*)t\tilde{y}_0} \setminus \lambda_v(\mathbb{C}^*)t\tilde{y}_0 = \{t\tilde{y}_{\tau_1}, t\tilde{y}_{\rho_4}\}.$$

Note that we have $t\tilde{y}_{\tau_1} \in \tilde{Y}_{\tau_1}$ and $t\tilde{y}_{\rho_4} \in \tilde{Y}_{\tau_2}$. Since Z' separated, we obtain $\tilde{\psi}(t\tilde{y}_{\tau_1}) = \tilde{\psi}(t\tilde{y}_{\rho_4})$. In particular, we obtain for $t \in \mathbb{T}_{y_{\tau_1}} = \mathbb{T}_{\tilde{y}_{\tau_1}}$ the equation

$$\tilde{\psi}(t\tilde{y}_{\rho_4}) = \tilde{\psi}(t\tilde{y}_{\tau_1}) = \tilde{\psi}(\tilde{y}_{\tau_1}).$$

Consequently, for every $t \in \mathbb{T}$, the set $\tilde{\psi}(\kappa^{-1}(ty_{\tau_1}))$ consists of a single point. Now we treat $\kappa^{-1}(ty_{\tau_1}) = \mathbb{T}_{\tilde{y}_{\tau_2}}$. First note that $\mathbb{T} = \mathbb{T}_{\tilde{y}_{\tau_1}}\mathbb{T}_{\tilde{y}_{\tau_2}}$ and hence any $t \in \mathbb{T}$ can be written as $t = t_{\tau_1}t_{\tau_2}$ with $t_{\tau_i} \in \mathbb{T}_{\tilde{y}_{\tau_i}}$. Consider again the one parameter subgroup $\lambda_v: \mathbb{C}^* \rightarrow \mathbb{T}$ corresponding to a lattice vector $v \in \mathbb{Z}^3$ in the relative interior of ρ_4 . Then we have:

$$\begin{aligned}
 \tilde{\psi}(t\tilde{y}_{\tau_2}) &= \lim_{s \rightarrow 0} \tilde{\psi}(t_{\tau_1}t_{\tau_2}\lambda_v(s)\tilde{y}_{\rho_4}) \\
 &= \lim_{s \rightarrow 0} \tilde{\psi}(t_{\tau_1}t_{\tau_2}\lambda_v(s)\tilde{y}_{\tau_1}) \\
 &= \lim_{s \rightarrow 0} \tilde{\psi}(t_{\tau_2}\lambda_v(s)\tilde{y}_{\tau_1}) \\
 &= \tilde{\psi}(t_{\tau_2}\tilde{y}_{\tau_2}) \\
 &= \tilde{\psi}(\tilde{y}_{\tau_2}).
 \end{aligned}$$

We show now that the map $\psi: Y \rightarrow Z$ is a morphism of dc-subsets. Consider the open affine toric subvarieties $Y_{\rho_3}, Y_{\tau_1} \subseteq \mathbb{C}^3$ and $\tilde{Y}_{\rho_3} \subseteq \tilde{Y}$ defined by the cones ρ_3 and τ_1 . Then we have $Y_{\tau_1}, Y_{\rho_3} \subseteq Y$ and, for $U := \tilde{Y}_{\tau_1} \cup \tilde{Y}_{\rho_3}$ and $V := Y_{\tau_1} \cup Y_{\rho_3}$, the restriction $\kappa|_U: U \rightarrow V$ is an isomorphism of varieties. Thus, we have $\psi = \tilde{\psi} \circ \kappa^{-1}$ on V , and hence ψ is a rational map.

We have to show that ψ defined near the point y_δ . For this we choose an affine open neighbourhood $W_0 \subseteq Z'$ of $\psi(y_\delta)$ in Z' and set $U_0 := \tilde{\psi}^{-1}(W_0)$ and $V_0 := \psi^{-1}(W_0)$. Then we have an open subset

$$V_0 \cap V = \kappa(U_0 \cap U) \subseteq \mathbb{C}^3,$$

on which ψ is a morphism. After realizing W_0 as a closed subset of some \mathbb{K}^r , we obtain functions $g_1, \dots, g_r \in \mathcal{O}(V_0 \cap V)$, such that the morphism ψ is given on $V_0 \cap V$ as

$$\psi|_{V_0 \cap V}: V_0 \cap V \rightarrow W_0, \quad y \mapsto (g_1(y), \dots, g_r(y)).$$

Suppose for the moment that each g_i is defined at y_δ . Then there are an open set $V' \subseteq \mathbb{C}^3$ containing $V_0 \cap V$ and the point y_δ such that g_1, \dots, g_r define a morphism $\psi': V' \rightarrow W_0$ extending $\psi|_{V_0 \cap V}$. This morphism fits into the diagram

$$\begin{array}{ccc} U_0 \cap U & \subseteq & \kappa^{-1}(V') \xrightarrow{\tilde{\psi}} W_0 \\ \kappa \downarrow & & \downarrow \kappa \nearrow \psi' \\ V_0 \cap V & \subseteq & V' \end{array}$$

Since $\tilde{\psi}$ and $\psi' \circ \kappa$ coincide on the dense open subset $U_0 \cap U$, they coincide on $\kappa^{-1}(V')$. Thus, by surjectivity of κ , we obtain $\psi' = \psi|_{V'}$ and see that ψ is a morphism on the neighbourhood $V' \subseteq \mathbb{C}^3$ of $y_\delta \in Y$. Hence, ψ is a morphism on the open set $V \cup V' \subseteq \mathbb{C}^3$. This shows that $\psi: Y \rightarrow Z$ is a morphism in the category of dc-subsets.

Thus, our task is to show that every g_i is defined in y_δ . For this, we regard g_i as a rational function on the normal ambient variety \mathbb{C}^3 of Y and show that $\text{div}(g_i)$ is nonnegative at y_δ . For this in turn, it suffices to show that any prime divisor on \mathbb{C}^3 containing y_δ meets the open set $V \cap V_0$.

Let D be a prime divisor on \mathbb{C}^3 passing through y_δ . We show that D meets the open set $V \cap V_0$. Since $\mathbb{C}^3 \setminus V$ is of codimension 2 in \mathbb{C}^3 , we obtain $D \cap V \neq \emptyset$. First consider the case that $\kappa^{-1}(D)$ contains the prime divisor $\overline{\mathbb{T}\tilde{y}_{\rho_4}} = \mathbb{T}\tilde{y}_{\rho_4} \cup \mathbb{T}\tilde{y}_{\tau_2}$. Then $\kappa^{-1}(D) \cap U_0 \neq \emptyset$, hence

$$D \cap V_0 \neq \emptyset \implies D \cap V \cap V_0 \neq \emptyset.$$

Next suppose that $\kappa^{-1}(D)$ doesn't contain $\overline{\mathbb{T}\tilde{y}_{\rho_4}} = \tilde{Y} \setminus U$. Then any component of $\kappa^{-1}(D)$ meets U . Since \mathbb{C}^3 is factorial, D is principal, and thus $\kappa^{-1}(D)$ is of pure codimension one in \tilde{Y} . Consequently, $\kappa^{-1}(D)$ equals the closure of $\kappa^{-1}(D) \cap U$ in \tilde{Y} . Since $\kappa^{-1}(D)$ intersects $\kappa^{-1}(y_\delta)$, we can conclude

$$\kappa^{-1}(D) \cap U \cap U_0 \neq \emptyset \implies D \cap V \cap V_0 \neq \emptyset.$$

□

REFERENCES

- [1] A'Campo-Neuen, Annette; Hausen, Jürgen: Quotients of toric varieties by the action of a subtorus. *Tohoku Math. J. (2)* 51 (1999), no. 1, 1–12.
- [2] A'Campo-Neuen, Annette; Hausen, Jürgen: Examples and counterexamples for existence of categorical quotients. *J. Algebra* 231 (2000), no. 1, 67–85.
- [3] A'Campo-Neuen, Annette; Hausen, Jürgen: Toric prevarieties and subtorus actions. *Geom. Dedicata* 87 (2001), no. 1-3, 35–64.
- [4] Białynicki-Birula, Andrzej: Categorical quotients. *J. Algebra* 239 (2001), no. 1, 35–55.
- [5] Fulton, William: Introduction to toric varieties. Princeton University Press, Princeton, NJ (1993).

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE
10, 72076 TÜBINGEN, GERMANY

E-mail address: celik@mail.mathematik.uni-tuebingen.de