

BESSEL TYPE INEQUALITIES IN HILBERT C^* -MODULES

S. S. DRAGOMIR¹, M. KHOSRAVI² AND M. S. MOSLEHIAN³

ABSTRACT. Regarding the generalizations of the Bessel inequality in Hilbert spaces which are due to Bombiari and Boas–Bellman, we obtain a version of the Bessel inequality and some generalizations of this inequality in the framework of Hilbert C^* -modules.

1. INTRODUCTION

The Bessel inequality states that if $(e_i)_{1 \leq i \leq n}$ is an orthonormal sequence of vectors in a Hilbert space $(\mathcal{H}; \langle \cdot, \cdot \rangle)$, then

$$\sum_{i=1}^n |\langle x, e_i \rangle|^2 \leq \|x\|^2 \quad (x \in \mathcal{H}).$$

A number of mathematicians have investigated the above inequality in various settings. One of the generalizations of the Bessel inequality was given by Bombieri [3] as follows.

Theorem 1.1. *If x, y_1, \dots, y_n are elements of a complex unitary space, then*

$$\sum_{i=1}^n |\langle x, y_i \rangle|^2 \leq \|x\|^2 \max_{1 \leq i \leq n} \sum_{j=1}^n |\langle y_i, y_j \rangle|.$$

In 1941, Boas [2] and in 1944, independently, Bellman [1] proved a result, which can be stated as follows.

Theorem 1.2. *If x, y_1, \dots, y_n are elements of a Hilbert space, then*

$$\sum_{i=1}^n |\langle x, y_i \rangle|^2 \leq \|x\|^2 \left[\max_{1 \leq i \leq n} \|y_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} |\langle y_i, y_j \rangle|^2 \right)^{1/2} \right].$$

Recently, Mitrinović, Pečarić and Fink [9] proved the following inequality and showed that it is equivalent with the Boas–Bellman theorem.

2000 *Mathematics Subject Classification.* Primary 46L08; Secondary 46L05, 47A63, 47B10, 47A30.

Key words and phrases. Bessel inequality; Boas–Bellman inequality; Hilbert C^* -module, C^* -algebra.

Theorem 1.3. *If x, y_1, \dots, y_n are elements of a Hilbert space and c_1, \dots, c_n are arbitrary complex numbers, then*

$$\left| \sum_{i=1}^n c_i \langle x, y_i \rangle \right|^2 \leq \|x\|^2 \sum_{i=1}^n |c_i|^2 \left[\max_{1 \leq i \leq n} \|y_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} |\langle y_i, y_j \rangle|^2 \right)^{1/2} \right].$$

In addition, Dragomir [4] obtained some other generalizations of the Bessel inequality, which are similar to Boas–Bellman inequality and Mitrinović–Pečarić–Fink inequality.

Our aim is to extend some of these generalizations of the Bessel inequality in the framework of Hilbert C^* -modules. A related concept to our work is the notion of frame. We would like to refer the interested reader to [6] for an extensive account on frames in Hilbert C^* -modules.

2. PRELIMINARIES

In this section we recall some fundamental definitions in the theory of Hilbert modules that will be used in the sequel.

Suppose that \mathcal{A} is a C^* -algebra and \mathcal{X} is a linear space which is an algebraic right \mathcal{A} -module satisfying $\lambda(xa) = x(\lambda a) = (\lambda x)a$ for all $x \in \mathcal{X}, a \in \mathcal{A}, \lambda \in \mathbb{C}$. The space \mathcal{X} is called a *pre-Hilbert \mathcal{A} -module* (or an *inner product \mathcal{A} -module*) if there exists an \mathcal{A} -valued inner product $\langle \cdot, \cdot \rangle : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{A}$ satisfying the following properties:

- (i) $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ if and only if $x = 0$;
- (ii) $\langle x, \lambda y + z \rangle = \lambda \langle x, y \rangle + \langle x, z \rangle$;
- (iii) $\langle x, ya \rangle = \langle x, y \rangle a$;
- (iv) $\langle x, y \rangle^* = \langle y, x \rangle$;

for all $x, y, z \in \mathcal{X}, a \in \mathcal{A}$ and $\lambda \in \mathbb{C}$.

By (ii) and (iv), $\langle \lambda x + ya, z \rangle = \bar{\lambda} \langle x, z \rangle + a^* \langle y, z \rangle$. It follows from the *Cauchy–Schwarz inequality* $\langle y, x \rangle \langle x, y \rangle \leq \|\langle x, x \rangle\| \|\langle y, y \rangle\|$ that $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$ is a norm on \mathcal{X} , where the latter norm denotes that in the C^* -algebra \mathcal{A} ; see [7, p. 5]. This norm makes \mathcal{X} into a right normed module over \mathcal{A} . A pre-Hilbert module X is called a *Hilbert \mathcal{A} -module* if it is complete with respect to its norm.

Two typical examples of Hilbert C^* -modules are as follows:

- (I) Every Hilbert space is a Hilbert \mathbb{C} -module.
- (II) Every C^* -algebra \mathcal{A} is a Hilbert \mathcal{A} -module via $\langle a, b \rangle = a^* b$ ($a, b \in \mathcal{A}$).

Notice that the inner product structure of a C^* -algebra is essentially more complicated than complex numbers. For instance, the concepts such as adjoint, orthogonality and theorems

such as Riesz' representation in the theory of complex Hilbert spaces cannot simply be generalized or transferred to that of Hilbert C^* -modules.

One may define an " \mathcal{A} -valued norm" $|\cdot|$ by $|x| = \langle x, x \rangle^{1/2}$. Clearly, $\| |x| \| = \|x\|$ for each $x \in \mathcal{X}$. It is known that $|\cdot|$ does not satisfy the triangle inequality in general. See [7, 8] for more detailed information on Hilbert C^* -modules.

3. MAIN RESULTS

We start our work by presenting a version of the Bessel inequality for Hilbert C^* -modules.

Theorem 3.1. *Let \mathcal{X} be a Hilbert \mathcal{A} -module and e_1, e_2, \dots, e_n be a sequence of unit vectors in X such that $\langle e_i, e_j \rangle = 0$ for $1 \leq i \neq j \leq n$. If $x \in X$, then*

$$\sum_{i=1}^n |\langle e_i, x \rangle|^2 \leq |x|^2. \quad (3.1)$$

Proof. The result follows from the following inequalities.

$$\begin{aligned} 0 \leq \left| x - \sum_{i=1}^n e_i \langle e_i, x \rangle \right|^2 &= \left\langle x - \sum_{i=1}^n e_i \langle e_i, x \rangle, x - \sum_{i=1}^n e_i \langle e_i, x \rangle \right\rangle \\ &= \langle x, x \rangle + \sum_{i=1}^n \sum_{j=1}^n \langle e_i, x \rangle^* \langle e_i, e_j \rangle \langle e_j, x \rangle - 2 \sum_{i=1}^n |\langle e_i, x \rangle|^2 \\ &= \langle x, x \rangle + \sum_{i=1}^n \langle e_i, x \rangle^* \langle e_i, e_i \rangle \langle e_i, x \rangle - 2 \sum_{i=1}^n |\langle e_i, x \rangle|^2 \\ &\leq |x|^2 + \sum_{i=1}^n \langle e_i, x \rangle^* \langle e_i, x \rangle - 2 \sum_{i=1}^n |\langle e_i, x \rangle|^2 \\ &= |x|^2 - \sum_{i=1}^n |\langle e_i, x \rangle|^2. \end{aligned}$$

□

The following lemma is useful to prove a Bombieri type inequality.

Lemma 3.2. *Let \mathcal{A} be a C^* -algebra and $a, b, c \in \mathcal{A}$. Then*

$$a^*cb + b^*c^*a \leq \|c\|(|a|^2 + |b|^2).$$

Proof. Using the universal representation of \mathcal{A} it is sufficient to prove that

$$T^*RS + S^*R^*T \leq \|R\|(|T|^2 + |S|^2)$$

for bounded linear operators T, R, S acting on a Hilbert space \mathcal{H} .

By the polar decomposition, there exists a partial isometry $U \in \mathbb{B}(\mathcal{H})$ such that $R = U|R|$. It follows from

$$\begin{aligned} 0 &\leq (S - U^*T)^*|R|(S - U^*T) \\ &= S^*|R|S - S^*|R|U^*T - T^*U|R|S + T^*U|R|U^*T, \end{aligned}$$

that

$$T^*RS + S^*R^*T \leq ||R|^{1/2}U^*T|^2 + ||R|^{1/2}S|^2.$$

Hence

$$\begin{aligned} T^*RS + S^*R^*T &\leq ||R|^{1/2}U^*T|^2 + ||R|^{1/2}S|^2 \\ &\leq ||R|^{1/2}U^*|^2 |T|^2 + ||R|^{1/2}|^2 |S|^2 \\ &\leq \|R\|(|T|^2 + |S|^2). \end{aligned}$$

□

Theorem 3.3. *If y_1, \dots, y_n are elements of a Hilbert C^* -module \mathcal{X} , then*

$$\left| \sum_{i=1}^n y_i a_i \right|^2 \leq \sum_{i=1}^n |a_i|^2 \max_{1 \leq i \leq n} \sum_{j=1}^n \|\langle y_i, y_j \rangle\|, \quad (3.2)$$

for all $a_1, \dots, a_n \in \mathcal{A}$.

Proof. By using Lemma 3.2, we have

$$\begin{aligned} \left| \sum_{i=1}^n y_i a_i \right|^2 &= \left\langle \sum_{i=1}^n y_i a_i, \sum_{i=1}^n y_i a_i \right\rangle = \sum_{1 \leq i, j \leq n} a_i^* \langle y_i, y_j \rangle a_j \\ &= \sum_{i=1}^n a_i^* \langle y_i, y_i \rangle a_i + \sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j \\ &= \sum_{i=1}^n a_i^* \langle y_i, y_i \rangle a_i + \sum_{1 \leq i < j \leq n} (a_i^* \langle y_i, y_j \rangle a_j + a_j^* \langle y_j, y_i \rangle a_i) \\ &\leq \sum_{i=1}^n \|\langle y_i, y_i \rangle\| |a_i|^2 + \sum_{1 \leq i < j \leq n} (\|\langle y_i, y_j \rangle\| |a_i|^2 + \|\langle y_i, y_j \rangle\| |a_j|^2) \\ &= \sum_{1 \leq i, j \leq n} \|\langle y_i, y_j \rangle\| |a_i|^2 \\ &\leq \sum_{i=1}^n |a_i|^2 \max_{1 \leq i \leq n} \sum_{j=1}^n \|\langle y_i, y_j \rangle\|. \end{aligned}$$

□

From Theorem 3.3 the following result of Bombieri type can be obtained.

Corollary 3.4. *Let $x, y_1, \dots, y_n \in \mathcal{X}$ and $a_1, \dots, a_n \in \mathcal{A}$. Then*

$$\left(\sum_{i=1}^n |\langle y_i, x \rangle|^2 \right)^2 \leq |x|^2 \left\| \sum_{i=1}^n |\langle y_i, x \rangle|^2 \right\| \max_{1 \leq i \leq n} \sum_{j=1}^n \|\langle y_i, y_j \rangle\|.$$

Proof. By using inequality (3.2), we have

$$\begin{aligned} \left(\sum_{i=1}^n |\langle y_i, x \rangle|^2 \right)^2 &= \left| \sum_{i=1}^n \langle y_i, x \rangle^* \langle y_i, x \rangle \right|^2 \\ &= \left| \left\langle \sum_{i=1}^n y_i \langle y_i, x \rangle, x \right\rangle \right|^2 \\ &\leq \left\| \sum_{i=1}^n y_i \langle y_i, x \rangle \right\|^2 |x|^2 \\ &\leq |x|^2 \left\| \sum_{i=1}^n |\langle y_i, x \rangle|^2 \right\| \max_{1 \leq i \leq n} \sum_{j=1}^n \|\langle y_i, y_j \rangle\| \\ &\leq |x|^2 \left\| \sum_{i=1}^n |\langle y_i, x \rangle|^2 \right\| \max_{1 \leq i \leq n} \sum_{j=1}^n \|\langle y_i, y_j \rangle\|. \end{aligned}$$

□

Now we state some other applications of inequality (3.2) for Hilbert space operators, although some of them can be deduced directly. Recall that the space $\mathbb{B}(\mathcal{H}, \mathcal{K})$ of all bounded linear operators between Hilbert spaces \mathcal{H} and \mathcal{K} can be regarded as a Hilbert C^* -module over the C^* -algebra $\mathbb{B}(\mathcal{H})$ via $\langle T, S \rangle = T^*S$.

Corollary 3.5. *Let \mathcal{H} and \mathcal{K} be Hilbert spaces and let $T_1, \dots, T_n \in \mathbb{B}(\mathcal{H}, \mathcal{K})$ be operators having orthogonal ranges. Then*

$$\left| \sum_{i=1}^n T_i S_i \right|^2 \leq \sum_{i=1}^n |S_i|^2 \max_{1 \leq i \leq n} \|T_i\|^2,$$

for any $S_1, \dots, S_n \in \mathbb{B}(\mathcal{H})$.

Proof. Clearly the operators T_i have orthogonal ranges if and only if $T_i^* T_j = 0$. Thus the result follows immediately from inequality (3.2). □

Corollary 3.6. *Let S_1, S_2 be operators on a Hilbert space \mathcal{H} and let T be an invertible operator on \mathcal{H} . Then*

$$|TS_1 + (T^*)^{-1}S_2|^2 \leq (|S_1|^2 + |S_2|^2) [1 + \max(\|T\|^2, \|T^{-1}\|^2)].$$

The next result is a refinement of one of the inequalities presented in Theorem 2.1 of [5].

Corollary 3.7. *If $\lambda_1, \dots, \lambda_n$ are complex numbers and T_1, \dots, T_n are operators on a Hilbert space \mathcal{H} , then*

$$\left| \sum_{i=1}^n \lambda_i T_i \right| \leq \max_{1 \leq i \leq n} |\lambda_i| \sum_{i=1}^n |\lambda_i| \sum_{i=1}^n |T_i|^2.$$

Proof. If we consider $\lambda_i T_i = (\lambda_i I) T_i$, we get from (3.2) that

$$\begin{aligned} \left| \sum_{i=1}^n \lambda_i T_i \right| &\leq \sum_{i=1}^n |T_i|^2 \max_{1 \leq i \leq n} \sum_{j=1}^n |\bar{\lambda}_i \lambda_j| \\ &= \sum_{i=1}^n |T_i|^2 \max_{1 \leq i \leq n} |\lambda_i| \sum_{j=1}^n |\lambda_j| \\ &= \max_{1 \leq i \leq n} |\lambda_i| \sum_{i=1}^n |\lambda_i| \sum_{i=1}^n |T_i|^2. \end{aligned}$$

□

The following theorem is similar to the Mitrinović–Pečarić–Fink theorem in the Hilbert space theory, with \mathcal{A} -valued norm instead of the usual norm.

Theorem 3.8. *If x, y_1, \dots, y_n are elements of a Hilbert \mathcal{A} -module \mathcal{X} and a_1, \dots, a_n are elements of \mathcal{A} , then*

$$\left| \sum_{i=1}^n a_i \langle y_i, x \rangle \right|^2 \leq |x|^2 \sum_{i=1}^n \|a_i\|^2 \left[\max_{1 \leq i \leq n} \|y_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{1/2} \right]. \quad (3.3)$$

Proof. By the Cauchy–Schwarz inequality,

$$\left| \sum_{i=1}^n a_i \langle y_i, x \rangle \right|^2 = \left| \left\langle \sum_{i=1}^n y_i a_i^*, x \right\rangle \right|^2 \leq \left\| \sum_{i=1}^n y_i a_i^* \right\|^2 |x|^2. \quad (3.4)$$

We also have

$$\begin{aligned}
 \left\| \sum_{i=1}^n y_i a_i^* \right\|^2 &= \left\| \left\langle \sum_{i=1}^n y_i a_i^*, \sum_{j=1}^n y_j a_j^* \right\rangle \right\| \\
 &= \left\| \sum_{1 \leq i, j \leq n} a_i \langle y_i, y_j \rangle a_j^* \right\| \\
 &\leq \sum_{1 \leq i, j \leq n} \|a_i\| \|\langle y_i, y_j \rangle\| \|a_j\| \\
 &= \sum_{i=1}^n \|a_i\|^2 \|y_i\|^2 + \sum_{1 \leq i \neq j \leq n} \|a_i\| \|a_j\| \|\langle y_i, y_j \rangle\| \\
 &\leq \max_{1 \leq i \leq n} \|y_i\| \sum_{i=1}^n \|a_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} \|a_i\|^2 \|a_j\|^2 \right)^{\frac{1}{2}} \left(\sum_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{\frac{1}{2}} \\
 &\leq \sum_{i=1}^n \|a_i\|^2 \left(\max_{1 \leq i \leq n} \|y_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{\frac{1}{2}} \right). \tag{3.5}
 \end{aligned}$$

Combining (3.4) and (3.5) we can get the desired result. \square

Corollary 3.9. *For x, y_1, \dots, y_n in a Hilbert \mathcal{A} -module \mathcal{X} ,*

$$\left(\sum_{i=1}^n |\langle y_i, x \rangle|^2 \right)^2 \leq |x|^2 \sum_{i=1}^n \|\langle y_i, x \rangle\|^2 \left[\max_{1 \leq i \leq n} \|y_i\|^2 + \left(\sum_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{\frac{1}{2}} \right]. \tag{3.6}$$

Proof. Set $a_i = \langle x, y_i \rangle$ in inequality (3.3). \square

Note that the inequality (3.6), can be considered as a generalization of Boas–Bellman inequality (1.1). However, for the case where $(y_i)_{1 \leq i \leq n}$ is an orthonormal sequence of vectors, the inequality (3.6) is a weaker result than (3.1).

Furthermore, all of the inequalities which are obtained by Dragomir in [4, Chapter 4] can be extended to Hilbert C^* -modules in a similar way. The details are left to the interested readers.

We can also prove some other Boas–Bellman type inequalities in Hilbert C^* -modules as follows.

Lemma 3.10. *Let \mathcal{A} be a C^* -algebra and \mathcal{X} be a Hilbert \mathcal{A} -module. Then*

$$\left| \sum_{i=1}^n y_i a_i \right|^2 \leq \max_{1 \leq i \leq n} \|y_i\|^2 \sum_{i=1}^n |a_i|^2 + B_n,$$

where

$$B_n = \begin{cases} (n-1)\sqrt{n} \max_{1 \leq i \leq n} \|a_i\| \max_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\| \left(\sum_{i=1}^n |a_i|^2 \right)^{\frac{1}{2}}, \\ \sqrt{n-1} \left(\max_{1 \leq i \leq n} \sum_{1 \leq j \neq i \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^n \|a_i\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^n |a_i|^2 \right)^{\frac{1}{2}}, \end{cases} \quad (3.7)$$

for any $y_i \in \mathcal{X}$ and $a_i \in \mathcal{A}$.

Proof. We observe that

$$\begin{aligned} \left| \sum_{i=1}^n y_i a_i \right|^2 &= \left\langle \sum_{i=1}^n y_i a_i, \sum_{i=1}^n y_i a_i \right\rangle \\ &= \sum_{1 \leq i, j \leq n} a_i^* \langle y_i, y_j \rangle a_j \\ &= \sum_{i=1}^n a_i^* \langle y_i, y_i \rangle a_i + \sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j. \end{aligned}$$

We also have

$$\sum_{i=1}^n a_i^* \langle y_i, y_i \rangle a_i = \sum_{i=1}^n a_i^* |y_i|^2 a_i \leq \sum_{i=1}^n \|y_i\|^2 |a_i|^2 \leq \max_{1 \leq i \leq n} \|y_i\|^2 \sum_{i=1}^n |a_i|^2.$$

To get the first inequality of (3.7) note that

$$\begin{aligned} \sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j &= 2 \sum_{1 \leq i < j \leq n} \operatorname{Re}(a_i^* \langle y_i, y_j \rangle a_j) \\ &\leq 2 \sum_{1 \leq i < j \leq n} \left(\frac{|a_i^* \langle y_i, y_j \rangle a_j|^2 + |a_j^* \langle y_j, y_i \rangle a_i|^2}{2} \right)^{1/2} \\ &\quad \left(\text{by } |\operatorname{Re}(c)| \leq \left(\frac{c^* c + c c^*}{2} \right)^{1/2}, \text{ where } c \in \mathcal{A} \right) \\ &= \sqrt{2} \sum_{1 \leq i < j \leq n} \left(|a_i^* \langle y_i, y_j \rangle a_j|^2 + |a_j^* \langle y_j, y_i \rangle a_i|^2 \right)^{1/2}. \end{aligned}$$

By operator convexity of the function $f(t) = t^2$, we conclude that

$$\begin{aligned}
 \sum_{i \neq j} a_i^* \langle y_i, y_j \rangle a_j &\leq \sqrt{2} \cdot \sqrt{\frac{n^2 - n}{2}} \left(\sum_{1 \leq i < j \leq n} |a_i^* \langle y_i, y_j \rangle a_j|^2 + |a_j^* \langle y_j, y_i \rangle a_i|^2 \right)^{1/2} \\
 &= \sqrt{n^2 - n} \left(\sum_{1 \leq i \neq j \leq n} |a_i^* \langle y_i, y_j \rangle a_j|^2 \right)^{1/2} \\
 &\leq \sqrt{n^2 - n} \max_{1 \leq i \leq n} \|a_i\| \max_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\| \left((n-1) \sum_{i=1}^n |a_i|^2 \right)^{1/2} \\
 &= (n-1) \sqrt{n} \max_{1 \leq i \leq n} \|a_i\| \max_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\| \left(\sum_{i=1}^n |a_i|^2 \right)^{1/2}.
 \end{aligned}$$

To obtain the second inequality of (3.7) notice that $\sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j$ is self-adjoint, hence

$$\begin{aligned}
 \left(\sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j \right)^2 &\leq \left| \sum_{1 \leq i \neq j \leq n} a_i^* \langle y_i, y_j \rangle a_j \right|^2 \\
 &= \left| \sum_{i=1}^n a_i^* \left(\sum_{1 \leq j \neq i \leq n} \langle y_i, y_j \rangle a_j \right) \right|^2 \\
 &\leq \sum_{i=1}^n \|a_i\|^2 \sum_{i=1}^n \left| \sum_{1 \leq j \neq i \leq n} \langle y_i, y_j \rangle a_j \right|^2 \\
 &\leq \sum_{i=1}^n \|a_i\|^2 \sum_{i=1}^n \left(\sum_{1 \leq j \neq i \leq n} \|\langle y_i, y_j \rangle\|^2 \sum_{1 \leq j \neq i \leq n} |a_j|^2 \right) \\
 &\leq \sum_{i=1}^n \|a_i\|^2 \left(\max_{1 \leq i \leq n} \sum_{1 \leq j \neq i \leq n} \|\langle y_i, y_j \rangle\|^2 \cdot \sum_{i=1}^n \sum_{1 \leq j \neq i \leq n} |a_j|^2 \right) \\
 &\leq (n-1) \sum_{i=1}^n |a_i|^2 \left(\sum_{i=1}^n \|a_i\|^2 \cdot \max_{1 \leq i \leq n} \sum_{1 \leq j \neq i \leq n} \|\langle y_i, y_j \rangle\|^2 \right).
 \end{aligned}$$

□

The following result can be stated as well.

Theorem 3.11. *Let $x, y_1, \dots, y_n \in \mathcal{X}$ and $a_1, \dots, a_n \in \mathcal{A}$. Then*

$$\left| \sum_{i=1}^n a_i \langle y_i, x \rangle \right|^2 \leq |x|^2 \left\| \sum_{i=1}^n |a_i^*|^2 \right\|^{1/2} \left[\max_{1 \leq i \leq n} \|y_i\|^2 \left\| \sum_{i=1}^n |a_i^*|^2 \right\|^{1/2} + B_n \right],$$

where

$$B_n := \begin{cases} (n-1)\sqrt{n} \max_{1 \leq i \leq n} \|a_i\| \max_{1 \leq i \neq j \leq n} \|\langle y_i, y_j \rangle\|, \\ \sqrt{n-1} \left(\max_{1 \leq i \leq n} \sum_{1 \leq j \neq i \leq n} \|\langle y_i, y_j \rangle\|^2 \right)^{1/2} \left(\sum_{i=1}^n \|a_i\|^2 \right)^{1/2}. \end{cases}$$

Remark 3.12. In the case where (y_i) is an orthogonal sequence, it follows from the theorem above that

$$\left(\sum_{i=1}^n |\langle y_i, x \rangle|^2 \right)^2 \leq |x|^2 \max_{1 \leq i \leq n} \|y_i\|^2 \left\| \sum_{i=1}^n |\langle y_i, x \rangle|^2 \right\|,$$

which is a stronger result than the inequality (3.6).

Acknowledgement.

This work was done when the second author was at the Research Group in Mathematical Inequalities and Applications (RGMIA) in Victoria University on her sabbatical leave from Tehran Teacher Training University. She thanks both universities for their support. The third author was supported by a grant from Ferdowsi University of Mashhad; (NO. MP88060MOS)

REFERENCES

- [1] R. Bellman, *Almost orthogonal series*, Bull. Amer. Math. Soc. **50** (1944), 517–519.
- [2] R.P. Boas, *A general moment problem*, Amer. J. Math. **63** (1941), 361–370.
- [3] E. Bombieri, *A note on the large sieve*, Acta Arith. **18** (1971), 401–404.
- [4] S.S. Dragomir, *Advances in Inequalities of the Schwarz, Grüss and Bessel Type in Inner Product Spaces*, Nova Science Publishers, Inc., Hauppauge, NY, 2005.
- [5] S.S. Dragomir, *Norm inequalities for sequences of operators related to the Schwarz inequality*, J. Inequal. Pure Appl. Math. **7** (2006), no. 3, Article 97.
- [6] M. Frank and D.R. Larson, *Frames in Hilbert C^* -modules and C^* -algebras*, J. Operator Theory **48** (2002), no. 2, 273–314.
- [7] E.C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, 210, Cambridge University Press, Cambridge, 1995.
- [8] V.M. Manuilov and E.V. Troitsky, *Hilbert C^* -modules*, Translations of Mathematical Monographs, 226. American Mathematical Society, Providence, RI, 2005.
- [9] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic, Dordrecht, 1993.

¹ RESEARCH GROUP IN MATHEMATICAL INEQUALITIES AND APPLICATIONS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, P. O. BOX 14428, MELBOURNE CITY, VIC, 8001, AUSTRALIA.

E-mail address: `sever.dragomir@vu.edu.au`

² DEPARTMENT OF MATHEMATICS, TEHRAN TEACHER TRAINING UNIVERSITY, P. O. BOX 15618, TAHRAN, IRAN

BANACH MATHEMATICAL RESEARCH GROUP (BMRG), MASHHAD, IRAN.

E-mail address: `khosravi_m@saba.tmu.ac.ir`

³ DEPARTMENT OF MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN;

CENTER OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, IRAN.

E-mail address: `moslehian@ferdowsi.um.ac.ir` and `moslehian@ams.org`