NOTES ON THE SPACES OF BILINEAR MULTIPLIERS

OSCAR BLASCO

ABSTRACT. A locally integrable function $m(\xi, \eta)$ defined on $\mathbb{R}^n \times \mathbb{R}^n$ is said to be a bilinear multiplier on \mathbb{R}^n of type (p_1, p_2, p_3) if

$$B_m(f,g)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \hat{f}(\xi) \hat{g}(\eta) m(\xi,\eta) e^{2\pi i (\langle \xi + \eta, x \rangle)} d\xi d\eta$$

defines a bounded bilinear operator from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^{p_3}(\mathbb{R}^n)$. The study of the basic properties of such spaces is investigated and several methods of constructing examples of bilinear multipliers are provided. The special case where $m(\xi, \eta) = M(\xi - \eta)$ for a given M defined on \mathbb{R}^n is also addressed.

1. Introduction.

Throughout the paper $C_{00}(\mathbb{R}^n)$ denotes the space of continuous functions defined in \mathbb{R}^n with compact support, $\mathcal{S}(\mathbb{R}^n)$ denotes the Schwartz class on \mathbb{R}^n , i.e. $f: \mathbb{R}^n \to \mathbb{C}$ such that $f \in C^{\infty}(\mathbb{R}^n)$ and $x^{\alpha} \frac{\partial^{|\beta|} f(x)}{\partial x_1^{\beta_1} \dots \partial x_n^{\beta_n}}$ is bounded for any $\beta = (\beta_1, \dots, \beta_n)$ and $\alpha = (\alpha_1, \dots, \alpha_n)$ where $x^{\alpha} = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and $|\beta| = \beta_1 + \dots + \beta_n$ and $\mathcal{P}(\mathbb{R}^n)$ stands for the set of functions in $\mathcal{S}(\mathbb{R}^n)$ such that $\hat{f} \in C_{00}(\mathbb{R}^n)$ where $\hat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-2\pi i \langle x, \xi \rangle} dx$.

We shall use the notation $\mathcal{M}_{p,q}(\mathbb{R}^n)$ (respect. $\tilde{\mathcal{M}}_{p,q}(\mathbb{R}^n)$), for $1 \leq p, q \leq \infty$, for the space of distributions $u \in \mathcal{S}'(\mathbb{R}^n)$ such that $u * \phi \in L^q(\mathbb{R}^n)$ for all $\phi \in L^p(\mathbb{R}^n)$ (respect. for the space of bounded functions m such that T_m defines a bounded operator from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ where $\widehat{T_m(\phi)}(\xi) = m(\xi)\widehat{f}(\xi)$.) We endow the space $\tilde{\mathcal{M}}_{p,q}(\mathbb{R}^n)$ with the "norm" of the operator T_m , that is $||m||_{p,q} = ||T_m||$.

Let us start off by mentioning some well known properties of the space of linear multipliers (see [1, 14]): $\mathcal{M}_{p,q}(\mathbb{R}^n) = \{0\}$ whenever q < p, $\mathcal{M}_{p,q}(\mathbb{R}^n) = \mathcal{M}_{q',p'}(\mathbb{R}^n)$ for $1 and for <math>1 \le p \le 2$, $\mathcal{M}_{1,1}(\mathbb{R}^n) \subset \mathcal{M}_{p,p}(\mathbb{R}^n) \subset \mathcal{M}_{2,2}(\mathbb{R}^n)$. We also have the identifications

$$\tilde{\mathcal{M}}_{2,2}(\mathbb{R}^n) = L^{\infty}(\mathbb{R}^n),$$

$$\mathcal{M}_{1,q}(\mathbb{R}^n) = \{ u \in \mathcal{S}'(\mathbb{R}^n) : u \in L^q(\mathbb{R}^n) \}, 1 < q < \infty,$$

$$\mathcal{M}_{1,1}(\mathbb{R}^n) = \{ u \in \mathcal{S}'(\mathbb{R}^n) : u = \mu \in M(\mathbb{R}^n) \}.$$

In this paper we shall be dealing with their bilinear analogues.

Definition 1.1. Let $1 \leq p_1, p_2 \leq \infty$ and $0 < p_3 \leq \infty$ and let $m(\xi, \eta)$ be a locally integrable function on $\mathbb{R}^n \times \mathbb{R}^n$. Define

$$B_m(f,g)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \hat{f}(\xi) \hat{g}(\eta) m(\xi,\eta) e^{2\pi i (\langle \xi + \eta, x \rangle)} d\xi d\eta$$

 $Key\ words\ and\ phrases.$ spaces of bilinear multipliers, bilinear Hilbert transform, bilinear fractional transform.

Partially supported by Proyecto MTM2008-04594/MTM.

for $f, g \in \mathcal{P}(\mathbb{R}^n)$.

m is said to be a bilinear multiplier on \mathbb{R}^n of type (p_1, p_2, p_3) if there exists C > 0 such that

$$||B_m(f,g)||_{p_3} \le C||f||_{p_1}||g||_{p_2}$$

for any $f, g \in \mathcal{P}(\mathbb{R}^n)$, i.e. B_m extends to a bounded bilinear operator from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^{p_3}(\mathbb{R}^n)$ (where we replace $L^{\infty}(\mathbb{R}^n)$ for $C_0(\mathbb{R}^n)$ in the case $p_i = \infty$ for i = 1, 2).

We write $\mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R}^n)$ for the space of bilinear multipliers of type (p_1,p_2,p_3) and $\|m\|_{p_1,p_2,p_3} = \|B_m\|$.

The study of bilinear multipliers for smooth symbols (where $m(\xi, \eta)$ is a "nice" regular function) goes back to the work by R.R. Coifman and Y. Meyer in [6].

Particularly simple examples are the following bilinear convolution-type operators: For a given $K \in L^1_{loc}(\mathbb{R}^n)$ we define

(1)
$$C_K(f,g)(x) = \int_{\mathbb{R}} f(x-y)g(x+y)K(y)dy$$

for f and g belonging to $C_{00}(\mathbb{R}^n)$.

If $K \in L^1(\mathbb{R}^n)$ then $m(\xi, \eta) = \hat{K}(\xi - \eta)$ defines a multiplier in $\mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R}^n)$ for $1/p_1 + 1/p_2 = 1/p_3$ if $p_3 \ge 1$ and $||m||_{p_1, p_2, p_3} \le ||K||_1$.

Indeed, for f and $g \in \mathcal{S}(\mathbb{R})$, one has $f(x-y) = \int_{\mathbb{R}^n} \hat{f}(\xi)e^{2\pi i\langle x-y,\xi\rangle}d\xi$ and $g(x+y) = \int_{\mathbb{R}^n} \hat{g}(\eta)e^{2\pi i\langle x+y,\eta\rangle}d\eta$. Hence we have

$$C_{K}(f,g)(x) = \int_{\mathbb{R}^{n}} f(x-y)g(x+y)K(y)dy$$

$$= \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} \hat{f}(\xi)\hat{g}(\eta)K(y)e^{2\pi i\langle x-y,\xi\rangle}e^{2\pi i\langle x+y,\eta\rangle}d\xi d\eta dy$$

$$= \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} \hat{f}(\xi)\hat{g}(\eta)(\int_{\mathbb{R}^{n}} K(y)e^{-2\pi i\langle \xi-\eta,y\rangle}dy)e^{2\pi i\langle \xi+\eta,x\rangle}d\xi d\eta$$

$$= \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} \hat{g}(\eta)\hat{f}(\xi)\hat{K}(\xi-\eta)e^{2\pi i\langle \xi+\eta,x\rangle}d\xi d\eta.$$

This motivates the introduction of the following class of multipliers.

Definition 1.2. Let $1 \leq p_1, p_2 \leq \infty$ and $0 < p_3 \leq \infty$. We denote by $\tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R}^n)$ the space of measurable functions $M : \mathbb{R}^n \to \mathbb{C}$ such that $m(\xi, \eta) = M(\xi - \eta) \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$, that is to say

$$B_M(f,g)(x) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \hat{f}(\xi)\hat{g}(\eta)M(\xi - \eta)e^{2\pi i\langle \xi + \eta, x \rangle}d\xi d\eta$$

extends to a bounded bilinear map from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ into $L^{p_3}(\mathbb{R}^n)$. We keep the notation $||M||_{p_1,p_2,p_3} = ||B_M||$.

It was only in the last decade that the cases $M_0(x) = \frac{1}{|x|^{1-\alpha}}$ were shown to define bilinear multipliers of type (p_1, p_2, p_3) for $1/p_3 = 1/p_1 + 1/p_2 - \alpha$ for $1 < p_1, p_2 < \infty$ and $0 < \alpha < 1/p_1 + 1/p_2$ (see (3) in Theorem 1.3) and, in the case n = 1, $M_1(x) = -isign(x)$ was shown to define a bilinear multiplier of type (p_1, p_2, p_3) for $1/p_3 = 1/p_1 + 1/p_2$ for $1 < p_1, p_2 < \infty$ and $p_3 > 2/3$ (see (2) in Theorem 1.3). These two main examples correspond to the following bilinear operators: the bilinear fractional integral defined by

$$I_{\alpha}(f,g)(x) = \int_{\mathbb{R}} \frac{f(x-y)g(x+y)}{|y|^{1-\alpha}} dy, \quad 0 < \alpha < 1$$

and the bilinear Hilbert transform defined by

$$H(f,g)(x) = \lim_{\varepsilon \to 0} \frac{1}{\pi} \int_{|y| > \varepsilon} \frac{f(x-y)g(x+y)}{y} dy$$

respectively.

Let us collect the results about their boundedness which are known nowadays.

Theorem 1.3. Let $1 < p_1, p_2 < \infty, 0 < \alpha < 1/p_1 + 1/p_2, 1/q = 1/p_1 + 1/p_2 - \alpha, 1/p_3 = 1/p_1 + 1/p_2 \text{ and } 2/3 < p_3 < \infty.$ Then there exist constants A and B such that

(2)
$$||H(f,g)||_{p_3} \le A||f||_{p_1}||g||_{p_2} (Lacey-Thiele, [12, 13]),$$

(3)
$$||I_{\alpha}(f,g)||_q \leq B||f||_{p_1}||g||_{p_2}$$
. (Kenig-Stein [11], Grafakos-Kalton [10]).

Our objective is to study the basic properties of the classes $\mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ and $\mathcal{M}_{p_1,p_2,p_3}(\mathbb{R})$, to find examples of bilinear multipliers in these classes and to get methods to produce new ones.

As usual, if $f \in L^1(\mathbb{R}^n)$ we denote by τ_x , M_x and D_t^p the translation $\tau_x f(y) = f(y-x)$ for $x \in \mathbb{R}^n$, the modulation $M_x f(y) = e^{2\pi i \langle x,y \rangle} f(y)$ and the dilation $D_t^p f(x) = t^{-n/p} f(\frac{x}{t})$ for $0 < p, t < \infty$.

With this notation out of the way one has, for $1 \le p \le \infty$ and 1/p + 1/p' = 1,

(4)
$$\widehat{(\tau_x f)}(\xi) = M_{-x} \widehat{f}(\xi), \quad \widehat{(M_x f)}(\xi) = \tau_x \widehat{f}(\xi), \quad \widehat{(D_t^p f)}(\xi) = D_{t^{-1}}^{p'} \widehat{f}(\xi).$$

Clearly τ_x, M_x and D_t^p are isometries on $L^p(\mathbb{R}^n)$ for any 0 .

Although most of the results presented in what follows have a formulation in $n \ge 1$ we shall restrict ourselves to the case n = 1 for simplicity. The reader is referred to [2, 3, 4, 5, 7] for several similar results on other groups, and to find same methods of transference.

2. Bilinear multipliers: The basics

Let us start by pointing out a characterization, for $p_3 \ge 1$, in terms of the duality, whose elementary proof is left to the reader.

Proposition 2.1. Let $1 \leq p_3 \leq \infty$. Then $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ if and only if there exists C > 0 such that

$$\left| \int_{\mathbb{R}^2} \hat{f}(\xi) \hat{g}(\eta) \hat{h}(\xi + \eta) m(\xi, \eta) d\xi d\eta \right| \le C \|f\|_{p_1} \|g\|_{p_2} \|h\|_{p_3'}$$

for all $f, g, h \in \mathcal{P}(\mathbb{R})$.

We now present a basic example of a bilinear multiplier. For a Borel regular measure in \mathbb{R} μ we denote $\hat{\mu}(\xi) = \int_{\mathbb{R}} e^{-2\pi i x \xi} d\mu(x)$ its Fourier transform.

Proposition 2.2. Let $p_3 \ge 1$ and $1/p_1 + 1/p_2 = 1/p_3$ and let $m(\xi, \eta) = \hat{\mu}(\alpha \xi + \beta \eta)$ where μ is a Borel regular measure in \mathbb{R} and $(\alpha, \beta) \in \mathbb{R}^2$. Then $m \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$ and $\|m\|_{p_1, p_2, p_3} \le \|\mu\|_1$.

Proof. Let us first rewrite the value $B_m(f,g)$ as follows:

$$\begin{split} B_m(f,g)(x) &= \int_{\mathbb{R}^2} \hat{f}(\xi) \hat{g}(\eta) \hat{\mu}(\alpha \xi + \beta \eta) e^{2\pi i (\xi + \eta) x} d\xi d\eta \\ &= \int_{\mathbb{R}^2} \hat{f}(\xi) \hat{g}(\eta) (\int_{\mathbb{R}} e^{-2\pi i (\alpha \xi + \beta \eta) t} d\mu(t)) e^{2\pi i (\xi + \eta) x} d\xi d\eta \\ &= \int_{\mathbb{R}} (\int_{\mathbb{R}^2} \hat{f}(\xi) \hat{g}(\eta) e^{2\pi i (x - \alpha t) \xi} e^{2\pi i (x - \beta t) \eta} d\xi d\eta) d\mu(t) \\ &= \int_{\mathbb{R}} f(x - \alpha t) g(x - \beta t) d\mu(t). \end{split}$$

Hence, using Minkowski's inequality, one has

$$||B_{m}(f,g)||_{p_{3}} \leq \int_{\mathbb{R}} ||f(\cdot - \alpha t)g(\cdot - \beta t)||_{p_{3}} d|\mu|(t)$$

$$\leq \int_{\mathbb{R}} ||f(\cdot - \alpha t)||_{p_{1}} ||g(\cdot - \beta t)||_{p_{2}} d|\mu|(t)$$

$$= ||f||_{p_{1}} ||g||_{p_{2}} \int_{\mathbb{R}} d|\mu|(t) = ||\mu||_{1} ||f||_{p_{1}} ||g||_{p_{2}}.$$

Let us start with some elementary properties of the bilinear multipliers when composing with translations, modulations and dilations.

Proposition 2.3. Let $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$.

(a) If $m_1 \in \tilde{\mathcal{M}}_{s_1,p_1}(\mathbb{R})$ and $m_2 \in \tilde{\mathcal{M}}_{s_2,p_2}(\mathbb{R})$ then $m_1(\xi)m(\xi,\eta)m_2(\eta) \in \mathcal{BM}_{(s_1,s_2,p_3)}(\mathbb{R})$.

$$||m_1mm_2||_{s_1,s_2,p_3} \le ||m_1||_{s_1,p_1} ||m||_{p_1,p_2,p_3} ||m_2||_{s_2,p_2}$$

(b)
$$\tau_{(\xi_0,\eta_0)} m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R}) \text{ for each } (\xi_0,\eta_0) \in \mathbb{R}^2 \text{ and }$$

$$\|\tau_{(\xi_0,\eta_0)}m\|_{p_1,p_2,p_3} = \|m\|_{p_1,p_2,p_3}.$$

(c) $M_{(\xi_0,\eta_0)}m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ for each $(\xi_0,\eta_0) \in \mathbb{R}^2$ and

$$||M_{(\xi_0,\eta_0)}m||_{p_1,p_2,p_3} = ||m||_{p_1,p_2,p_3}$$

(d) If
$$\frac{2}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3}$$
 and $0 < t < \infty$ then $D_t^q m \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$ and

$$||D_t^q m||_{p_1,p_2,p_3} = ||m||_{p_1,p_2,p_3}.$$

Proof. Use (4) to deduce the following formulas

(5)
$$B_{m_1 m m_2}(f, g) = B_m(T_{m_1} f, T_{m_2} g).$$

(6)
$$B_{\tau_{(\xi_0,\eta_0)}m}(f,g) = M_{\xi_0+\eta_0}B_m(M_{-\xi_0}f,M_{-\eta_0}g).$$

(7)
$$B_{M_{(\xi_0,\eta_0)}m}(f,g) = B_m(\tau_{-\xi_0}f,\tau_{-\eta_0}g).$$

(8)
$$B_m(D_t^{p_1}f, D_t^{p_2}g) = D_t^{p_3}B_{D_t^qm}(f, g).$$

Let us check only the validity of last one. The other ones follow easily from the previous facts.

$$\begin{split} B_m(D_t^{p_1}f,D_t^{p_2}g)(x) &= \int_{\mathbb{R}^2} t^{\frac{1}{p_1'}} \hat{f}(t\xi) t^{\frac{1}{p_2'}} \hat{g}(t\eta) m(\xi,\eta) e^{2\pi i (\xi+\eta)x} d\xi d\eta \\ &= \int_{\mathbb{R}^2} t^{\frac{1}{p_1'}} \hat{f}(\xi) t^{\frac{1}{p_2'}} \hat{g}(\eta) m(\frac{\xi}{t},\frac{\eta}{t}) e^{2\pi i (\xi+\eta)\frac{x}{t}} t^{-2} d\xi d\eta \\ &= t^{-\frac{1}{p_3}} \int_{\mathbb{R}^2} \hat{f}(\xi) \hat{g}(\eta) t^{-\frac{1}{p_1} - \frac{1}{p_2} + \frac{1}{p_3}} m(\frac{\xi}{t},\frac{\eta}{t}) e^{2\pi i (\xi+\eta)\frac{x}{t}} d\xi d\eta \\ &= D_t^{p_3} B_{D_t^{q_1}m}(f,g)(x). \end{split}$$

From (8) we can see that the condition $1/p_1 + 1/p_2 = 1/p_3$ is also connected to the homogeneity of the symbol.

Proposition 2.4. Let $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ such that $m(t\xi,t\eta) = m(\xi,\eta)$ for any t > 0. Then $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3}$.

Proof. From assumption $D_t^{\infty} m = m$. Using (8) we have

$$B_m(D_t^{p_1}f, D_t^{p_2}g) = t^{1/p_3 - (1/p_1 + 1/p_2)}D_t^{p_3}B_m(f, g)$$

and therefore

$$||B_{m}(f,g)||_{p_{3}} = ||D_{t}^{p_{3}}B_{m}(f,g)||_{p_{3}}$$

$$= t^{-1/p_{3}+(1/p_{1}+1/p_{2})}||B_{m}(D^{p_{1}}f,D_{t}^{p_{2}}g)||_{p_{3}}$$

$$\leq t^{-1/p_{3}+(1/p_{1}+1/p_{2})}||B_{m}|||f||_{p_{1}}||g||_{p_{2}}.$$

For this to hold for any $0 < t < \infty$ one needs $1/p_1 + 1/p_2 = 1/p_3$.

Let us combine the previous results to get new bilinear multipliers from a given one.

Proposition 2.5. Let $p_3 \geq 1$ and $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$.

- (a) If $Q = [a, b] \times [c, d]$ and $1 < p_1, p_2 < \infty$ then $m\chi_Q \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$ and $||m\chi_{Q}||_{p_{1},p_{2},p_{3}} \leq C||m||_{p_{1},p_{2},p_{3}}.$ (b) If $\Phi \in L^{1}(\mathbb{R}^{2})$ then $\Phi*m \in \mathcal{BM}_{(p_{1},p_{2},p_{3})}(\mathbb{R})$ and $||\Phi*m||_{p_{1},p_{2},p_{3}} \leq ||\Phi||_{1}||m||_{p_{1},p_{2},p_{3}}.$
- (c) If $\Phi \in L^1(\mathbb{R}^2)$ then $\hat{\Phi}m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ and $\|\hat{\Phi}m\|_{p_1,p_2,p_3} \leq \|\Phi\|_1 \|m\|_{p_1,p_2,p_3}$.
- (d) If $\psi \in L^1(\mathbb{R}^+, t^{\frac{1}{p_3} (\frac{1}{p_1} + \frac{1}{p_2})})$ then $m_{\psi}(\xi, \eta) = \int_0^{\infty} m(t\xi, t\eta) \psi(t) dt \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$. Moreover $\|m_{\psi}\|_{p_1, p_2, p_3} \leq \|\psi\|_1 \|m\|_{p_1, p_2, p_3}$.

Proof. (a) Use that $\chi_{[a,b]} \in \tilde{\mathcal{M}}_{p_1,p_1}$ for $1 < p_1 < \infty$ and $\chi_{[c,d]} \in \tilde{\mathcal{M}}_{p_2,p_2}$ for $1 < p_2 < \infty$ together with Proposition 2.3 part (a).

(b) Note that

$$B_{\Phi*m}(f,g)(x) = \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{\mathbb{R}^2} m(\xi - u, \eta - v)\Phi(u, v)dudv\right) e^{2\pi i(\xi + \eta)x} d\xi d\eta$$

$$= \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta)m(\xi - u, \eta - v)e^{2\pi i(\xi + \eta)x} d\xi d\eta\right) \Phi(u, v)dudv$$

$$= \int_{\mathbb{R}^2} B_{\tau(u,v)}m(f,g)(x)\Phi(u, v)dudv.$$

From the vector-valued Minkowski inequality and Proposition 2.3 part (b), we have

$$||B_{\Phi*m}(f,g)||_{p_3} \leq \int_{\mathbb{R}^2} ||B_{\tau_{(u,v)}m}(f,g)||_{p_3} |\Phi(u,v)| du dv$$

$$\leq ||m||_{p_1,p_2,p_3} ||f||_{p_1} ||g||_{p_2} ||\Phi||_1.$$

(c) Observe that

$$B_{\hat{\Phi}m}(f,g)(x) = \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{\mathbb{R}^2} M_{(-u,-v)}m(\xi,\eta)\Phi(u,v)dudv\right) e^{2\pi i(\xi+\eta)x}d\xi d\eta$$
$$= \int_{\mathbb{R}^2} B_{M_{(-u,-v)}m}(f,g)(x)\Phi(u,v)dudv.$$

Argue as above, using now Proposition 2.3 part (c), to conclude the result. (d) Use now Proposition 2.3 part (d), for $\frac{1}{p_3} - (\frac{1}{p_1} + \frac{1}{p_2}) = -\frac{2}{q}$,

$$B_{m_{\psi}}(f,g)(x) = \int_{\mathbb{R}^{2}} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{0}^{\infty} D_{t^{-1}}^{q} m(\xi,\eta) t^{-2/q} \psi(t) dt \right) e^{2\pi i (\xi+\eta)x} d\xi d\eta$$
$$= \int_{0}^{\infty} B_{D_{t^{-1}}^{q} m}(f,g)(x) t^{-2/q} \psi(t) dt.$$

With all these procedures we have several useful methods to produce multipliers in $\mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$. Let us mention one application of each of them.

xample 2.6. (1) If $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3}$, $m_1 \in \tilde{\mathcal{M}}_{(p_1,p_1)}$ and $m_2 \in \tilde{\mathcal{M}}_{(p_2,p_2)}$ then $m(\xi,\eta) = m_1(\xi)m_2(\eta) \in \mathcal{BM}_{p_1,p_2,p_3}$. (2) If $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$, $p_3 \geq 1$ and Q_1,Q_2 are bounded measurable sets in Example 2.6.

$$\frac{1}{|Q_1||Q_2|}\int_{Q_1\times Q_2}m(\xi+u,\eta+v)dudv\in\mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R}).$$

- (3) If $\Phi \in L^1(\mathbb{R}^2)$ then $\hat{\Phi} \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ for $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3}$, $p_3 \ge 1$. (4) If $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$, $|\frac{1}{p_1} + \frac{1}{p_2} \frac{1}{p_3}| < 1$ then

$$m_1(\xi, \eta) = \int_0^\infty m(t\xi, t\eta) \frac{dt}{1 + t^2} \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R}).$$

A combination of the previous results gives the following examples of bilinear multipliers in $\mathcal{B}M_{(1,1,p_3)}(\mathbb{R})$ whose proof is left to the reader.

Corollary 2.7. Let $\Phi \in L^1(\mathbb{R}^2)$, $\psi_1 \in L^{p_1}(\mathbb{R})$ and $\psi_2 \in L^{p_2}(\mathbb{R})$ and $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_2}$ $\frac{1}{p_3} \leq 1$ then

$$m(\xi, \eta) = \hat{\psi}_1(\xi)\hat{\Phi}(\xi, \eta)\hat{\psi}_2(\eta) \in \mathcal{B}M_{(1,1,p_3)}(\mathbb{R}).$$

Let us use Proposition 2.1 and interpolation to get a sufficient integrability condition to guarantee that $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$.

Theorem 2.8. Let $1 \le p_1, p_2 \le p \le 2$ and $p_3 \ge p'$ such that $\frac{1}{p_1} + \frac{1}{p_2} - \frac{2}{p} = \frac{1}{p_3}$. If $m \in L^p(\mathbb{R}^2)$ then $m \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$.

Proof. Let us show first that $m \in \mathcal{BM}_{(p,p,\infty)}(\mathbb{R})$. Let $f,g \in L^p(\mathbb{R})$ and $h \in L^1(\mathbb{R})$. Using Hölder and Hausdorff-Young's inequalities one gets

$$|\int_{\mathbb{R}^{2}} \hat{f}(\xi)\hat{g}(\eta)\hat{h}(\xi+\eta)m(\xi,\eta)d\xi d\eta| \leq ||m||_{L^{p}(\mathbb{R}^{2})}||\hat{h}||_{\infty}||\hat{f}||_{p'}||\hat{g}||_{p'}$$
$$\leq ||m||_{L^{p}(\mathbb{R}^{2})}||h||_{1}||f||_{p}||g||_{p}.$$

Similarly, changing the variables $\xi + \eta = u$, $\xi = -v$, one has

$$\int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta)\hat{h}(\xi+\eta)m(\xi,\eta)d\xi d\eta = \int_{\mathbb{R}^2} \hat{f}(-v)\hat{g}(u+v)\hat{h}(u)m(-v,u+v)dv du.$$

An argument as above gives also the estimate

$$\left| \int_{\mathbb{R}^2} \hat{f}(-v)\hat{g}(u+v)\hat{h}(u)m(-v,u+v)dvdu \right| \leq \|m\|_{L^p(\mathbb{R}^2)} \|g\|_1 \|f\|_p \|h\|_p.$$

This shows that $m \in \mathcal{BM}_{(p,1,p')}(\mathbb{R})$. A similar argument shows also that $m \in$ $\mathcal{BM}_{(1,p,p')}(\mathbb{R}).$

Given $1 \leq \tilde{p}_1 \leq p$ and $p' \leq \tilde{p}_3 \leq \infty$ with $\frac{1}{\tilde{p}_1} - \frac{1}{\tilde{p}_3} = \frac{1}{p}$ we have $0 \leq \theta \leq 1$ such that $\frac{1}{\tilde{p}_1} = \frac{1-\theta}{p} + \frac{\theta}{1}$ and $\frac{1}{\tilde{p}_3} = \frac{1-\theta}{\infty} + \frac{\theta}{p'}$. Hence, by interpolation, $m \in \mathcal{BM}_{(\tilde{p}_1, p, \tilde{p}_3)}(\mathbb{R})$. Similarly $m \in \mathcal{BM}_{(p, \tilde{p}_2, \tilde{q}_3)}(\mathbb{R})$ whenever $1 \leq \tilde{p}_2 \leq p$ and $p' \leq \tilde{q}_3 \leq \infty$ with

 $\frac{1}{\bar{p}_2} - \frac{1}{\bar{q}_3} = \frac{1}{p}$. To finish the proof we observe that if $1 < p_1 < p$ and $1 < p_2 < p$ then for each

 $0 < \theta < 1$ there exist $1 \le \tilde{p}_1 \le p_1 < p$ and $1 \le \tilde{p}_2 \le p_2 < p$ such that

$$\frac{1}{p_1} - \frac{1}{p} = (1 - \theta)(\frac{1}{\tilde{p}_1} - \frac{1}{p}), \quad \frac{1}{p_2} - \frac{1}{p} = \theta(\frac{1}{\tilde{p}_2} - \frac{1}{p}).$$

Denoting \tilde{p}_3 , \tilde{q}_3 the values such that $\frac{1}{\tilde{p}_2} - \frac{1}{p} = \frac{1}{\tilde{p}_3}$ and $\frac{1}{\tilde{p}_2} - \frac{1}{p} = \frac{1}{\tilde{q}_3}$ one obtains that

$$\frac{1}{p_1} = \frac{(1-\theta)}{\tilde{p}_1} + \frac{\theta}{p}, \quad \frac{1}{p_2} = \frac{(1-\theta)}{p} + \frac{\theta}{\tilde{p}_1}, \quad \frac{1}{p_3} = \frac{(1-\theta)}{\tilde{p}_3} + \frac{\theta}{\tilde{q}_3}.$$

Hence the result follows again from interpolation between the last ones.

3. Bilinear multipliers defined by functions in one variable

Let us restrict ourselves to a smaller family of multipliers where $m(\xi, \eta) = M(\xi - 1)$ η) for some M defined in \mathbb{R} . These multipliers satisfy

(9)
$$B_m(M_x f, M_x g) = M_{2x} B_m(f, g).$$

As in the introduction we use the notation $\tilde{\mathcal{M}}_{p_1,p_2,p_3}(\mathbb{R})$ for the space of functions $M: \mathbb{R} \to \mathbb{C}$ such that $m(\xi, \eta) = M(\xi - \eta) \in \mathcal{BM}_{(p_1, p_2, p_3)}(\mathbb{R})$, that is to say

$$B_M(f,g)(x) = \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta)M(\xi - \eta)e^{2\pi i(\xi + \eta)x}d\xi d\eta,$$

defined for \hat{f} and \hat{g} compactly supported, extends to a bounded bilinear map from $L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R})$ into $L^{p_3}(\mathbb{R})$. We keep the notation $||M||_{p_1,p_2,p_3} = ||B_M||$.

The reader should be aware that the starting assumption on the function M is only relevant for the definition of the bilinear mapping to make sense when acting on certain classes of "nice" functions. Then a density argument allows to extend functions belonging to Lebesgue spaces. We would like to point out the following observation.

Remark 3.1. If $M_n \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ are functions such that $M_n(x) \to M(x)$ a.e and $\sup_n \|M_n\| < \infty$ then $M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ and $\|M\|_{p_1,p_2,p_3} \leq \sup_n \|M_n\|_{p_1,p_2,p_3}$. Indeed, this fact follows from Fatou's lemma, since

$$||B_M(f,g)||_{p_3} \le \liminf ||B_{M_n}(f,g)||_{p_3} \le \sup_n ||M_n||_{p_1,p_2,p_3} ||f||_{p_1} ||g||_{p_2}.$$

Remark 3.2. The case $M(x) = \frac{1}{|x|^{1-\alpha}}$ (and even the n-dimensional case) corresponds to the bilinear fractional integral. This was first shown by C. Kenig and E. Stein in [11] to belong to $\tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ for any $1 < p_1, p_2 < \infty, 0 < \alpha < 1/p_1+1/p_2$ and $1/p_1+1/p_2=1/p_3-\alpha$. Another very important and non trivial example is the bilinear Hilbert transform, given by $M(x)=-i \operatorname{sign}(x)$, which was shown by M. Lacey and C. Thiele in [12, 13, ?] to belong to $\tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ for any $1 < p_1, p_2 < \infty, 1/p_1+1/p_2=1/p_3$ and $p_3>2/3$. These results were extended to other cases in [10] and [8, 9] respectively.

We start reformulating the definition of this class of bilinear multipliers.

Proposition 3.3. Let $M \in L^1_{loc}(\mathbb{R})$, $f, g \in \mathcal{P}(\mathbb{R})$. Then

(10)
$$B_M(f,g)(x) = \frac{1}{2} \int_{\mathbb{R}^2} \hat{f}(\frac{u+v}{2}) \hat{g}(\frac{u-v}{2}) M(v) e^{2\pi i u x} du dv$$

(11)
$$B_M(f,g)(-x) = \int_{\mathbb{R}} (\widehat{\tau_x g} * M)(\xi) \widehat{\tau_x f}(\xi) d\xi.$$

(12)
$$\widehat{B_M(f,g)}(x) = \frac{1}{2} C_M(\widehat{D_{1/2}^1 f}, \widehat{D_{1/2}^1 g})(x).$$

Proof. (10) follows changing variables.

To show (11) observe that

$$B_{M}(f,g)(-x) = \int_{\mathbb{R}^{2}} \widehat{\tau_{x}f}(\xi)\widehat{\tau_{x}g}(\eta)M(\xi-\eta)d\xi d\eta$$
$$= \int_{\mathbb{R}} (\int_{\mathbb{R}} \widehat{\tau_{x}g}(\eta)M(\xi-\eta)d\eta)\widehat{\tau_{x}f}(\xi)d\xi$$
$$= \int_{\mathbb{R}} (\widehat{\tau_{x}g}*M)(\xi)\widehat{\tau_{x}f}(\xi)d\xi$$

Finally, using (10), we have

$$\begin{split} B_M(f,g)(x) &= \frac{1}{2} \int_{\mathbb{R}} \Big(\int_{\mathbb{R}} \hat{f}(\frac{u+v}{2}) \hat{g}(\frac{u-v}{2}) M(v) dv \Big) e^{2\pi i u x} dv \\ &= \frac{1}{2} \int_{\mathbb{R}} C_M(D_{1/2}^{\infty} \hat{f}, D_{1/2}^{\infty} \hat{g})(u) e^{2\pi i u x} du. \end{split}$$

This implies (12).

For symbols M which are integrable we can write B_M in terms C_K for some kernel K.

Proposition 3.4. Let $M \in L^1(\mathbb{R})$ and set $K(t) = \hat{M}(-t)$. Then $B_M = C_K$, i.e

$$B_M(f,g) = \int_{\mathbb{R}} f(x-t)g(x+t)K(t)dt$$

Proof.

$$C_K(f,g)(x) = \int_{\mathbb{R}} f(x-t)g(x+t)\hat{M}(-t)dt$$

$$= \int_{\mathbb{R}} (\int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta)e^{2\pi i(x-t)\xi}e^{2\pi i(x+t)\eta}d\xi d\eta)\hat{M}(-t)dt$$

$$= \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta)(\int_{\mathbb{R}} \hat{M}(t)e^{2\pi i(\xi-\eta)t}dt)e^{2\pi i(\xi+\eta)x}d\xi d\eta$$

$$= B_M(f,g)(x).$$

This class does have much richer properties than $\mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$. As above use the notation $f_t(x) = D_t^1 f(x) = \frac{1}{t} f(\frac{x}{t})$ for a function f defined in \mathbb{R} . The following facts are immediate.

(13)
$$\tau_y B_M(f,g) = B_M(\tau_y f, \tau_y g), y \in \mathbb{R}.$$

(14)
$$M_{2y}B_M(f,g) = B_M(M_y f, M_y g), y \in \mathbb{R}.$$

(15)
$$(B_M(f,g))_t = B_{D_{t-1}^1 M}(f_t, g_t), t > 0.$$

When specializing the properties obtained for $m(\xi, \eta)$ to the case $M(\xi - \eta)$ we get the following facts:

(16)
$$B_M(\tau_{-y}f, \tau_y g) = B_{M_y M}(f, g), y \in \mathbb{R}.$$

(17)
$$B_M(M_yf, M_{-y}g) = B_{\tau_{2y}M}(f, g), y \in \mathbb{R}.$$
 For $\frac{1}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3}$ we have
$$B_M(D_t^{p_1}f, D_t^{p_2}g) = D_t^{p_3}B_{D_t^qM}(f, g), t > 0.$$

As in the previous section we can generate new multipliers in $\tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$.

Proposition 3.5. Let $p_3 \geq 1$, $\phi \in L^1(\mathbb{R})$ and $M \in \mathcal{M}_{(p_1,p_2,p_3)}(\mathbb{R})$. Then

- (a) $\phi * M \in \tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R})$ and $\|\phi * M\|_{p_1, p_2, p_3} \le \|\phi\|_1 \|M\|_{p_1, p_2, p_3}$.
- (b) $\hat{\phi}M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R}) \text{ and } \|\hat{\phi}M\|_{p_1,p_2,p_3} \leq \|\phi\|_1 \|M\|_{p_1,p_2,p_3}.$ (c) $If \psi \in L^1(\mathbb{R}^+, t^{\frac{1}{p_3} (\frac{1}{p_1} + \frac{1}{p_2})}) \text{ then } M_{\psi}(\xi) = \int_0^{\infty} M(t\xi) \psi(t) dt \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R}).$ Moreover $||M_{\psi}||_{p_1,p_2,p_3} \leq ||\psi||_1 ||M||_{p_1,p_2,p_3}$.

Proof. (a) Apply Minkowski's inequality to the following fact:

$$B_{\phi*M}(f,g)(x) = \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{\mathbb{R}} M(\xi - \eta - u)\phi(u)du \right) e^{2\pi i(\xi + \eta)x} d\xi d\eta$$

$$= \int_{\mathbb{R}} \left(\int_{\mathbb{R}^2} \widehat{M_{-u}f}(\xi)\hat{g}(\eta)M(\xi - \eta)e^{2\pi i(\xi + \eta)x} d\xi d\eta \right) e^{2\pi i ux} \phi(u)du$$

$$= \int_{\mathbb{R}} M_u B_M(M_{-u}f,g)(x)\phi(u)du.$$

(b) Observe that

$$B_{\hat{\phi}m}(f,g)(x) = \int_{\mathbb{R}^2} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{\mathbb{R}} (M_{-u}m)(\xi - \eta)\phi(u)du\right) e^{2\pi i(\xi + \eta)x} d\xi d\eta$$
$$= \int_{\mathbb{R}^2} B_{M_{-u}m}(f,g)(x)\phi(u)du.$$

Use now Minkowski's again and (16).

(c) Write
$$\frac{1}{p_3} - (\frac{1}{p_1} + \frac{1}{p_2}) = -\frac{1}{q}$$
,

$$B_{M_{\psi}}(f,g)(x) = \int_{\mathbb{R}^{2}} \hat{f}(\xi)\hat{g}(\eta) \left(\int_{0}^{\infty} D_{t^{-1}}^{q} M(\xi) t^{-1/q} \psi(t) dt\right) e^{2\pi i (\xi+\eta)x} d\xi d\eta$$
$$= \int_{0}^{\infty} B_{D_{t^{-1}}^{q} M}(f,g)(x) t^{-1/q} \psi(t) dt.$$

The result follows from (18) and Minkowski's again.

Proposition 3.6. Let $p_3 \geq 1$, $\phi \in L^1(\mathbb{R})$ and $M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$. Then $m(\xi,\eta) = M(\xi - \eta)\hat{\phi}(\xi + \eta) \in \mathcal{BM}_{(p_1,p_2,p_3)}(\mathbb{R})$ and $\|m\|_{p_1,p_2,p_3} \leq \|\phi\|_1 \|M\|_{p_1,p_2,p_3}$.

Proof. Apply Young's inequality to the following fact:

$$B_{m}(f,g)(x) = \int_{\mathbb{R}^{2}} \hat{f}(\xi)\hat{g}(\eta)M(\xi-\eta)(\int_{\mathbb{R}} \phi(y)e^{-2\pi i(\xi+\eta)y}dy)e^{2\pi i(\xi+\eta)x}d\xi d\eta$$
$$= \int_{\mathbb{R}} (\int_{\mathbb{R}^{2}} \hat{f}(\xi)\hat{g}(\eta)M(\xi-\eta)e^{2\pi i(\xi+\eta)(x-y)}d\xi d\eta)\phi(y)dy$$
$$= \phi * B_{M}(f,g)(x).$$

Let us show that the classes $\tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ are reduced to $\{0\}$ for some values of the parameters.

Theorem 3.7. Let $p_3 \geq 1$ such that $\frac{1}{p_1} + \frac{1}{p_2} < \frac{1}{p_3}$. Then $\tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R}) = \{0\}$.

Proof. Let $M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$. Using Proposition 3.5 we have that $\phi * M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ for any ϕ continuous with compact support. Hence we may assume that $M \in L^1(\mathbb{R})$. Using Proposition 3.4 one has that

$$B_M(f,g)(x) = \int_{(x+B_R)\cap(-x+B_R)} f(x-t)g(x+t)\hat{M}(-t)dt$$

for any f and g continuous functions supported in a ball $B_R = \{|x| \leq R\}$. Therefore one concludes that $supp(B_M(f,g)) \subset B_{2R}$ in such a case. On the other hand for any compactly supported function h, 0 and <math>g big enough one can say that $||h \pm \tau_y f||_p = 2^{1/p} ||f||_p$.

Consider $\{r_k\}$ the Rademacher system in [0,1] and observe that, for each $N \in \mathbb{N}$ and $y \in \mathbb{R}$, the orthonormality of the system gives

$$\int_{0}^{1} B_{M}(\sum_{k=0}^{N} r_{k}(t)\tau_{ky}f, \sum_{k=0}^{N} r_{k}(t)\tau_{ky}f)dt = \sum_{k=0}^{N} B_{M}(\tau_{ky}f, \tau_{ky}g)$$

Therefore, since $\sum_{k=0}^{N} B_M(\tau_{ky}f, \tau_{ky}g) = \sum_{k=0}^{N} \tau_{ky}B_M(f,g)$, we conclude that for g big enough

$$\|\sum_{k=0}^{N} \tau_{ky} B_M(f,g)\|_{p_3}^{p_3} = (N+1) \|B_M(f,g)\|_{p_3}^{p_3}.$$

On the other hand, for $p_3 \geq 1$,

$$\| \int_{0}^{1} B_{M}(\sum_{k=0}^{N} r_{k}(t)\tau_{ky}f, \sum_{k=0}^{N} r_{k}(t)\tau_{ky}g)dt \|_{p_{3}}$$

$$\leq \int_{0}^{1} \|B_{M}(\sum_{k=0}^{N} r_{k}(t)\tau_{ky}f, \sum_{k=0}^{N} r_{k}(t)\tau_{ky}g) \|_{p_{3}}dt$$

$$\leq \int \|B_{M}\| \|\sum_{k=0}^{N} r_{k}(t)\tau_{ky}f \|_{p_{1}} \|\sum_{k=0}^{N} r_{k}(t)\tau_{ky}g) \|_{p_{2}}dt$$

$$\leq \|B_{M}\| \sup_{0 < t < 1} \|\sum_{k=0}^{N} r_{k}(t)\tau_{ky}f \|_{p_{1}} \sup_{0 < t < 1} \|\sum_{k=0}^{N} r_{k}(t)\tau_{ky}g \|_{p_{2}}$$

$$\leq \|B_{M}\| \|(N+1)^{1/p_{1}} \|f\|_{p_{1}} (N+1)^{1/p_{2}} \|g\|_{p_{2}}.$$

This implies that $(N+1)^{1/p_3} \|B_M(f,g)\|^{p_3} \le C(N+1)^{1/p_1+1/p_2} \|f\|_{p_1} \|g\|_{p_2}$. Hence $1/p_1 + 1/p_2 \ge 1/p_3$.

The following elementary lemma is quite useful to get necessary conditions on multipliers.

Lemma 3.8. Let $M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$. If $\frac{1}{q} = \frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3}$ then there exists C > 0 such that

$$\left| \int_{\mathbb{R}} e^{-\lambda^2 \xi^2} M(\xi) d\xi \right| \le C \|M\|_{p_1, p_2, p_3} \lambda^{\frac{1}{q} - 1}$$

for any $\lambda > 0$.

Proof. Let $\lambda > 0$ and denote G_{λ} such that $\hat{G}_{\lambda}(\xi) = e^{-2\lambda^2 \xi^2}$. Using (10) one concludes that

$$B_{M}(G_{\lambda}, G_{\lambda})(x) = \frac{1}{2} \int_{\mathbb{R}^{2}} e^{-\lambda^{2}v^{2}} e^{-\lambda^{2}u^{2}} M(v) e^{2\pi i u x} du dv$$

$$= \frac{1}{2} \left(\int_{\mathbb{R}} e^{-\lambda^{2}v^{2}} M(v) dv \right) \left(\frac{1}{\lambda} \int_{\mathbb{R}} e^{-u^{2}} e^{2\pi i u \frac{x}{\lambda}} du \right)$$

$$= C \frac{1}{\lambda} e^{-\pi^{2} \frac{x^{2}}{\lambda^{2}}} \left(\int_{\mathbb{R}} e^{-\lambda^{2}v^{2}} M(v) dv \right).$$

Since $\|G_{\lambda}\|_p = C_p \lambda^{\frac{1}{p}-1}$ and $M \in \tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ one gets that

$$||B_M(G_\lambda, G_\lambda)||_{p_3} = C|\int_{\mathbb{R}} e^{-\lambda^2 v^2} M(v) dv |\lambda^{\frac{1}{p_3} - 1} \le C||M||_{p_1, p_2, p_3} \lambda^{\frac{1}{p_1} - 1} \lambda^{\frac{1}{p_2} - 1}.$$

Therefore
$$|\int_{\mathbb{R}} e^{-\lambda^2 \xi^2} M(\xi) d\xi| \le C ||M||_{p_1, p_2, p_3} \lambda^{\frac{1}{q} - 1}$$
.

Theorem 3.9. If there exists a non-zero continuous and integrable function M belonging to $\tilde{\mathcal{M}}_{(p_1,p_2,p_3)}(\mathbb{R})$ then

$$\frac{1}{p_3} \le \frac{1}{p_1} + \frac{1}{p_2} \le \frac{1}{p_3} + 1.$$

Proof. Assume first that $\frac{1}{p_1} + \frac{1}{p_2} < \frac{1}{p_3}$. Use Lemma 3.8 applied to $\tau_{-2y}M$ for any $y \in \mathbb{R}$ together with (16) to obtain

$$|\lambda \int_{\mathbb{R}} e^{-\lambda^2 \xi^2} M(\xi + 2y) d\xi| \le C ||M||_{p_1, p_2, p_3} \lambda^{\frac{1}{q}}.$$

Therefore, using the continuity of M and q < 0 one gets

$$\lim_{\lambda \to \infty} |\lambda \int_{\mathbb{R}} e^{-\lambda^2 \xi^2} M(\xi + 2y) d\xi| = |M(2y)| = 0.$$

Hence M = 0.

Assume now that $\frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3} > 1$. Using again Lemma 3.8, applied to M_yM , together with (17) we obtain

$$\left| \int_{\mathbb{R}} e^{-\lambda^2 \xi^2} M(\xi) e^{2\pi i y \xi} d\xi \right| \le C \|M\|_{p_1, p_2, p_3} \lambda^{\frac{1}{q} - 1}.$$

Therefore, taking limits again as $\lambda \to 0$, since 1/q-1>0 we get $|\hat{M}(y)|=0$. Hence M=0.

Corollary 3.10. (see [16, Prop 3.1]) Let $p_3 \ge 1$ such that $\frac{1}{p_1} + \frac{1}{p_2} < \frac{1}{p_3}$ or $\frac{1}{p_1} + \frac{1}{p_2} > \frac{1}{p_3}$ $\frac{1}{p_3} + 1$. Then $\tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R}) = \{0\}$.

Proof. Let $M \in \mathcal{M}_{(p_1,p_2,p_3)}(\mathbb{R})$. From Proposition 3.5 we have that $\phi * M \in$ $\mathcal{M}_{(p_1,p_2,p_3)}(\mathbb{R})$ for any ϕ compactly supported and continuous. Now use Theorem 3.9 to conclude that $\phi * M = 0$ for any compactly supported and continuous ϕ . This implies that M = 0.

Let us now use some interpolation methods to get more examples of multipliers in $\mathcal{M}_{(p_1,p_2,p_3)}(\mathbb{R})$. First note that, selecting $\alpha=1$ and $\beta=-1$ in Proposition 2.2 we obtain the following simple example.

Proposition 3.11. If $\mu \in M(\mathbb{R})$ then $M = \hat{\mu} \in \tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R})$ for $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_2} \leq 1$ and $||M|| \leq ||\mu||_1$.

Theorem 3.12. Let $\frac{1}{p_3} \leq \frac{1}{p_1} + \frac{1}{p_2} \leq \min\{2, \frac{1}{p_3} + 1\}$. If $M \in L^1(\mathbb{R})$ and $M = \hat{K}$ for some $K \in L^q(\mathbb{R})$ where $\frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3} = 1 - \frac{1}{q}$ then $M \in \tilde{\mathcal{M}}_{(p_1, p_2, p_3)}(\mathbb{R})$ with $||M||_{p_1,p_2,p_3} \le C||K||_q.$

Proof. Consider the trilinear form

$$T(K, f, g) = \int_{\mathbb{R}} f(x - t)g(x + t)K(t)dt.$$

From Proposition 3.4 we have $B_M(f,g) = T(K,f,g)$ for $M = \hat{K}$. Now use Proposition 3.11 to conclude that T is bounded in $L^1(\mathbb{R}) \times L^{q_1}(\mathbb{R}) \times L^{q_2}(\mathbb{R}) \to L^{s_1}(\mathbb{R})$ where $\frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{s_1} \le 1$ and it has norm bounded by 1.

Assume first that $\frac{1}{p_1} + \frac{1}{p_2} \le 1$. Hence T is bounded in $L^1(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to$ $L^p(\mathbb{R})$ for $\frac{1}{p_1} + \frac{1}{p_2} = \hat{1}$. On the other hand, using Hölder's inequality

$$\sup_{x} \left| \int_{\mathbb{R}} f(x-t)g(x+t)K(t)dt \right| \le ||f||_{p_1} ||g||_{p_2} ||K||_{p'}.$$

This shows that T is also bounded in $L^{p'}(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to L^{\infty}(\mathbb{R})$. Therefore, by interpolation, selecting $0 < \theta < 1$ such that $\frac{1}{p_3} = \frac{1-\theta}{p}$, one obtains that T is bounded in $L^q(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to L^{p_3}(\mathbb{R})$ for $\frac{1}{p_1} + \frac{1}{p_2} - \frac{1}{p_3} = 1 - \frac{1}{q}$.

Assume now that $1 < \frac{1}{p_1} + \frac{1}{p_2} \le 2$. Using that $\int_{\mathbb{R}} f(x-t)g(x+t)dt = f * g(2x)$, Young's inequality implies that

$$\| \int_{\mathbb{R}} f(x-t)g(x+t)K(t)dt \|_{r_3} \le \|K\|_{\infty} \|D_{1/2}^{\infty}(|f| * |g|)\|_{r_3} \le C\|f\|_{r_1} \|g\|_{r_2} \|K\|_{\infty}$$

whenever $\frac{1}{r_1} + \frac{1}{r_2} \ge 1$ and $\frac{1}{r_1} + \frac{1}{r_2} - 1 = \frac{1}{r_3}$. Hence T is bounded in $L^{\infty}(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to L^p(\mathbb{R})$ where $\frac{1}{p_1} + \frac{1}{p_2} - 1 = \frac{1}{p_2}$

Using duality, $\langle T(K, f, q), h \rangle = \langle T(h, \bar{f}, q), K \rangle$, where $\bar{f}(x) = f(-x)$, that is

$$\int_{\mathbb{R}^2} f(x-t)g(x+t)K(t)h(x)dtdx = \int_{\mathbb{R}} (\int_{\mathbb{R}} \bar{f}(t-x)g(x+t)h(x)dx)K(t)dt.$$

Therefore T is also bounded in $L^{p'}(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to L^1(\mathbb{R})$.

Select $0 \le \theta \le 1$ such that $\frac{1}{p_3} = \frac{1}{p} + \frac{\theta}{p'}$. Now using interpolation T will be bounded in $L^q(\mathbb{R}) \times L^{p_1}(\mathbb{R}) \times L^{p_2}(\mathbb{R}) \to L^{p_3}(\mathbb{R})$ for $\frac{1}{q} = \frac{\theta}{p'} = \frac{1}{p_3} - \frac{1}{p} = \frac{1}{p_3} - \frac{1}{p_1} - \frac{1}{p_2} + \frac{1}{p_3}$.

References

- [1] Bergh, J., Löfström, J., Interpolation spaces. Springer Verlag [1970].
- [2] Blasco, O. Bilinear multipliers and transference. Int. J. Math. Math. Sci. 2005(4) [2005], 545-
- Berkson, E., Blasco, O., Carro, M. and Gillespie, A. Discretization versus transference for bilinear operators. Banach Spaces and their Applications in Analysis, 11-30, De Gruyter
- [4] Blasco, O., Carro, M., and Gillespie, A. Bilinear Hilbert transform on measure spaces. J. Fourier Anal. and Appl. 11 [2005], 459-470.
- [5] Blasco, O., Villarroya, F., Transference of bilinear multipliers on Lorentz spaces. Illinois J. Math. 47(4),[2005], 1327-1343
- [6] Coifman R.R., Meyer, Y. Fourier Analysis of multilinear convolution, Calderón theorem and analysis of Lipschitz curves. Euclidean Harmonic Analysis (Proc. Sem. Univ. Maryland, College Univ., Md) Lecture Notes in Maths. 779, [1979], 104-122.
- [7] Fan, D., Sato, S., Transference of certain multilinear multipliers operators. J. Austral. Math. Soc. 70, [2001], pp. 37-55.
- [8] Gilbert, J, Nahmod, A., Bilinear operators with non-smooth symbols, J. Fourier Anal. Appl., **7**[2001], pp. 435-467.
- [9] Gilbert, J, Nahmod, A., Boundedness Bilinear operators with non-smooth symbols, Mat. Res. Letters, **7**[2000], pp. 767-778.
- [10] Grafakos, L., Kalton, N., Some remarks on multilinear maps and interpolation. Math. Annalen **319** [2001], pp. 151-180
- [11] Kenig, C. E., Stein, E.M., Multilinear estimates and fractional integration. Math. Res. Lett. 6 [1999], pp. 1-15.
- [12] Lacey M., Thiele C., L^p estimates on the bilinear Hilbert transform for 2 AnnalsMath. 146, [1997], pp. 693-724.
- [13] Lacey, M., Thiele, C., On Calderón's conjecture. Ann. Math. 149 2 [1999] pp. 475-496.
- [14] Stein, E. M., Weiss, G., Introduction to Fourier Analysis on Euclidean spaces. Princeton Univ. Press [1971].
- [15] Villarroya, P., La transformada de Hilbert bilineal. Doctoral Dissertation. Univ. of Valencia
- [16] Villarroya, P., Bilinear multipliers on Lorentz spaces. Czechoslovac Math. J. (to appear) [2007].

DEPARTMENT OF MATHEMATICS, UNIVERSITAT DE VALENCIA, BURJASSOT 46100 (VALENCIA)

E-mail address: oscar.blasco@uv.es