

NATURAL EXTENSIONS FOR α -ROSEN CONTINUED FRACTIONS

CORNELIS KRAAIKAMP, THOMAS A. SCHMIDT, AND IONICA SMEETS

ABSTRACT. We give natural extensions for the α -Rosen continued fractions of Dajani *et al* for a set of small α values by appropriately adding and deleting rectangles from the region of the natural extension for the standard Rosen fractions. It follows that the underlying maps have equal entropy.

1. INTRODUCTION

Dajani, Kraaikamp and Steiner [3] introduced a generalization of both the α -continued fractions of Nakada [8] and the Rosen continued fractions [13]. Using direct methods, similar to those of [2] for the classical Rosen fractions, they determine in [3] natural extensions for certain of their α -Rosen fractions.

Here we give a method that begins with the explicit region of the natural extension of a Rosen fraction (as already determined in [2]) and determines the regions for the natural extensions for various α -Rosen fractions. One advantage of this approach is that one easily sees that these various α -Rosen fraction maps determine isomorphic dynamical systems; in particular the associated one-dimensional maps have the same entropy. This can be compared to results on the entropy of Nakada's α -continued fractions, obtained by Nakada [8] and others [7], [6], [11].

We define all notation below, here we simply note that for each $q \in \mathbb{Z}, q \geq 3$ and all appropriate values of α , one can define an α -Rosen continued fraction map T_α on $[\lambda(\alpha-1), \lambda\alpha)$. When $\alpha = 1/2$, the map is the original map of Rosen; [3] showed that the domain of the natural extension is connected for all $\alpha \in [1/2, 1/\lambda]$. We determine the least α_0 such that for all $\alpha \in (\alpha_0, 1/\lambda]$ the natural extension is connected.

We prove the following.

Theorem 1. Fix $q \in \mathbb{Z}, q \geq 4$ and $\lambda = \lambda_q = 2 \cos \frac{\pi}{q}$.

(i.) Let

$$\alpha_0 := \begin{cases} \frac{\lambda^2 - 4 + \sqrt{(4 - \lambda^2)^2 + 4\lambda^2}}{2\lambda^2} & \text{if } q \text{ is even,} \\ \frac{\lambda - 2 + \sqrt{2\lambda^2 - 4\lambda + 4}}{\lambda^2} & \text{otherwise.} \end{cases}$$

Date: 11 March, 2009.

2000 Mathematics Subject Classification. 11K50, 37A45, 37A35.

Key words and phrases. Rosen fractions, natural extensions, entropy.

The second author was supported in part by Bezoekersbeurs **B 040.11.083** of the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO) and also thanks TU Delft for providing a stimulating atmosphere.

Then $(\alpha_0, 1/\lambda]$ is the largest interval containing $1/2$ for which each domain of the natural extension of T_α is connected.

(ii.) Furthermore, let

$$\omega_0 := \begin{cases} 1/\lambda & \text{if } q \text{ is even,} \\ \frac{\lambda - 2 + \sqrt{\lambda^2 - 4\lambda + 8}}{2\lambda} & \text{otherwise.} \end{cases}$$

Then the entropy of the α -Rosen map for each $\alpha \in [\alpha_0, \omega_0]$ is equal to the entropy of the standard Rosen map.

(Here, “the domain of the natural extension of T_α ” refers to the largest region on which the standard number theoretic map — defined below in Equation (3) — is bijective.) The value of the entropy of the standard Rosen map was given by H. Nakada [9], and is equal to

$$C \cdot \frac{(q-2)\pi^2}{2q},$$

where C is the normalizing constant (which depends on the parity of the index q) found by [2]. We recall the value of C in Equations (6) and (8).




FIGURE 1. Some values of α_0 and of ω_0 . Even index on left, odd index on right.

Our approach is to determine the domain of the natural extension using a standard “number theoretic” two-dimensional operator, here $\mathcal{T}_\alpha$ (depending on both λ and α). The domain for Rosen fractions, denoted here by $\Omega_{1/2}$, was obtained in [2]. For fixed λ and given α , we determine our domain by adding and deleting various regions to $\Omega_{1/2}$ by a process that we informally refer to as “quilting”. The regions are determined from appropriate orbits of the strips in $\Omega_{1/2}$ above intervals where the T_α –“digits” (see below) differ from the $T_{1/2}$ –digits. These strips are mapped by $\mathcal{T}_{1/2}$ to a region that must be deleted, their image under $\mathcal{T}_\alpha$ must be added; thereafter their further $\mathcal{T}_\alpha$ –orbits are deleted and added, respectively. Our approach succeeds without great difficulty exactly because in a small number of steps the orbits of the added regions agree with the orbits of the deleted regions — the infinitely many potential holes are quilted over by the added regions.

1.1. Outline of Paper. The following two subsections complete this Introduction by giving basic notation and then defining our building blocks, the basic deleted and added regions. In Section 2 we sketch a main argument of our approach — when the orbits of these basic regions agree after the same number of steps, entropy is preserved. We give an example of our techniques in Section 3 by re-establishing known results for certain classical Nakada α -fractions. In Sections 4 and 5 we give the proof of Theorem 1, in the even and odd index case, respectively. Finally, in

Section 6 we indicate how our results can be extended to show that in the odd index case, the entropy of T_α decreases when $\alpha > \omega_0$.

1.2. Basic Notation.

1.2.1. *One dimensional maps.* Let $q \in \mathbb{Z}, q \geq 3$ and $\lambda = \lambda_q = \cos \frac{\pi}{q}$. For $\alpha \in [0, \frac{1}{\lambda}]$, we let $\mathbb{I}_\alpha := [\lambda(\alpha - 1), \lambda\alpha]$ and define the map $T_\alpha : \mathbb{I}_\alpha \rightarrow \mathbb{I}_\alpha$ by

$$(1) \quad T_\alpha(x) := \left\lfloor \frac{1}{x} \right\rfloor - \lambda \left\lfloor \left\lfloor \frac{1}{\lambda x} \right\rfloor + 1 - \alpha \right\rfloor, \text{ for } x \neq 0; \quad T_\alpha(0) := 0.$$

For $x \in \mathbb{I}_\alpha$, put $d(x) := d_\alpha(x) = \lfloor \frac{1}{\lambda x} \rfloor + 1 - \alpha$ and $\varepsilon(x) := \text{sgn}(x)$. Furthermore, for $n \geq 1$ with $T_\alpha^{n-1}(x) \neq 0$ put

$$\varepsilon_n(x) = \varepsilon_n = \varepsilon(T_\alpha^{n-1}(x)) \text{ and } d_n(x) = d_n = d(T_\alpha^{n-1}(x)).$$

This yields the α -Rosen continued fraction of x :

$$x = \frac{\varepsilon_1}{d_1\lambda + \frac{\varepsilon_2}{d_2\lambda + \dots}} =: [\varepsilon_1 : d_1, \varepsilon_2 : d_2, \dots],$$

where $\varepsilon \in \{\pm 1\}$ and $d_i \in \mathbb{N}$. Fixing $\alpha = \frac{1}{2}$, results in the Rosen fractions. On the other hand, fixing $q = 3$ and considering general α , we have Nakada's α -expansions. These include the regular continued fractions, given by $\alpha = \frac{1}{\lambda} = 1$ and the nearest integer continued fractions, given by $\alpha = \frac{1}{2}$.

1.2.2. *Dynamics.* When studying the dynamics of these maps, the orbits of the interval endpoints of $\mathbb{I}_\alpha$ are of utmost importance. We define

$$\begin{aligned} l_0 &= (\alpha - 1)\lambda & \text{and } l_n &= T_\alpha^n(l_0), \text{ for } n \geq 1, \\ r_0 &= \alpha\lambda & \text{and } r_n &= T_\alpha^n(r_0), \text{ for } n \geq 1. \end{aligned}$$

The *cylinders* for the map T_α are

$$\Delta(\varepsilon : r) = \{x \mid \text{sgn}(x) = \varepsilon \text{ and } d_\alpha(x) = r\}.$$

Letting

$$(2) \quad \delta_r := \frac{1}{\lambda(r + \alpha)},$$

for r sufficiently large, we have *full* cylinders (each mapped surjectively by T_α onto $\mathbb{I}_\alpha$) of the form

$$\Delta(+1 : r) = (\delta_r, \delta_{r-1}] \text{ and } \Delta(-1 : r) = [-\delta_{r-1}, -\delta_r).$$

For future reference, we introduce notation for strips that fiber over cylinders: let

$$\mathcal{D}(\varepsilon : r) := \{(x, y) \mid x \in \Delta(\varepsilon : r)\}.$$

The standard number theoretic planar map associated to continued fractions gives here

$$(3) \quad \mathcal{T}_\alpha(x, y) = \left(T_\alpha(x), \frac{1}{r\lambda + \varepsilon y} \right);$$

it is easily checked that this map has invariant measure

$$d\mu = \frac{dx dy}{(1 + xy)^2}.$$

1.2.3. *Hecke groups.* Rosen defined his continued fractions in order to study aspects of the Hecke groups, $G_q \subset \mathrm{PSL}(2, \mathbb{R})$. With fixed index q as above, let

$$(4) \quad S = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix}, T = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \text{ and } U = \begin{bmatrix} \lambda & -1 \\ 1 & 0 \end{bmatrix}.$$

Then G_q is generated by any two of these, as $U = ST$. In fact, $U^q = \mathrm{Id}$. It turns out that the $T_{1/2}$ orbit of $\lambda/2$ is exactly the orbit of ∞ under powers of U : $r_j = U^{j+1}(\infty)$, where as usual we use the Möbius (or, fractional linear) action of 2×2 matrices on the reals (extended to include ∞ , as necessary). From [2] we know $U^n = \begin{bmatrix} B_{n+1} & -B_n \\ B_n & -B_{n-1} \end{bmatrix}$ where the sequence B_n is

$$(5) \quad B_0 = 0, \quad B_1 = 1, \quad B_n = \lambda B_{n-1} - B_{n-2}, \quad \text{for } n = 2, 3, \dots$$

1.3. Regions of changed digits; basic deletion and addition regions. Fix λ and choose some $\alpha \in [0, \frac{1}{\lambda}]$. For α in the range that we consider, our intention is to *solve* for the region of the natural extension of T_α , using the operator $\mathcal{T}_\alpha$. Of course, for many points $(x, y) \in \Omega_{1/2}$, we have $\mathcal{T}_\alpha(x, y) = \mathcal{T}_{1/2}(x, y)$. It is this last fact that we exploit, and find that it is important to understand $\mathcal{T}_\alpha(x, y)$ for all points where this value differs from $\mathcal{T}_{1/2}(x, y)$.

Definition 1. The *region of changed digits* is

$$C = \{ (x, y) \in \Omega_{1/2} \mid x \in \mathbb{I}_\alpha \cap \mathbb{I}_{1/2} \text{ and } d_\alpha(x) \neq d_{1/2}(x) \}.$$

The region C is a disjoint union of rectangles; in general, for each digit d , the subset of C determined by x whose digit has changed to $d_\alpha(x)$ consists of two connected components, one with x negative, the other with positive x values. See Figure 2 for a schematic representation of this in the classical $\lambda = 1$ setting.

We also identify a region of $\Omega_{1/2}$ that obviously cannot be part of the new natural extension, as its marginal projection lies outside of $\mathbb{I}_\alpha$.

Definition 2. The *basic deleted region* is the $\mathcal{T}_{1/2}$ -image of the region of changed digits:

$$D_0 := \mathcal{T}_{1/2}(C).$$

Lemma 1. For $\alpha < 1/2$, the *basic deleted region* is

$$D_0 = \{ (x, y) \in \Omega_{1/2} \mid x > \alpha\lambda \}.$$

Proof. The projection of C to the real line is a disjoint union of intervals. The boundaries of the cylinders for the map T_z are determined by the values $\frac{1}{\lambda(d+z)}$; thus, each of the connected components of this projection lies at the end of a cylinder for $T_{1/2}$. Since for fixed d the function $z \mapsto \frac{1}{\lambda(d+z)}$ is decreasing, the connected components lie at the right end of cylinders for negative x , and at the left end of cylinders of positive x values. But, $T_{1/2}$ itself is an increasing function for negative x and a decreasing function for positive x . Thus, this projection is exactly the $T_{1/2}$ -preimage of $(\alpha\lambda, \lambda/2]$. From this, it easily follows that $\mathcal{T}_{1/2}(C) = \{ (x, y) \in \Omega_{1/2} \mid x > \alpha\lambda \}$. $\square$

Remark 1. For $\alpha > 1/2$, one finds that $D_0 = \{ (x, y) \in \Omega_{1/2} \mid x < (\alpha - 1)\lambda \}$.

We are ready to define the region that in fact is the domain for the natural extension of T_α .

Definition 3. Let

$$\Omega_\alpha := \left(\Omega_{1/2} \bigcup \bigcup_{k=1}^{\infty} \mathcal{T}_\alpha^k(C) \right) \setminus \left(\bigcup_{k=0}^{\infty} \mathcal{T}_\alpha^k(D_0) \setminus \left(\bigcup_{k=0}^{\infty} \mathcal{T}_\alpha^k(D_0) \cap \bigcup_{k=1}^{\infty} \mathcal{T}_\alpha^k(C) \right) \right).$$

Thus, Ω_α is created by adding to $\Omega_{1/2}$ all images of the regions of changed digits and then deleting all images of the basic deleted region that are not contained in the added regions.

By analogy informed by Lemma 1, we define the *basic added region* to be

$$A_0 := \mathcal{T}_\alpha(C).$$

2. SUCCESSFUL QUILTING RESULTS IN EQUAL ENTROPY

We sketch here a main argument for our approach. Indeed, the results of this section show that we will have proven part (ii) of Theorem 1 once we show that for each $\alpha \in [\alpha_0, \omega_0]$ there exists some k satisfying the hypotheses of Proposition 1.

We mildly informally let μ denote the probability measure on Ω_α induced by $d\mu = \frac{dx dy}{(1+xy)^2}$. Let $\bar{\mathcal{B}}_\alpha$ denote the Borel σ -algebra of Ω_α .

Proposition 1. Fix $q \in \mathbb{Z}, q \geq 3$ and $\lambda = \lambda_q = 2 \cos \frac{\pi}{q}$, and choose some $\alpha \in [0, 1/\lambda]$. Let A_0 and D_0 be defined as above. Suppose that there is some natural number k such that

$$\mathcal{T}_\alpha^k(A_0) = \mathcal{T}_\alpha^k(D_0).$$

Then $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ is isomorphic to $(\mathcal{T}_{1/2}, \Omega_{1/2}, \bar{\mathcal{B}}_{1/2}, \mu)$.

Proof. By [2], $\mathcal{T}_{1/2}$ is bijective (up to μ -measure zero) on $\Omega_{1/2}$; one easily verifies that $\mathcal{T}_\alpha$ is injective onto $\bigcup_{j=0}^{k-1} \mathcal{T}_\alpha^j(D_0)$. Therefore, we can define $f : \Omega_{1/2} \rightarrow \Omega_\alpha$ by f being the identity off of $\bigcup_{j=0}^{k-1} \mathcal{T}_\alpha^j(D_0)$ and $f := \mathcal{T}_\alpha^{j+1} \circ \mathcal{T}_{1/2}^{-1} \circ \mathcal{T}_\alpha^{-j}$ on $\mathcal{T}_\alpha^j(D_0)$. Since each of $\mathcal{T}_\alpha$ and $\mathcal{T}_{1/2}$ preserves the measure μ , one easily shows that this is an isomorphism. $\square$

Proposition 2. With notation as above, let μ_α denote the marginal measure obtained by integrating μ on the fibers of $\pi : \Omega_\alpha \rightarrow \mathbb{I}_\alpha$. Further let $\mathcal{B}_\alpha$ denote the Borel σ -algebra of $\mathbb{I}_\alpha$. Then $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ is the natural extension of $(\mathcal{T}_\alpha, \mathbb{I}_\alpha, \mathcal{B}_\alpha, \mu_\alpha)$.

This is perhaps best proven using F. Schweiger's formalization of the ideas of [10], see in particular Section 22 of Schweiger's textbook [14]. For similar applications, see [3] or [5]. We leave this verification to the reader.

Corollary 1. Under the hypotheses of Proposition 1, the systems $(\mathcal{T}_\alpha, \mathbb{I}_\alpha, \mathcal{B}_\alpha, \mu_\alpha)$ have the same entropy.

Proof. It is known that a system and its natural extension have the same entropy [12]. Since the natural extensions here are all isomorphic, they certainly have the same entropy. $\square$

Remark 2. (1.) Rohlin [12] introduced the notion of natural extension explicitly in order to treat entropy.

- (2.) As we display in each of Sections 3, 4 and 5, the key to successful quilting is equality of orbits of r_0 and of l_0 after a fixed number of steps. Compare this with the discussion of [11] relating eventual equality of these orbits and behavior of the entropy in the classical case.

3. CLASSICAL CASE, $\lambda = 1$: NAKADA'S α -CONTINUED FRACTIONS

Aiming to maximize expository clarity, as an example we re-establish the form of the natural extension for Nakada's α -continued fraction [8] (thus with $\lambda = 1$) in the case of $\sqrt{2} - 1 \leq \alpha \leq 1/2$.

It is easily verified that $\alpha = 1/2$ gives the classical nearest integer continued fractions (NICF), whose natural extension was given by [8], $(\mathcal{T}_{1/2}, \Omega_{1/2}, \bar{\mathcal{B}}_{1/2}, \mu)$, with

$$\Omega_{1/2} = \left([-1/2, 0) \times [0, g^2) \right) \cup \left([0, 1/2) \times [0, g) \right),$$

where $g = \frac{-1+\sqrt{5}}{2}$ is the small golden number. (Here, the non-full cylinders of $T_{1/2}$ are $\Delta(+1 : 2) = [\delta_2, 1/2)$ and $\Delta(-1 : 2) = [-1/2, -\delta_2)$.)

Our goal is to re-establish the following result.

Theorem 2. *For all $\alpha \in [\sqrt{2} - 1, 1/2]$, the system $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ is isomorphic to $(\mathcal{T}_{1/2}, \Omega_{1/2}, \bar{\mathcal{B}}_{1/2}, \mu)$, where*

$$\begin{aligned} \Omega_\alpha := & [\alpha - 1, l_1) \times [0, g^2) \cup \\ & [l_1, r_1) \times \left([0, g^2) \cup [1/2, g) \right) \cup \\ & [r_1, \alpha) \times [0, g), \end{aligned}$$

and $\mathcal{B}_\alpha$ denotes the σ -algebra of μ -Borel subsets of Ω_α .

Furthermore, the entropy of T_α with respect to the marginal measure of the above system equals $\frac{1}{\ln(1+g)} \frac{\pi^2}{6}$.

Remark 3. The constancy of the entropy in this setting was first established by Moussa, Cassa and Marmi [7], the value at $\alpha = 1/2$ having been determined by [8]. We also note that Nakada and Natsui [11] explicitly show the isomorphism of these natural extensions (see their Appendix).

3.1. Explicit form of the basic addition region. Fix $(\mathcal{S}, \Omega) := (\mathcal{T}_{1/2}, \Omega_{1/2})$ and consider also a fixed $\alpha \in [\sqrt{2} - 1, 1/2)$. Recall that our intention is to solve for the region of the natural extension of T_α , by finding the $\mathcal{T} := \mathcal{T}_\alpha$ -orbits of basic added and deleted regions.

Remark 4. Other than $\mathcal{S}$ and Ω , all notation refers to values dependent upon α unless explicit dependence upon $1/2$ is indicated.

Note that with r fixed, δ_r as defined in Equation (2) is a decreasing function in α . Note also that for $\alpha \in (\sqrt{2} - 1, 1/2)$ one has $\delta_2(\alpha) < \alpha$.

Lemma 2. *The basic added region is given by*

$$A_0 = [\alpha - 1, -1/2) \times [0, g^2).$$




FIGURE 2. Regions of change of digit and basic deletion. Here each γ_m represents $\delta_m(1/2)$. To aid visualization, we use here $C_{\varepsilon r} = C \cap \Delta(\varepsilon : r)$.

Proof. Similarly to the proof of Lemma 1, we note that for $x < 0$ each connected component of the marginal projection of C lies at the left end of a T_α -cylinder and is sent by (the locally increasing function) T_α to $[\alpha - 1, -1/2)$. When $x > 0$, the component lies at the right end of its cylinder and is also sent by T_α to $[\alpha - 1, -1/2)$. One trivially checks that no other points of $\mathbb{I}_\alpha$ are sent to this subinterval.

We now discuss the y -coordinates of points in $\mathcal{T}(A_0)$. The fibers of $\Omega_{1/2}$ above cylinders where $x < 0$ are of the form $[0, g^2)$, while above cylinders with $x > 0$ the fibers are of the form $[0, g)$. Now, for $(x, y) \in \mathcal{D}(\varepsilon : r)$, the y -coordinate of $\mathcal{T}(x, y)$ is $1/(r + \varepsilon y)$. Thus, $\mathcal{T}$ sends $\mathcal{D}(-1 : r)$ and $\mathcal{D}(+1 : r)$ to horizontal strips whose y -values lie in $[\frac{1}{r+g}, \frac{1}{r-g^2}]$. But, $g^2 = 1 - g$ and thus this image lies directly above that of the horizontal strip given by $\mathcal{T}$ applied to $\mathcal{D}(-1 : r+1)$ and $\mathcal{D}(+1 : r+1)$.

Now, the greatest y -value of $\mathcal{T}(C)$ comes from the intersection of the projection of C with $\mathcal{D}(-1 : 3)$. Thus, $A_0 = [\alpha - 1, -1/2) \times [0, 1/(3 - g^2))$. Since $1/(3 - g^2) = g^2$, the result follows. $\square$

3.2. Quilting. We now show that the $\mathcal{T}$ -orbits of the added regions eventually match the orbits of the deleted regions, and thus $\mathcal{T}$ is bijective (modulo μ -measure zero) on Ω_α .

Lemma 3. *The following equality holds:*

$$l_2 = r_2.$$

Furthermore, there is a $d \in \mathbb{N}$ such that

$$l_1 \in \Delta(-1 : d) \text{ and } r_1 \in \Delta(1 : d - 1).$$

Proof. We have that $l_0 \in \Delta(-1 : 2)$ and $r_0 \in \Delta(+1 : 2)$, giving

$$l_1 = \frac{2\alpha - 1}{1 - \alpha} \text{ and } r_1 = \frac{1 - 2\alpha}{\alpha}.$$

Therefore,

$$l_2 = \frac{1 - \alpha}{1 - 2\alpha} - d \text{ and } r_2 = \frac{\alpha}{1 - 2\alpha} - d',$$




FIGURE 3. Representative region Ω_α for $\alpha \in (\sqrt{2} - 1, 1/2)$; $q = 3$.

with d, d' the appropriate T_α -digits. Now, $l_2 - r_2 = 1 + d' - d$ and is the difference of two elements in $\mathbb{I}_\alpha$, an interval of length one. We thus conclude both that $d' = d - 1$ and $l_2 = r_2$. $\square$

The orbit of the basic addition region, A_0 , is quickly synchronized with that of the basic deletion region, D_0 . Recall that $\mathcal{D}(\varepsilon : r)$ fibers over $\Delta(\varepsilon : r)$.

Lemma 4. *We have*

$$\mathcal{T}^2(A_0) = \mathcal{T}^2(D_0).$$

Proof. Let $A_1 := \mathcal{T}(A_0)$; since $A_0 \subset \mathcal{D}(-1 : 2)$, an elementary calculation shows that this is

$$A_1 = [l_1, 0) \times [1/2, g).$$

Similarly, defining $D_1 := \mathcal{T}(D_0)$, one has $D_1 \subset \mathcal{D}(1 : 2)$, and finds

$$D_1 = [0, r_1) \times [g^2, 1/2).$$

With d as in Lemma 3, let

$$A'_1 := A_1 \cap \mathcal{D}(-1 : d) \quad \text{and} \quad D'_1 := D_1 \cap \mathcal{D}(+1 : d - 1).$$

By elementary calculation, Lemma 3, and an application of the identity $g^2 = 1 - g$, one finds that

$$\begin{aligned} \mathcal{T}(A'_1) &= \mathcal{T}(D'_1) \\ &= [l_2, \alpha) \times \left[\frac{2}{2d-1}, d-g \right). \end{aligned}$$

Each of $A_1 \setminus A'_1$ and $D_1 \setminus D'_1$ projects to the union of full cylinders:

$$A_1 \setminus A'_1 = \bigcup_{m=d+1}^{\infty} \Delta(-1 : -m) \times [1/2, g) \quad \text{and} \quad D_1 \setminus D'_1 = \bigcup_{m=d}^{\infty} \Delta(+1 : m) \times [1/2, g^2).$$

Since $\mathcal{T}(\Delta(-1 : -m) \times [1/2, g)) = \mathcal{T}(\Delta(+1 : m - 1) \times [1/2, g^2))$, we conclude that $\mathcal{T}(A_1) = \mathcal{T}(D_1)$, and the result follows. $\square$

3.3. Isomorphic systems. In this subsection, we complete the proof of Theorem 2.

Corollary 2. *We have*

$$\begin{aligned} \Omega_\alpha = & [\alpha - 1, l_1) \times [0, g^2) \cup \\ & [l_1, r_1) \times \left([0, g^2) \cup [1/2, g) \right) \cup \\ & [r_1, \alpha) \times [0, g). \end{aligned}$$

Furthermore, the μ -area of Ω_α equals that of $\Omega_{1/2}$.

Proof. From the above,

$$\Omega_\alpha = \left(\Omega \cup A_0 \cup A_1 \right) \setminus \left(D_0 \cup D_1 \right).$$

The explicit shape of Ω_α follows.

By definition of A_0 and by Lemma 1, A_0 and D_0 are the images of the union of the change of digit regions under $\mathcal{T}$ and $\mathcal{S}$, respectively. Since both $\mathcal{S}$ and $\mathcal{T}$ are μ -measure preserving, thus $\mu(D_0) = \mu(A_0)$. Again since $\mathcal{T}$ preserves measure, we also have $\mu(D_1) = \mu(A_1)$. $\square$

Of course, the equality of the areas is already implied by the arguments of Section 2. Indeed, since the entropy of the NICF is known, those arguments can be easily adapted to complete the proof of the Theorem 2.

4. EVEN q ; $\alpha \in (\alpha_0, 1/2]$

The natural extension for the Rosen fractions was determined in [2]. The exact form of the domain depends on the parity of the index q .

4.1. Natural extensions for Rosen fractions. Let $q = 2p$ for $p \in \mathbb{N}$ and $p \geq 2$. The domain, see Figure 4, given by [2] is

$$\Omega = \bigcup_{j=1}^p J_j \times K_j,$$


here J_j is defined as follows: Let $\varphi_j = T_{1/2}^j(-\frac{\lambda}{2})$, then $J_j = [\varphi_{j-1}, \varphi_j)$ for $j \in \{1, 2, \dots, p-1\}$ and $J_p = [0, \frac{\lambda}{2})$. Further, $K_j = [0, L_j]$ for $j \in \{1, 2, \dots, p-1\}$ and $K_p = [0, 1]$, where the values of the L_j are given by the following relations:

$$\begin{aligned} L_1 &= 1/(\lambda + 1), \\ L_j &= 1/(\lambda - L_{j-1}) \quad \text{for } j \in \{2, \dots, p-1\}. \end{aligned}$$

The normalizing constant C such that $Cd\mu$ gives a probability measure on Ω_α is

$$(6) \quad C = \frac{1}{\ln[(1 + \cos \pi/q)/\sin \pi/q]},$$

see Lemma 3.2 of [2].

FIGURE 4. The natural extension for $q = 8$ and $\alpha = \frac{1}{2}$.

In this case, using that U is of order q (in the projective group) for the B_j as defined in (5), we have that

$$(7) \quad B_{p-1} = B_{p+1} = \frac{\lambda}{2} B_p \quad \text{and} \quad B_{p-2} = \left(\frac{\lambda^2}{2} - 1 \right) B_p.$$

The key to understanding the system for $\mathcal{T}_\alpha$ is that for all $q = 2p$, one has that $l_p = r_p$ for all of our α . In proving this, it is convenient to use the fact that the orbits of r_0 and $-r_0$ coincide after one application of T_α .

Lemma 5. *For any $\alpha > \alpha_0$, the T_α -expansion of both l_0 and $-r_0$ starts as $[(-1 : 1)^{p-1}, \dots]$.*

Proof. A point that lies to the left of the appropriate $p - 2$ nd pre-image of $-\delta_1$ has ones for its first $p - 1$ digits. This pre-image is given by $(S^{-1}T)^{-p+2}(-\delta_1)$, where we are using the matrices defined in Equation (4).

But,

$$(S^{-1}T)^{-p+2} = (S^{-1}U^{-1}S)^{-p+2}$$

$$= S^{-1}U^{p-2}S$$

$$= \begin{bmatrix} B_{p-1} - \lambda B_{p-2} & -B_{p-2} \\ B_{p-2} & B_{p-1} \end{bmatrix}.$$

Now, the relations of Equation (7) give

$$(S^{-1}T)^{-p+2} = \frac{B_p}{2} \begin{bmatrix} -\lambda^3 + 3\lambda & -\lambda^2 + 2 \\ \lambda^2 - 2 & \lambda \end{bmatrix}.$$

Thus the $p - 2$ nd pre-image of $-\delta_1$ is given by

$$T_\alpha^{-p+2}(-\delta_1) = \frac{(\lambda^3 - 3\lambda)\delta_1 - \lambda^2 + 2}{(-\lambda^2 + 2)\delta_1 + \lambda} = -\frac{\alpha(\lambda^3 - 2\lambda) + \lambda}{\alpha\lambda^2 + 2}.$$

In particular, $-r_0$ starts with $p - 1$ digits one if

$$-r_0 = -\alpha\lambda < -\frac{\alpha(\lambda^3 - 2\lambda) + \lambda}{\alpha\lambda^2 + 2}.$$

Rewriting this inequality yields that it holds whenever

$$\alpha > \alpha_0 = \frac{\lambda^2 - 4 + \sqrt{(4 - \lambda^2)^2 + 4\lambda^2}}{2\lambda^2}.$$

Finally, if $\alpha > \alpha_0$, then it immediately follows that l_0 also starts with $p - 1$ ones since $l_0 = (\alpha - 1)\lambda < -\alpha\lambda = -r_0$. $\square$

Lemma 6. For $\alpha \in (\alpha_0, 1/2)$, $r_p = l_p$. Furthermore, there is $d \in \mathbb{N}$ such that

$$l_{p-1} \in \Delta(-1 : d) \text{ and } r_{p-1} \in \Delta(1 : d - 1) .$$

Proof. Using the previous lemma, we find

$$\begin{aligned} l_{p-1} &= T_\alpha^{p-1}(l_0) = (S^{-1}T)^{p-1}((\alpha - 1)\lambda) \\ &= \begin{bmatrix} -\lambda & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \lambda & \lambda^2 - 2 \\ -\lambda^2 + 2 & -\lambda^3 + 3\lambda \end{bmatrix} ((\alpha - 1)\lambda) \\ &= \frac{(1 - 2\alpha)}{\alpha\lambda^2 - 2} , \end{aligned}$$

and similarly, since $T_\alpha(r_0) = T_\alpha(-r_0)$,

$$r_{p-1} = T_\alpha^{p-1}(-r_0) = \frac{(2\alpha - 1)\lambda}{(1 - \alpha)\lambda^2 - 2} .$$

Therefore,

$$\left| \frac{1}{l_{p-1}} \right| - \left| \frac{1}{r_{p-1}} \right| = \lambda .$$

But, then $T_\alpha(l_{p-1}) - T_\alpha(r_{p-1})$ is an integer multiple of λ . However, this is the difference of two elements of $\mathbb{I}_\alpha$, and thus this multiple must be zero. We conclude that $r_p = l_p$. That the digits of these points is as claimed follows as in the classical case. $\square$




FIGURE 5. The added and deleted rectangles for $\alpha = 0.48$ in the natural extensions for $q = 8$. Here A_i, D_i denote $T^i(A_0), T^i(D_0)$, respectively.

To describe the initial orbits of A_0 and D_0 , we use the following sequence.

$$H_1 = \frac{1}{\lambda} \text{ and } H_i = \frac{1}{\lambda - H_{i-1}} \text{ for } i \geq 2 .$$

Lemma 7. For $\alpha \in (\alpha_0, 1/2)$, $T^p(A_0) = T^p(D_0)$.

Proof. One easily checks that the basic added region and basic deleted regions are

$$A_0 = [l_0, -\lambda/2) \times [0, L_1) \text{ and } D_0 = (r_0, \lambda/2] \times [0, R) .$$

Since $-\lambda/2 < -r_0$, from Lemma 5 we find that all $x \in [l_0, -\lambda/2)$ share the same first $p-1$ of their T_α -digits, and in fact that $T_\alpha^{p-1}(-\lambda/2) = T_{1/2}^{p-1}(-\lambda/2)$. But, $T_{1/2}^{p-1}(-\lambda/2) = 0$. Thus, recalling that $1 = 1/(\lambda - L_{p-1})$, we find that

$$\mathcal{T}^{p-1}(A_0) = [l_{p-1}, 0) \times [H_{p-1}, 1).$$

Paying attention to sign and orientation, one finds that $\mathcal{T}(A_0) = [\varphi_1, r_1) \times [L_1, H_1)$, and thus

$$\mathcal{T}^{p-1}(D_0) = [0, r_{p-1}) \times [L_{p-1}, H_{p-1}).$$

Analogously to the classical case, with d from Lemma 6, we let

$$A'_{p-1} := \mathcal{T}^{p-1}(A_0) \cap \mathcal{D}(-1 : d),$$

and

$$D'_{p-1} := \mathcal{T}^{p-1}(D_0) \cap \mathcal{D}(+1 : d-1).$$

Arguing analogously to the classical case, we find that the $\mathcal{T}$ images of A'_{p-1} and D'_{p-1} agree, and then that the various layers from each of $\mathcal{T}^{p-1}(A_0) \setminus A'_{p-1}$ and of $\mathcal{T}^{p-1}(D_0) \setminus D'_{p-1}$ also agree. The result follows. $\square$




FIGURE 6. Change of topology at $\alpha = \alpha_0$: Simulations of the natural extension for $q = 8$ with on the left $\alpha = \alpha_1 - 0.001$ and on the right $\alpha = \alpha_1 + 0.001$.

Lemma 8. *The region Ω_{α_0} is not connected.*

Proof. The proof of Lemma 5 shows that if $\alpha = \alpha_0$, then $r_{p-2} = -\delta_1$. But then

$$\mathcal{T}^{p-1}(D_0) = [0, \alpha\lambda) \times [L_{p-1}, H_{p-1}).$$

Thus, since the $\mathcal{T}$ -orbit of A_0 can never fill in this deleted strip, we see that Ω_{α_0} is indeed disconnected. $\square$

The arguments of Section 2 then finish the proof of Theorem 1 in this case of even index q .

5. ODD q ; $\alpha \in (\alpha_0, 1/2]$

Let $q = 2h + 3$, for $h \geq 1$.

5.1. Natural extensions for Rosen fractions. We recycle notation, now using φ_j and L_j as follows (all necessary calculations are in [2]):

$$\varphi_0 = -\frac{\lambda}{2}, \quad \text{and} \quad \varphi_j = T_{1/2}^j \left(-\frac{\lambda}{2} \right), \quad 0 \leq j \leq 2h+1.$$

We recall that

$$-\frac{\lambda}{2} \leq \varphi_j < -\frac{2}{3\lambda} \quad \text{for} \quad j \in \{0, 1, \dots, h-1\} \cup \{h+1, \dots, 2h\},$$

$$-\frac{2}{3\lambda} < \varphi_h < -\frac{2}{5\lambda}, \quad \text{and}$$

$$\varphi_{2h+1} = 0.$$

Also we put, with R the positive root of $R^2 + (2 - \lambda)R - 1 = 0$,

$$L_1 = \frac{1}{2\lambda - L_{2h}},$$

$$L_2 = \frac{1}{2\lambda - L_{2h+1}},$$

$$L_j = \frac{1}{\lambda - L_{j-2}}, \quad 2 < j \leq 2h+2.$$

The domain, see Figure 7, given in [2] is $\Omega = \bigcup_{j=1}^{2h+2} J_j \times K_j$, where

$$J_{2k} = [\varphi_{h+k}, \varphi_k), \quad \text{for } k \in \{1, \dots, h\},$$

$$J_{2k+1} = [\varphi_k, \varphi_{h+k+1}), \quad \text{for } k \in \{0, 1, \dots, h\},$$

and $J_{2h+2} = [0, \frac{\lambda}{2})$. Let $K_j = [0, L_j]$ for $j \in \{1, \dots, 2h+1\}$ and let $K_{2h+2} = [0, R]$.

Here, the normalizing constant C such that $Cd\mu$ gives a probability measure on Ω_α is

$$(8) \quad C = \frac{1}{\ln(1 + R)},$$

see Lemma 3.4 of [2].




FIGURE 7. The natural extension for $q = 7$ and $\alpha = \frac{1}{2}$.

Similarly to the even case, for each of our α , the basic added region and basic deleted regions are given by

$$A_0 = [l_0, -\lambda/2) \times [0, L_1), \quad D_0 = (r_0, \lambda/2] \times [0, R).$$

In this case, using that U is of order $2h+3$ we have that

$$(9) \quad B_{h+1} = B_{h+2}, \quad B_h = (\lambda - 1)B_{h+1} \quad \text{and} \quad B_{h-1} = (\lambda^2 - \lambda - 1)B_{h+1}.$$

In this section we prove that for $q = 2h + 3$ one has that $l_{2h+2} = r_{2h+2}$ for all $\alpha \in (\alpha_0, \frac{1}{2})$ and consequently that the added blocks coincide with the deleted blocks after $2h + 2$ steps. Here also, we use the fact that the orbits of r_0 and $-r_0$ coincide after one application of T_α .

Lemma 9. *For any $\alpha > \alpha_0$, the T_α -expansion of both l_0 and $-r_0$ starts as $[(-1 : 1)^h, (-1, 2), (-1, 1)^h, \dots]$.*

Proof. We recall that $-\delta_1 < \varphi_h < -\delta_2$. We certainly have that $l_0 < \varphi_0 < r_0$; it immediately follows that $T_\alpha^h(l_0) < \varphi_h < -\delta_2$ and that $T_\alpha^h(-r_0) > \varphi_h > -\delta_1$. To prove the lemma it suffices to show that $T_\alpha^h(l_0) > -\delta_1$, $T_\alpha^h(r_0) < -\delta_2$ and $T_\alpha^{2h}(-r_0) < \delta_1$. We only show the last of these inequalities, because one can easily check that it imposes the strongest restriction on the value of α .

Assuming the first two conditions are met we find that

$$T_\alpha^{2h}(-r_0) = (S^{-1}T)^{h-1}S^{-2}T(S^{-1}T)^h(-r_0).$$

Again,

$$\begin{aligned} (S^{-1}T)^h &= (S^{-1}U^{-1}S)^h \\ &= S^{-1}U^{-h}S \\ &= \begin{bmatrix} -B_{h+2} & -B_{h+3} \\ B_{h+3} & B_{h+4} \end{bmatrix}. \end{aligned}$$

It easily follows that

$$(S^{-1}T)^{h-1} = \begin{bmatrix} -B_{h+3} & -B_{h+4} \\ B_{h+4} & B_{h+5} \end{bmatrix}.$$

We find that

$$\begin{aligned} (S^{-1}T)^{h-1}S^{-2}T(S^{-1}T)^h &= \begin{bmatrix} -B_{h+3} & -B_{h+4} \\ B_{h+4} & B_{h+5} \end{bmatrix} \begin{bmatrix} -2\lambda & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} -B_{h+2} & -B_{h+3} \\ B_{h+3} & B_{h+4} \end{bmatrix} \\ &= \begin{bmatrix} -B_{h+2}^2 - B_{h+1}B_{h+3} & -2B_{h+2}B_{h+3} \\ B_{h+2}B_{h+3} + B_{h+1}B_{h+4} & B_{h+2}B_{h+4} + B_{h+3}^2 \end{bmatrix}. \end{aligned}$$

Using the relations (9) yields that

$$T_\alpha^{2h}(-r_0) = B_{h+1}^2 \begin{bmatrix} \lambda & 2(\lambda - 1) \\ 2 - \lambda^2 & (3 - 2\lambda)\lambda \end{bmatrix}.$$

Substituting $r_0 = \alpha\lambda$ gives

$$T_\alpha^{2h}(-r_0) = \frac{-\alpha\lambda^2 + 2(\lambda - 1)}{\alpha\lambda^3 - 2\alpha\lambda + (3 - 2\lambda)\lambda}.$$

A calculation shows that $\frac{-\alpha\lambda^2 + 2(\lambda - 1)}{\alpha\lambda^3 - 2\alpha\lambda + (3 - 2\lambda)\lambda} < -\delta_1$ whenever

$$\alpha > \alpha_0 = \frac{\lambda - 2 + \sqrt{2\lambda^2 - 4\lambda + 4}}{\lambda^2}.$$

□

Lemma 10. *For $\alpha \in (\alpha_0, 1/2)$ one has that $\mathcal{T}^{2h+2}(A_0) = \mathcal{T}^{2h+2}(D_0)$.*

Proof. Thanks to Lemma 9 it is easy to compute l_{2h+1} and r_{2h+1} . We have

$$\begin{aligned} l_{2h+1} &= T^{2h+1}(l_0) = \begin{bmatrix} -2 & -\lambda \\ \lambda & 2\lambda - 2 \end{bmatrix} ((\alpha - 1)\lambda) \\ &= \frac{-2(\alpha - 1)\lambda - \lambda}{(\alpha - 1)\lambda^2 + 2\lambda - 2}, \\ r_{2h+1} &= T^{2h+1}(r_0) = \begin{bmatrix} -2 & -\lambda \\ \lambda & 2\lambda - 2 \end{bmatrix} (-\alpha\lambda) \\ &= \frac{-2\alpha\lambda - \lambda}{\alpha\lambda^2 + 2\lambda - 2}. \end{aligned}$$

We thus find that

$$\left| \frac{1}{r_{2h+1}} \right| - \left| \frac{1}{l_{2h+1}} \right| = \frac{-2\alpha\lambda^2 + \lambda^2}{(2\alpha - 1)\lambda} = -\lambda.$$

Arguments completely analogous to the even case now give that that $l_{2h+2} = r_{2h+2}$ and in fact that $T^{2h+2}(A_0) = T^{2h+2}(D_0)$. $\square$




FIGURE 8. Change of topology at $\alpha = \alpha_0$: Simulations of the natural extension for $q = 9$; on left $\alpha = \alpha_1 - 0.001$, on right $\alpha = \alpha_0 + 0.001$.

Arguing as in Lemma 8 and completing with the arguments of Section 2 finish the proof of Theorem 1 in this case of odd index q .

6. LARGE α , BY WAY OF DAJANI *et al*

We finish the proof of Theorem 1 by appropriately interpreting results of [3].

6.1. Successful quilting for $\alpha \in (1/2, \omega_0]$. As already stated in our introduction, [3] shows that the domain Ω_α is connected for all q and all $\alpha \in (1/2, 1/\lambda]$. Thus with the above, part (i) of Theorem 1 follows.

For even q and $\alpha \in (1/2, 1/\lambda]$, Theorem 2.2 of [3] shows that $r_p = l_p$ and that the digits of r_{p-1} and l_{p-1} agree up to sign and a shift (of the correct sign) by one. From this, just as above, one can in fact show that $T^p(A_0) = T^p(D_0)$ in these cases as well.

For odd q and $\alpha \in (1/2, \omega_0)$, Theorem 2.9 of [3] shows that $r_{2h+2} = l_{2h+2}$ and that the digits of their orbit predecessors agree up to sign and a shift by one. Thus, here one can show that $\mathcal{T}^{2h+2}(A_0) = \mathcal{T}^{2h+2}(D_0)$.

Using the results of Section 2, we thus have that part (ii) of Theorem 1 holds.

6.2. Nearly successful quilting and unequal entropy. We now show that the entropy of T_α is *not* equal to that of $T_{1/2}$ for $\alpha_0 > \omega_0$ in the case of odd q . Indeed, for these values, the results of [3] show that although the natural extensions remain connected, the conditions for successful quilting are not fully satisfied.

Lemma 11. *With notation as above, suppose that there are distinct natural numbers k, k' such that*

$$\mathcal{T}_\alpha^k(A_0) = \mathcal{T}_\alpha^{k'}(D_0).$$

Then the entropy of $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ differs from that of $(\mathcal{T}_{1/2}, \Omega_{1/2}, \bar{\mathcal{B}}_{1/2}, \mu)$.

Proof. (Sketch) If $k < k'$, then we can produce a new system $(\Omega'_\alpha, T'_\alpha, \bar{\mathcal{B}}'_\alpha, \mu)$ by inducing past one “copy” of A_0 . This system can be shown to be isomorphic to $(\mathcal{T}_{1/2}, \Omega_{1/2}, \bar{\mathcal{B}}_{1/2}, \mu)$. But, by the Abramov formula [1], the induced system has entropy differing from that of the full system by a multiplicative factor equal to the measure of A_0 . Thus, the entropy of $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ is less than that of the system of index $1/2$.

Similarly, if $k > k'$, we form a new system by inducing past a copy of D_0 in the index $1/2$ system. This allows us to conclude that the entropy of $(\mathcal{T}_\alpha, \Omega_\alpha, \bar{\mathcal{B}}_\alpha, \mu)$ is greater than that of the system of index $1/2$. $\square$

The following is part of Theorem 2.9 of [3].

Lemma 12. (Dajani *et al* [3]) For odd q and $\alpha \in (\omega_0, 1/\lambda]$ there are distinct natural numbers k, k' such that $l_k = r_{k'}$ and that the T_α -digits of $l_{k-1} = r_{k'-1}$ differ by one.

Corollary 3. *For odd q and $\alpha \in (\omega_0, 1/\lambda]$, the maps T_α and $T_{1/2}$ have distinct entropy values.*

Proof. From Lemma 12, one can show that $\mathcal{T}_\alpha^k(A_0) = \mathcal{T}_\alpha^{k'}(D_0)$. Lemma 11 then applies. $\square$

In fact, Theorem 2.9 of [3] shows that our $k' = k + 1$; thus, from the proof of Lemma 11, we see that the entropy of T_α for $\alpha \in (\omega_0, 1/\lambda]$ is decreasing, confirming the results for the case $q = 3$ of [8], see also [6].

REFERENCES

- [1] L. M. Abramov, *The entropy of a derived automorphism. (Russian)*, Dokl. Akad. Nauk SSSR 128 (1959), 647–650. [Amer. Math. Soc. Transl. Series 2, **49** (1966), 162–166.]
- [2] R. Burton, C. Kraaikamp, and T.A. Schmidt, *Natural extensions for the Rosen fractions*, TAMS 352 (2000), 1277–1298.
- [3] K. Dajani, C. Kraaikamp and W. Steiner, *Metrical theory for α -Rosen fractions*, to appear in J. Eur. Math. Soc. (JEMS). Available at <http://arxiv.org/abs/math/0702516>, 2008.
- [4] C. Kraaikamp, *A new class of continued fraction expansions*, Acta Arith. 57 (1991), no. 1, 1–39.
- [5] C. Kraaikamp, H. Nakada and T.A. Schmidt, *Metrical theory for α -Rosen fractions*, to appear in Acta Arithmetica. Available at <http://arxiv.org/abs/math/0812.0548>, 2008.

- [6] L. Luzzi and S. Marmi, *On the entropy of Japanese continued fractions*, Discrete Contin. Dyn. Syst. **20** (2008), no. 3, 673–711.
- [7] P. Moussa, A. Cassa, and S. Marmi, *Continued fractions and Bruno functions*, J. Comput. Appl. Math. **105** (1995), no. 3, 403–415.
- [8] H. Nakada, *Metrical theory for a class of continued fraction transformations and their natural extensions*, Tokyo J. Math. **4** (1981), 399–426.
- [9] ———, *On the Lenstra constant associated to the Rosen continued fractions*, to appear in J. Eur. Math. Soc. (JEMS). Available at <http://arxiv.org/abs/math/0705.3756>, 2007.
- [10] H. Nakada, S. Ito, and S. Tanaka, *On the invariant measure for the transformations associated with some real continued-fractions*, Keio Engrg. Rep. **30** (1977), no. 13, 159–175.
- [11] H. Nakada, and R. Natsui, *The non-monotonicity of the entropy of α -continued fraction transformations*, Nonlinearity **21** (2008), no. 6, 1207–1225.
- [12] V.A. Rohlin, *Exact endomorphisms of Lebesgue spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **25** (1961), 499–530. Amer. Math. Soc. Transl. Series 2, **39** (1964), 1–36.
- [13] D. Rosen, *A class of continued fractions associated with certain properly discontinuous groups*, Duke Math. J. **21** (1954), 549–563.
- [14] F. Schweiger, *Ergodic theory of fibred systems and metric number theory*. Oxford: Clarendon Press, 1995.

CORNELIS KRAAIKAMP, EWI, DELFT UNIVERSITY OF TECHNOLOGY, MEKELWEG 4, 2628 CD DELFT, THE NETHERLANDS

E-mail address: `c.kraaikamp@tudelft.nl`

THOMAS A. SCHMIDT, DEPARTMENT OF MATHEMATICS, OREGON STATE UNIVERSITY, CORVALLIS, OR 97331, USA

E-mail address: `toms@math.orst.edu`

IONICA SMEETS, MATHEMATICAL INSTITUTE, LEIDEN UNIVERSITY, NIELS BOHRWEG 1, 2333 CA LEIDEN, THE NETHERLANDS

E-mail address: `ionica.smeets@gmail.com`