Quantum correlations and dynamics from classical random fields valued in complex Hilbert spaces

Andrei

School of Mathematics and Systems Engineering University of Växjö, S-35195, Sweden

October 29, 2018

Abstract

One of the crucial differences between mathematical models of classical and quantum mechanics is the use of the tensor product of the state spaces of subsystems as the state space of the corresponding composite system. (To describe an ensemble of classical composite systems one uses random variables taking values in the Cartesian product of the state spaces of subsystems.) We show that, nevertheless, it is possible to establish a natural correspondence between the classical and quantum probabilistic descriptions of composite systems. Quantum averages for composite systems (including entangled) can be represented as averages with respect to classical random fields. It is essentially what Albert Einstein was dreamed of. Quantum mechanics is represented as classical statistical mechanics with infinite-dimensional phase space. While the mathematical construction is completely rigorous, its physical interpretation is a complicated problem (which will not be discussed in this paper).

1 Introduction

Nowadays it is commonly accepted that the use of the *tensor product of* the state spaces of subsystems as the state space of the corresponding composite system is one of the main distinguishing features of QM. It is especially important in quantum information theory where systems in entangled states play an fundamental role.¹ In this paper we do not discuss extremely complicated problems related to interpretations of quantum mechanics. We proceed in the framework of mathematical physics.

There are known two models [8] for computations of averages for ensembles of composite systems: a) classical probability model (due to Kolmogorov [9]) based on integrals, b) quantum probability model (due to von Neumann [10], see, e.g., [11], [12], [13] for the modern treatment of the problem) based on traces of self-adjoint operators. We show that, in

¹While it is sufficiently well studied mathematically, entanglement is still quite mysterious physically. Its widely used interpretation as the evidence of "nonlocal correlations" can not be considered as completely satisfactory, see, e.g., [1]- [5] and especially [6]-[8] for recent debates. Therefore clarification of the structure of its mathematical description may have important consequences. This paper is a step in this direction.

spite of a rather common opinion, quantum correlations for observables on subsystems of a composite systems can be represented as correlations with respect to a classical (Gaussian) random fields. Moreover, dynamics of quantum correlations induced by Schrödinger's equation can be reduced to dynamics of correlations for a classical "prequantum" stochastic processes.

For non-composite systems theory has been developed in a series of author's papers [14]- [20]. It is known under the name Prequantum Classical Statistical Field Theory (PCSFT). In this theory ensembles of quantum particles are represented by classical random fields, probability distributions on a Hilbert space. We remark that appealing to classical random fields is rather common in various attempts to create a kind of classical statistical mechanics which reproduces predictions of QM. We can mention stochastic electrodynamics, e.g., [21], [25], or the semiclassical model, e.g., [26]–[28]. Bohmian mechanics also contains a kind of classical field, the pilot wave. However, in this model randomness is coupled to particles and not to fields. The same can be said about Nelson's stochastic QM [29] and its generalization due to Mark Davidson [30], [31] as well as the recent prequantum model of 't Hooft [32], [33], see also Thomas Elze [35]. Physically the present paper belongs to the domain of "quantum mechanics as emergent phenomenon", cf. mentioned papers of 't Hooft and Elze as well as models which were recently created Rusov et al. [36] and Kisil [37].

However, as was already pointed, we prefer not to go in the debate on a physical meaning of the proposed mathematical construction. Our aim was to unify two mathematical descriptions of averages, classical and quantum. This aim was approached via representation of quantum correlations by Gaussian integrals over the Hilbert space $H_1 \times H_2$, where H_{i} = 1, 2, are the (Hilbert) state spaces of the subsystems. On the other hand, we could not totally escape the interpretation problem. The main message from our mathematical construction is that a quantum pure state of a composite system should be considered not as a "state vector" belonging to the tensor product $H_1 \otimes H_2$, but as non-diagonal block of an operator acting in the Cartesian product $H_1 \times H_2$ see (35). This operator, say D, is the covariance operator of the prequantum classical random field. We remind that consideration of a *density operator as the* covariance operator of the corresponding prequantum random process is the crucial point of PCSFT, see [14]- [20]. In this paper we extend this approach to composite systems.

We point out to another approach providing a possibility to represent quantum averages by operating only with classical probability distributions, namely, quantum tomographic approach, see Manko et al. [38]-[42].

Results of these paper were shortly announced in [43], where they were presented on the physical level of rigorousness. Unfortunately, such a formal presentation induced (to be totally honest) a mistake – unfortunately, in the basic equality (22) coupling the quantum correlation with the nontrivial term of the classical field correlation. In this paper we proceed in the rigorous mathematical framework, the basic formula and its consequences were corrected. Formulas of the present theory are more complicated than of one given in [43], but they are mathematically correct.

2 Statistical models of classical and quantum mechanics

Everywhere in this papers Hilbert spaces are separable. Let H_j , j = 1, 2, be (real or complex) Hilbert spaces. We denote the space of bounded linear operators from H_1 to H_2 by the symbol $\mathcal{L}(H_1, H_2)$. Let H be a (real or complex) Hilbert space. We denote the space of self-adjoint bounded operators in H by the symbol $\mathcal{L}_s(H)$.

2.1 Classical model

a). States are represented by points of some set M (state space).

b). Physical variables are represented by functions $f:M\to {\bf R}$ belonging to some functional space $V(M).^2$

c). Statistical states are represented by probability measures on M belonging to some class S(M).

d). The average of a physical variable (which is represented by a function $f \in V(M)$) with respect to a statistical state (which is represented by a probability measure $p \in S(M)$) is given by

$$\langle f \rangle_p \equiv \int_M f(\phi) dp(\phi).$$
 (1)

By using the language of probability theory we can say that there is given a random vector $\phi(\omega)$, where ω is a random parameter, taking values in M. Then $\langle f \rangle_{\phi} = Ef(\phi(\omega)) = \langle f \rangle_p$. Here and everywhere below E denotes classical mathematical expectation (average).

If the state space M is a space of functions, e.g., $M = L_2(\mathbf{R}^3)$, then M-valued random vectors are called *random fields*. For each ω , $\phi(\omega)$ is a function of $x \in \mathbf{R}^3 : \phi(x, \omega)$.

e). If systems $S_i, i = 1, 2, ..., k$, have state spaces M_i , respectively, then the composite system $S = (S_i)_{i=1}^k$ has state space $M = M_1 \times ... \times M_k$, the Cartesian product of the state spaces M_i . Ensembles of S-systems are described by random vectors in $M : \phi(\omega) = (\phi_1(\omega), ..., \phi_k(\omega))$ (or equivalently by probability measures on M.) A trivial, but important, remark is that in general components of $\phi(\omega)$ are not independent. There are nontrivial correlations between them. The best way to describe these correlations is to use the *covariance operator* (it will be defined little bit later).

A classical statistical model is a pair M = (S(M), V(M)).

2.2 Quantum case

Let H be a complex Hilbert space.

- a). States (pure) are represented by classes of normalized vectors of H with respect to the equivalence relation: $\psi_1 = e^{i\theta}\psi_2$.
- a). Physical observables are represented by operators $\widehat{A} : H \to H$

of the class $\mathcal{L}_{s}(H)$. (To simplify considerations, we shall consider only quantum observables represented by bounded operators.)

²The choice of a concrete functional space V(M) depends on various physical and mathematical factors. In classical mechanics for systems with the finite number of degrees of freedom M is chosen as the phase space $\mathbf{R}^{2n}; V(M)$ is the space of smooth functions.

b). Statistical states are represented by density operators. The class of such operators is denoted by $\mathcal{D}(H)$.

d). Average of a physical observable (which is represented by the operator $\widehat{A} \in \mathcal{L}_{s}(H)$) with respect to a statistical state (which is represented by the density operator $\rho \in \mathcal{D}(H)$) is given by von Neumann's formula

$$\langle A \rangle_{\rho} \equiv \operatorname{Tr} \rho \widehat{A}$$
 (2)

e). If quantum systems $S_i, i = 1, 2, ..., k$, have the state spaces H_i , respectively, then the system $S = (S_i)_{i=1}^k$ has the state space $H_1 \otimes ... \otimes H_k$, the tensor product of state spaces H_i .

The quantum statistical model is the pair $N_{\text{quant}} = (\mathcal{D}(H), \mathcal{L}_s(H)).$

At the first sight the gap between classical statistical mechanics and quantum mechanics is huge [10]. Impossibility to reduce quantum averages to classical averages is the main source of the great ideological difference between classical and quantum probabilistic descriptions.

3 Gaussian measures on real and complex Hilbert spaces

3.1 Real case

Let W be a real Hilbert space. Let $A \in \mathcal{L}_s(W)$.

We start with derivation of the basic mathematical formula which was used in [14]- [20]. We will calculate the Gaussian integral of the quadratic form

$$f_A(\phi) = (A\phi, \phi). \tag{3}$$

Consider a σ -additive Gaussian measure p on the σ -field of Borel subsets of W. This measure is determined by its covariance operator $B: W \to W$ and mean value $m \in W$. For example, B and m determine the Fourier transform of p

$$\tilde{p}(y) = \int_{W} e^{i(y,\phi)} dp(\phi) = e^{\frac{1}{2}(By,y) + i(m,y)}, y \in W.$$

In what follows we restrict our considerations to Gaussian measures with zero mean value: $(m, y) = \int_W (y, \psi) dp(\psi) = 0$ for any $y \in W$. Sometimes there will be used the symbol p_B to denote the Gaussian measure with the covariance operator B and m = 0. We recall that the covariance operator B is defined by its bilinear form

$$(By_1, y_2) = \int (y_1, \phi)(y_2, \phi) dp(\phi), y_1, y_2 \in W,$$
(4)

and it has the following properties: a) $B \ge 0$, i.e., $(By, y) \ge 0, y \in W$; b) B is a self-adjoint operator, $B \in \mathcal{L}_s(W)$; c) B is a trace-class operator and $\operatorname{Tr} B = \int_W ||\phi||^2 dp(\phi)$. It is *dispersion* of the probability p. Thus for Gaussian probability we have $\sigma^2(p) = \operatorname{Tr} B$. We remark that the list of properties of the covariation operator of a Gaussian measure differs from the list of properties of a von Neumann density operator only by one condition: $\operatorname{Tr} \rho = 1$, for a density operator ρ . Thus, for any covariance operator B, its scaling $B/\operatorname{Tr} B$ can be considered as a density operator. By using (4) we can easily find the Gaussian integral of the quadratic form $f_A(\phi)$ defined by (3):

$$\int_{W} f_A(\phi) dp_B(\phi) = \int_{W} (A\phi, \phi) dp_B(\phi)$$
$$= \sum_{i,j=1}^{\infty} (Ae_i, e_j) \int_{W} (e_i, \phi) (e_j, \phi) dp_B(\psi) = \sum_{i,j=1}^{\infty} (Ae_i, e_j) (Be_i, e_j),$$

where $\{e_i\}$ is some orthonormal basis in W. Thus $\int_W f_A(\phi) dp_B(\phi) = \text{Tr } BA$.

3.2 Complex case

Let Q and P be two copies of a real Hilbert space. Let us consider their Cartesian product $H = Q \times P$, "phase space," endowed with the symplectic operator $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Consider the class of Gaussian measures (with zero mean value) which are invariant with respect to the action of the operator J; denote this class S(H). It is easy to show that $p \in S(H)$ if and only if its covariance operator commutes with the symplectic operator, [44].

As always, we consider complexification of H (which will be denoted by the same symbol), $H = Q \oplus iP$. The complex scalar product is denoted by the symbol $\langle \cdot, \cdot \rangle$. We will also use the operation of complex congugation * in complex Hilbert space H, for $\phi = \phi_1 + i\phi_2, \phi_1 \in Q, \phi_2 \in P$, we set * $(\phi) = \overline{\phi} = \phi_1 - i\phi_2$. We will use the following trivial fact:

$$\langle \overline{u}, \overline{v} \rangle = \langle v, u \rangle.$$
 (5)

We introduce the complex covariance operator of a measure p on the complex Hilbert space H

$$\langle Dy_1, y_2 \rangle = \int_H \langle y_1, \phi \rangle \langle \phi, y_2 \rangle dp(\phi).$$

We also consider the complex Fourier transform of p

$$\tilde{p}(y) = \int_{H} exp\{i(\langle y, \phi \rangle + \langle \phi, y \rangle)\}dp(\phi).$$
(6)

Any J-invariant Gaussian measure on H is determined by its complex Fourier transform,[44]: $\tilde{p}(y) = exp\{-\langle Dy, y\rangle\}$.

We remark that J-invariance is a strong constraint on the class of Gaussian measures under consideration. Consider a measure p on the Cartesian product $H = Q \times P$. Its real covariance operator has the block structure³

 $B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}$, where $B_{11}^* = B_{11}, B_{22}^* = B_{22}, B_{12}^* = B_{21}$. Consider also its complex covariance operator D. It can be realized as acting in the Cartesian product of two real Hilbert spaces and in such a representation it also has the block structure $D_{\text{real}} = \begin{pmatrix} L & C \\ -C & L \end{pmatrix}$. It was shown in [44] that $L = B_{11} + B_{22}$ and $C = B_{12} - B_{21}$. It also was shown that

 $^{^{3}}$ Little bit later we will use the block structure of the complex covariance operator for a measure defined on the Cartesian product of two *complex Hilbert spaces*. The reader should be careful and not mix these two totally different block structures!

if measure is symplectically invariant then $B_{11} = B_{22}, B_{21} = -B_{12}$. Thus in the latter case the complex and real covariance operators are coupled in a simple way: $D_{\text{real}} = 2B$.

Lemma 1. For any measure $p \in S(H)$ the following representation takes place

$$\int_{H} \langle \xi_1, \phi \rangle \langle \eta_1, \phi \rangle \langle \phi, \xi_2 \rangle \langle \phi, \eta_2 \rangle dp(\phi) = \langle D\xi_1, \eta_2 \rangle \langle D\eta_1, \xi_2 \rangle + \langle D\xi_1, \xi_2 \rangle \langle D\eta_1, \eta_2 \rangle$$
(7)

To prove this formula, one should differentiate the Fourier transform (6) four times.

Let H_1 and H_2 be two complex Hilbert spaces and let $D_{21} \in \mathcal{L}(H_1, H_2), D_{12} \in \mathcal{L}(H_2, H_1)$. Then $D_{21} \otimes D_{12} \in \mathcal{L}(H_1 \otimes H_2, H_2 \otimes H_1)$. Let us consider the permutation operator $\sigma : H_2 \otimes H_1 \to H_1 \otimes H_2, \sigma(\phi_2 \otimes \phi_1) = \phi_1 \otimes \phi_2$. We remark that $\sigma \in \mathcal{L}(H_2 \otimes H_1, H_1 \otimes H_2)$.

Let p be a measure on the Cartesian product $H_1 \times H_2$ of two Hilbert spaces. Then its covariance operator has the block structure

$$D = \begin{pmatrix} D_{11} & D_{12} \\ D_{21} & D_{22} \end{pmatrix}, \tag{8}$$

where $D_{ii}: H_i \to H_i$ and $D_{ij}: H_j \to H_i$. The operator is self-adjoint. Hence $D_{ii}^* = D_{ii}$, and $D_{12}^* = D_{21}$.

Let H be a complex Hilbert space and let $\widehat{A} \in \mathcal{L}(H, H)$. We consider its quadratic form (which will play an important role in our further considerations)

$$\phi \to f_A(\phi) = \langle A\phi, \phi \rangle.$$

We make a trivial, but ideologically important remark: $f_A: H \to H$, is a "usual function" which is defined point wise.

In the same way as in the real case we prove the equality

$$\int_{H} f_A(\phi) dp_D(\phi) = \text{Tr } DA$$
(9)

Theorem 1. Let $p \in S(H_1 \times H_2)$ with the (complex) covariance operator D and let $\widehat{A}_i \in \mathcal{L}(H_i, H_i), i = 1, 2$. Then

$$\int_{H_1 \times H_2} f_{A_1}(\phi_1) f_{A_2}(\phi_2) dp(\phi) = \operatorname{Tr} D_{11} \widehat{A}_1 \operatorname{Tr} D_{22} \widehat{A}_2 + \operatorname{Tr} D_{12} \widehat{A}_2 D_{21} \widehat{A}_1$$
(10)

This theorem is a consequence of the following general result:

Lemma 2. Let $p \in S(H)$ with the (complex) covariance operator D and let $\widehat{A}_i \in \mathcal{L}(H,H), i = 1, 2$. Then

$$\int_{H} f_{A_1}(\phi) f_{A_2}(\phi) dp(\phi) = \operatorname{Tr} D\widehat{A}_1 \operatorname{Tr} D\widehat{A}_2 + \operatorname{Tr} D\widehat{A}_2 D\widehat{A}_1.$$
(11)

Proof. By Lemma 1 the integral can be represented as

$$I = \sum_{i_1 j_1} \sum_{i_2 j_2} \langle \hat{A}_1 e_{i_1}, e_{j_1} \rangle \langle \hat{A}_2 e_{i_2}, e_{j_2} \rangle$$

$$\times [\langle De_{j_1}, e_{i_2} \rangle \langle De_{j_2}, e_{i_1} \rangle + \langle De_{j_1}, e_{i_1} \rangle \langle De_{j_2}, e_{i_2} \rangle] = I_1 + I_2$$

where $\{e_i\}$ is an orthonormal basis in H. Here

$$\begin{split} I_1 &= \sum_{i_1 i_2} \sum_{j_1} \langle \hat{A}_1 e_{i_1}, e_{j_1} \rangle \langle e_{j_1}, D e_{i_2} \rangle \sum_{j_2} \langle \hat{A}_2 e_{i_2}, e_{j_2} \rangle \langle e_{j_2}, D e_{i_1} \rangle \\ &= \sum_{i_1 i_2} \langle D \hat{A}_1 e_{i_1}, e_{i_2} \rangle \langle e_{i_2}, \hat{A}_2^* D e_{i_1} \rangle \\ &= \sum_i \langle D \hat{A}_2 D \hat{A}_1 e_{i_1}, e_{i_1} \rangle = \operatorname{Tr} D \hat{A}_2 D \hat{A}_1. \\ I_2 &= \sum_{i_1 i_2} \sum_{j_2} \langle \hat{A}_1 e_{i_1} e_{j_1} \rangle \langle e_{j_1}, D e_{i_1} \rangle \sum_{j_2} \langle \hat{A}_2 e_{i_2} e_{j_2} \rangle \langle e_{j_2}, D e_{i_2} \rangle \\ &= \sum_{i_1 i_2} \langle \hat{A}_1 e_{i_1}, D e_{i_1} \rangle \langle \hat{A}_2 e_{i_2}, D e_{i_2} \rangle = \operatorname{Tr} D \hat{A}_1 \operatorname{Tr} D_{A_2}. \end{split}$$

Proposition 1. Let conditions of Theorem 1 hold. Then

$$\int_{H_1 \times H_2} f_{A_1}(\phi_1) f_{A_2}(\phi_2) dp(\phi) = \operatorname{Tr}(D_{11} \otimes D_{22} + \sigma(D_{21} \otimes D_{12})) \widehat{A}_1 \otimes \widehat{A}_2.$$
(12)

Proof. It is sufficient to prove that

$$\operatorname{Tr} \sigma(D_{21} \otimes D_{12})\widehat{A}_1 \otimes \widehat{A}_2 = \operatorname{Tr} D_{12}\widehat{A}_2 D_{21}\widehat{A}_1.$$

We have

$$\operatorname{Tr}\sigma(D_{21}\otimes D_{12})\widehat{A}_{1}\otimes A_{2} = \sum_{ij} \langle \sigma(D_{21}\otimes D_{12})\widehat{A}_{1}\otimes \widehat{A}_{2}e_{i}\otimes f_{j}, e_{i}\otimes f_{j} \rangle$$
$$= \sum_{ij} \langle D_{12}\widehat{A}_{2}f_{j}\otimes D_{21}\widehat{A}_{1}e_{i}, e_{i}\otimes f_{j} \rangle = \sum_{ij} \langle D_{12}\widehat{A}_{2}f_{j}, e_{i} \rangle \langle D_{21}\widehat{A}_{1}e_{i}, f_{j} \rangle.$$
On the other hand, TrD, \widehat{A} , D, \widehat{A}

On the other hand, $TrD_{12}A_2D_{21}A_1$

$$=\sum_{i} \langle D_{12}\widehat{A}_2 D_{21}\widehat{A}_1 e_i, e_i \rangle \sum_{i} \langle D_{21}\widehat{A}_1 e_i, A_2^* D_{21} e_i \rangle \sum_{ij} \langle D_{21}\widehat{A}_1 e_i, f_j \rangle \langle f_j, A_2^* D_{21} e_i \rangle$$

4 Vectors, operators, traces

4.1 Vector and operator realizations of the tensor product

In quantum theory a pure state of a composite system is represented by a normalized vector belonging to the tensor product $H_1 \otimes H_2$. On the other hand, in functional analysis it is common to use elements of $H_1 \otimes H_2$ as operators. The standard construction provides the realization of any vector $\Psi \in H_1 \otimes H_2$ by a linear operator from $H_2^* \to H_1$, where H_2^* is the space dual to H_2 . For a vector $Z = u \otimes v$, one puts $\widehat{Z}x^* = x^*(v)u$ for $x^* \in H_2^*$. This correspondence $Z \to \widehat{Z}$ is extended to isomorphism of $H_1 \otimes H_2$ with the space of Hilbert-Smidt operators $\mathcal{HS}(H_2^*, H_1)$ or (equivalently) with the space of anti-linear HS-operators from H_2 to H_1 . The main compication in coming considerations is related to our need to represent vectors from the tesor product by *linear* operators from H_2 to H_1 . We present a new construction which seems to be unknown, so it may be interesting even from the purely mathematical viewpoint. We remind the definition of the *HS*-norm corresponding to the trace scalar product in $\mathcal{HS}(H_2, H_1)$. Take an arbitrary orthonormal basis $\{f_k\}$ in H_2 . Then

$$\langle \widehat{L_1}, \widehat{L_2} \rangle = \sum_k \langle \widehat{L_1} f_k, \widehat{L_2} f_k \rangle = \mathrm{Tr} \widehat{\mathbf{L}_2}^* \widehat{\mathbf{L}_1}, \ \|\widehat{\mathbf{L}}\|_2^2 = \mathrm{Tr} \widehat{\mathbf{L}}^* \widehat{\mathbf{L}}$$

(index 2 is typically used for the Hilbert-Schmidt norm).

Let $\{e_j\}$ and $\{f_j\}$ be two orthonormal bases in H_1 and H_2 , respectively. Then

$$\Psi = \sum_{ij} \psi_{ij} e_i \otimes f_j, \psi_{ij} \in \mathbf{C},$$

and $||\Psi||^2 = \sum_{ij} |\psi_{ij}|^2$. We remark that, for an orthonormal basis, say $\{f_j\}$, the system of complex conjugate vectors $\{\overline{f}_j\}$ is also an orthonormal basis. Really, by (5) we get $\langle \overline{f}_j, \overline{f}_i \rangle = \langle f_i, f_j \rangle = \delta_{ij}$.

We set, for $\phi \in H_2$,

$$\widehat{\Psi}\phi = \sum_{ij} \psi_{ij} \langle \phi_2, \overline{f}_j \rangle e_i.$$
(13)

We emphasize that, for the vector Ψ , the expansion with respect to the basis $\{e_i \otimes f_j\}$ was used. In cotrast to this, in the expansion of the operator $\widehat{\Psi}$ the basis $\{e_i \otimes \overline{f_j}\}$ was used. If the basis is real than the definition is essentially simplified:

$$\widehat{\Psi}\phi = \sum_{ij} \psi_{ij} \langle \phi_2, f_j \rangle e_i.$$
(14)

Take now a factorizable vector $\Psi = u \otimes v$. It defines the rank one operator

$$\widehat{\Psi}\phi = \langle \phi, \overline{v} \rangle u = \langle v, \overline{\phi} \rangle u.$$

To play with the definition, take $\Psi = cu \otimes v = (cu) \otimes v = u \otimes (cv)$, where $c \in \mathbf{C}$. On the one hand, we have $\widehat{\Psi}\phi = c\langle\phi, \overline{v}\rangle u$; on the other hand, $\widehat{\Psi}\phi = \langle\phi, \overline{cv}\rangle u = \langle\phi, \overline{cv}\rangle u = c\langle\phi, \overline{v}\rangle u$. Thus our definition is consistent with scaling by a complex constant. Consider now very special, but at the same time very important case: $H_i = L_2(\mathbf{R}^{n_i}), i = 1, 2$. Here our definition gives the following representation (a special case of (14)):

$$\widehat{\Psi}\phi(x) = \int \Psi(x, y)\phi(y)dy \tag{15}$$

We proceed formally and use the real basis $\{e_x \otimes f_y\}$, where $e_x(t) = \delta(t-x)$ and $f_y(s) = \delta(s-y)$. Finally, we remind a result of theory of integral operators. The operator given by (15) is of the *HS*-type in *L*₂-spaces if and only if its kernel $\Psi(x, y)$ is square integrable. Thus the condition $\Psi \in L_2(\mathbf{R}^{n_1} \times \mathbf{R}^{n_2})$ for a function Ψ is equivalent to the *HS*-condition for the operator $\widehat{\Psi}$. In the abstract form this fact will be formulated in coming lemma.

Lemma 3. Each vector $\Psi \in H_1 \otimes H_2$ determines (uniquely) operator $\widehat{\Psi} \in \mathcal{L}(H_2, H_1)$ and, moreover, $\widehat{\Psi} \in \mathcal{HS}(H_1, H_2)$ and $||\widehat{\Psi}||_2 = ||\Psi||$ (the norm of a vector coicides with the Hilbert-Schmidt norm of the corresponding operator)

Proof. We start with proof of corectness of definition (13), i.e., it does not depend on the choice of orthonormal bases. Let $\{e'_j\}$ and $\{f'_j\}$ be two orthonormal bases in H_1 and H_2 , respectively, which are in general different from bases $\{e_j\}$ and $\{f_j\}$. Consider the expansion $\Psi = \sum_{ij} \psi'_{ij} e'_i \otimes f'_j$. We remark that $\overline{f'_j} = \sum_k \langle \overline{f'_j}, \overline{f_k} \rangle \overline{f_k} = \sum_k \langle f_k, f'_j \rangle \overline{f_k}$, which can also be obtained by complex conjugation from the expansion $f'_j = \sum_k \langle f'_j, f_k \rangle f_k$. In this basis the vector Ψ defines the operator

$$\widehat{\Psi}'\phi = \sum_{ij} \psi_{ij}' \langle \phi, \overline{f_j'} \rangle e_i'.$$
(16)

We have

$$\begin{aligned} \widehat{\Psi}'\phi &= \sum_{ij} \sum_{nm} \psi_{nm} \langle e_n, e_i' \rangle \langle f_m, f_j' \rangle \langle \phi, \sum_k \langle \overline{f_j'}, \overline{f_k} \rangle \overline{f_k} \rangle \sum_p \langle e_i', e_p \rangle e_p \\ &= \sum_{nm} \psi_{nm} \sum_{kp} \langle \phi, \overline{f_k} \rangle \sum_i \langle e_n, e_i' \rangle \langle e_i', e_p \rangle \sum_j \langle \overline{f_k}, \overline{f_j'} \rangle \langle f_m, f_j' \rangle e_p. \end{aligned}$$

We remark

$$\sum_{j} \langle \overline{f_k}, \overline{f'_j} \rangle \langle f_m, f'_j \rangle = \sum_{j} \langle f_m, f'_j \rangle \langle f'_j, f_k \rangle = \delta_{km}$$

Thus $\widehat{\Psi}'\phi = \sum_{nm} \psi_{nm} \langle \phi, \overline{f_m} \rangle e_n$. It coincides with (13). We now apply the Cauchy-Bunyakovsky inequality to the expression (13) and obtain: $\|\widehat{\Psi}\phi\|^2 = \sum_i |\sum_j \psi_{ij} \langle \phi, \overline{f_j} \rangle|^2 \leq \sum_i \sum_j |\psi_{ij}|^2 \sum_j |\langle \phi, \overline{f_j} \rangle|^2 = \|\Psi\|^2 \|\phi\|^2$. Thus this operator belongs to the space $\mathcal{L}(H_2, H_1)$. Finally, we show that it belongs even to the space $\mathcal{HS}(H_2, H_1) : \|\widehat{\Psi}\|_2^2 = \text{Tr}\widehat{\Psi}^*\widehat{\Psi} = \sum_k \langle \widehat{\Psi}\overline{f_k}, \widehat{\Psi}\overline{f_k} \rangle = \sum_k \langle \sum_{i_1} \psi_{i_1k} e_{i_1}, \sum_{i_1} \psi_{i_2k} e_{i_2} \rangle = \sum_{ki} \psi_{ik} \overline{\psi_{ik}}.$

Moreover, it is easy to show that any operator $\widehat{L} \in \mathcal{HS}(H_2, H_1)$ can be represented as (13) for some vector $\Psi \in H_1 \otimes H_2$. In the combination with Lemma 3 this remark implies:

Corollary 1. The equality (13) establishes the isomorphism of Hilbert spaces $H_1 \otimes H_2$ and $\mathcal{HS}(H_2, H_1)$.

We now find the adjoint operator. We have:

$$\langle \widehat{\Psi}y, x \rangle = \sum_{nm} \psi_{nm} \langle y, \overline{f_m} \rangle \langle e_n, x \rangle = \langle y, \sum_{nm} \overline{\psi_{nm}} \langle x, e_n \rangle \overline{f_m} \rangle.$$

Thus

$$\widehat{\Psi}^* x = \sum_{nm} \overline{\psi_{nm}} \langle x, e_n \rangle \overline{f_m}.$$

4.2 Operation of the complex conjugation in the space of self-adjoint operators

Let $\widehat{A} \in \mathcal{L}_s(W)$, where W is Hilbert space. We define "complex conjugate operator" $\widehat{\overline{A}}$ by its bilinear form:

$$\langle \bar{A}u, v \rangle = \langle \bar{v}, \hat{A}\bar{u} \rangle.$$
 (17)

Let $\{f_j\}$ be an orthormal basis in W. We find the matrix of the operator $\widehat{B} \equiv \widehat{A}$ with respect to this basis: $b_{ij} = \langle \widehat{A}f_i, f_j \rangle = \langle \overline{f}_j, \widehat{A}\overline{f}_i \rangle$. In the special case of the real basis, i.e., $\overline{f}_j = f_j$, we have:

$$b_{ij} = \overline{a_{ij}} = a_{ji}.$$

Thus its matrix is given by the transposition of the matrix of \widehat{A} . We will use the fact that the operator \widehat{A} is self-adjoint (we remind that \widehat{A} is self-ajoint):

$$\langle \hat{\bar{A}}u, v \rangle = \langle \hat{A}\bar{v}, \bar{u} \rangle = \overline{\langle \bar{u}, \hat{A}\bar{v} \rangle} = \overline{\langle \hat{\bar{A}}v, u \rangle} = \langle u, \hat{\bar{A}}v \rangle.$$
(18)

We will also use the fact that, for a positiv operator \hat{N} , the operator \hat{N} is also positiv:

$$\langle \bar{N}u, u \rangle = \langle \bar{u}, \hat{N}\bar{u} \rangle = \langle \hat{N}\bar{u}, \bar{u} \rangle.$$
(19)

Consider the quadratic form of the complex conjugate operator $\widehat{\bar{A}}$ of a self-adjoint operator \widehat{A} :

$$f_{\bar{A}}(\phi) = \langle \overline{\hat{A}}\phi, \phi \rangle = \langle \overline{\phi}, \widehat{A}\overline{\phi} \rangle = f_A(\overline{\phi}).$$
(20)

Consider the group $\{e, *\}$, where * is the operation of complex conjugation in a compex Hilbert space W. It induces the action in the space of real valued functions on $W : f \to \overline{f}$, where $\overline{f}(\phi) = f(\overline{\phi})$. We hope that the symbol \overline{f} will not be misleading. Only real valued functions are under consideration. Thus it cannot be mixed with the operation of complex conjugation on the range of values.

Theorem 1 implies (for $\widehat{A}_i \in \mathcal{L}_s(H_i, H_i), i = 1, 2$):

$$\int_{H_1 \times H_2} f_{A_1}(\phi_1) \bar{f}_{A_2}(\phi_2) dp(\phi_1, \phi_2) = \text{Tr} D_{11} \hat{A}_1 \text{Tr} D_{22} \hat{\bar{A}}_2 + \text{Tr} D_{12} \hat{\bar{A}}_2 D_{21} \hat{A}_1$$
(21)

4.3 The basic operator equality

Lemma 4. Let $\Psi \in H_1 \otimes H_2$. Then, for any pair of operators $\widehat{A}_j \in \mathcal{L}_s(H_j), j = 1, 2$,

$$\mathrm{Tr}\widehat{\Psi}\widehat{A}_{2}\widehat{\Psi}^{*}\widehat{A}_{1} = \langle \widehat{A}_{1} \otimes \widehat{A}_{2} \rangle_{\Psi} \equiv \langle \widehat{A}_{1} \otimes \widehat{A}_{2} \Psi, \Psi \rangle.$$
(22)

Proof. We have:

$$\begin{aligned} \mathrm{Tr}\widehat{\Psi}\widehat{\mathbf{A}}_{2}\widehat{\Psi}^{*}\widehat{\mathbf{A}}_{1} &= \sum_{\mathbf{k}} \langle \widehat{\Psi}\widehat{\mathbf{A}}_{2}\widehat{\Psi}^{*}\widehat{\mathbf{A}}_{1}\mathbf{e}_{\mathbf{k}}, \mathbf{e}_{\mathbf{k}} \rangle = \sum_{\mathbf{k}} \sum_{\mathbf{ij}} \psi_{\mathbf{ij}} \langle \widehat{\mathbf{A}}_{2}\widehat{\Psi}^{*}\widehat{\mathbf{A}}_{1}\mathbf{e}_{\mathbf{k}}, \overline{\mathbf{f_{j}}} \rangle \delta_{\mathbf{ik}} \\ &= \sum_{ij} \psi_{ij} \langle \widehat{A}_{2}\widehat{\Psi}^{*}\widehat{A}_{1}\mathbf{e}_{i}, \overline{f_{j}} \rangle = \sum_{i_{1}j_{1}} \sum_{i_{2}j_{2}} \psi_{i_{1}j_{1}} \overline{\psi_{i_{2}j_{2}}} \langle \widehat{A}_{1}\mathbf{e}_{i_{1}}, \mathbf{e}_{i_{2}} \rangle \langle \overline{f_{j_{2}}}, \widehat{\overline{A}}_{2}\overline{f_{j_{1}}} \rangle. \end{aligned}$$

By (18) and (17) we obtain $\langle \overline{f_{j_2}}, \widehat{\overline{A}}_2 \overline{f_{j_1}} \rangle = \langle \widehat{\overline{A}}_2 \overline{f_{j_2}}, \overline{f_{j_1}} \rangle = \langle f_{j_1}, \widehat{\overline{A}}_2 f_{j_2} \rangle = \langle \widehat{A}_2 f_{j_1}, f_{j_2} \rangle$. Thus

$$\mathrm{Tr}\widehat{\Psi}\widehat{A}_{2}\widehat{\Psi}^{*}\widehat{A}_{1} = \sum_{i_{1}j_{1}}\sum_{i_{2}j_{2}}\psi_{i_{1}j_{1}}\overline{\psi_{i_{2}j_{2}}}\langle\widehat{A}_{1}e_{i_{1}},e_{i_{2}}\rangle\langle\widehat{A}_{2}f_{j_{1}},f_{j_{2}}\rangle.$$

On the other hand, we obtain:

$$\langle \hat{A}_1 \otimes \hat{A}_2 \Psi, \Psi \rangle = \sum_{i_1 j_1} \sum_{i_2 j_2} \psi_{i_1 j_1} \overline{\psi_{i_2 j_2}} \langle \hat{A}_1 e_{i_1}, e_{i_2} \rangle \langle \hat{A}_2 f_{j_1}, f_{j_2} \rangle = \operatorname{Tr} \widehat{\Psi} \widehat{\bar{A}}_2 \widehat{\Psi}^* \widehat{A}_1.$$

If Ψ is normalized by 1, then the right-hand side of equality (22) is nothing else than average of the observable $\hat{C} = \hat{A}_1 \otimes \hat{A}_2$ describing correlations between measurement of observables \hat{A}_1 and \hat{A}_2 on subsystems S_1 and S_2 of a composite system $S = (S_1, S_2)$ which is prepared in the state Ψ . On the other hand, the left-hand side of equality (22) has the form of the second term in the right-hand side of formula (10) giving Guassian integral of the product of two quadratic forms corresponding to operators \hat{A}_1 and \hat{A}_2 . These mathematical coincidences provide a possibility to couple quantum correlations with classical Gaussian correlations, by selecting the covariance operator of the prequantum Gaussian distribution (corresponding to quantum state Ψ) in the right way.

4.4 Operator representation of reduced density operators

Lemma 5. For any vector $\Psi \in H_1 \otimes H_2$, the following equality holds:

$$\mathrm{Tr}_{H_2}\Psi\otimes\Psi=\widehat{\Psi}\widehat{\Psi}^*.$$
 (23)

Proof. The operator $\widehat{\Psi}\widehat{\Psi}^*$ acts on vector $x \in H_1$ as

$$\begin{split} \widehat{\Psi}\widehat{\Psi}^*x &= \sum_{nm} \overline{\psi_{nm}} \langle x, e_n \rangle \widehat{\Psi}\overline{f_m} = \sum_{nm} \sum_{kl} \psi_{kl} \overline{\psi_{nm}} \langle \overline{f_m}, \overline{f_l} \rangle \langle x, e_n \rangle e_k \\ &= \sum_{nk} [\sum_m \psi_{km} \overline{\psi_{nm}}] \langle x, e_n \rangle e_k. \end{split}$$

Thus its bilinear form is given by

$$\langle \widehat{\Psi} \widehat{\Psi}^* x, y \rangle = \sum_{nk} [\sum_{m} \psi_{km} \overline{\psi_{nm}}] \langle x, e_n \rangle \langle e_k, y \rangle$$

Take an orthogonormal basis $\{f_m\}$ in H_2 . The bilinear form of the operator $\text{Tr}_{H_2}\Psi\otimes\Psi$ is given by

$$\langle \mathrm{Tr}_{H_2}\Psi\otimes\Psi x, y\rangle = \sum_m \langle \mathrm{Tr}_{H_2}\Psi\otimes\Psi x\otimes f_m, y\otimes f_m\rangle$$

$$=\sum_{m}\sum_{kl}\langle x,e_k\rangle\langle e_l,y\rangle\langle e_k\otimes f_m,\Psi\rangle\langle\Psi,e_l\otimes f_m\rangle=\sum_{kl}[\sum_{m}\psi_{lm}\overline{\psi_{km}}]\langle x,e_k\rangle\langle e_l,y\rangle.$$

By setting $k \to n, l \to k$ we obtain the coincidence of two bilinear forms and hence the operators.

If the vector Ψ is normalized, then $\rho_{\Psi} = \Psi \otimes \Psi$ is the corresponding density operator and $\rho_{\Psi}^{(1)}$ is the H_1 -reduced density operator. By (23) we obtain

$$\rho_{\Psi}^{(1)} = \widehat{\Psi}\widehat{\Psi}^*. \tag{24}$$

Unfortunately, for the H_1 -reduced density operator $\rho_{\Psi}^{(2)},$ similar statement is not true; in general,

$$\rho_{\Psi}^{(2)} \neq \widehat{\Psi}^* \widehat{\Psi}.$$
 (25)

Let find the bilinear form of the operator $\widehat{\Psi}^*\widehat{\Psi}: H_2 \to H_2:$

$$\widehat{\Psi}^* \widehat{\Psi} y = \sum_{nm} \psi_{nm} \langle y, \overline{f_m} \rangle \widehat{\Psi}^* e_n = \sum_{nm,ij} \psi_{nm} \overline{\psi_{ij}} \langle y, \overline{f_m} \rangle \langle e_n, e_i \rangle \overline{f_j}$$

$$= \sum_{mj} [\sum_i \psi_{im} \overline{\psi_{ij}}] \langle y, \overline{f_m} \rangle \overline{f_j}.$$

$$\langle \widehat{\Psi}^* \widehat{\Psi} y, u \rangle = \sum_{mj} [\sum_i \psi_{im} \overline{\psi_{ij}}] \langle y, \overline{f_m} \rangle \langle \overline{f_j}, u \rangle.$$
(26)

However, for any orthonormal basis $\{e_m\}$ in H_1 , we obtain:

$$\langle \operatorname{Tr}_{H_1} \Psi \otimes \Psi y, u \rangle = \sum_m \langle \Psi \otimes \Psi e_m \otimes y, e_m \otimes u \rangle$$
$$= \sum_{mjl} \langle e_m \otimes f_j, \Psi \rangle \langle \Psi, e_m \otimes f_l \rangle \langle y, f_j \rangle \langle f_l, u \rangle$$
$$\sum_{jl} [\sum_m \psi_{ml} \overline{\psi_{mj}}] \langle y, f_j \rangle \langle f_l, u \rangle.$$
(27)

By setting $m \to i, j \to m, l \to j$ in (27) we obtain

$$\langle \operatorname{Tr}_{H_1} \Psi \otimes \Psi y, u \rangle = \sum_{mj} [\sum_i \psi_{ij} \overline{\psi_{im}}] \langle y, f_m \rangle \langle f_j, u \rangle.$$
 (28)

Comparing (26) and (28) we see that in general they do not coincide. To show this, let us take the real basis $\{f_m\}$. In this case (26) has the form:

$$\langle \widehat{\Psi}^* \widehat{\Psi} y, u \rangle = \sum_{mj} \left[\sum_i \psi_{im} \overline{\psi_{ij}} \right] \langle y, f_m \rangle \langle f_j, u \rangle.$$
⁽²⁹⁾

Thus in this special basis the matrix elements of operators $\widehat{\Psi}^*\widehat{\Psi}$ and $\operatorname{Tr}_{H_2}\Psi\otimes\Psi$ are coupled via the complex conjugation. In general, we have:

Lemma 5^{*}. For any vector $\Psi \in H_1 \otimes H_2$, the following equality holds for the operator $\widehat{T} = \operatorname{Tr}_{H_1} \Psi \otimes \Psi$:

$$\widehat{\overline{T}} = \widehat{\Psi}^* \widehat{\Psi}.$$
(30)

Proof. We now present proof which is not based on matrix elements. In particular, it illustrates well features of complex conjugate operators. For \hat{T} defined in the formulation, we obtain:

$$\langle \widehat{\bar{T}}y, u \rangle = \langle \bar{u}, \widehat{T}\bar{y} \rangle = \overline{\langle \widehat{T}\bar{y}, \bar{u} \rangle}.$$

By (27) we get:

$$\langle \hat{\overline{T}}y, u \rangle = \sum_{mj} [\sum_{i} \overline{\psi_{ij}} \psi_{im}] \langle f_m, \overline{y} \rangle \langle \overline{u}, f_j \rangle.$$

By (5) $\langle \bar{u}, v \rangle = \langle \bar{v}, u \rangle$ and hence $\langle \bar{u}, f_j \rangle = \langle \bar{f}_j, u \rangle$ and $\langle u, \bar{v} \rangle = \langle v, \bar{u} \rangle$ and hence $\langle f_m, \bar{y} \rangle = \langle y, \bar{f}_m \rangle$. Thus

$$\langle \widehat{\bar{T}}y, u \rangle = \sum_{mj} [\sum_{i} \overline{\psi_{ij}} \psi_{im}] \langle y, \bar{f_m} \rangle \langle \bar{f_j}, u \rangle = \langle \widehat{\Psi}^* \widehat{\Psi}y, u \rangle.$$

In particular, if Ψ is normalized (pure quantum state) then we obtain

$$\overline{\rho_{\Psi}^{(2)}} = \widehat{\Psi}^* \widehat{\Psi}.$$
(31)

Lemma 6. Let ρ be a density operator in a Hilbert space W. Then, for any $\widehat{A} \in \mathcal{L}_s(W)$, the following equality holds:

$$\operatorname{Tr} \overline{\rho}\overline{A} = \operatorname{Tr} \rho \widehat{A}. \tag{32}$$

Proof. Let $\{e_k\}$ be an orthonormal basis in W. Then

$$\operatorname{Tr} \overline{\rho} \widehat{\overline{A}} = \sum_{\mathbf{k}} \langle \overline{\rho} \widehat{\overline{A}} \mathbf{e}_{\mathbf{k}}, \mathbf{e}_{\mathbf{k}} \rangle = \sum_{\mathbf{k}} \langle \overline{e}_{\mathbf{k}}, \rho \overline{\widehat{\overline{A}} \mathbf{e}_{\mathbf{k}}} \rangle$$
$$= \sum_{k} \langle \rho \overline{e}_{k}, \overline{\widehat{A}} \overline{e}_{k} \rangle = \sum_{k} \langle \widehat{A} e_{k}, \overline{\rho \overline{e}_{k}} \rangle =$$
$$= \sum_{k} \langle \rho \overline{e}_{k}, \widehat{A} \overline{e}_{k} \rangle == \sum_{k} \langle \widehat{A} \rho \overline{e}_{k}, \overline{e}_{k} \rangle.$$

Since the trace does not depend on the choice of a basis, we can select the real basis. Hence, it was proved that

$$\operatorname{Tr} \overline{\rho} \widehat{\overline{A}} = \operatorname{Tr} \widehat{A} \rho.$$

Finally, we will prove that

$$\operatorname{Tr} \widehat{A}\rho = \operatorname{Tr} \rho \widehat{A}.$$

Take the basis consisting of the eigenvectors of the density operator: $\rho = \sum_{k} p_k e_k \otimes e_k$. Then

$$\operatorname{Tr} \rho \widehat{A} = \sum_{k} \langle \rho \widehat{A} e_{k}, e_{k} \rangle = \sum_{k} p_{k} \langle \widehat{A} e_{k}, e_{k} \rangle = \sum_{k} \langle \widehat{A} \rho e_{k}, e_{k} \rangle.$$

Corollary 2. Let $\Psi \in H_1 \otimes H_2$ be normalized (pure quantum state). Then

$$\operatorname{Tr} \widehat{\Psi}^* \widehat{\Psi} \overline{A} = \operatorname{Tr} \rho_{\Psi}^{(2)} \widehat{A}.$$
(33)

5 Classical random field description

5.1 Ensemble of noncomposite quantum systems

In what-follows random vectors taking values in a Hilbert space are called *random fields*. This definition is motivated by consideration of the Hilbert space $H = L_2(\mathbf{R}^m)$ of square integrable functions.

Let $\phi(\omega)$ denote a Gaussian random field in a complex Hilbert space H. Everywhere below we consider Gaussian random fields with probability distributions of the class S(H). The covariance operator of a random field is defined as the covariance operator of its probability distribution.

The correspondence between QM and PCSFT in the case of a single quantum system with the state space H is established in the following way:

1). Density operators (statistical states of QM) are identified with covariance operators of prequantum random fields, $\rho \mapsto D$.

2). Self-adjoint operators (quantum observables) are identified with quadratic functionals, $\widehat{A} \mapsto f_A$.

The equality (9) can be written as

$$Ef_A(\phi(\omega)) = \operatorname{Tr}\rho\widehat{A} \equiv \langle \widehat{A} \rangle_{\rho}.$$

It establishes the correspondence between PCSFT-averages and QM-averages. This story was presented in [14]- [20]. Now we modify it. Originally the source of coming modification was purely mathematical – to solve the problem of positive definiteness in theory for composite systems, see section 5.3. However, it happens that a natural physical interpretation can be provided.

To escape measure-theoretic difficulties, at the moment we proceed in the finite-dimensional Hilbert space. We will come back to the real physical case (for infinite dimension) in section 5.3, see Proposition 4. Let ρ be a density operator. Set $D = \rho + \alpha I$, where I is the unit operator and $\alpha > 0$. Consider the Gaussian random vector $\phi(\omega)$ with the covariance operator D. The additional term αI we can consider as (α -scaling of) the Gaussian normal distribution. It describes *spatial white noise* when the dimension of the space goes to infinity.

We have $Ef_A(\phi(\omega)) = \text{Tr}\rho \hat{A} + \alpha \text{Tr} \hat{A}$. In this model (modification of PCSFT created in [15]–[20]) quantum average can be obtained as a shift of classical average:

$$\langle \hat{A} \rangle_{\rho} = E f_A(\phi(\omega)) - \alpha \operatorname{Tr} \widehat{A}.$$
 (34)

The shift is generated by the presence of the background Gaussian noise (say "zero point field", cf. SED, [21]-[25]). Thus QM-average can be considered as simply normalization of average with respect to a prequantum random field. Normalization consists of substraction of the contribution of the background field. While in the finite-dimensional case the use of such a normalization is just a matter of test, in the infinite-dimensional case it becames very important. If quantum observable is represented by an operator A which is not of the trace class, then the normalization $TrA = \infty$. In other words the quadratic form $f_A(\phi)$ is not integrable, cf. Proposition 4, with respect to the probability distribution p_D , where $\rho + \alpha I$. Of course, it is a pure theoretical problem. In real experimental practice we are able to measure only observables represented by operators of finite ranks, see von Neumann [10]. Other observables (in particular, all observables given by operators with continuous spectra) are just mathematical idealizations. Nevertheless, it is convenient to have a theory which is able to operate with such quantities as well. From the PCSFT-viewpoint QM is such a theory. Thus in our approach QM has some analogy with QFT, but all divergences are regularized from the very beginning by choosing a special representation of classical averages.

5.2 Ensemble of composite quantum systems

Consider a composite quantum system $S = (S_1, S_2)$. Here S_j has the state space H_j , a complex Hilbert space. Let $\phi_1(\omega)$ and $\phi_2(\omega)$ be two Gaussian random fields, in Hilbert spaces H_1 and H_2 , respectively. Consider the Cartesian product of these Hilbert spaces, $H_1 \times H_2$, and the vector Gaussian random field $\phi(\omega) = (\phi_1(\omega), \phi_2(\omega)) \in H_1 \times H_2$. In the case under consideration its covariance operator has the block structure given by (8). Set

$$\langle f_{A_1}, f_{A_2} \rangle == E f_{A_1} \bar{f}_{A_2} = \int_{H_1 \times H_2} f_{A_1}(\phi_1) \bar{f}_{A_2}(\phi_2) dp(\phi_1, \phi_2).$$

Set also

$$\operatorname{cov}\left(f_{A_{1}}, f_{A_{2}}\right) = \langle f_{A_{1}}, f_{A_{2}} \rangle - \langle f_{A_{1}} \rangle \langle \bar{f}_{A_{2}} \rangle$$

Equalities (21) and (22) imply

Proposition 2. Let $\widehat{A}_i \in \mathcal{L}_s(H_i)$, i = 1, 2 and let $\Psi \in H_1 \otimes H_2$ with the unit norm. Then, for any Gaussian random field $\phi(\omega)$ in $H_1 \times H_2$ with the covariance matrix D such that the non-diagonal block

$$D_{12} = \widehat{\Psi} \tag{35}$$

the following equality takes place:

$$\operatorname{cov}\left(f_{A_1}, f_{A_2}\right) = \left(\widehat{A}_1 \otimes \widehat{A}_2 \Psi, \Psi\right) \equiv \langle \widehat{A}_1 \otimes \widehat{A}_2 \rangle_{\Psi}.$$
(36)

This equality establishes coupling between quantum and classical correlations. In the next section we will unify classical descriptions for a single system, section 5, and a composite system.

5.3 Making consistent PCSFT-models for ensembles of noncomposite and composite systems

Operators D_{ii} are responsible for averages of functionals depending only on one of components of the vector random field $\phi(\omega)$. In particular, $Ef_{A_1}(\phi_1)(\omega)) = \operatorname{Tr} D_{11}\widehat{A}_1$ and $E\overline{f}_{A_2}(\phi_2)(\omega)) = \operatorname{Tr} D_{22}\overline{A}_2$. We will construct such a random field that these "marginal averages" will match those given by QM. For the latter, we have:

$$\langle \widehat{A}_1 \rangle_{\Psi} = (\widehat{A}_1 \otimes I_2 \Psi, \Psi) = \operatorname{Tr} \rho_{\Psi}^{(1)} \widehat{A}_1, \langle \widehat{A}_2 \rangle_{\Psi} = (I_1 \otimes \widehat{A}_2 \Psi, \Psi) = \operatorname{Tr} \rho_{\Psi}^{(2)} \widehat{A}_2,$$

where I_i denotes the unit operator in H_i , i = 1, 2. By equality (24) the first average can written as

$$\langle \widehat{A}_1 \rangle_{\Psi} = \operatorname{Tr}(\widehat{\Psi}\widehat{\Psi}^*)\widehat{A}_1.$$

By equality (32) the second average can be written as

$$\langle \hat{A}_2 \rangle_{\Psi} = \mathrm{Tr} \overline{\rho_{\Psi}^{(2)}} \widehat{\bar{A}}_2$$

and, finally, by (31)

$$\langle \hat{A}_2 \rangle_{\Psi} = \operatorname{Tr}(\widehat{\Psi}^* \widehat{\Psi}) \widehat{\bar{A}}_2,$$

Thus it would be natural to take

$$D = \begin{pmatrix} \widehat{\Psi}\widehat{\Psi}^* & \widehat{\Psi} \\ \widehat{\Psi}^* & \widehat{\Psi}^*\widehat{\Psi} \end{pmatrix}.$$

Its off-diagonal block reproduces correct quantum correlations between systems S_1 and S_2 and its diagonal blocks produce correct quantum averages for system S_1 and system S_2 .

However, in general (i.e., for an arbitrary pure state Ψ) this operator is not positively defined. Therefore (in general) it could not be chosen as the covariance operator of a random field. Let us consider a modification which will be positively defined and such that quantum and classical averages will be coupled by a simple rule. Thus from quantum averages one can easily find classical averages and vice versa.

Proposition 3. For any normalized vector $\Psi \in H_1 \otimes H_2$, the operator

$$D = \begin{pmatrix} (\widehat{\Psi}\widehat{\Psi}^* + I_1/4) & \widehat{\Psi} \\ \widehat{\Psi}^* & (\widehat{\Psi}^*\widehat{\Psi} + I_2/4) \end{pmatrix}$$
(37)

is positively defined.

Proof. For any vector $\phi = (\phi_1, \phi_2) \in \Psi \in H_1 \times H_2$, we have: $(\tilde{D}_{\Psi}\phi, \phi) = ||\widehat{\Psi}^*\phi_1||^2 + \frac{||\phi_1||^2}{4} + (\widehat{\Psi}\phi_2, \phi_1) + (\widehat{\Psi}^*\phi_1, \phi_2) + ||\widehat{\Psi}\phi_2||^2 + \frac{||\phi_2||^2}{4} \ge (||\Psi\phi_1||^2 - ||\Psi^*\phi_1||||\phi_2|| + \frac{||\phi_2||^2}{4}) + (||\widehat{\Psi}\phi_2||^2 - ||\phi_1||||\widehat{\Psi}\phi_2|| + \frac{||\phi_1||^2}{4}) \ge 0.$ Thus operator D is positively defined.⁴

We continue to proceed in the finite-dimensional case (to escape the problem of existence of σ -additive Gaussian measure on infinite-dimensional space).

For the Gaussian measure with covariance operator (37), we have: $\langle f_{A_1} \rangle = \langle \widehat{A}_1 \rangle_{\Psi} - \text{Tr}\widehat{A}_1/4, \langle \overline{f}_{A_2} \rangle = \langle \widehat{A}_2 \rangle_{\Psi} - \text{Tr}\widehat{A}_2/4$. These relations for averages together with relation for correlations (36) provide coupling between PCSFT and QM.

In the infinite-dimensional case Gaussian distribution with he covariance operator given by (37) is not σ -additive.

To make it σ -additive one should consider a rigged Hilbert space: $\mathbf{H}_+ \subset \mathbf{H} \subset \mathbf{H}_-$, where $H = H_1 \times H_2$, and both embedding operators are of the Hilbert-Schmidt class.

⁴Of course, the same effect can be approached by adding αI for $\alpha \geq 1/4$.

Proposition 4. For any normalized vector $\Psi \in H_1 \otimes H_2$, the operator (37) determines the σ -additive Gaussian distribution on \mathbf{H}_- or equivalently the random field $\phi(\omega)$ valued in \mathbf{H}_- . For trace class operators $\widehat{A}_i : H_i \to H_i, i = 1, 2$, equalities (36) and (11) take place.

To prove this proposition, one should repeat the previous proofs, existence of traces is based on the trace class condition for of operators A_i (and not the trace class feature of the covariance operator of a Gaussian measure). The crucial difference with the finite dimensional case is that the prequantum random field takes values not in the Cartesian product $H = H_1 \otimes H_2$, but in its Hilbert-Schmidt extension. For mathematical details, I would like to recommend the excellent short book of A. V. Skorohod [45], see also [46]-[48] for applications to mathematical physics.

6 Classical (Hilbert valued) stochastic process corresponding to Schrödinger's evolution

We again start our considerations by considering the finite-dimensional case. Since we do not try to go beyond QM, but only reproduce its predictions, we use Schrödinger's equation for dynamics of the "wave function"⁵. We only change the interpretation of the Ψ -function of the composite system. Thus we start with Schrödinger's equation for a composite system $S = (S_1, S_2)$:

$$i\frac{d\Psi}{dt}(t) = \widehat{H}\Psi(t), \ \Psi(0) = \Psi_0, \tag{38}$$

where \hat{H} is Hamiltonian of S.

Hence, at the instant t, the covariance matrix of the prequantum random field (vector in the finite-dimensional case) $\phi(t, \omega)$ has the form:

$$D(t) = \begin{pmatrix} (\widehat{\Psi(t)}\widehat{\Psi(t)}^* + I_1/4) & \widehat{\Psi(t)} \\ \widehat{\Psi(t)}^* & (\widehat{\Psi(t)}^*\widehat{\Psi(t)} + I_2/4) \end{pmatrix}$$
(39)

The following fundamental question (having both mathematical and physical counterparts) immediately arises:

"Can one construct a stochastic process (valued in the Cartesian product $H_1 \times H_2$) such that at each $t \in [0, \infty)$ its covariance matrix coincides with D(t)?"

6.1 Bernoulli type process

The formal mathematical answer is yes! It is easy to construct such a stochastic process. Take space $\Omega = \prod_{t \in [0,\infty)} H_1 \times H_2$ as the space of random parameters, points of this space $\omega = (\omega_t)$ can be considered as functions $\omega : [0, \infty) \to H_1 \times H_2$, trajectories. Consider the family of Gaussian measures p_t on $H_1 \times H_2$ having zero mean value and covariance operators D_t , $t \in [0, \infty)$. Consider now (on Ω) the direct product of these measures, $P = \prod_{t \in [0,\infty)} p_t$.

Proposition 5. Let $\phi(t, \omega)$ be a stochastic process having the probability distribution P on Ω . Then, for any pair of vectors $y_1, y_2 \in H_1 \times H_2$

 $^{{}^{5}}$ So, we do not try to modify this equation, cf. [17]

and any instant of time $t \geq 0$,

$$E\langle y_1, \phi(t,\omega) \rangle \langle \phi(t,\omega), y_2 \rangle = \langle D(t)y_1, y_2 \rangle, \tag{40}$$

where D(t) is given by (40).

Existence of this stochastic process is a consequence of famous Kolmogorov's theorem. The equality (40) is a consequence of the definition of probability P on Ω .

Thus there exists a prequantum classical stochastic process inducing the Schrödinger evolution for any composite system prepared initially in a pure state. One may say that, for a composite system, Schrödinger's equation describes dynamics of the nondiagonal block of the covariance matrix of such a prequantum stochastic process.

This story becomes essentially more complicated after the remark that such a *prequantum process is not uniquely determined* by the D(t)! To determine uniquely a Gaussian process (up to natural equivalence), one should define not only covariance for each instant of time, i.e,

 $E\langle y_1, \phi(t,\omega) \rangle \langle \phi(t,\omega), y_2 \rangle$, but so called *covariance kernel* D(t,s):

$$E\langle y_1, \phi(t,\omega) \rangle, \langle \phi(s,\omega), y_2 \rangle = \langle D(t,s)y_1, y_2 \rangle.$$

However, the formalism of QM does not provide such a possibility. It is a consequence of the trivial fact (but of the great importance, cf. von Neumann [10]) that Schrödinger's equation for a composite system is dynamics with respect to a *single time parameter* t, common for both subsystems, and not with respect to a pair of time parameters (t, s) corresponding to internal times of subsystems.

Nevertheless, one may feel that the process existing due to Proposition 5 is not adequate to the real physical situation. Since its probability distribution P is the direct product of probabilities corresponding to different instances of time, it is the *Bernoulli process*. Its value at the instance of time t is totally independent from the previous behavior. Although this process provides right averages for each instance of time, it is hard to believe that real physical dynamics of e.g. an electron is of the Bernoulli-type (and for any Hamiltonian \hat{H}). We are looking for more realistic stochastic processes.

6.2 Stochastic (local) dynamics in the absence of interaction

We restrict our consideration to dynamics in the absence of interactions between S_1 and S_2 after the preparation procedure. Thus we are interested in propagation of initially correlated random fields (vectors in the finitedimensional case). Although it is a rather special dynamics, it plays an important role in quantum foundations. In particular, it describes the evolution of entanglement in the EPR-Bohm type experiments. Thus we consider Hamiltonian

$$\widehat{H} = \widehat{H_1} \otimes I_2 + I_1 \otimes \widehat{H_2},\tag{41}$$

where \widehat{H}_j is Hamiltonian of S_j (here we use QM terminology).

Lemma 7. Let Hamiltonian have the form (41). Then

$$\widehat{\Psi(t)} = e^{-i\widehat{H}_1 t} \widehat{\Psi_0} e^{-i\widehat{H}_2 t}.$$
(42)

$$\widehat{\Psi(t)}^* = e^{i\widehat{H_2}t}\widehat{\Psi_0}^* e^{-i\widehat{H_1}t}.$$
(43)

 $\mathbf{Proof.}$ In this case

$$\Psi(t) = e^{-it(\widehat{H_1} \otimes I_2 + I_1 \otimes \widehat{H_2})} \Psi_0.$$

We expand the initial state $\Psi_0: \Psi_0 = \sum_{ij}^k \psi_{ij} e_i \otimes f_j$. Then

$$\Psi(t) = \sum_{ij} \psi_{ij} e^{-i\widehat{H}_1 t} e_j \otimes e^{-i\widehat{H}_2 t} f_j$$

Thus, for $v \in H_2$, we get

$$\widehat{\Psi(t)}v = \sum_{ij}^{k} \psi_{ij} \langle v, e^{i\widehat{H_2}t} \bar{f}_j \rangle e^{-i\widehat{H_1}t} e_j = e^{-i\widehat{H_1}t} [\sum_{ij}^{k} \psi_{ij} \langle e^{-i\widehat{H_2}t}v, \bar{f}_j \rangle e_j].$$

By using Lemma 7 we prove: Lemma 8. Let the condition of Lemma 7 hold. Then

$$\widehat{\Psi(t)}\widehat{\Psi(t)}^* = e^{-i\widehat{H}_1 t}\widehat{\Psi_0}\widehat{\Psi_0}^* e^{i\widehat{H}_1 t}.$$
(44)

$$\widehat{\Psi(t)}^* \widehat{\Psi(t)} = e^{i\widehat{H_2}t} \widehat{\Psi_0}^* \widehat{\Psi_0} e^{-i\widehat{H_2}t}.$$
(45)

Finally, we obtain:

Lemma 9. Let the condition of Lemma 7 hold. Then the operator D(t) given by (39) can be represented in the form:

$$D(t) = \begin{pmatrix} (e^{-i\widehat{H_1}t}\widehat{\Psi_0}\widehat{\Psi_0}^*e^{i\widehat{H_1}t} + I/4) & e^{-i\widehat{H_1}t}\widehat{\Psi_0}e^{-i\widehat{H_2}t} \\ e^{i\widehat{H_2}t}\widehat{\Psi_0}^*e^{i\widehat{H_1}t} & (e^{i\widehat{H_2}t}\widehat{\Psi_0}^*\widehat{\Psi_0}e^{-i\widehat{H_2}t} + I/4) \end{pmatrix}$$
(46)

By using this representation it is easy to prove:

Proposition 6. Let the condition of Lemma 7 hold. Then the operator D(t) given by (39) is the covariance operator (for each instance of time t) of the vector process with coordinates

$$\phi_1(t,\omega) = e^{-i\overline{H_1}t}\xi_{01}(\omega), \phi_2(t,\omega) = e^{i\overline{H_2}t}\xi_{02}(\omega), \tag{47}$$

where the initial random vector $\xi_0(\omega) = (\xi_{01}(\omega), \xi_{02}(\omega))$ is Gaussian with zero mean value and the covariance operator D(0).

Proof. We will find not only the covariance operator for a fixed instant of time, but even the covariance kernel. We have, for any pair of vectors $u, w \in H_1$,

$$\begin{split} E\langle u,\phi_1(t,\omega)\rangle\langle\phi_1(s,\omega),w\rangle &= E\langle e^{i\widehat{H_1}t}u,\xi_{01}(\omega)\rangle\langle\xi_{01}(\omega)\rangle, e^{i\widehat{H_1}s}w\rangle\\ &= \langle e^{-i\widehat{H_k}s}(\widehat{\Psi_0}^*\widehat{\Psi_0}+I_1/4)e^{i\widehat{H_1}t}u,w\rangle. \end{split}$$

The same calculations can be done for the second diagonal block. Thus diagonal blocks of the covariance operator of the stochastic process given by (47) coincide with diagonal blocks of the operator D(t). We now consider nondiagonal blocks. Let now $u \in H_1, v \in H_2$. We have:

$$E\langle u, \phi_1(t,\omega) \rangle \langle \phi_2(s,\omega), v \rangle = E\langle e^{i\widehat{H_1}t}u, \xi_{01}(\omega) \rangle \langle \xi_{02}(\omega) \rangle, e^{-i\widehat{H_2}s}v \rangle$$
$$= \langle \widehat{\Psi_0}^* e^{i\widehat{H_1}t}u, e^{-i\widehat{H_2}s}v \rangle = \langle e^{i\widehat{H_2}s}\widehat{\Psi_0}^* e^{i\widehat{H_1}t}u, v \rangle.$$

The same calculations can be done for the second nondiagonal block. Thus the covariance kernel has the form D(t, s) =

$$\begin{pmatrix} e^{-i\widehat{H_1}s}\widehat{\Psi_0}\widehat{\Psi_0}^*e^{i\widehat{H_1}t} + e^{i\widehat{H_1}(t-s)}I/4 & e^{-i\widehat{H_1}s}\widehat{\Psi_0}e^{i\widehat{-H_2}t} \\ e^{i\widehat{H_2}s}\widehat{\Psi_0}^*e^{i\widehat{H_1}t} & e^{i\widehat{H_2}s}\widehat{\Psi_0}^*\widehat{\Psi_0}e^{-i\widehat{H_2}t} + e^{-i\widehat{H_2}(t-s)}I/4 \end{pmatrix}$$

$$(48)$$

And hence, for t = s, D(t, t) = D(t).

We emphasize that dynamics (47) by itself is purely deterministic, stochasticity is generated by initial conditions. One might say that this process describes propagation of uncertainty of preparation.

We remark that Propositions 5 and 6 provide two different stochastic processes. The covariance kernel (48) differs from the covariance kernel of the process which has been constructed by considering the product of Gaussian distributions p_t . The latter has the covariance kernel

$$D(t,s) = D(t)\delta(t-s).$$

For this process, its realization at different instants of time are independent. The process defined by (47) contains nontrivial dependence between its realizations at different times. I think that it is closer to the real physical situation.

The following interesting problem arises:

To construct a stochastic process for an arbitrary Hamiltonian, such that in the case of the absence of interactions this construction gives the process (47).

At the moment I am not able to solve this problem.

6.3 Stochastic nonlocal dynamics

We now consider another classical stochastic process reproducing dynamics of quantum correlations.

Proposition 7. Let operator D(t) be defined by (39). Then the stochastic process

$$\xi(t,\omega) = \sqrt{D(t)}\eta_0(\omega), \qquad (49)$$

where $\eta_0(\omega) \in H_1 \times H_2$ is distributed N(0, I), has the covariance operator D(t) for any $t \ge 0$.

Proof. Let $y_1, y_2 \in H_1 \times H_2$. Then

$$E\langle y_1, \xi(t,\omega)\rangle\langle \xi(t,\omega), y_2\rangle = E\langle \sqrt{D(t)}y_1, \eta_0(\omega)\rangle\langle \eta_0(\omega),$$

$$\sqrt{D(t)}y_2\rangle = \langle \sqrt{D(t)}y_1, \sqrt{D(t)}y_2\rangle = \langle D(t)y_1, y_2\rangle.$$

It is clear that the covariance kernel is given by

$$D(t,s) = \sqrt{D(s)D(t)}.$$
(50)

We remind that, for Hamiltonian without interaction, we constructed the stochastic process $\phi(t, \omega)$ given by (47). In general stochastic processes, $\xi(t, \omega)$ and $\phi(t, \omega)$ given by (49) and (47) do not coincide:

We can write the process (47) as $\phi(t) = V(t)\sqrt{D(0)}\eta_0$, where $V(t) = \text{diag}(e^{-it\hat{H}_1}, e^{it\hat{H}_2})$. Hence,

$$D_{\phi}(t,s) = V(s)D(0)V(t)^*.$$

On the other hand, the covariance kernel of process (49) is given by $D_{\xi}(t,s) = \sqrt{D(s)D(t)}$. We remark that $D(t) = V(t)D(0)V(t)^*$. Hence, $\sqrt{D(t)} = V(t)\sqrt{D(0)}V(t)^*$. Thus

$$D_{\xi}(t,s) = V(s)\sqrt{D(0)}V^{*}(s)V(t)\sqrt{D(0)}V^{*}(t).$$

We remark that $V^*(s)V(t) \neq I, t \neq s$.

^

The process $\xi(t,\omega)$ is nonlocal in the following sense. Its component $\xi_1(t,\omega)$ is guided not only by the Hamiltonian of S_1 , but also of S_2 ; the same is valid for $\xi_2(t,\omega)$. Thus PCSFT (at least at the moment) cannot provide a definite answer on locality of the prequantum world. Quantum correlations can be produced by local as well as nonlocal prequantum stochastic processes.

Proposition 8. In the case of Hamiltonian without interaction, see (41), the stochastic process (49) can be represented in the form:

$$\xi_1(t) = e^{-it\hat{H}_1} Q_{11}^0 e^{it\hat{H}_1} \eta_{01} + e^{-it\hat{H}_1} Q_{12}^0 e^{-it\hat{H}_2} \eta_{02}, \qquad (51)$$

~

$$\xi_2(t) = e^{itH_2} Q_{21}^0 e^{itH_1} \eta_{01} + e^{itH_2} Q_{22}^0 e^{-itH_2} \eta_{02}, \tag{52}$$

where

$$\sqrt{D(0)} = \begin{pmatrix} Q_{11}^0 & Q_{12}^0 \\ Q_{21}^0 & Q_{22}^0 \end{pmatrix},$$

and $\eta_0(\omega) \in N(0, I)$.

Proof. For example, take $y_1, y_2 \in H_1$ and consider average

$$E \langle y_{1}, \xi(t, \omega) \rangle \langle \xi(s, \omega), y_{2} \rangle$$

$$= E \langle e^{-it\hat{H}_{1}} Q_{11}^{0} e^{it\hat{H}_{1}} y_{1}, \eta_{02}(\omega) \rangle \langle \eta_{01}(\omega), e^{-is\hat{H}_{1}} Q_{11}^{0} e^{is\hat{H}_{1}} y_{2} \rangle$$

$$+ E \langle e^{it\hat{H}_{2}} (Q_{12}^{0})^{*} e^{it\hat{H}_{1}} y_{1}, \eta_{02}(\omega) \rangle \langle \eta_{02}(\omega), e^{is\hat{H}_{2}} (Q_{12}^{0})^{*} e^{is\hat{H}_{1}} y_{2} \rangle$$

$$+ E \langle e^{-it\hat{H}_{1}} Q_{11}^{0} e^{it\hat{H}_{1}} y_{1}, \eta_{10}(\omega) \rangle \langle \eta_{20}(\omega), e^{-is\hat{H}_{2}} (Q_{12}^{0})^{*} e^{is\hat{H}_{1}} y_{2} \rangle$$

$$+ E \langle e^{it\hat{H}_{2}} (Q_{12}^{0})^{*} e^{it\hat{H}_{1}} y_{1}, \eta_{10}(\omega) \rangle \langle \eta_{20}(\omega), e^{-is\hat{H}_{1}} Q_{11}^{0} e^{is\hat{H}_{1}} y_{2} \rangle.$$

Two last terms are equal to zero, since

 $E\langle z_1, \eta_{10}(\omega) \rangle \langle \eta_{20}(\omega), z_2 \rangle = 0$

for any pair $z_1 \in H_1, z_2 \in H_2$. The first two give us

$$\langle e^{-it\hat{H}_1} Q^0_{11} e^{it\hat{H}_1} y_1, e^{-is\hat{H}_1} Q^0_{11} e^{is\hat{H}_1} y_2 \rangle$$

+ $\langle e^{it\hat{H}_2} (Q^0_{12})^* e^{it\hat{H}_1} y_1, e^{is\hat{H}_2} (Q^0_{12})^* e^{is\hat{H}_1} y_2 \rangle$

Thus

$$D_{11}(t,s) = e^{-is\hat{H}_1} Q_{11}^0 e^{i(s-t)\hat{H}_1} Q_{11}^0 e^{it\hat{H}_1} + e^{-is\hat{H}_1} Q_{12}^0 e^{i(t-s)\hat{H}_2} (Q_{12}^0)^* e^{it\hat{H}_1}.$$

Representation (51), (52) implies that even in the absence of interaction

between the subsystems S_1 and S_2 of the system S the dynamics of S_1 depends on the Hamiltonian \hat{H}_2 and vice versa. It can be interpreted as a sign of "action at the distance". Thus the stochastic process $\xi(t)$ can be considered as "nonlocal" – opposite to the process $\phi(t)$ given by (47).

6.4 Infinite-dimensional case

To proceed in the infinite-dimensional case, one should consider a rigged Hilbert space: $\mathbf{H}_+ \subset \mathbf{H} \subset \mathbf{H}_-$, where $H = H_1 \times H_2$, and both embedding operators are of the Hilbert-Schmidt class. Stochastic processes take values in the Hilbert space H_- (and not $H = H_1 \times H_2$). All previous results are valid for any unitary dynamics $\Psi(t) = U(t)\Psi_0$.

This paper was written under support of the grant "Mathematical Modeling" of Växjö university and the grant QBIC of Tokyo University of Science. It was presented at "Feynman Festival", June, 2009; the author would like to thank Vladimir Manko for his critical comments which improved understanding of the model.

References

- Khrennikov, A.(ed): Foundations of Probability and Physics. Series PQ-QP: Quantum Probability and White Noise Analysis 13. WSP, Singapore (2001)
- [2] Khrennikov, A.(ed): Quantum Theory: Reconsideration of Foundations. Ser. Math. Model. 2, Växjö University Press, Växjö (2002); electronic volume: http://www.vxu.se/msi/forskn/publications.html
- [3] Adenier, G., Khrennikov, A. and Nieuwenhuizen, Th.M. (eds.): Quantum Theory: Reconsideration of Foundations-3. American Institute of Physics, Ser. Conference Proceedings 810, Melville, NY (2006)
- [4] Adenier, G., Fuchs, C. and Khrennikov, A.(eds): Foundations of Probability and Physics-3. American Institute of Physics, Ser. Conference Proceedings 889, Melville, NY (2007)
- [5] Accardi, L., G. Adenier, C.A. Fuchs, G. Jaeger, A. Yu. Khrennikov, J.-A. Larsson, S. Stenholm (eds.): Foundations of Probability and Physics-5, American Institute of Physics, Ser. Conference Proceedings, 1101, Melville, NY (2009)
- [6] Hess, K., Michielsen, K., De Raedt, H.: Possible experience: From Boole to Bell. *EPL*, 87, 60007 (2009).
- [7] Hess, K.: Modeling experiments using quantum and Kolmogorov probability. J. of Physics: Condensed Matter, 20, 454207 (2009).
- [8] A. Yu. Khrennikov, Interpretations of Probability. De Gruyter, Berlin (2009), second edition (completed).
- Kolmogoroff, A. N.: Grundbegriffe der Wahrscheinlichkeitsrechnung. Springer Verlag, Berlin (1933); English translation: Kolmogorov, A.N.: Foundations of the Probability Theory. Chelsea Publishing Company, New York (1956)
- [10] J. von Neumann, Mathematical Foundations of Quantum Mechanics, Princeton Univ. Press, Princeton, N.J., 1955.
- [11] P. Busch, M. Grabowski, P. Lahti, Operational Quantum Physics, Springer Verlag, Berlin, 1995.
- [12] M. Ozawa, Conditional probability and a posteriori states in quantum mechanics, *Publ. Res. Inst. Math. Sci.* 21, 279-295 (1985).
- [13] A. Yu. Khrennikov, Contextual approach to quantum formalism, Springer, Berlin-Heidelberg-New York, 2009.

- [14] A. Yu. Khrennikov, Einstein's dream. Proceedings of Conference The nature of light: What are photons? C. Roychoudhuri, A. F. Kracklauer, K. Creath. Proceedings of SPIE, 6664, 2007, 666409-1 – 666409-9.
- [15] A. Yu. Khrennikov, "Prequantum classical statistical model with infinite dimensional phase-space," J. Phys. A: Math. Gen. 38, pp. 9051-9073, 2005.
- [16] A. Yu. Khrennikov, "Generalizations of quantum mechanics induced by classical statistical field theory," *Found. Phys. Letters* 18, pp. 637-650, 2005.
- [17] A. Yu. Khrennikov, "Nonlinear Schrödinger equations from prequantum classical statistical field theory," *Physics Letters A* 357, pp. 171-176, 2006.
- [18] A. Yu. Khrennikov, "Prequantum classical statistical field theory: Complex representation, Hamilton-Schrödinger equation, and interpretation of stationary states," *Found. Phys. Lett.* **19**, pp. 299-319, 2006.
- [19] A. Yu. Khrennikov, "On the problem of hidden variables for quantum field theory," *Nuovo Cimento* B 121, pp. 505-515, 2006.
- [20] A. Yu. Khrennikov, Born's rule from classical random fields, *Physics Letters A*, **372**, N 44, 6588-6592 (2008).
- [21] De la Pena, L. and Cetto, A. M.: The quantum dice: An introduction to stochastic electrodynamics. Kluwer, Dordrecht (1996)
- [22] A. Casado, T. Marshall, E. Santos, J. Opt. Soc. Am. B 14, pp. 494-205, 1997.
- [23] G. Brida, M. Genovese, M. Gramegna, C. Novero and E. Predazzi, *Phys. Lett* A **299**, pp. 121-141, 2002.
- [24] Nieuwenhuizen, Th. M.: Classical phase space density for relativistic electron. In: Adenier, G., Khrennikov, A. and Nieuwenhuizen, Th.M. (eds.) Quantum Theory: Reconsideration of Foundations-3. American Institute of Physics, Ser. Conference Proceedings, vol. 810, pp. 198-210. Melville, NY (2006)
- [25] Boyer, T. H.: A brief survey of stochastic electrodynamics. In: Barut, A. O. (ed) Foundations of Radiation Theory and Quantum Electrodynamics, pp. 141-162. Plenum, New York (1980)
- [26] Scully, M. O. and Zubairy, M. S.: Quantum optics. Cambridge University Press, Cambridge (1997)
- [27] Louisell, H. H.: Quantum statistical properties of radiation. J. Wiley, New York (1973)
- [28] Mandel, L. and Wolf, E.: Optical coherence and quantum optics. Cambridge University Press, Cambridge (1995)
- [29] Nelson, E: Quantum fluctuation Princeton Univ. Press, Princeton (1985)
- [30] Davidson, M.: J. Math. Phys. 20, 1865-1870 (1979)
- [31] Davidson, M.: Stochastic models of quantum mechanics a perspective. In: Adenier, G., Fuchs, C. and Khrennikov, A. (eds.) Foundations of Probability and Physics-4. American Institute of Physics, Ser. Conference Proceedings, vol. 889, pp. 106–119. Melville, NY (2007)

- [32] 't Hooft, G.: Quantum mechanics and determinism. hep-th/0105105 (2001)
- [33] 't Hooft, G.: Determinism beneath quantum mechanics. quant-ph/0212095 (2002)
- [34] 't Hooft, G.: The free-will postulate in quantum mechanics. quant-ph/0701097 (2007)
- [35] Elze, T.: The attractor and the quantum states. arXiv: 0806.3408 (2008)
- [36] Rusov, V. D. , Vlasenko, D. S., and Mavrodiev, S.Cht.: Quantization in classical mechanics and reality of Bohm's psi-field. arXiv:0906.1723 (2009)
- [37] Kisil, V.: A quantum-classical brackets from p-mechanics. Europhys. Lett. 72 (6), 873-?879 (2005)
- [38] V. I. Manko, J. of Russian Laser Research, 17, 579-584 (1996).
- [39] V. I. Manko and E. V. Shchukin, J. Russian Laser Research, 22, 545-560 (2001).
- [40] M. A. Manko, V. I. Manko, R. V. Mendes, J. Russian Laser Research, 27, 507-532.
- [41] S. De Nicola, R. Fedele, M. A. Man'ko and V. I. Man'ko, Quantum tomography, wave packets, and solitons. J. of Russian Laser Research, 25, 1071-2836, 2004.
- [42] O. V. Manko and V. I. Manko, J. Russian Laser Research, 25, 477-492 (2004).
- [43] A. Khrennikov, Entanglement's dynamics from classical stochastic process. *Europhysics Letters*, 88, 40005.1-6 (2009).
- [44] A. Khrennikov, To quantum averages through asymptotic expansion of classical averages on infinite-dimensional space. J. Math. Phys., 48 (1), Art. No. 013512 (2007).
- [45] A. V. Skorohod, Integration in Hilbert space. Springer-Verlag, Berlin, 1974.
- [46] A. L. Daletski and S. V. Fomin, Measures and differential equations in infinite-dimensional spaces. Kluwer, Dordrecht, 1991.
- [47] S. Albeverio, and M. Röckner, Prob. Theory and Related Fields 89, 347 (1991).
- [48] S. Albeverio, R. Hoegh-Krohn, Phys. Lett. B 177, 175 (1989).