

A model for sexed coagulation

Raoul Normand*

*Laboratoire de Probabilités et Modèles Aléatoires, UPMC,
175 rue du Chevaleret, 75013 Paris, France*

Abstract

We consider in this work a model for aggregation, where the coalescing particles initially have a certain number of potential links (called arms) which are used to perform coagulations. This model is sexed, in the sense that there are male and female arms: two particles may coagulate only if one has an available male arm, and the other has an available female arm. After a coagulation, the used arms are no longer available. We are interested in the concentrations of the different types of particles, which are governed by a modification of Smoluchowski's coagulation equation — that is, an infinite system of nonlinear differential equations. Using generating functions and solving a nonlinear PDE, we show that, up to some critical time, there is a unique solution to this equation. The Lagrange Inversion Formula allows in some cases to obtain explicit solutions, and to relate our model to two recent models for limited aggregation. We also show that, whenever the critical time is infinite, the concentrations converge to a state where all arms have disappeared, and the distribution of the masses is related to the law of the size of some two-type Galton-Watson tree. Finally, we consider a microscopic model for coagulation: we construct a sequence of Marcus-Luschnikov processes, and show that it converges, before the critical time, to the solution of our modified Smoluchowski's equation.

MSC: Primary 34A34, 60K35; Secondary 60B12, 82C23, 82D60

Keywords: Coagulation equations; Gelation; Generating functions; Method of characteristics; Marcus-Luschnikov Process; Hydrodynamic limit

1 Introduction

In this work, we study a model for coagulation of particles, generalizing the original model of Smoluchowski [24], and a recent model of Bertoin [2]. We consider particles which are initially given a certain number of male and female arms. These arms are used to perform the coagulations: two particles coagulate when a male arm of one and a female arm of another bind. This can be used to model the formation of polymers. For instance, consider male particles (which have only male arms), and female particles. Then a coagulation between a male and a female particle can be thought of as an ionic bond between a cation and an anion.

In our model, a particle is characterised by a triple (a, b, m) , $a \in \mathbb{N}$ being its number of male arms, $b \in \mathbb{N}$ its number of female arms, and $m \in \mathbb{N}^*$ its mass. Two particles may coagulate

*E-mail address: raoul.normand@upmc.fr.

when one has an available female arm and the other has an available male arm, and when a coagulation occurs, the used arms disappear. Hence, we may only observe the transition

$$\{(a, b, m), (a', b', m')\} \rightarrow (a + a' - 1, b + b' - 1, m + m').$$

We will assume that this transition occurs with a rate given by the number of pairs formed of a female arm and of a male arm, that is $a'b + ab'$. We wish to study how the concentration of each type of particle evolves when time passes. The precise mathematical formulation is given in Section 2.

This model is a modification of the well-known model of Smoluchowski [24]. Recall that Smoluchowski's coagulation equations [24] describe the evolution of the concentrations of particles in a medium, where particles are characterised only by their masses. When two particles of masses m and m' coagulate, they merge into a single particle of mass $m+m'$. Such a coagulation occurs with rate $\kappa(m, m')$, where κ is some symmetric nonnegative kernel. In Smoluchowski's original model, the mass are assumed to be positive integers. The concentration $c_t(m)$ of particles of mass m is governed by the following infinite system of nonlinear differential equations

$$\frac{d}{dt}c_t(m) = \frac{1}{2} \sum_{m'=1}^{m-1} c_t(m')c_t(m-m')\kappa(m', m-m') - c_t(m) \sum_{m'=1}^{+\infty} c_t(m')\kappa(m, m'),$$

for $m \in \mathbb{N}^*$. The first term accounts for the creation of particles of mass m by coagulation of particles of mass m' and $m-m'$; the second for disappearance of particles of mass m by coagulation with other particles.

For general kernels κ , explicit solutions are not known. However, some have been obtained in different cases, notably whenever the kernel is constant [24], additive [12] or multiplicative [19]. In the multiplicative case, solutions are obtained up to a critical time, known as the gelation time. This is interpreted as the time when a particle of infinite mass appears. It absorbs some of the particles, and the total mass starts to decrease.

Smoluchowski's equation (and some variations) have been extensively studied, both from an analytical (e.g. [5], [9], [14]) and a probabilistic point of view (e.g. [6], [15], [17], [23], and see also the review by Aldous [1]). In general, little is known after the gelation time, and most results are obtained before the gelation time (see however [10], [11]). The existence and uniqueness of a solution before gelation has been obtained only in 1999 by Norris ([23]), under the assumption that κ is sublinear, i. e. $\kappa(n, m)/(nm)$ is bounded.

From a probabilistic point of view, some microscopic models have been studied, beginning with Marcus [22] and Lushnikov [20]. Heuristically, one considers a finite number of particles, and each couple of particles with masses m and m' coalesces with rate $\kappa(m, m')$. After suitable change of time and renormalization, one expects this system to converge to a solution of Smoluchowski's equation. This has been shown by Jeon in 1998 [17] (up to extraction of a subsequence), provided the rate is strictly sublinear (i.e. $\kappa(n, m)/n \rightarrow 0$ when $n \rightarrow +\infty$). In particular, there is no gelation in this case. Norris [23] extended his results one year later by showing the convergence of the model before gelation, whenever the rate is sublinear. Other points of view are also considered; e.g. in [15], the authors show that coagulating Brownian particles form clusters whose size evolves according to Smoluchowski's equation.

An interesting question is to deal with the case when the coagulations are restricted by some device. Typically, one may think of covalent bonds: a given atom can only perform a given number of bonds. In this direction, Bertoin [2] studied two models where a particle is characterised by its number of arms and by its mass, and it uses its arms to perform aggregations.

The concentrations of each type of particle is governed by a modification of Smoluchowski's equation. In [2], he obtains solution up to some time T , and shows that whenever gelation does not occur (i.e. $T = +\infty$), there is a limit state where all the arms have disappeared: the concentrations converge to limiting concentrations which bear a striking resemblance with the law of the size of some Galton-Watson tree. This fact is explained in [3] and [4]. It is also worth noticing that Bertoin's model can be related to Smoluchowski's for the constant, additive and multiplicative kernels. We will also see that our model contains Bertoin's: the oriented model corresponds indeed to the sexed model if each particle is given precisely one female arm, and the symmetric model corresponds to the sexed one if the particles are given a gender uniformly at random.

This paper is divided in two parts. In the first one (Sections 2 to 5), we shall study the sexed Smoluchowski's equation, which is an infinite system of nonlinear differential equations. We first (Section 2) introduce the problem, and prove some physically intuitive facts. Then (Section 3), we prove our main result: up to some critical time, there exists a unique solution to the system, and its moment generating function can be expressed explicitly in terms of the initial data. The tools used are analogous to those in [2], but since we are dealing with a two-dimensional problem, several technical issues need to be addressed. The outline of the proof is as follows. First, we transform the system into a PDE problem by considering the generating functions of the concentrations. This PDE is not quasilinear, but it may however be solved by the method of characteristics. This method requires the inversion of a two-dimensional mapping, and this can be done precisely up to the critical time. Unfortunately, even for monodisperse initial conditions (i.e. there are only particles of mass 1 at time 0), the inversion is not explicit (one could use the two-variable Lagrange inversion formula, but in general, the expression it provides is too cumbersome). Nonetheless, in some specific cases (Section 4), the Lagrange Inversion Formula yields explicit results. In particular, we recover the solutions obtained in [2]. Finally, we show (Section 5) that there exist limiting concentrations when $t \rightarrow +\infty$, and that they are related to the distribution of the total progeny of some two-type Galton-Watson process.

In the second part (Section 6), we study a microscopic model. Given a finite number of particles, we let them coagulate and observe the evolution of the concentrations of the different types of particles. This is a Marcus-Luschnikov process, and we show that it converges, before the critical time, to a process solving Smoluchowski's equation (1). As pointed out earlier, this kind of convergence had already been proved by Norris ([23], see as well [17]). The difference here is that we consider a model with male or female arms. Moreover, the proof is made much easier by the fact that the rate of coagulation is explicit. In particular, we will appeal to the PDE obtained in the first part. This discrete model provides a justification to the sexed Smoluchowski's equation (1).

Finally, note that our construction can also provide a model for random oriented graphs, since a coagulation can be seen as the creation of an oriented edge between two vertices in a graph, whose orientation is given e.g. from the male arm to the female arm. Hence, we can consider a large number n of particles and let them coagulate. When all the coagulations are performed, we obtain a set of oriented graphs. When $n \rightarrow +\infty$, we may wonder what the distribution of their sizes is, what a typical graph looks like, etc. A heuristic answer, motivated by the works [3], [4], and by the results obtained in this paper (Section 5), is that a typical graph would be a two-type Galton-Watson tree (with the convention of orientation above), provided there are few arms (with the notations of this paper, this means $T_c = +\infty$ and μ is not degenerate). This configuration model is to be investigated in another article.

2 Setting and results

2.1 Notations

Let us first introduce some notations and Smoluchowski's equation, and state our main result.

- $\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{N}^* = \{1, 2, \dots\}$.
- $S = \mathbb{N} \times \mathbb{N} \times \mathbb{N}^*$ is the set of the different types of particles. A generic element of S will be denoted by p , and if $p = (a, b, m)$, we will call a p -particle a particle with a male arms, b female arms, and mass m .
- For $p = (a, b, m) \in S$ and $p' = (a', b', m') \in S$, we will denote

$$p.p' = a'b + ab'$$

the *rate* of coagulation and

$$p \circ p' = (a + a' - 1, b + b' - 1, m + m')$$

the type of the particle resulting from such a coagulation. We say that $p' \preceq p$ if $a' \leq a + 1$, $b' \leq b + 1$ and $m' \leq m - 1$. When $p' \preceq p$, we write

$$p \setminus p' = (a + 1 - a', b + 1 - b', m - m')$$

the type of particle such that $p' \circ (p \setminus p') = p$.

- For two functions $c, f : S \rightarrow \mathbb{R}$, we will denote, when the series converge absolutely,

$$\langle c, f \rangle := \sum_{p \in S} c(p)f(p).$$

When using this notation, we will write, with a slight abuse of notation, a for the function $(a, b, m) \mapsto a$, b for $(a, b, m) \mapsto b$, etc.

Let us recall our goal. We are interested in a system of coagulating particles with male and female arms. We assume that each couple formed of a p -particle and of a p' -particle coagulate with rate $p.p'$, to form a $p \circ p'$ -particle. This means that if we denote $c_t(p)$ the concentration of p -particles then $(c_t(p), p \in S)$ solves the following infinite system of nonlinear differential equations

$$\frac{d}{dt}c_t(p) = \frac{1}{2} \sum_{p' \preceq p} p'.(p \setminus p')c_t(p')c_t(p \setminus p') - c_t(p) \sum_{p' \in S} p.p'c_t(p'). \quad (1)$$

The first term accounts for the creation of p -particles by coagulation of p' - and $p \setminus p'$ -particles (the factor $1/2$ comes from an obvious symmetry). The second accounts for the disappearance of p -particles by coagulation with other particles. Let us once write down this formula explicitly. For all $(a, b, m) \in S$, the concentration of (a, b, m) -particles verifies

$$\begin{aligned} \frac{d}{dt}c_t(a, b, m) &= \frac{1}{2} \sum_{m'=1}^{m-1} \sum_{a'=0}^{a+1} \sum_{b'=0}^{b+1} (a'(b+1-b') + b'(a+1-a')) \times \\ &\quad c_t(a', b', m')c_t(a+1-a', b+1-b', m-m') \\ &\quad - \sum_{m' \geq 1} \sum_{a' \geq 0} \sum_{b' \geq 0} (ab' + a'b)c_t(a, b, m)c_t(a', b', m'). \end{aligned}$$

Let us now define what we call a *solution* to Smoluchowski's equation.

Definition 1. We call a family $((c_t(p))_{p \in S}, t \in [0, T])$ of differentiable functions a solution of Smoluchowski's equation (or of system (1)), if

1. For every $t \in [0, T]$, $\sum_{(a,b,m) \in S} (a+b)|c_t(a,b,m)| < +\infty$,
2. The family $(c_t(p))$ solves the system (1) for $t \in [0, T]$.

Remark 1. • We will always assume that at time 0, $\langle a+b+1, c_0 \rangle < +\infty$, and that the mean number of male arms $\langle a, c_0 \rangle$ and the mean number of female arms $\langle b, c_0 \rangle$ are equal. Physically, it is then obvious that they will remain equal as time passes. This shall be proven later on, in Lemma 3.

- It is easy to see that if $(c_t)_{t \in [0, T]}$ is a solution to (1) with initial conditions c_0 , and $\lambda > 0$, then $(\lambda c_t / \lambda^2)_{t \in [0, T]}$ is a solution to (1) with initial conditions λc_0 . Hence, it is enough to assume that $\langle a, c_0 \rangle = \langle b, c_0 \rangle = 1$, what will always be the case from now on.

2.2 Main result

Our main result is existence and uniqueness of a solution to (1) up to a critical time. In all the statements and proofs, we are given nonnegative initial concentrations c_0 such that $\langle 1, c_0 \rangle < +\infty$, $\langle a, c_0 \rangle = \langle b, c_0 \rangle = 1$ and $\langle a^2 + b^2, c_0 \rangle < +\infty$. We can then define the critical time T_c .

Definition 2. Let

$$M = \langle ab, c_0 \rangle + \sqrt{\langle a^2 - a, c_0 \rangle \langle b^2 - b, c_0 \rangle}$$

and

$$T_c = \begin{cases} +\infty & \text{if } M \leq 1 \\ \frac{1}{M-1} & \text{if } M > 1. \end{cases}$$

We will also constantly use the generating function of (c_0)

$$g_0(x, y, z) := \sum_{(a,b,m) \in S} c_0(a, b, m) x^a y^b z^m.$$

Since $\langle 1, c_0 \rangle < +\infty$, g_0 is well-defined on $[0, 1]^3$. Using the assumption $\langle a+b, c_0 \rangle = 1$, and e.g. monotone convergence, we also see that its partial derivatives with respect to x and y are well-defined and continuous on $[0, 1]^3$. For the same reason, they remain in $[0, 1]$. We shall prove the following result.

Theorem 1. (i) Smoluchowski's equation (1) with initial conditions c_0 has a unique solution (c_t) defined on $[0, T_c)$.

(ii) Let

$$\Gamma_r = \inf\{t \geq 0, \langle a^2 + b^2, c_t \rangle > r\} \quad \text{and} \quad \Gamma_\infty = \sup_{r>0} \Gamma_r. \quad (2)$$

Then $\Gamma_\infty = T_c$.

(iii) For $t \in [0, T_c)$ and $z \in [0, 1]$, the mapping $\phi_t(\cdot, \cdot, z)$, given for $(x, y) \in [0, 1]^2$ by

$$\phi_t(x, y, z) = \left((1+t)x - t \frac{\partial g_0}{\partial y}(x, y, z), (1+t)y - t \frac{\partial g_0}{\partial x}(x, y, z) \right),$$

has a right inverse $h_t = (h_t^{(1)}, h_t^{(2)})$ which is well-defined and analytic on $(0, 1)^2$. Then the generating function g_t of (c_t) is given by

$$g_t(x, y, z) = \frac{1}{1+t} \left(\tilde{H}_t^{(2)}(x, y, z) + \tilde{H}_t^{(1)}(0, y, z) \right) + G_t(z). \quad (3)$$

where for $t > 0$

$$\tilde{h}_t^{(1)} = \frac{1+t}{t} h_t^{(1)}(x, y, z) - \frac{x}{t} \quad ; \quad \tilde{h}_t^{(2)} := \frac{1+t}{t} h_t^{(2)}(x, y, z) - \frac{y}{t} \quad (4)$$

and

- $\tilde{H}_t^{(1)}$ is the antiderivative of $\tilde{h}_t^{(1)}$ with respect to y , vanishing at $y = 0$,
- $\tilde{H}_t^{(2)}$ is the antiderivative of $\tilde{h}_t^{(2)}$ with respect to x , vanishing at $x = 0$,
- $G_t(z)$ is the antiderivative of

$$\frac{\partial g_0}{\partial z} \left(h_t^{(1)}(0, 0, z), h_t^{(2)}(0, 0, z), z \right) \quad (5)$$

with respect to z , vanishing at 0.

(iv) The total mass $\langle m, c_t \rangle$ is constant on $[0, T_c)$.

2.3 Preliminary results

In this section, we give some physically intuitive results, and deduce the “weak” form of the equation. Let us start with the following lemma (recall that $c_t(p)$ is meant to model a concentration).

Lemma 1. *Any solution to Smoluchowski’s equation remains nonnegative, i.e. if $(c_t)_{t \in [0, T)}$ is a solution to (1), then for all $t \in [0, T)$ and $p \in S$, $c_t(p) \geq 0$.*

Proof. Take some $t \in [0, T)$. System (1) gives

$$\frac{d}{dt} c_t(a, b, 1) = -c_t(a, b, 1) \sum_{m' \geq 1} \sum_{a' \geq 0} \sum_{b' \geq 0} (ab' + a'b) c_t(a', b', m') := -\gamma(t) c_t(a, b, 1).$$

Let $G(t) = \int_0^t \gamma(s) ds$. Then $c_t(a, b, 1) = c_0(a, b, 1) e^{-G(t)}$, so it remains nonnegative. Let now $m \geq 1$, and suppose that the $c_t(a, b, m')$ are nonnegative for $a, b \geq 0$ and $1 \leq m' \leq m$. For some $p = (a, b, m+1)$, we have

$$\begin{aligned} \frac{d}{dt} c_t(p) &= \frac{1}{2} \sum_{p' \preceq p} p' \cdot (p \setminus p') c_t(p') c_t(p \setminus p') - c_t(p) \sum_{p' \in S} p \cdot p' c_t(p') \\ &= \beta(t) - c_t(p) \gamma(t). \end{aligned}$$

So we may write

$$c_t(p) = \left(c_0(p) + \int_0^t \beta(s) e^{G(s)} ds \right) e^{-G(t)}.$$

But $\beta(t) \geq 0$ since it is a linear combination with nonnegative coefficients of the $c_t(a, b, m')$ for $a, b \geq 0$ and $1 \leq m' \leq m$. So $c_t(a, b, m + 1)$ is nonnegative, what gives the result by induction. \square

The following two lemmas are straightforward generalizations of Lemma 1 and 2 in [2]. Note however that the monotone convergence used in the proofs requires that the coefficients (c_t) be nonnegative.

Lemma 2. (i) *If (c_t) is a solution to Smulochowski's equation (1), then $t \mapsto \langle 1, c_t \rangle$, $t \mapsto \langle a, c_t \rangle$ and $t \mapsto \langle b, c_t \rangle$ are decreasing.*

(ii) *A family (c_t) is a solution to (1) if and only if it solves*

$$\frac{d}{dt} \langle c_t, f \rangle = \frac{1}{2} \sum_{p, p' \in S} p \cdot p' c_t(p) c_t(p') (f(p \circ p') - f(p) - f(p')) \quad (6)$$

for every bounded $f : S \rightarrow \mathbb{R}$.

Remark 2. • The derivative in this lemma has to be understood in the weak sense, i.e. the formula actually holds in the integral form. It is anyway easy to check, using the decreasing monotonicity of $t \mapsto \langle a, c_t \rangle$ and $t \mapsto \langle b, c_t \rangle$, that if $f(a, b, m) \rightarrow 0$ when $(a, b, m) \rightarrow \infty$, then the series actually converges uniformly, and hence that $\langle c_t, f \rangle$ is C^1 and that the formula holds in the strong sense.

- Consider in particular, the generating function of c_t

$$g_t(x, y, z) = \langle x^a y^b z^m, c_t \rangle = \sum_{(a, b, m) \in S} c_t(a, b, m) x^a y^b z^m.$$

Since $t \mapsto \langle c_t, 1 \rangle$ decreases, it is well-defined. Fix $z \in [0, 1]$. For every $(x, y) \in (0, 1)^2$, $(t, x, y) \mapsto g_t(x, y, z)$ is C^1 in t because of the remark above, and formula (6) shows that its partial derivative with respect to t is C^1 . Moreover, it is C^2 in (x, y) (even analytic), and the partial derivatives with respect to x and y are C^1 in t (it is the remark above with $f(a, b, m) = ax^{a-1}y^b z^m$ and $f(a, b, m) = bx^a y^{b-1} z^m$). For the same reason, the second order partial derivatives are continuous in t . Finally, $g(\cdot, \cdot, z)$ is regular, in the sense of Definition 3 below.

Definition 3. *We say that a function $(t, x, y) \mapsto g_t(x, y)$ defined on $[0, T] \times (0, 1)^2$ is regular if*

- $t \mapsto g_t(x, y)$ is C^1 ,
- $(x, y) \mapsto \frac{\partial g_t}{\partial t}(x, y)$ is C^1 ,
- $(x, y) \mapsto g_t(x, y)$ is C^2 ,
- $t \mapsto \frac{\partial g_t}{\partial x}(x, y)$ and $t \mapsto \frac{\partial g_t}{\partial y}(x, y)$ are C^1 .

We may now prove the fact pointed out earlier that the mean number of male and female arms remains equal as time passes.

Lemma 3. *Let (c_t) be a solution to Smoluchowski's equation, and let Γ_∞ be defined as in (2). Consider the mean numbers of male and female arms $A_t = \langle a, c_t \rangle$ and $B_t = \langle b, c_t \rangle$, and assume $A_0 = B_0 = 1$. Then for all $t \in [0, T \wedge \Gamma_\infty)$*

$$A_t = B_t = \frac{1}{1+t}. \quad (7)$$

Proof. For $t < \Gamma_\infty$, we can use dominated convergence in (6) to choose $f = a$ and $f = b$. We obtain that

$$\frac{d}{dt}A_t = \frac{d}{dt}B_t = -A_tB_t$$

for $t < \Gamma_\infty$. Since $A_0 = B_0 = 1$, the result follows. \square

3 Proof of the theorem

3.1 Overview of the method

In this section, we give a sketch of the proof which contains all the important ideas. The rigorous proof however requires some care, and it is given in detail afterwards. So, consider a solution $(c_t)_{t \in [0, T]}$ to Smoluchowski's equation (1), and

$$g_t(x, y, z) = \langle x^a y^b z^m, c_t \rangle = \sum_{a \geq 0} \sum_{b \geq 0} \sum_{m \geq 1} c_t(a, b, m) x^a y^b z^m.$$

Using (6) and Lemma 3, it is easy to see that g_t solves the following PDE

$$\frac{\partial g_t}{\partial t} = \frac{\partial g_t}{\partial x} \frac{\partial g_t}{\partial y} - \frac{1}{1+t} \left(x \frac{\partial g_t}{\partial x} + y \frac{\partial g_t}{\partial y} \right). \quad (8)$$

Now, we can solve this PDE using the method of characteristics: we want to find a trajectory $(x(t), y(t))$ starting from some $(x, y) \in [0, 1]^2$ such that $g_t(x(t), y(t), z)$ is easy to compute. So let

$$(p_1(t), p_2(t)) = \left(\frac{\partial g_t}{\partial x}(x(t), y(t), z), \frac{\partial g_t}{\partial y}(x(t), y(t), z) \right).$$

An easy calculation shows that

$$\dot{p}_1(t) = \frac{\partial^2 g_t}{\partial x^2} \left(\dot{x}(t) + p_2(t) - \frac{x(t)}{1+t} \right) + \frac{\partial^2 g_t}{\partial x \partial y} \left(\dot{y}(t) + p_1(t) - \frac{y(t)}{1+t} \right) - \frac{p_1(t)}{1+t}, \quad (9)$$

and a similar formula for p_2 . Now, if we require

$$\dot{x}(t) + p_2(t) - \frac{x(t)}{1+t} = \dot{y}(t) + p_1(t) - \frac{y(t)}{1+t} = 0,$$

then

$$\dot{p}_i(t) = -\frac{p_i(t)}{1+t}, \quad i = 1, 2.$$

These ODE's are readily solved, with $p_1(0) = \frac{\partial g_0}{\partial x}(x, y)$ and $p_2(0) = \frac{\partial g_0}{\partial y}(x, y)$, and we obtain

$$p_i(t) = \frac{p_i(0)}{1+t} \quad (10)$$

and

$$x(t) = x + (x - p_2(0))t \quad ; \quad y(t) = y + (y - p_1(0))t.$$

Using the PDE, we now see that

$$\frac{d}{dt} g_t(x(t), y(t), z) = -\frac{p_1(0)p_2(0)}{(1+t)^2}, \quad (11)$$

so by integrating

$$g_t(x(t), y(t), z) = g_t(\phi_t(x, y, z), z) = g_0(x, y, z) - \frac{t}{1+t} \frac{\partial g_0}{\partial x}(x, y, z) \frac{\partial g_0}{\partial y}(x, y, z). \quad (12)$$

To obtain g_t , it only remains to invert ϕ_t , for, if $\phi_t(h_t) = \text{Id}$, then

$$g_t(x, y, z) = g_0(h_t(x, y, z)) - \frac{t}{1+t} \frac{\partial g_0}{\partial x}(h_t(x, y, z)) \frac{\partial g_0}{\partial y}(h_t(x, y, z)).$$

We may now start a rigorous proof, which consists mainly of 3 steps

- Study the map ϕ_t ,
- Solve the PDE (18),
- Show that the generating function of a family (c_t) solves (18) if and only if (c_t) solves Smoluchowski's equation (1).

3.2 Inversion of the mapping

In this section, we study the map ϕ_t , which is useful both for solving theoretically the PDE, and for obtaining explicit solutions. We will need two preliminary lemmas.

Lemma 4. *Let $\alpha > 0$, $\beta, \gamma \geq 0$ and $K = [0, \alpha] \times [0, \beta] \times [0, \gamma]$. For $(r, s, t) \in K$, denote*

$$A(r, s, t) := \begin{pmatrix} r & s \\ t & r \end{pmatrix}.$$

Then for every $\epsilon > 0$, there is a norm $\|\cdot\|$ on \mathbb{R}^2 such that

$$\max_{(r,s,t) \in K} \|A(r, s, t)\| \leq \alpha + \sqrt{\beta\gamma} + \epsilon,$$

where we also denote by $\|\cdot\|$ the induced norm on the 2×2 matrices.

Remark 3. This is a uniform version of the well known result which states that

- For every (square) matrix A and norm $\|\cdot\|$, one has $\|A\| \geq \rho(A)$, where $\rho(A)$ is the spectral radius of A ,

- For every matrix A and $\epsilon > 0$, there is a norm $\|\cdot\|$ such that $\|A\| \leq \rho(A) + \epsilon$.

Note indeed that $\alpha + \sqrt{\beta\gamma}$ is the spectral radius of $A(\alpha, \beta, \gamma)$.

Proof. 1. First assume that β and γ are positive. We can diagonalize $A := A(\alpha, \beta, \gamma)$. If we let $a := \alpha$, $b := \sqrt{\beta}$ and $c := \sqrt{\gamma}$ then

$$A = P \begin{pmatrix} a + bc & 0 \\ 0 & a - bc \end{pmatrix} P^{-1},$$

where

$$P = \begin{pmatrix} b & -b \\ c & c \end{pmatrix} \quad ; \quad P^{-1} = \frac{1}{2bc} \begin{pmatrix} c & b \\ -c & b \end{pmatrix}.$$

Now, consider the following norm: for $x \in \mathbb{R}^2$, let $\|x\| = \|P^{-1}x\|_\infty$, where $\|(x_1, x_2)\|_\infty = \max(|x_1|, |x_2|)$. Then for any 2×2 matrix M ,

$$\|M\| = \max_{x \neq 0} \frac{\|Mx\|}{\|x\|} = \max_{x \neq 0} \frac{\|P^{-1}Mx\|_\infty}{\|P^{-1}x\|_\infty} = \max_{y \neq 0} \frac{\|P^{-1}MPy\|_\infty}{\|y\|_\infty} = \|P^{-1}MP\|_\infty.$$

An easy computation shows that for $(r, s, t) \in K$,

$$P^{-1}A(r, s, t)P = \begin{pmatrix} r + \frac{bt}{2c} + \frac{cs}{2b} & -\frac{bt}{2c} + \frac{cs}{2b} \\ \frac{bt}{2c} - \frac{cs}{2b} & r - \frac{bt}{2c} - \frac{cs}{2b} \end{pmatrix}.$$

Recall that for a matrix M ,

$$\|M\|_\infty = \max_i \sum_j |M_{i,j}|,$$

so that, since $r \geq 0$,

$$\|P^{-1}A(r, s, t)P\|_\infty = r + \frac{bt}{2c} + \frac{cs}{2b} + \left| \frac{bt}{2c} - \frac{cs}{2b} \right| := F(r, s, t).$$

It remains to find the maximum of F on K . First, note that for $(r, s, t) \in K$,

$$0 \leq F(r, s, t) \leq F(\alpha, s, t).$$

Then, for every $(s, t) \in [0, \beta] \times [0, \gamma]$, we can write $t = ps$, $p \geq 0$. If $p \leq c^2/b^2$, then $cs/(2b) \geq bt/(2c)$, so that $F(\alpha, s, t) = \alpha + cs/b$. But $s \leq b^2$, so $F(\alpha, s, t) \leq \alpha + bc = \alpha + \sqrt{\beta\gamma}$. And if $p > c^2/b^2$, then $cs/(2b) \leq bt/(2c)$, so that $F(\alpha, s, t) = \alpha + bt/c$. But $t \leq c^2$, so $F(\alpha, s, t) \leq \alpha + bc = \alpha + \sqrt{\beta\gamma}$. Finally, the maximum of F on K , i.e. the maximum of $\|A(r, s, t)\|$ on K , is $\alpha + \sqrt{\beta\gamma}$.

2. Assume now that β or γ is zero, say e.g. $\gamma = 0$. Take $\epsilon > 0$, and $M > 0$ such that $\beta/M < \epsilon$. Consider the norm $\|x\| = \|Px\|_\infty$, where P is a diagonal matrix with diagonal $(1, M)$. For $(r, s, 0) \in K$, we have as before

$$\|A(r, s, 0)\| = \|PA(r, s, 0)P^{-1}\|_\infty = \left\| \begin{pmatrix} r & s/M \\ 0 & r \end{pmatrix} \right\|_\infty.$$

Since $s \leq \beta$, this shows that $\|A(r, s, 0)\| \leq \alpha + \epsilon$.

□

We will deal often with real-analytic functions in the remaining of the proofs. For the definitions and results on this topic, we refer to [21]. In particular, we will use the following result to study the regularity of ϕ_t .

Lemma 5. *Let $f_i(x_1, \dots, x_l, u_1, \dots, u_n)$, $i = 1 \dots n$, be real analytic functions in a neighbourhood of a point $(p, q) \in \mathbb{R}^l \times \mathbb{R}^n$. Consider the system of equations in u_1, \dots, u_n*

$$f_i(x_1, \dots, x_l, u_1, \dots, u_n) = 0, \quad i = 1, \dots, n. \quad (13)$$

Assume that $f(p, q) = 0$, and that the matrix

$$\left(\frac{\partial f_i}{\partial u_j}(p, q) \right)_{1 \leq i, j \leq n} \quad (14)$$

is invertible. Then there is a neighbourhood $U \subset \mathbb{R}^l$ of p and real-analytic functions $h_j : U \rightarrow \mathbb{R}$, $j = 1, \dots, n$, such that for all $j = 1, \dots, n$, $h_j(p) = q_j$ (where $q = (q_1, \dots, q_n)$), and for all $x \in U$ $f_i(x, h_1(x), \dots, h_n(x)) = 0$. Moreover, $(h_1(x), \dots, h_n(x))$ is locally the unique solution to (13).

We will show the following result.

Proposition 1. *For $t \in [0, T_c)$ and $z \in [0, 1]$, define $\phi_t(\cdot, \cdot, z) : [0, 1]^2 \rightarrow \mathbb{R}^2$ by*

$$\phi_t(x, y, z) = \left((1+t)x - t \frac{\partial g_0}{\partial y}(x, y, z), (1+t)y - t \frac{\partial g_0}{\partial x}(x, y, z) \right), \quad (15)$$

and let $K_t(z)$ be the closed subset of $[0, 1]^2$: $K_t(z) = \phi_t(\cdot, \cdot, z)^{-1}([0, 1]^2)$. Then

- (i) $\phi_t(\cdot, \cdot, z) : K_t(z) \rightarrow [0, 1]^2$ is a homeomorphism. Denote $h_t(\cdot, \cdot, z) = (h_t^{(1)}(\cdot, \cdot, z), h_t^{(2)}(\cdot, \cdot, z))$ its inverse.
- (ii) For $i = 1, 2$, $(x, y, z, t) \mapsto h_t^{(i)}(x, y, z)$ is an analytic function on $(0, 1)^3 \times (0, T_c)$.

Proof. (i) Fix some $z \in [0, 1]$ and some $t \in (0, T_c)$, and keep the notations of the statement. For notational simplicity, we omit the parameter z . Let $0 \leq t < T_c$. We first want to show that $\phi_t : K_t \rightarrow [0, 1]^2$ is one-to-one and onto. Fix $(u, v) \in [0, 1]^2$ and let us check that there is a unique couple $(x, y) \in [0, 1]^2$ such that $\phi_t(x, y) = (u, v)$. This requirement is equivalent to finding a unique fixed point to

$$F_t(x, y) = \left(\frac{u}{1+t} + \frac{t}{1+t} \frac{\partial g_0}{\partial y}(x, y, z), \frac{v}{1+t} + \frac{t}{1+t} \frac{\partial g_0}{\partial x}(x, y, z) \right).$$

Because of the remark above, F_t is a mapping from $[0, 1]^2$ to $[0, 1]^2$. It remains to check that it is contracting. Its differential is

$$DF_t(x, y) = \frac{t}{1+t} \begin{pmatrix} \frac{\partial^2 g_0}{\partial x \partial y} & \frac{\partial^2 g_0}{\partial y^2} \\ \frac{\partial^2 g_0}{\partial x^2} & \frac{\partial^2 g_0}{\partial x \partial y} \end{pmatrix} := \frac{t}{1+t} \begin{pmatrix} \alpha(x, y, z) & \beta(x, y, z) \\ \gamma(x, y, z) & \alpha(x, y, z) \end{pmatrix}. \quad (16)$$

Let $\alpha = \alpha(1, 1, 1) = \langle ab, c_0 \rangle$, $\beta = \beta(1, 1, 1) = \langle b^2, c_0 \rangle$ and $\gamma = \gamma(1, 1, 1) = \langle a^2, c_0 \rangle$. Since $t < T_c$, then $\frac{t}{1+t}(\alpha + \sqrt{\beta\gamma} + \epsilon) < 1$ for some small enough $\epsilon > 0$. Hence, by Lemma 4, there is a norm $\|\cdot\|$ such that

$$\max_{(x,y) \in [0,1]^2} \|DF_t(x, y)\| \leq \frac{t}{1+t}(\alpha + \sqrt{\beta\gamma} + \epsilon) < 1,$$

so that F_t is contracting. Hence it has a unique fixed point. As a consequence, there is a unique couple $(x, y) \in [0, 1]^2$ such that $\phi_t(x, y) = (u, v)$. Moreover, since F_t is continuous with respect to (u, v) and uniformly contracting in (u, v) , then the mapping $(u, v) \mapsto (x, y)$ is continuous, that is $h_t : [0, 1]^2 \rightarrow K_t$ is a homeomorphism.

(ii) Let $t_0 \in (0, T_c)$, $z_0 \in (0, 1)$ and $(x_0, y_0) \in U_{t_0}$. We want to invert the function ϕ_t above, that is, we want to find $(u_1, u_2) \in [0, 1]^2$ such that

$$\begin{cases} (1+t)u_1 - t \frac{\partial g_0}{\partial y}(u_1, u_2, z) - x := f_1(x, y, z, t, u_1, u_2) = 0 \\ (1+t)u_2 - t \frac{\partial g_0}{\partial x}(u_1, u_2, z) - y := f_2(x, y, z, t, u_1, u_2) = 0. \end{cases} \quad (17)$$

We have just shown that there exists $(u_1^0, u_2^0) \in (0, 1)^2$ such that $f_i(x_0, y_0, z_0, t_0, u_1^0, u_2^0) = 0$, $i = 1, 2$. Now, the matrix (14) is precisely

$$D\phi_{t_0}(x_0, y_0, z_0) = \begin{pmatrix} 1 + t_0 - t_0\alpha(x_0, y_0, z_0) & -t_0\beta(x_0, y_0, z_0) \\ -t_0\gamma(x_0, y_0) & 1 + t_0 - t_0\alpha(x_0, y_0) \end{pmatrix},$$

and it is easy to see that since $t_0 < T_c$, it is invertible. So Lemma 5 says that there is a neighbourhood $U \subset (0, 1)^3 \times (0, T_c)$ of (x_0, y_0, z_0, t_0) , and real-analytic functions $u_1, u_2 : U \rightarrow \mathbb{R}$ such that $u_1(x_0, y_0, z_0, t_0) = u_1^0$, $u_2(x_0, y_0, z_0, t_0) = u_2^0$, and for $(x, y, z, t) \in U$

$$\begin{cases} (1+t)u_1(x, y, z, t) - t \frac{\partial g_0}{\partial y}(u_1(x, y, z, t), u_2(x, y, z, t), z) = x \\ (1+t)u_2(x, y, z, t) - t \frac{\partial g_0}{\partial x}(u_1(x, y, z, t), u_2(x, y, z, t), z) = y. \end{cases}$$

Because of the uniqueness of the solution to (17), then $(u_1, u_2) = (h^{(1)}, h^{(2)})$. So the functions $h_t^{(i)}$ are real-analytic functions on $(0, 1)^3 \times (0, T_c)$. □

3.3 Study of the PDE

The following (non-quasilinear) PDE is a central feature of our discussion

$$\frac{\partial g_t}{\partial t} = \frac{\partial g_t}{\partial x} \frac{\partial g_t}{\partial y} - \frac{1}{1+t} \left(x \frac{\partial g_t}{\partial x} + y \frac{\partial g_t}{\partial y} \right). \quad (18)$$

Proposition 2. (i) For every $z \in [0, 1]$, the PDE (18) with initial conditions $g_0 = g_0(\cdot, \cdot, z)$ has a unique regular (in the sense of Definition 3) solution $(t, x, y) \mapsto g_t(x, y, z)$ defined on $[0, T_c) \times (0, 1)^2$. Moreover, $(t, x, y, z) \mapsto g_t(x, y, z)$ is analytic on $(0, T_c) \times (0, 1)^3$.

(ii) The solution of the PDE is given by

$$g_t(x, y, z) = g_0(h_t(x, y, z), z) - \frac{t}{1+t} \frac{\partial g_0}{\partial x}(h_t(x, y, z), z) \frac{\partial g_0}{\partial y}(h_t(x, y, z), z), \quad (19)$$

where h_t is defined in Proposition 1.

(iii) We have the alternative expression

$$g_t(x, y, z) = \frac{1}{1+t} \left(\tilde{H}_t^{(2)}(x, y, z) + \tilde{H}_t^{(1)}(0, y, z) \right) + G_t(z) \quad (20)$$

in the notations of Theorem 1.

Remark 4. Formula (20) will be useful to compute explicit solutions. Indeed, what we aim at obtaining is the analytic expansion of g_t , to extract its coefficients and get (c_t) . But assume for instance that we have found an explicit analytic expression for h_t . Substituting it in (19) will not provide any explicit power series expansion, since we need to compound g_0 and h_t . That is when formula (20) comes in handy, since then, one just needs the analytic expansion of $h_t^{(1)}$ and $h_t^{(2)}$ to obtain the expansion of g_t . Note however that G_0 may be tedious to compute in general, but since it is a function of z only, it is relevant only when we wish to compute the concentrations of particles with no arms. Nonetheless, their concentrations can be obtained thanks to the system (1), since

$$\frac{d}{dt} c_t(0, 0, m) = \sum_{m'=1}^{m-1} c_t(1, 0, m') c_t(0, 1, m - m'). \quad (21)$$

Proof. We will prove the statement in three steps. First we will show that a solution has to be written as in (19). Next that this formula does provide a solution. Proving formula (20) is then an easy matter. In all the proof, some $z \in [0, 1]$ is fixed.

1. Let $U_t = \phi_t(\cdot, \cdot, z)^{-1}((0, 1)^2)$, and consider g_t a regular solution of (18) on $[0, T) \times (0, 1)^2$. Fix $t_0 \in (0, T)$ and $(x, y) \in U_{t_0}$, and let

$$(p_1(t), p_2(t)) := \left(\frac{\partial g_t}{\partial x}(\phi_t(x, y, z), z), \frac{\partial g_t}{\partial y}(\phi_t(x, y, z), z) \right).$$

It is easy to see that U_t decreases with t , so for $t \leq t_0$, this definition makes sense and we can differentiate p_i . The regularity assumptions on g_t are just those needed to allow the use of Schwarz's theorem, and an easy computation shows that

$$\begin{cases} \dot{p}_1(t) &= p_1(t) \left(\frac{\partial^2 g_t}{\partial x \partial y} - \frac{1}{1+t} \right) + p_2(t) \frac{\partial^2 g_t}{\partial x^2} - \frac{1}{1+t} \left(\frac{\partial g_0}{\partial y} \frac{\partial^2 g_t}{\partial x^2} - \frac{\partial g_0}{\partial x} \frac{\partial^2 g_t}{\partial x \partial y} \right) \\ \dot{p}_2(t) &= p_1(t) \frac{\partial^2 g_t}{\partial y^2} + p_2(t) \left(\frac{\partial^2 g_t}{\partial x \partial y} - \frac{1}{1+t} \right) - \frac{1}{1+t} \left(\frac{\partial g_0}{\partial x} \frac{\partial^2 g_t}{\partial y^2} - \frac{\partial g_0}{\partial y} \frac{\partial^2 g_t}{\partial x \partial y} \right) \end{cases}$$

for every $0 \leq t \leq t_0$ (the partial derivatives of g_t are taken at ϕ_t). This system of linear differential equations, with initial conditions

$$\left(\frac{\partial g_0}{\partial x}(x, y, z), \frac{\partial g_0}{\partial y}(x, y, z) \right),$$

has continuous coefficients (see Remark 2) so it has a unique solution which is given by (10). Hence

$$\frac{\partial g_t}{\partial x}(\phi_t(x, y, z), z) = \frac{\partial g_0}{\partial x}(x, y, z)/(1+t) \quad ; \quad \frac{\partial g_t}{\partial y}(\phi_t(x, y, z), z) = \frac{\partial g_0}{\partial y}(x, y, z)/(1+t) \quad (22)$$

for all $(x, y) \in U_t$. Then, it is easy to check that for all $(x, y) \in U_t$, (11) and (12) hold. Replacing (x, y) by $h_t(x, y, z)$ (recall $h_t : (0, 1)^2 \rightarrow U_t$ is the right-inverse of ϕ_t), we finally obtain (19). This shows that the PDE has at most one solution.

2. The existence of a solution is now straightforward. Let g_t be defined as in (19). Because of the regularity of h_t and of g_0 , g has the required regularity properties. It then suffices to show that it is actually a solution. To this end, let us first compute

$$(p_1(t), p_2(t)) := \left(\frac{\partial g_t}{\partial x}(\phi_t), \frac{\partial g_t}{\partial y}(\phi_t) \right)$$

for some fixed $t \in [0, T_c)$ and $(x, y) \in U_t$. But by differentiating $g_t(\phi_t)$ with respect to x and y , we see that it solves the following system

$$\begin{cases} p_1(t) \frac{\partial \phi_t^{(1)}}{\partial x} + p_2(t) \frac{\partial \phi_t^{(2)}}{\partial x} = \frac{\partial g_t(\phi_t)}{\partial x} = -\frac{t}{1+t} \left(\frac{\partial^2 g_0}{\partial x^2} \frac{\partial g_0}{\partial y} + \frac{\partial g_0}{\partial x} \frac{\partial^2 g_0}{\partial x \partial y} \right) \\ p_1(t) \frac{\partial \phi_t^{(1)}}{\partial y} + p_2(t) \frac{\partial \phi_t^{(2)}}{\partial y} = \frac{\partial g_t(\phi_t)}{\partial y} = -\frac{t}{1+t} \left(\frac{\partial^2 g_0}{\partial x \partial y} \frac{\partial g_0}{\partial y} + \frac{\partial g_0}{\partial x} \frac{\partial^2 g_0}{\partial y^2} \right). \end{cases} \quad (23)$$

For $t < T_c$, this linear system has a nonzero determinant, and hence a unique solution, given by equation (22). To conclude, let $k_t = g_t(\phi_t)$. On the one hand, thanks to (19) and (22), we have

$$\frac{d}{dt} k_t = -\frac{1}{(1+t)^2} \frac{\partial g_0}{\partial x} \frac{\partial g_0}{\partial y} = -\frac{\partial g_t}{\partial x}(\phi_t) \frac{\partial g_t}{\partial y}(\phi_t).$$

On the other hand, still with (22), we obtain

$$\begin{aligned} \frac{d}{dt} k_t &= \frac{d g_t}{dt}(\phi_t) + \frac{\partial g_t}{\partial x}(\phi_t) \frac{d}{dt} \phi_t^{(1)} + \frac{\partial g_t}{\partial y}(\phi_t) \frac{d}{dt} \phi_t^{(2)} \\ &= \frac{d g_t}{dt}(\phi_t) - 2 \frac{\partial g_t}{\partial x}(\phi_t) \frac{\partial g_t}{\partial y}(\phi_t) + \frac{\partial g_t}{\partial x}(\phi_t) \left(x - t \frac{\partial g_t}{\partial x}(\phi_t) \right) + \frac{\partial g_t}{\partial y}(\phi_t) \left(y - t \frac{\partial g_t}{\partial y}(\phi_t) \right). \end{aligned}$$

By using this equality, compounding by h_t , and using (22) again, we finally observe that g_t solves the PDE (18) for $t \in [0, T_c)$, $(x, y) \in (0, 1)^2$.

3. Let $g_0(h_t(x, y, z), z) = f(x, y, z)$. We have

$$\frac{\partial f}{\partial x} = \frac{\partial g_0}{\partial x}(h_t, z) \frac{\partial h_t^{(1)}}{\partial x} + \frac{\partial g_0}{\partial y}(h_t, z) \frac{\partial h_t^{(2)}}{\partial x}.$$

Recall that $\phi_t(h_t) = \text{Id}$, so with the notations of Theorem 1, we get

$$\frac{\partial g_0}{\partial x}(h_t, z) = \tilde{h}_t^{(2)} \quad ; \quad \frac{\partial g_0}{\partial y}(h_t, z) = \tilde{h}_t^{(1)}.$$

Hence, with $q = t/(1+t)$, we obtain

$$\begin{aligned}\frac{\partial f}{\partial x} &= \tilde{h}_t^{(2)} \frac{\partial h_t^{(1)}}{\partial x} + \tilde{h}_t^{(1)} \frac{\partial h_t^{(2)}}{\partial x} \\ &= q \left(\tilde{h}_t^{(2)} \frac{\partial \tilde{h}_t^{(1)}}{\partial x} + \tilde{h}_t^{(1)} \frac{\partial \tilde{h}_t^{(2)}}{\partial x} \right) + \frac{q}{t} \tilde{h}_t^{(2)}.\end{aligned}$$

Integrating this equality gives

$$\begin{aligned}f(x, y, z) &= q \tilde{h}_t^{(1)}(x, y, z) \tilde{h}_t^{(2)}(x, y, z) - q \tilde{h}_t^{(1)}(0, y, z) \tilde{h}_t^{(2)}(0, y, z) \\ &\quad + \frac{q}{t} \tilde{H}_t^{(2)}(x, y, z) + f(0, y, z).\end{aligned}\tag{24}$$

Just in the same way, we can differentiate $f(0, y, z)$ with respect to y , and $f(0, 0, z)$ with respect to z to obtain (20).

□

3.4 Equivalence between the system and the PDE

Smoluchowski's equation is solved thanks to the PDE (18).

Proposition 3. (i) *Let $(c_t)_{t \in [0, T]}$ be a solution to Smoluchowski's equation, and let $g_t(x, y, z) := \langle c_t, x^a y^b z^m \rangle$ be its generating function. Then for all $z \in [0, 1]$, $(t, x, y) \mapsto g_t(x, y, z)$ is a regular solution to the PDE (18) on $[0, T \wedge \Gamma_\infty) \times (0, 1)^2$, with initial conditions $g_0(\cdot, \cdot, z)$.*

(ii) *Conversely, let $(c_t(p))_{p \in S}$, $t \in [0, T]$ be a family of differentiable functions. Let $g_t(x, y, z)$ be its generating function and assume it is defined for $t \in [0, T]$, $(x, y) \in (0, 1)^2$ and $z \in [0, 1]$. Assume that for every $z \in [0, 1]$, $g_t(\cdot, \cdot, z)$ is a regular solution to the PDE (18) with initial conditions $g_0(\cdot, \cdot, z)$. Then*

- For all $p \in S$ and $t \in [0, T)$, $c_t(p) \geq 0$,
- (c_t) is a solution to Smoluchowski's equation for $t \in [T \wedge T_c)$, with initial conditions c_0 .

Remark 5. An important feature of this result is that the PDE (18) and the system (1) are equivalent only before the critical time (T_c or Γ_∞). A priori, a solution of Smoluchowski's equation may be defined after Γ_∞ , but then its generating function may not solve the PDE after this time. Conversely, a generating function may solve the PDE after T_c , but its coefficients may not solve Smoluchowski's equation after this time. This fact is crucial when we study the microscopic model. We indeed obtain a family of coefficients whose generating function solves the PDE (on $[0, +\infty)$), but we cannot ensure that they solve Smoluchowski's equation after T_c (actually, we believe that they do not).

A preliminary result to the proof is the following. Its proof is exactly the same as the one of Lemma 1.

Lemma 6. Let $(c_t)_{t \in [0, T]}$ be a solution to the system

$$\frac{d}{dt}c_t(p) = \frac{1}{2} \sum_{p' \preceq p} p' \cdot (p \setminus p') c_t(p') c_t(p \setminus p') - \frac{a+b}{1+t} c_t(p) \quad (25)$$

for $p = (a, b, m) \in S$, with nonnegative initial conditions. Then for all $t \in [0, T)$ and $p \in S$, $c_t(p) \geq 0$.

Proof of Proposition 3. (i) Take the same notations as in the proposition, part (i). First note that g is regular according to Remark 2. If one takes $f(a, b, m) = x^a y^b z^m$ in (6), for some fixed $(x, y, z) \in (0, 1)^2 \times [0, 1]$, then one gets

$$\frac{d}{dt}g_t(x, y, z) = \frac{\partial g_t}{\partial x} \frac{\partial g_t}{\partial y} - A_t y \frac{\partial g_t}{\partial y} - B_t x \frac{\partial g_t}{\partial x}.$$

Recall from Lemma 3 that when $t < \Gamma_\infty$, $A_t = B_t = 1/(1+t)$. Replacing in the equation above shows that g_t solves (18) for $(x, y) \in (0, 1)^2$ and $0 \leq t < T \wedge \Gamma_\infty$.

(ii) Take now the notations of the proposition, part (ii).

- For every $(x, y, z) \in (0, 1)^3$, the series converges normally in t . Since g_t solves the PDE (18), we may write, using the normal convergence and a Cauchy product, that

$$\begin{aligned} g_t(x, y, z) = & g_0(x, y, z) + \frac{1}{2} \sum_{p=(a,b,m) \in S} x^a y^b z^m \sum_{p' \preceq p} p' \cdot (p \setminus p') \int_0^t c_s(p') c_s(p \setminus p') ds \\ & - \sum_{p=(a,b,m) \in S} x^a y^b z^m \int_0^t \frac{a+b}{1+s} c_s(p) ds \end{aligned} \quad (26)$$

for $(x, y, z) \in (0, 1)^3$ and $t \in [0, T)$. Identifying the coefficients now shows that (c_t) solves the system (25), so the nonnegativity of the $c_t(p)$'s follows from Lemma 6.

- Let $t < T_c \wedge T$, $U_t = \phi_t^{-1}(\cdot, \cdot, 1)((0, 1)^2)$, $K_t = K_t(1) = \phi_t^{-1}(\cdot, \cdot, 1)([0, 1]^2)$, and recall from (22) that since g is a regular solution to (18), then for all $(x, y) \in U_t$

$$\frac{\partial g_t}{\partial x}(\phi_t(x, y, 1), 1) = \frac{\partial g_0}{\partial x}(x, y, 1) \frac{1}{1+t},$$

what we can write

$$\sum_{(a,b,m) \in S} a c_t(a, b, m) \phi_t^{(1)}(x, y, 1)^a \phi_t^{(2)}(x, y, 1)^b = \sum_{(a,b,m) \in S} a c_0(a, b, m) x^a y^b \times \frac{1}{1+t} \quad (27)$$

Note now that since $t < T_c$, then $\phi_t(\cdot, \cdot, 1) : K_t \rightarrow [0, 1]^2$ is a homeomorphism, so $\overline{U}_t = K_t$. Since $(1, 1) \in K_t$, we can pass to the limit in the equality above when $(x, y) \rightarrow (1, 1)$. Using monotone convergence and the continuity of ϕ_t , we obtain

$$\sum_{(a,b,m) \in S} a c_t(a, b, m) = \sum_{(a,b,m) \in S} a c_0(a, b, m) \times \frac{1}{1+t} = \frac{1}{1+t}.$$

The same reasoning shows that $\langle b, c_t \rangle = 1/(1+t)$ for $t < T_c$. Hence, we may re-write (26) before T_c by substituting

$$\frac{a}{1+t} = a \sum_{(a', b', m') \in S} b' c_t(a', b', m') \quad ; \quad \frac{b}{1+t} = b \sum_{(a', b', m') \in S} a' c_t(a', b', m').$$

Identifying the coefficients now shows that (c_t) solves Smoluchowski's equation for $t \in [0, T \wedge T_c)$. □

3.5 Existence and uniqueness of a solution

With these results, proving Theorem 1 is now an easy matter.

Proof of theorem 1. 1. Let us first prove that $\Gamma_\infty = T_c$. Take $(c_t(a, b, m))_{t \in [0, T]}$ a solution to the system (1), and g_t its generating function. We have

$$\frac{\partial^2 g_t}{\partial x^2}(1, 1, 1) = \langle c_t, a^2 - a \rangle,$$

the left-hand side being understood as a monotone limit. Note that by Lemma 3 $\langle c_t, a \rangle$ is bounded by 1, so $\langle c_t, a^2 \rangle$ explodes if and only if $\frac{\partial^2 g_t}{\partial x^2}(1, 1, 1)$ explodes. Let us compute the latter. Recall from (22) that for $(x, y) \in U_t$

$$\frac{\partial g_t}{\partial x}(\phi_t(x, y, 1), 1) = \frac{\partial g_0}{\partial x}(x, y, 1)/(1+t).$$

Differentiating this equality with respect to x , and having (x, y) tend to $(1, 1)$, we obtain

$$\frac{\partial^2 g_t}{\partial x^2}(1, 1, 1) \frac{\partial \phi_t^{(1)}}{\partial x}(1, 1, 1) + \frac{\partial^2 g_t}{\partial x \partial y}(1, 1, 1) \frac{\partial \phi_t^{(2)}}{\partial x}(1, 1, 1) = \frac{\partial^2 g_0}{\partial x^2}(1, 1, 1)/(1+t),$$

what we write

$$a \frac{\partial \phi_t^{(1)}}{\partial x}(1, 1, 1) + c \frac{\partial \phi_t^{(2)}}{\partial x}(1, 1, 1) = \frac{\gamma}{1+t}, \quad (28)$$

where

$$\alpha = \frac{\partial^2 g_0}{\partial x \partial y} = \langle ab, c_0 \rangle \quad ; \quad \beta = \frac{\partial^2 g_0}{\partial y^2} = \langle b^2 - b, c_0 \rangle \quad ; \quad \gamma = \frac{\partial^2 g_0}{\partial x^2} = \langle a^2 - a, c_0 \rangle.$$

By differentiating with respect to y , we get

$$a \frac{\partial \phi_t^{(1)}}{\partial y}(1, 1) + c \frac{\partial \phi_t^{(2)}}{\partial y}(1, 1) = \frac{\alpha}{1+t}. \quad (29)$$

Hence a and c verify

$$\begin{pmatrix} 1+t-t\alpha & -t\gamma \\ -t\beta & 1+t-t\alpha \end{pmatrix} \begin{pmatrix} a \\ c \end{pmatrix} = \frac{1}{1+t} \begin{pmatrix} \gamma \\ \alpha \end{pmatrix},$$

so that

$$a = \frac{\partial^2 g_t}{\partial x^2}(1, 1, 1) = \sum_{(a,b,m) \in S} a(a-1)c_t(a, b, m) = \frac{\gamma}{(1+t-t\alpha)^2 - t^2\gamma\beta}.$$

This expression is valid as long as $t < T_c$, since the determinant of the matrix is then nonzero. In the same way, we also have

$$\frac{\partial^2 g_t}{\partial y^2}(1, 1, 1) = \sum_{(a,b,m) \in S} b(b-1)c_t(a, b, m) = \frac{\beta}{(1+t-t\alpha)^2 - t^2\gamma\beta}.$$

If γ or β is nonzero, then $\langle c_t, a^2 + b^2 \rangle \rightarrow +\infty$ when $t \rightarrow T_c$, i.e. $\Gamma_\infty = T_c$. If $\gamma = \beta = 0$, then $\Gamma_\infty = +\infty$. But the conditions $\gamma = \beta = 0$ and $A_0 = B_0 = 1$ impose $\sum_m c_0(1, 1, m) = 1$. So $M = 1$ and $T_c = +\infty = \Gamma_\infty$.

2. Uniqueness is now easy to obtain: assume $(c_t^{(1)})$ and $(c_t^{(2)})$ solve the system 1 on $[0, T)$, $T \leq T_c$, with initial conditions c_0 . Let $g_t^{(1)}$ and $g_t^{(2)}$ be their generating functions. Since $\Gamma_\infty = T_c$ and $T \leq T_c$, then by Proposition 3, for every $z \in [0, 1]$, they are regular solutions to the PDE (18) on $[0, T) \times (0, 1)^2$, with initial conditions $g_0(\cdot, \cdot, z)$. But by Proposition 2 there is a unique regular solution to the PDE on $[0, T_c)$, so $g_t^{(1)} = g_t^{(2)}$ on $[0, T)$, so that $(c_t^{(1)}) = (c_t^{(2)})$.
3. Conversely, for every $z \in [0, 1]$, let $g_t(x, y, z)$, $(t, x, y) \in [0, T_c) \times (0, 1)^2$, be the regular (analytic in (t, x, y, z)) solution to the PDE (18) obtained in Proposition 2. In particular, it is analytic for every t as a function of (x, y, z) . Consider $g_t(x, y, z) = \sum_{(a,b,m) \in S} c_t(a, b, m)x^a y^b z^m$ its power series expansion around 0. Since g solves the PDE (18), it is easy to check that (c_t) solves (25), so by Lemma 6, the coefficients $c_t(p)$ are nonnegative. Hence (see e.g. the proof of Bernstein's theorem in [21]) the series actually converges on $(0, 1)^3$. So g_t is the generating function of a family (c_t) of differentiable functions, it is defined for $(x, y, z, t) \in (0, 1)^2 \times [0, 1] \times [0, T_c)$, and for every $z \in [0, 1]$ $g(\cdot, \cdot, z)$ is a regular solution to (18) with initial conditions $g_0(\cdot, \cdot, z)$. By Proposition 2, this means that (c_t) is a solution to Smoluchowski's equation with initial conditions c_0 .
4. Let us finally prove that the total mass is conserved. Consider $\psi_t(x, y, z) = (\phi_t(x, y, z), z)$, $U'_t = \psi_t^{-1}((0, 1)^3)$ and $K'_t = \psi_t^{-1}([0, 1]^3)$. For $(x, y, z) \in U'_t$, we can differentiate $g_t(\psi_t(x, y, z))$ with respect to z , and using (22), we obtain

$$\frac{\partial g_t}{\partial z}(\psi_t(x, y, z)) = \frac{\partial g_0}{\partial z}(\psi_t(x, y, z)).$$

Now, ψ_t is a homeomorphism from K'_t to $[0, 1]^3$, so $\overline{U'_t} = K_t$. But $(1, 1, 1) \in K'_t$, so we may pass to the limit when $(x, y, z) \rightarrow (1, 1, 1)$ in the equality above, to obtain

$$\frac{\partial g_t}{\partial z}(1, 1, 1) = \frac{\partial g_0}{\partial z}(1, 1, 1),$$

what precisely means $\langle m, c_t \rangle = \langle m, c_0 \rangle$.

□

4 Explicit formulae

We give in this short section some explicit solutions, without giving the full details of the computations. We will always assume that at time 0, there are only particles of size 1 in the medium. So given a (finite) measure μ on $\mathbb{N} \times \mathbb{N}$, we assume

$$c_0(a, b, m) = \mu(a, b)1_{\{m=1\}}$$

and as usual $\langle a, c_0 \rangle = \langle b, c_0 \rangle = 1$. To obtain the solutions, we need to invert ϕ_t , what can be done using the (two-variable) Lagrange inversion formula (a statement is given by Good [13]). But it is much more involved than the one-dimensional formula, and the expressions it would provide can hardly be called explicit. Let us however study three easy cases. Only the last one requires the two-variable formula.

4.1 Particles with one female arm

The first case is when each particle has exactly one female arm, and a number of male arms distributed according to a measure μ_1 . So

$$\mu(a, b) = \begin{cases} 0 & \text{if } b \neq 1 \\ \mu_1(a) & \text{if } b = 1 \end{cases}$$

and we will assume that $A_0 = B_0 = 1$, i.e. μ_1 is a probability measure with unit mean. In this case, we obtain, for every $a, b \geq 0$ and $m \geq 1$

$$c_t(a, b, m) = \begin{cases} 0 & \text{if } b \neq 1 \\ \frac{t^{m-1}}{(1+t)^{m+a}} \frac{1}{m} \binom{m+a-1}{a} \mu_1^{*m}(m+a-1) & \text{if } b = 1 \end{cases}.$$

In particular there exists only particles with one female arms, what is physically obvious. Moreover, the concentration $c_t(a, 1, m)$ is exactly the concentration of particles with a arms and mass m obtained in the ‘‘oriented model’’ of [2], with initial distribution μ_1 . This is also natural, since in this case, $(a, 1, m)$ - and $(a', 1, m')$ -particles indeed coagulate with rate $a + a'$, which is the rate of the oriented model. Note also that $T_c = +\infty$, like in the oriented model.

4.2 Arms with uniform random genders

In this model, the total number of arms of a particle is chosen according to a measure μ_1 , then each arm is given a gender independently, with probability $1/2$. That is, we let

$$\mu(a, b) = \mu_1(a+b) \binom{a+b}{b} \frac{1}{2^{a+b}}.$$

We will assume that μ_1 has mean 2, so that $A_0 = B_0 = 1$. Let $\nu(j) = (j+1)\mu(j+1)$. Then we obtain, for $(a, b) \neq (0, 0)$

$$c_t(a, b, m) = \frac{1}{2} \frac{t^{m-1}}{(1+t)^{a+b+m-1}} \frac{(m+a+b-2)!}{m!a!b!} \left(\frac{\nu_1}{2}\right)^{*m} (m+a+b-2)$$

and

$$c_t(0, 0, m) = \frac{1}{m(m-1)} \frac{1}{2} \frac{1}{(1+1/t)^{m-1}} \left(\frac{\nu_1}{2}\right)^{*m} (m-2)$$

provided $\nu_1(0) > 0$. If $T_c = +\infty$, this condition means that $\nu_1 \neq \delta_1$. In particular, one easily checks that

$$\sum_{a+b=k} c_t(a, b, m) = 2c_t^{sym}(k, m)$$

where $c_t^{sym}(k, m)$ is the concentration of particles with k arms and mass m in the symmetric model of [2], with initial arm distribution $\mu_1/2$. The factor 2 comes from the normalisation: in our model, the total concentration of arms in the medium is 2, when it is 1 in the symmetric model. It is also worth stressing the stronger fact that for $a, b \geq 0$, we have

$$c_t(a, b, m) = \frac{1}{2} \binom{a+b}{b} \frac{1}{2^{a+b}} c_t(a+b, m).$$

Hence, at any given time, the distribution of the number of male (or female) arms is still binomial. So, if at some time we chose to reassign to each arm a gender uniformly and independently, and let the system evolve on from this state, no difference would be observed. Or we could watch a system evolve like the symmetric model starting from an arm distribution $\mu_1/2$, and then at some time give the arms a gender uniformly at random and independently. The evolution afterwards will be the evolution of the sexed model with initial arm distribution μ . Note as before that the critical time is the same than in the symmetric model with initial distribution $\mu_1/2$.

More generally, consider initial concentrations such that for all $a, b \geq 0$ $\mu(a, b) = \mu(b, a)$, and the solution $(c_t)_{t \in [0, T_c]}$ to Smoluchowski's equation (1). Then it is easy to check (by uniqueness) that for all $t \in [0, T_c]$ $c_t(a, b, m) = c_t(b, a, m)$, and that, if we denote

$$k_t(l, m) := \sum_{a+b=l} c_t(a, b, m),$$

then (k_t) is governed (up to a factor 1/2) by the symmetric Smoluchowski equation of [2]. Hence, in this case too, $k_t(l, m) = 2c_t^{sym}(l, m)$.

4.3 Particles with one gender

Let us finally consider the more intricate case where at time 0, the arms of each particle have all the same gender. This is motivated by the idea of ionic bonds: a particle with only male (resp. female) arms can be considered as a cation (resp. an anion), and cations can only bond with anions. Hence, consider, for $i = 1, 2$, μ_i two measures with mean 1 such that $\mu_1(0) = \mu_2(0)$, and take

$$\mu(a, b) = \begin{cases} \mu_1(a) & \text{if } b = 0 \\ \mu_2(b) & \text{if } a = 0 \\ 0 & \text{else} \end{cases}$$

and $\nu_i(j) = (j+1)\mu_i(j+1)$. The two-variable Lagrange inversion formula gives, for $(a, b) \neq (0, 0)$

$$c_t(a, b, m) = \frac{t^{m-1}}{(1+t)^{m+a+b-1}} \sum_{k=0}^m \frac{(m-k+b-1)!(k+a-1)!}{(m-k)!k!a!b!} \nu_1^{*(m-k)}(k+a-1) \nu_2^{*k}(m-k-1+b).$$

If we also let

$$\nu_1 \diamond \nu_2(m) = (m-1) \sum_{k=1}^{m-1} \frac{1}{k} \nu_1^{*(m-k)}(k-1) \frac{1}{m-k} \nu_2^{*k}(m-k-1)$$

then, for $m \geq 2$,

$$c_t(0, 0, m) = \frac{1}{m-1} \frac{1}{(1+1/t)^{m-1}} \nu_1 \diamond \nu_2(m),$$

provided $\nu_1(0)\nu_2(0) > 0$ (which means, if $T_c = +\infty$, that ν_1 and ν_2 are not δ_1). In particular, we see that if $T_c = +\infty$ and $\nu_1, \nu_2 \neq \delta_1$, then for $m \geq 2$,

$$c_t(a, b, m) \rightarrow \begin{cases} 0 & \text{if } (a, b) \neq (0, 0) \\ \frac{1}{m-1} \nu_1 \diamond \nu_2(m) & \text{if } a = b = 0. \end{cases}$$

Hence all the arms are used to coagulate. Chemically, this means that there are no more ions in the medium. The limiting distribution of the sizes is given by $(m-1)^{-1} \nu_1 \diamond \nu_2(m)$. We will generalize this fact in the following section, and give a probabilistic interpretation of the measure $\nu_1 \diamond \nu_2$. Also, if M_i is the mean of ν_i , then $M = \sqrt{M_1 M_2}$ and $T = 1/(M-1)$, or $+\infty$ if $M \leq 1$. If $\mu_1 = \mu_2 = \mu$, then the critical time is the same as in the symmetric model with initial distribution μ .

5 Limiting concentrations and Galton-Watson processes

5.1 Convergence of the concentrations

In this section, we will study the limiting concentrations. Similarly to what happens in the oriented and symmetric model of [2], we expect the concentrations to converge when the time tends to $+\infty$, whenever gelation does not occur. Physically, this would mean that the system converges to a terminal state where all arms have been used (otherwise, further coagulations “should” occur). This is actually true, and this is an easy consequence of the preceding results.

Corollary 1. *Assume $T_c = +\infty$, and let $(c_t)_{t \geq 0}$ be the solution to Smoluchowski’s equation (1).*

(i) *When $t \rightarrow +\infty$, there exists limiting concentrations $c_\infty(m)$ such that*

$$c_t(a, b, m) \rightarrow c_\infty(m) 1_{\{a=b=0\}}$$

in $\ell^1(\mathbb{N} \times \mathbb{N} \times \mathbb{N}^)$.*

(ii) *For $z \in [0, 1)$, the generating function $g_\infty(z)$ of $(c_\infty(m))_{m \geq 1}$ is the antiderivative vanishing at 0 of*

$$\frac{\partial g_0}{\partial z} (h_\infty^{(1)}(z), h_\infty^{(2)}(z), z)$$

where $(h_\infty^{(1)}, h_\infty^{(2)})$ is characterised by

$$\begin{cases} h_\infty^{(1)}(z) &= \frac{\partial g_0}{\partial y} (h_\infty^{(1)}(z), h_\infty^{(2)}(z), z) \\ h_\infty^{(2)}(z) &= \frac{\partial g_0}{\partial x} (h_\infty^{(1)}(z), h_\infty^{(2)}(z), z). \end{cases} \quad (30)$$

Proof. (i) Since $\Gamma_\infty = T_c = +\infty$, then (7) holds for all $t \geq 0$, so

$$\sum_{(a,b) \neq (0,0)} c_t(a, b, m) \rightarrow 0$$

Then, using (21), we get for all $t \geq 0$

$$c_t(0, 0, m) = c_0(0, 0, m) + \sum_{m'=1}^{m-1} \int_0^t c_s(1, 0, m') c_s(0, 1, m - m') ds$$

But the integrand is bounded by $A_s B_s = 1/(1+s)^2$. Hence the integral has a finite limit when $t \rightarrow +\infty$, and so does $c_t(0, 0, m)$. Finally, $\langle m, c_t \rangle$ is bounded by Theorem 1, and $\langle m, c_\infty \rangle < +\infty$ by Fatou's lemma, so the Cauchy-Schwarz inequality shows that

$$\sum_{m \geq 1} |c_t(0, 0, m) - c_\infty(m)| \rightarrow 0$$

and the result follows.

(ii) By ℓ^1 -convergence, we have

$$g_\infty(z) = \lim_{t \rightarrow +\infty} g_t(0, 0, z),$$

so, using (3) and the fact that $\tilde{H}_t^{(1)}$ and $\tilde{H}_t^{(2)}$ are bounded by 1,

$$g_\infty(z) = \lim_{t \rightarrow +\infty} G_t(z).$$

It just remains to check that $h_t^{(1)}(0, 0, z)$ and $h_t^{(2)}(0, 0, z)$ do have a limit when $t \rightarrow +\infty$. From their definition (4), they have the same limit (if any) as $k_t^{(1)}(z) := \tilde{h}_t^{(1)}(0, 0, z)$ and $k_t^{(2)}(z) := \tilde{h}_t^{(2)}(0, 0, z)$. Now fix $Z \in [0, 1)$, and consider $k_t^{(1)}$ and $k_t^{(2)}$ as (continuous) maps on $[0, Z]$. But h_t is the right-inverse of ϕ_t , so

$$\frac{\partial g_0}{\partial x}(k_t^{(1)}(z), k_t^{(2)}(z), z) = k_t^{(2)}(z) \quad ; \quad \frac{\partial g_0}{\partial y}(k_t^{(1)}(z), k_t^{(2)}(z), z) = k_t^{(1)}(z) \quad (31)$$

and, since $\langle a^2 + b^2, c_0 \rangle < +\infty$, $\frac{\partial g_0}{\partial x}(x, y, z)$ has a bounded differential on $[0, 1]^2 \times [0, Z]$. Hence $k_t^{(2)}$, and for the same reason $k_t^{(1)}$, are Lipschitz-continuous on $[0, Z]$, with a constant independent of t . Ascoli's theorem thus shows that the families $(k_t^{(1)})$ and $(k_t^{(2)})$, $t \geq 0$ lie in a compact set (for the uniform topology on $[0, Z]$). So the family $(k_t^{(1)}, k_t^{(2)})$ lies in a compact set, and passing to the limit in (31) shows that any of its limit points solves (30). But since $T_c = +\infty$, the application

$$(x, y) \mapsto \left(\frac{\partial g_0}{\partial y}(x, y, z), \frac{\partial g_0}{\partial x}(x, y, z) \right)$$

is contracting for every $z \in [0, Z]$. So there is a unique solution to (30), and $(k_t^{(1)}, k_t^{(2)})$ converges to this solution. □

5.2 Connection with two-type Galton-Watson processes

In [2], Bertoin shows that for monodisperse initial conditions (i.e. $c_0(a, m) = \mu(a)1_{\{m=0\}}$ for some measure μ) and when gelation does not occur, the limiting concentrations can be described in terms of Galton-Watson processes. The same kind of analogy is observed in our case. Precisely, consider a Galton-Watson tree with two genders, constructed as follows. We start from a male or a female ancestor. It gives birth to a number a of male children, and a number b of female children, where (a, b) is distributed according to a law $\mu_m(a, b)$ if the ancestor is a male, $\mu_f(a, b)$ if it is a female. Then each child gives birth to a certain number of children, distributed according to μ_m or μ_f , depending on his gender, and so on. Consider $T_f(\mu_m, \mu_f)$ (resp. $T_m(\mu_m, \mu_f)$) the total population of such a Galton-Watson process, starting from a female (resp. male) ancestor. Let for $r \in [0, 1)$

$$g_f(r) = \mathbb{E}(r^{T_f}) \quad ; \quad g_m(r) = \mathbb{E}(r^{T_m})$$

their generating functions. It is an easy exercise to check that they solve the following system, where ϕ_f (resp. ϕ_m) is the generating function of μ_f (resp. μ_m)

$$\begin{cases} g_m(r) &= r\phi_m(g_m(r), g_f(r)) \\ g_f(r) &= r\phi_f(g_m(r), g_f(r)). \end{cases} \quad (32)$$

Besides, if $T(\mu_m, \mu_f)$ is the size of a Galton-Watson tree started from a male and a female ancestor (each tree growing independently), then

$$\mathbb{P}(T(\mu_m, \mu_f) = m) = [z^m]g_m(r)g_f(r), \quad (33)$$

where $[z^m]h(z)$ is the coefficient of z^m in the expansion around 0 of an analytic function h .

Now, let us go back to our study. Assume monodisperse initial conditions, i.e. there is a finite measure μ on $\mathbb{N} \times \mathbb{N}$ such that $c_0(a, b, m) = 1_{\{m=0\}}\mu(a, b)$, and assume $T_c = +\infty$. We will use the same notations as in the previous section. Using (30), we obtain

$$g'_\infty(z) = \frac{1}{z}h_\infty^{(1)}(z)h_\infty^{(2)}(z),$$

so that for $n \geq 2$

$$[z^n]g_\infty(z) = \frac{1}{n-1}[z^n]h_\infty^{(1)}(z)h_\infty^{(2)}(z). \quad (34)$$

Let

- $\nu_m(a, b) = (b+1)\mu(a, b+1)$ the probability measure with generating function $\phi_m := \frac{\partial g_0}{\partial y}$,
- $\nu_f(a, b) = (a+1)\mu(a+1, b)$ the probability measure with generating function $\phi_f := \frac{\partial g_0}{\partial x}$

(they are probability measures because of the assumption $\langle a, c_0 \rangle = \langle b, c_0 \rangle = 1$), and consider the two-type Galton-Watson process as above with these reproduction laws. Because of (30) and (32), (g_m, g_f) and $(h_\infty^{(1)}, h_\infty^{(2)})$ solve the same equation, which has a unique solution by Corollary 1, so $h_\infty^{(1)} = g_m$, $h_\infty^{(2)} = g_f$. Hence by (33) and (34)

$$\mathbb{P}(T(\nu_m, \nu_f) = m) = [z^m]h_\infty^{(1)}(z)h_\infty^{(2)}(z) = (m-1)[z^m]g_\infty(z) = (m-1)c_\infty(m).$$

Finally, let us call a measure μ degenerate if $\mu = \delta_{(1,1)}$ or $\mu = \frac{1}{2}(\delta_{(2,0)} + \delta_{(0,2)})$, or $\mu(a, b) = 0$ for $a \neq 1$, or $\mu(a, b) = 0$ for $b \neq 1$. We let the reader check (using e.g. Theorem 10.1 in [16]) that under the assumptions $T_c = +\infty$, and ruling out the degenerate cases, $T_m(\nu_m, \nu_f)$ and $T_f(\mu_m, \mu_f)$ are finite a.s., and that the process is supercritical if $T_c < +\infty$.

Corollary 2. *The limiting concentrations verify for $m \geq 2$*

$$\mathbb{P}(T(\nu_m, \nu_f) = m) = (m - 1)c_\infty(m).$$

Moreover, if μ is not degenerate, then $T(\nu_m, \nu_f) < +\infty$ a.s.

Hence, the law $\nu_1 \diamond \nu_2$ defined in section 4.4 is the law of the total population of a two-type Galton Watson process started from one male and one female ancestors, where the males give birth to females according to the law ν_2 , and the females give birth to males according to the law ν_1 . In particular, if $\nu_1 = \nu_2 = \nu$, then $\nu \diamond \nu$ is the distribution of the size of a Galton-Watson tree with reproduction law ν and starting from two ancestors. So we get (what is not obvious from the formula for \diamond), that for $m \geq 2$

$$\nu \diamond \nu(m) = \frac{2}{m} \nu^{*m}(m - 2).$$

This corollary answers another question about gelation. By Theorem 1, the total mass $\langle m, c_t \rangle$ is conserved as time passes, so gelation does not occur before T_c . But if $T_c = +\infty$, it may occur at infinity: some mass may be lost then. For monodisperse initial conditions, Corollary 2 proves that this cannot happen, except in the degenerate cases. Denote indeed $C_t = \langle c_t, 1 \rangle$. Because of the ℓ^1 -convergence in Corollary 1, $C_t \rightarrow \sum c_\infty(m)$, and Equation (6) yields $C_t = C_0 - t/(1 + t)$. Hence

$$C_0 - 1 = \sum_{m \geq 1} c_\infty(m).$$

So, whenever μ is not degenerated, Corollary 2 gives

$$\begin{aligned} \sum_{(a,b,m) \in S} m c_\infty(m) &= \sum_{(a,b,m) \in S} ((m - 1)c_\infty(m) + c_\infty(m)) = \mathbb{P}(T(\mu_m, \mu_f) < +\infty) + \sum c_\infty(m) \\ &= 1 + \sum c_\infty(m) = 1 + C_0 - 1 = C_0, \end{aligned}$$

which is precisely the total mass at time 0. For the degenerate cases, we can get explicit expressions for the concentrations, and these show that the mass at infinity is 0.

6 Microscopic model

6.1 Notations and preliminary results

The goal of this section is to construct a sequence of random processes modeling the coagulation of particles with male and female arms. We will start with n particles (and then let $n \rightarrow +\infty$). Let us first set some notations.

- Recall $S = \mathbb{N} \times \mathbb{N} \times \mathbb{N}^*$.

- $\llbracket 0, n \rrbracket = \{0, \dots, n\}$.
- $M > 0$ is a fixed real number. The number of arms and the total mass are assumed to grow at most like Mn (see the definition of E_n).
- For $p = (a, b, m) \in S$ and $p' = (a', b', m') \in S$, we will denote $p.p' = a'b + ab'$ the *rate* of coagulation, and $p \circ p' = (a + a' - 1, b + b' - 1, m + m')$ the type of the particle resulting from such a coagulation.
- The sequence of the number of p -particles is an element of

$$E_n = \left\{ N \in \llbracket 0, n \rrbracket^S, \sum_{(a,b,m) \in S} (a + b + m)N(a, b, m) \leq Mn \right\}$$

which is a finite set.

- $\frac{1}{n}E_n$ is a subset of

$$E = \left\{ C \in [0, 1]^S, \sum_{(a,b,m) \in S} (a + b + m)C(a, b, m) \leq M \right\}.$$

An element of E represents the sequence of concentrations of p -particles. E is a metric space endowed with the distance

$$d(C^{(1)}, C^{(2)}) = \sum_{p \in S} |C^{(1)}(p) - C^{(2)}(p)|.$$

- We will call C-convergence the compact convergence (i.e. uniform convergence on every compact set) for functions from \mathbb{R}^+ to E .
- $\mathbb{D}([0, +\infty), H)$ is the space of càdlàg functions from $[0, +\infty)$ to a metric space (H, d) , endowed with the Skorokhod distance. We will call S-convergence the convergence for Skorokhod's distance. For the basic facts about Skorokhod distance for functions with value in a (complete separable) metric space, see [8].

It is easy to check the following result.

Lemma 7. *(E, d) is a compact metric space. In particular, it is a Polish space.*

We will also need the following result about tightness of probability measures on $\mathbb{D}([0, +\infty), H)$ (cf [8] or [18]).

Lemma 8. *Let (H, d) be a Polish space and $(Y^{(n)})_{n \geq 0}$ a sequence of processes on $\mathbb{D}([0, +\infty), H)$, defined on their respective probability space $(\Omega^n, \mathcal{A}^n, \mathbb{P}^n)$. Let $P^{(n)}$ be the law of $Y^{(n)}$, and $P_t^{(n)}$ the law of $Y_t^{(n)}$. Then the sequence (P_n) is tight if and only if*

1. *There is a dense subset T of \mathbb{R}^+ such that for every $t \in T$, $(P_t^{(n)})_{n \geq 0}$ is tight.*

2. For every $N > 0$, $\alpha > 0$, $\epsilon > 0$, there exists a $\delta > 0$ such that

$$\limsup_n \mathbb{P}_n (\omega \in \Omega^n, w^N(Y^{(n)}(\omega), \delta) > \alpha) < \epsilon$$

where, for $Y \in \mathbb{D}([0, +\infty), H)$

$$w^N(Y, \delta) := \inf_{\pi \in \Pi_\delta} \max_{t_i \in \pi} \sup_{t_i \leq s < t < t_{i+1}} d(Y_t, Y_s)$$

Π_δ being the set of all subdivisions $0 = t_0 < t_1 < \dots < t_n = N$ of $[0, N]$ such that $t_{i+1} - t_i \geq \delta$ for all i .

6.2 Model

Let us now introduce the model. Informally, we consider a finite number n of particles with integer mass, and assume that at time 0, the total mass of the system plus the total number of arms is less than Mn . Then, each pair formed of a p -particle and of a p' -particle may coagulate with rate $\frac{1}{2}p \cdot p'$, independently of the other pairs, to form a $p \circ p'$ -particle, that is, the time one has to wait to see them coagulate is exponential with parameter $\frac{1}{2}p \cdot p'$. In other words, assume the system in in the state η at a given time, that is $\eta \in E_n$ and $\eta(p)$ is the number of p -particles. There are $\eta(p)\eta(p')$ (or $\eta(p)(\eta(p) - 1)$ if $p = p'$) pairs formed of a p -particle and of a p' -particle. Let

$$\lambda_\eta(p, p') = \begin{cases} \frac{1}{2}p \cdot p' \eta(p) \eta(p') & \text{if } p \neq p' \\ \frac{1}{2}p \cdot p \eta(p) (\eta(p) - 1) & \text{if } p = p'. \end{cases}$$

We set independently on each couple (p, p') an exponential clock with parameter $\lambda_\eta(p, p')$ (an exponential random variable with parameter 0 is assumed to be a.s. infinite). There is a.s. one and only clock which rings first. If it is the clock on the couple (p, p') , then the system jumps to the state $\eta + \Delta_{p, p'}$ where

$$\begin{aligned} \Delta_{p, p'}(p) &= \Delta_{p, p'}(p') = -1 & \text{if } p \neq p' \\ \Delta_{p, p'}(p) &= -2 & \text{if } p = p' \\ \Delta_{p, p'}(p \circ p') &= +1. \end{aligned}$$

Then restart the construction afresh from the new state. Note that only finitely many $\eta(p)$ are nonzero, so the first jump occurs after an exponential time with parameter

$$\lambda_\eta = \sum_{p, p' \in S} \lambda_\eta(p, p') < +\infty.$$

We will consider the Markov chain constructed according to this rule. That is, we fix for every $n \geq 1$

- An element $X_0^{(n)}$ of E_n , which is the initial number of particles.
- A pure-jump Markov process $X^{(n)}$ on E_n , defined on some probability space $(\Omega^n, \mathcal{A}^n, \mathbb{P}^n)$, starting from $X_0^{(n)}$, and with generator

$$Gf(\eta) = \sum_{(p, p') \in S^2} (f(\eta + \Delta_{p, p'}) - f(\eta)) \lambda_\eta(p, p')$$

for every bounded function $f : E_n \rightarrow \mathbb{R}$. The construction of such a process is obvious since E_n is finite.

- The rescaled and time-changed process

$$C_t^{(n)} = \frac{1}{n} X_{t/n}^{(n)}.$$

Note that $C^{(n)}$ is a pure-jump Markov process on $\frac{1}{n}E_n \subset E$, starting from $C_0^{(n)} = X_0^{(n)}/n$, and with generator

$$G^{(n)}f(\eta) = \sum_{(p,p') \in S^2} \left(f\left(\eta + \frac{1}{n}\Delta_{p,p'}\right) - f(\eta) \right) \lambda_\eta^{(n)}(p,p')$$

where

$$\lambda_\eta^{(n)}(p,p') = \frac{1}{n} \lambda_{n\eta}(p,p').$$

The law P_n of the process $C^{(n)}$ is a probability measure on $\mathbb{D}([0, +\infty), E)$. We will prove that the sequence (P_n) is tight, and that for every limit point P , and almost every process (C_t) with law P , (C_t) solves some system, which is Smoluchowski's equation (1) before the critical time. Because of the uniqueness of such a solution, this will show that (P_n) itself converges to the solution of Smoluchowski's equation before the critical time. The proof of tightness is analogous to the one in [17], up to some slight modifications.

6.3 Tightness

Lemma 9. *The sequence $(P_n)_{n \geq 0}$ is tight.*

Proof. We will use Lemma 8. For $t \geq 0$, let $P_t^{(n)}$ be the law of $C_t^{(n)}$, which is a probability measure on E . Since E is compact by Lemma 7, the tightness of the sequence $(P_t^{(n)})_{n \geq 0}$ is obvious.

Now, $C^{(n)}$ is a pure-jump process on $\frac{1}{n}E_n \subset \mathbb{E}$, with generator $G^{(n)}$. Hence, when the process is in the state η , then the time before the next jump is exponential with parameter

$$\lambda_\eta^{(n)} := \sum_{p,p' \in S} \lambda_\eta^{(n)}(p,p') = \frac{1}{2} \left(\sum_{p,p' \in S} np.p'\eta(p)\eta(p') - \sum_{p \in S} p.p\eta(p) \right)$$

and, since $\eta \in E$, $\lambda_\eta^{(n)} \leq M^2 n := cn$. Now take $N > 0$, $\beta > 0$, $\epsilon > 0$, and let $\delta > 0$ such that $N = \delta l$ for some $l \in \mathbb{N}^*$, and $3c\delta e/\beta < 1$. Consider the partition $t_0 = 0 < t_1 = \delta < \dots < t_l = \lambda = N$ of $[0, N]$. Let $Z_i := \sup_{t_i \leq s < t_{i+1}} d(C_s^{(n)}, C_t^{(n)})$ for $0 \leq i \leq l-1$. Then

$$\mathbb{P}_n(w^N(C^{(n)}, \delta) > \beta) \leq \mathbb{P}_n \left(\max_{0 \leq i \leq l-1} Z_i > \beta \right) \leq l \max_{0 \leq i \leq l-1} \mathbb{P}_n(Z_i > \beta).$$

But the size of a jump, that is $d(C_{t^-}^{(n)}, C_t^{(n)})$, is $3/n$. Hence, if $Z_i > \beta$, then the process has jumped more than $k := \lceil \beta n/3 \rceil := \lceil c'n \rceil$ times between t_i and t_{i+1} (where $\lceil x \rceil$ is the first integer strictly greater than x). If S_k is the time of the k -th jump, the Markov property tell us that

$$\mathbb{P}_n(Z_i > \beta) \leq \mathbb{P}_n(S_k \leq \delta).$$

But S_k is the sum of k independent exponential random variables, with parameter smaller than cn . So, if S'_k is the sum of k independent exponential random variables with parameter cn (on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$), then S_k is stochastically dominated by S'_k , that is

$$\mathbb{P}_n(S_k \leq \delta) \leq \mathbb{P}(S'_k \leq \delta).$$

To conclude, note that the last term is the probability that a Poisson process with parameter cn jumps more than k times on $[0, \delta]$. Hence

$$\mathbb{P}(S_k \leq \delta) = e^{-cn\delta} \sum_{i \geq k} \frac{(cn\delta)^i}{i!} \leq \frac{(cn\delta)^{k+1}}{(k+1)!} = \frac{(cn\delta)^{\lceil cn \rceil + 1}}{(\lceil cn \rceil + 1)!}$$

and Stirling's formula shows that this tends to zero for $(c\delta e/c') < 1$. \square

6.4 Convergence

In this section, we prove the convergence of $C^{(n)}$ to a process solving the system (25), and deduce that it solves Smoluchowski's equation (1) before T_c .

Proposition 4. *Assume that the following convergences in distribution hold*

- For every $p \in S$, $C_0^{(n)}(p) \rightarrow c_0(p)$ for some non random $c_0(p) \geq 0$,
- $\sum_{(a,b,m) \in S} aC_0^{(n)}(a,b,m) \rightarrow 1$,
- $\sum_{(a,b,m) \in S} bC_0^{(n)}(a,b,m) \rightarrow 1$.

Let P be a limit point of (P_n) , and let (c_t) be a process with law P . Then a.s. (c_t) solves the system (25), with initial conditions (c_0) .

In the following proofs, we take a subsequence of $(C^{(n)})$ which converges in law to some possibly random $c \in \mathbb{D}([0, +\infty), E)$. For notational simplicity, we will assume that $(C^{(n)})$ itself tends to c . Since E is compact, it is separable, and hence, so is $\mathbb{D}([0, +\infty), E)$. Skorokhod's representation theorem (cf e.g. [8]) now allows us to assume that the $C^{(n)}$ are defined on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$, that $C^{(n)} \rightarrow c$ a.s. (that is, for almost every $\omega \in \Omega$, the function $C^{(n)}(\omega)$ tends to $c(\omega)$ for Skorokhod's distance), and that in the statement, there is a.s. convergence. We will also constantly use the fact that for every bounded Borel function $f : E \rightarrow \mathbb{R}$, the processes

$$C_t^{(n)} - C_0^{(n)} - \int_0^t G^{(n)} f(C_s^{(n)}) ds := M_t^{(n)} \quad (35)$$

and

$$\left(M_t^{(n)} \right)^2 - \int_0^t \left(G^{(n)}(f^2)(C_s^{(n)}) - 2f(C_s^{(n)})G^{(n)}f(C_s^{(n)}) \right) ds \quad (36)$$

are martingales. Note also that if $f : E \rightarrow \mathbb{R}$ is "linear", then for all $\eta \in E$

$$G^{(n)}f(\eta) = \frac{1}{2} \sum_{p,p' \in S} f(\Delta_{p,p'}) p \cdot p' \eta(p) \eta(p') - \frac{1}{2n} \sum_{p \in S} f(\Delta_{p,p}) \eta(p) \quad (37)$$

and that

$$(G^{(n)}(f^2) - 2fG^{(n)}f)(\eta) = \frac{1}{2n} \left(\sum_{p,p' \in S} f(\Delta_{p,p'})^2 \eta(p)\eta(p')p \cdot p' - \frac{1}{n} \sum_{p \in S} f(\Delta_{p,p})^2 \eta(p)p \cdot p \right) \quad (38)$$

We will also need the following convergence result.

Lemma 10. *Let*

$$A_s^{(n)} = \sum_{(a,b,m) \in S} aC_s^{(n)}(a,b,m) \quad \text{and} \quad B_s^{(n)} = \sum_{(a,b,m) \in S} bC_s^{(n)}(a,b,m).$$

Then $(A^{(n)})$ and $(B^{(n)})$ C-converge a.s. to $t \mapsto 1/(1+t)$.

Proof. Obviously, we cannot pass to the limit immediately in these expressions. So consider the maps from E to \mathbb{R} : $C \mapsto \langle a, C \rangle$ and $C \mapsto \langle b, C \rangle$, which are measurable and bounded (by M). By (35), there are martingales $M^{A,(n)}$ and $M^{B,(n)}$ such that

$$A_t^{(n)} = A_0^{(n)} - \int_0^t A_s^{(n)} B_s^{(n)} ds + M_t^{A,(n)} \quad ; \quad B_t^{(n)} = B_0^{(n)} - \int_0^t A_s^{(n)} B_s^{(n)} ds + M_t^{B,(n)}. \quad (39)$$

Now, (36) and (38) show that the quadratic variation of $M^{A,(n)}$ verifies

$$\langle M^{A,(n)} \rangle_t \leq \frac{1}{n} \int_0^t A_s B_s ds \leq \frac{M^2 t}{n}.$$

By Doob's inequality,

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} M_t^{A,(n)} \right)^2 \right) \rightarrow 0$$

for all $T > 0$. Hence there is a subsequence of $(M^{A,(n)})$ which C-converges a.s. to 0. In particular, it S-converges. For notational simplicity, we will assume that $(M^{A,(n)})$ itself converges. For the same reason, we may assume that $(M^{B,(n)})$ also S-converges a.s. to 0.

Now, note that the proof of Lemma 9 still works for $(A^{(n)})$, since the size of its jumps is bounded by $1/n$. So $(A^{(n)})$ is tight, and by Prokhorov's theorem, this means that for almost every $\omega \in \Omega$, $(A^{(n)}(\omega))_{n \geq 0}$ lies in a compact of $\mathbb{D}([0, +\infty), E)$ (actually, this is a consequence of the proof of Lemma 9, and we do not need this implication of Prokhorov's theorem). The same works for $(B^{(n)})$, so we can find $\Omega' \subset \Omega$ with $\mathbb{P}(\Omega') = 1$, such that for every $\omega \in \Omega'$, $C_n(\omega) \rightarrow c(\omega)$ (for Skorokhod's distance), $M^{A,(n)}(\omega) \rightarrow 0$ and $M^{B,(n)}(\omega) \rightarrow 0$ (compactly), and $(A^{(n)}(\omega))_{n \geq 0}$ and $(B^{(n)}(\omega))_{n \geq 0}$ lie in a compact.

Next, fix $\omega \in \Omega'$, and let us find the limit of $(A^{(n)}(\omega), B^{(n)}(\omega))$ (for the product topology — which is not Skorokhod's topology on $\mathbb{D}([0, +\infty), E^2)$). Since it lies in a compact set, it is enough to show that it has only one limit point. So assume $(A^{(n)}(\omega), B^{(n)}(\omega))$ converges to some (A, B) . Then $(A_t^{(n)}(\omega))$ converges to A_t for every $t \in K$, the set of continuity points of A . But A is càdlàg, so it has only countably many points of discontinuity, and hence K^c has Lebesgue-measure 0. Hence $(A^{(n)}(\omega))$ converges to A Lebesgue-a.s., and ditto for $(B^{(n)}(\omega))$. Also, $(A^{(n)}(\omega))$ and $(B^{(n)}(\omega))$ are bounded by M , so using dominated convergence in (39) and recalling that $A_0^{(n)}(\omega)$ and $B_0^{(n)}(\omega) \rightarrow 1$ by assumption, we obtain

$$A_t = 1 - \int_0^t A_s B_s ds \quad ; \quad B_t = 1 - \int_0^t A_s B_s ds.$$

Hence

$$A_t = B_t = \frac{1}{1+t}.$$

Finally there is only one limit point to $(A^{(n)}(\omega), B^{(n)}(\omega))$. So $(A^{(n)}(\omega))$ and $(B^{(n)}(\omega))$ both S-converge to $t \mapsto 1/(1+t)$, and, since this function is continuous, they C-converge. \square

Remark 6. As pointed out in the proof, the convergence of $A^{(n)}$ and $B^{(n)}$ to the actual number of arms

$$A_t = \sum_{(a,b,m) \in S} ac_t(a,b,m) \quad \text{and} \quad B_t = \sum_{(a,b,m) \in S} bc_t(a,b,m)$$

is not obvious. There is no such problem for a strictly sublinear coagulation rate (as in Jeon's proof [17]). In our (linear) case, we prove below that this convergence holds before the critical time (we also refer to Norris [23] for general sublinear rates in a model with no arms). In fact, if there is a solution (c_t) to (1) defined after T_c , we believe that $A_t^{(n)}$ and $B_t^{(n)}$ do not converge to A_t and B_t after T_c (and that this number of arms is then strictly lesser than $1/(1+t)$). This would suggest that T_c is actually a gelation time: some of the arms are lost in a "gel" (a particle with an infinite mass and infinitely many arms).

Proof of Proposition 4. 1. Take some $p_0 = (a_0, b_0, m_0) \in S$, and let for $C \in \mathbb{D}([0, +\infty), E)$, $f(C) = C(p_0)$. According to (35),

$$C_t^{(n)}(p_0) - C_0^{(n)}(p_0) - \int_0^t G^{(n)} f(C_s^{(n)}) ds := M_t^{p_0, (n)} \quad (40)$$

is a martingale. Note also that for $p, p' \in S$, $f(\Delta_{p,p'})$ is 0, except if p or p' or $p \circ p'$ is p_0 . Hence, it is easy to check using (35) that

$$\begin{aligned} G^{(n)} f(C_s^{(n)}) &= - \sum_{p \in S} C_s^{(n)}(p) C_s^{(n)}(p_0) p_0 \cdot p + \frac{1}{2} \sum_{p \preceq p_0} p \cdot (p_0 \setminus p) C_s^{(n)}(p) C_s^{(n)}(p_0 \setminus p) \\ &\quad - \frac{1}{n} \sum_{p \in S} f(\Delta_{p,p}) C_s^{(n)}(p). \end{aligned} \quad (41)$$

The last term is due to the difference between $\lambda_\eta(p, p')$ when $p \neq p'$ and when $p = p'$. In any case, it tends to 0 uniformly on \mathbb{R}^+ and uniformly in p_0 .

2. Let us now study the martingale term. By Doob's inequality, we have for every $T > 0$

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} M_t^{p_0, (n)} \right)^2 \right) \leq 4 \mathbb{E} \left(\left(M_T^{p_0, (n)} \right)^2 \right)$$

and by (36), this last term is

$$\mathbb{E} \left(\int_0^T (G^{(n)} f^2 - 2fG^{(n)} f)(C_s^{(n)}) ds \right).$$

But by (38), and since $f(\Delta_{p,p'}) \leq M$ for all $p, p' \in S$, then $(G^{(n)} f^2 - 2fG^{(n)} f)(C_s^{(n)}) \leq M^4/n$, so that

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} M_t^{p_0, (n)} \right)^2 \right) \rightarrow 0.$$

Hence, there is a subsequence of $(M^{p_0, (n)})$ which a.s. converges to 0 uniformly on \mathbb{R}^+ . For notational simplicity, we will now assume that $(M^{p_0, (n)})$ itself C-converges to 0. Using the diagonal method, we may as well assume that $(M^{p_0, (n)})$ C-converges to 0 for every $p_0 \in S$.

3. We have already seen in the proof of Lemma 9 that $d(C_t^{(n)}, C_{t-}^{(n)}) \leq 3/n$ a.s. Hence

$$\int_0^{+\infty} e^{-s} (\sup_{0 \leq t \leq s} d(X_t, X_{t-}) \wedge 1) ds \rightarrow 0 \quad \text{a.s.}$$

This is enough to ensure that c is almost surely continuous (cf [8]), so $C^{(n)}$ actually C-converges to c . From the definition of d , it is also obvious that $C^{(n)}(p)$ C-converges to $c(p)$ for every $p \in S$.

4. With these results, we may now pass to the limit in (40) and (41). Write (41) in the form

$$G^{(n)} f(C_s^{(n)}) = -\alpha_n(s) + \frac{1}{2} \beta_n(s) + \epsilon_n(s).$$

Equation (40) shows that

$$C_t^{(n)}(p_0) = C_0^{(n)}(p_0) - \int_0^t \alpha_n(s) ds + \frac{1}{2} \int_0^t \beta_n(s) ds + \int_0^t \epsilon_n(s) ds + M_t^{(n)}. \quad (42)$$

By Point 3, $C_t^{(n)}(p_0)$ C-converges a.s. to $c(p_0)$. $C_0^{(n)}(p_0)$ tends to $c_0(p_0)$ by assumption. $\beta_n(t)$ is a finite sum, so

$$\beta_n(t) \rightarrow \sum_{p \leq p_0} p \cdot (p_0 \setminus p) C_s^{(n)}(p) C_s^{(n)}(p_0 \setminus p)$$

compactly. Finally, note that

$$\begin{aligned} \alpha_n(s) &= C_s^{(n)}(p_0) \left(a_0 \sum_{(a,b,m) \in S} b C_s^{(n)}(a, b, m) + b_0 \sum_{(a,b,m) \in S} a C_s^{(n)}(a, b, m) \right) \\ &= C_s^{(n)}(p_0) (a_0 B_s^{(n)} + b_0 A_s^{(n)}). \end{aligned}$$

By Lemma 10, $A_t^{(n)}$ and $B_t^{(n)}$ converge compactly to $t \mapsto 1/(1+t)$, so

$$\lim_{n \rightarrow +\infty} \alpha_n(s) = \frac{a+b}{1+t} c_s(p_0) \quad \text{a.s.}$$

compactly. Since these are all compact convergences, we can pass to the limit in (42), for all $p_0 \in S$. This readily shows that $(c(p))$ solves (25), with initial conditions (c_0) . \square

Theorem 2. *Assume the same hypothesis as in Proposition 4, and assume as well*

$$\sum_{(a,b,m) \in S} (a^2 + b^2) c_0(a, b, m) < +\infty.$$

Let T_c be defined as in Definition 2. Then $(C_t^{(n)})_{t \in [0, T_c]}$ converges (in distribution) to the unique solution of Smoluchowski's equation (1).

Remark 7. Obviously, convergence has to be understood with respect to Skorokhod's topology on $[0, T)$ (which is the trace topology of Skorokhod's topology on $[0, +\infty)$). In particular the sequence of the laws of $(C_t^{(n)})_{t \in [0, T)}$ is tight.

Proof. Let Q_n be the law of $(C_t^{(n)})_{t \in [0, T_c)}$. The sequence (Q_n) is tight. Let Q one of its limit points, and let c a process with law Q . By Proposition 4 above, c solves a.s. the system (25), with initial conditions (c_0) . Now, let $g_t(x, y, z)$ the (a priori random) generating function of c . It is easy to see that g is well defined for $(t, x, y, z) \in [0, T_c) \times (0, 1)^2 \times [0, 1]$, and that $g_t(\cdot, \cdot, z)$ is regular for every $z \in [0, 1]$. Moreover, we see as in the proof of Proposition 3 that for every $z \in [0, 1]$, $g_t(\cdot, \cdot, z)$ solves the PDE (18) with initial conditions $(x, y) \mapsto g_0(x, y, z) = \sum c_0(a, b, m) x^a y^b z^m$. Hence by Proposition 3, (c_t) solves Smoluchowski's equation 1 until T_c . But by Theorem 1, there is a unique solution to this equation on $[0, T_c)$. Hence there is a unique limit point to (Q_n) , so the sequence itself converges to the solution of Smoluchowski's equation on $[0, T_c)$. \square

Acknowledgments This is a part of the author's PhD thesis. I would like to thank my advisors Jean Bertoin and Lorenzo Zambotti for introducing this subject, for their useful advice, and for their encouragement.

References

- [1] D. J. Aldous, *Deterministic and stochastic models for coalescence (aggregation and coagulation): a review of the mean-field theory for probabilists*, Bernoulli 5 (1999), no. 1, 3–48
- [2] J. Bertoin, *Two solvable systems of coagulation equations with limited aggregations*, Ann. Inst. H. Poincaré Anal. Non Linéaire AN (2008), doi:10.1016/j.anihpc.2008.10.007
- [3] J. Bertoin, V. Sidoravicius, M. E. Varès, *A system of grabbing particles related to Galton-Watson trees*, To appear in Random Structures Algorithms. Preprint available at <http://arxiv.org/abs/0804.0726>
- [4] J. Bertoin, V. Sidoravicius, *The structure of typical clusters in large sparse random configurations*. J. Stat. Phys. 135 (2009), 87–105
- [5] J. Carr, F. P. da Costa, *Instantaneous gelation in coagulation dynamics*, Z. Angew. Math. Phys. 43 (1992), no. 6, 974–983
- [6] M. Deaconu, E. Tanré, *Smoluchowski's coagulation equation: probabilistic interpretation of solutions for constant, additive and multiplicative kernels*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 29 (2000), no. 3, 549–579
- [7] M. Dwass, *The total progeny in a branching process and a related random walk*, J. Appl. Prob. 6 (1969), 682–686
- [8] S. N. Ethier, T.G. Kurtz, *Markov processes, characterisation and convergence*, Wiley and Sons, 2005

- [9] M. Escobedo, S. Mischler, B. Perthame, *Gelation in coagulation and fragmentation models*, Comm. Math. Phys. 231 (2002), no. 1, 157–188
- [10] N. Fournier, J-S. Giet, *Convergence of the Marcus-Lushnikov process*, Methodol. Comput. Appl. Probab. 6 (2004), no. 2, 219–231
- [11] N. Fournier, P. Laurençot, *Marcus Lushnikov processes, Smoluchowski's and Flory's models*, Stochastic Process. Appl. 119 (2009), no. 1, 167–189
- [12] A. M. Golovin, *The solution of the coagulation equation for cloud droplets in a rising air current*, Izv. Geophys. Ser. 5 (1963), 482–487
- [13] I. J. Good, *Generalizations to several variables of Lagrange's expansion, with applications to stochastic processes*, Proc. Cambridge Philos. Soc. 56 (1960), 367–380.
- [14] A. Hammond, F. Rezakhanlou, *Moment bounds for the Smoluchowski equation and their consequences*, Comm. Math. Phys. 276 (2007), no. 3, 645–670
- [15] A. Hammond, F. Rezakhanlou, *The kinetic limit of a system of coagulating Brownian particles*, Arch. Ration. Mech. Anal. 185 (2007), no. 1, 1–67
- [16] T.E. Harris, *The Theory of Branching Processes*, Dover Publications, 2002
- [17] I. Jeon, *Existence of gelling solutions for coagulation-fragmentation equations*, Comm. Math. Phys. 194 (1998), no. 3, 541–567.
- [18] A. Joffe, M. Métivier, *Weak convergence of sequences of semimartingales with applications to multitype branching processes*, Adv. in Appl. Probab. 18 (1986), no. 1, 20–65
- [19] J.B. McLeod, *On an infinite set of nonlinear differential equations*, Quart. J. Math. Oxford 13 (1962), 119–128
- [20] A. A. Lushnikov, *Certain new aspects of the coagulation theory*, Izv. Atm. Ok. Fiz. 14 (1978), 738–743
- [21] S. G. Krantz, H.R. Parks, *A primer of real analytic functions*, 2nd ed., Birkhäuser, 2002
- [22] A. H. Marcus, *Stochastic coalescence*, Technometrics 10 (1968), 133–143.
- [23] J. R. Norris, *Smoluchowski's coagulation equation: uniqueness, nonuniqueness and a hydrodynamic limit for the stochastic coalescent*, Ann. Appl. Probab. 9 (1999), no. 1, 78–109.
- [24] M. von Smoluchowski, *Drei Vorträge über Diffusion, Brownsche Molekularbewegung und Koagulation von Kolloidteilchen*, Phys. Zeit. 17 (1916), 557–571 and 585–599