Nonmass Eigenstates of Fermion and Boson Fields

Xin-Bing Huang*

Shanghai United Center for Astrophysics (SUCA), Shanghai Normal University, No.100 Guilin Road, Shanghai 200234, China

Abstract

It appears natural to consider the four dimensional relativistic massive field as a five dimensional massless field. If the fifth coordinate is interpreted as the proper time, then the fifth moment can be understood as the rest mass. After introducing the rest mass operator, we define the mass eigenstate and the nonmass eigenstate. The general equations of spin-0, spin- $\frac{1}{2}$ and spin-1 fields are obtained respectively. It is shown that the Klein-Gordon equation, the Dirac equation and the Proca equation describe the mass eigenstates only. The rest mass of spin- $\frac{1}{2}$ field and the rest mass squared of Boson fields are calculated. The U(1) gauge field that couples to the nonmass eigenstates is studied carefully, whose gauge boson can be massive.

PACS numbers: 12.15.Ff, 11.10.Kk, 12.60.Cn

huangxb@shnu.edu.cn

What is the origin of the rest mass? How to distinguish the massive field and the massless field explicitly from the mathematical point of view? What is the essential difference between the flavor eigenstates and the mass eigenstates? All those problems are fundamental, in which the problem of the rest mass has been studied widely from different viewpoints. In the standard model of electroweak interactions [1, 2], the rest masses of leptons and quarks originate from their Yukawa couplings with the Higgs field [3], and the mismatch between the flavor eigenstates and the mass eigenstates of leptons and quarks is caused by the Higgs interactions [4]. In the 5-dimensional Kaluza-Klein theory [5, 6, 7] and the 11-dimensional string theory (or called M-theory) [8], the quantum field can obtain the rest mass via the compactification of extra dimensional massless particle and interpret the fifth coordinate as the particle's proper time while the fifth moment can be understood as the mass. In this letter, we use the proper time to define the rest mass operator and discuss the nonmass eigenstates of Boson and Fermion fields.

Consider a Minkowskian spacetime with the following metric tensor (covariant components)

$$\eta_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} , \quad (\mu, \nu = 0, 1, 2, 3) .$$
 (1)

In this letter we will use the contravariant three-vector

$$x^{i} = \{x^{1}, x^{2}, x^{3}\} \equiv \{x, y, z\}, \quad (i = 1, 2, 3),$$
 (2)

and four-vector

$$x^{\mu} = \{x^0, x^1, x^2, x^3\} \equiv \{ct, x, y, z\} , \qquad (3)$$

for the description of the spacetime coordinates, where the timelike component is denoted as zero component. The proper time s can be given by

$$s^{2} = x^{\mu}x_{\mu} = c^{2}t^{2} - x^{2} - y^{2} - z^{2} , \qquad (4)$$

where c is the speed of light in vacuum, which is invariant under the Lorentz transformations. From (4) one can acquire that

$$sds = ctd(ct) - xdx - ydy - zdz , \qquad (5)$$

therefore

$$\frac{1}{c}\frac{\partial s}{\partial t} = \frac{ct}{s} , \quad \frac{\partial s}{\partial x} = -\frac{x}{s} , \quad \frac{\partial s}{\partial y} = -\frac{y}{s} , \quad \frac{\partial s}{\partial z} = -\frac{z}{s} , \quad (6)$$

furthermore

$$\frac{\partial}{\partial s} = \frac{s}{c^2 t} \frac{\partial}{\partial t} - \frac{s}{x} \frac{\partial}{\partial x} - \frac{s}{y} \frac{\partial}{\partial y} - \frac{s}{z} \frac{\partial}{\partial z} = \mathbf{n}^{\mu} \partial_{\mu} , \quad \partial_{\mu} \equiv \frac{\partial}{\partial x^{\mu}} , \quad (7)$$

where the contravariant vector \mathbf{n}^{μ} has been defined by

$$\mathbf{n}^{\mu} = \left\{ \frac{s}{ct}, -\frac{s}{x}, -\frac{s}{y}, -\frac{s}{z} \right\} .$$
(8)

In order to discuss the rest mass we start by considering a free particle with the relativistic relation

$$\frac{E^2}{c^2} = p_1^2 + p_2^2 + p_3^2 + m_0^2 c^2 , \qquad (9)$$

here m_0 is the rest mass of the particle. Obviously the rest mass plays the equal role with the component of momentum in above equation, they therefore should have the similar quantization. In elementary quantum mechanics [10], the energy E and the component of momentum p_i are quantized by

$$\hat{E} = i\hbar \frac{\partial}{\partial t} , \quad \hat{p}_i = -i\hbar \frac{\partial}{\partial x^i} , \qquad (10)$$

which are Hermitian operators, namely, $\hat{E}^{\dagger} = \hat{E}$ and $\hat{p}_{i}^{\dagger} = \hat{p}_{i}$. If the rest mass is quantized by a Hermitian operator also, then the real number m_{0} in (9) must be one of the eigenvalues of this Hermitian operator. According to the relativistic relation (9) and the operators of energy and momentum (10), we define the rest mass operator as follows

$$\hat{m} = -i\frac{\hbar}{c}\frac{\partial}{\partial s} , \qquad (11)$$

which is invariant under the Lorentz transformations. It is easy to prove that the rest mass operator is Hermitian if and only if s is timelike.

Assume that a set of eigenfunctions $\sigma_j(x^{\mu})$ $(j = 1, 2, \dots, n)$ constitutes an *n*dimensional complete Hilbert space, which are the eigenfunctions of \hat{m} , and m_j are the corresponding eigenvalues, which are real and nonnegative—as the rest mass, of course, must be. We reexpress this assumption in the mathematical language

$$\hat{m}\sigma_j(x^\mu) = -i\frac{\hbar}{c}\frac{\partial\sigma_j(x^\mu)}{\partial s} = m_j\sigma_j(x^\mu) , \quad (j = 1, 2, \cdots, n) .$$
(12)

Above assumption shows that the mass eigenstate has been defined by the eigenfunction of the rest mass operator. According to quantum mechanics [10], we can then define the nonmass eigenstate $\sigma(x^{\mu})$ by

$$\sigma(x^{\mu}) = \sum_{j=1}^{n} a_j \sigma_j(x^{\mu}) , \qquad \sum_{j=1}^{n} a_j a_j^* = 1 , \qquad (13)$$

where a_j is complex and a_j^* the complex conjugate of a_j .

In quantum field theories, the mass operator had been introduced and the nonmass eigenstates had also been used to denote the off-shell states [11, 12]. But we give a quite different definition for the rest mass operator. The nonmass eigenstate discussed by us will devote it to a better understanding of particle mixing.

From elementary quantum mechanics it is known that the schrödinger equation corresponds to the non-relativistic energy relation in operator form. we therefore replace the energy, the momentum and the rest mass in the relativistic relation (9) by their corresponding operators (10) and (11) to get a relativistic equation

$$\left(\frac{\partial^2}{c^2\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} - \frac{\partial^2}{\partial s^2}\right)\phi(x^{\mu}) = 0 , \qquad (14)$$

here $\phi(x^{\mu})$ is a nonmass eigenstate of the free scalar fields. This equation is invariant under the Lorentz transformations. For a mass-squared eigenstate $\phi(x^{\mu})$, we obtain

$$\hat{m}^2 \phi(x^{\mu}) = -\frac{\hbar^2}{c^2} \frac{\partial^2 \phi(x^{\mu})}{\partial s^2} = m_{\phi}^2 \phi(x^{\mu}) .$$
(15)

Then (14) reduces to

$$\left(\frac{\partial^2}{c^2\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} + \frac{m_\phi^2 c^2}{\hbar^2}\right)\phi(x^\mu) = 0.$$
(16)

The definition of the rest mass operator \hat{m} shows that (15) is the eigen equation of the operator \hat{m}^2 . Therefore, the function $\phi(x^{\mu})$ should be called the mass-squared eigenstate of a scalar field. (16) is the Klein-Gordon equation.

Assume that a set of mass-squared eigenstates $\phi_j(x^{\mu})$ given by (15) constitutes an *n*-dimensional complete Hilbert space, where m_j is the corresponding rest mass. Then the nonmass eigenstate of spin-0 fields is generally given by

$$\phi(x^{\mu}) = \sum_{j=1}^{n} a_j \phi_j(x^{\mu}) , \qquad \sum_{j=1}^{n} a_j a_j^* = 1 .$$
(17)

Since $\phi_j(x^{\mu})$ are mass-squared eigenstates, one can only obtain the square rest mass of a nonmass eigenstate defined by (17) as follows

$$m^2 = \sum_{j=1}^n a_j a_j^* m_j^2 . (18)$$

Following the historical approach of Dirac who, in 1928, obtained a relativistic covariant wave equation for spin- $\frac{1}{2}$ field, we give the relativistic covariant equation for the nonmass eigenstate of spin- $\frac{1}{2}$ fields with a general potential $V(x^{\mu})$

$$i\hbar \frac{\partial \psi(x^{\mu})}{\partial t} = \left[-i\hbar c\hat{\alpha}^{i} \frac{\partial}{\partial x^{i}} - i\hbar c\hat{\beta} \frac{\partial}{\partial s} + V \right] \psi(x^{\mu}) , \qquad (19)$$

where $\psi(x^{\mu})$ is a 4×1 matrix, and $\hat{\alpha}^{i}, \hat{\beta}$ are 4×4 Hermitian matrices defined by Dirac. One can prove the covariance of this equation by noticing that the operator $\frac{\partial}{\partial s}$ is invariant under the Lorentz transformations.

For a mass eigenstate $\psi(x^{\mu})$, we get

$$\hat{m}\psi(x^{\mu}) = -i\frac{\hbar}{c}\frac{\partial\psi(x^{\mu})}{\partial s} = m_{\psi}\psi(x^{\mu}) , \qquad (20)$$

and

$$\left(i\hbar\gamma^{\mu}\frac{\partial}{\partial x^{\mu}} - m_{\psi}c\right)\psi(x^{\mu}) = \frac{V}{c}\gamma^{0}\psi(x^{\mu}) , \qquad (21)$$

here we have adopted the definitions of $\gamma^{\mu} = \{\gamma^0, \gamma^1, \gamma^2, \gamma^3\}$ and $\gamma^0 = \hat{\beta}, \ \gamma^i = \hat{\beta}\hat{\alpha}^i$. From above analysis, we can draw a conclusion that the mass eigenstate of the spin- $\frac{1}{2}$ field is described by the well-known Dirac equation (21).

From (19) we find that the Lagrange density of a nonmass eigenstate of free spin- $\frac{1}{2}$ particles has the form[†]

$$\mathcal{L}_{1n} = \bar{\psi}(x^{\mu}) \left(i\hbar c\gamma^{\mu} \frac{\partial}{\partial x^{\mu}} + i\hbar c \frac{\partial}{\partial s} \right) \psi(x^{\mu}) .$$
(22)

 $\bar{\psi} \equiv \psi^{\dagger} \gamma^{0}$ is called the spinor adjoint to ψ . The variation of above \mathcal{L}_{1n} with respect to $\bar{\psi}(x, z)$ yields the general equation for a nonmass eigenstate of free spin- $\frac{1}{2}$ Fermions

$$\left(\gamma^{\mu}\frac{\partial}{\partial x^{\mu}} + \frac{\partial}{\partial s}\right)\psi(x^{\mu}) = 0.$$
(23)

The above equation is (19) with V = 0. The Dirac equation in the models of flat 1 + 4 dimensional spacetime is generally given by [6, 7, 13, 14]

$$\left(i\hbar\gamma^{\mu}\frac{\partial}{\partial x^{\mu}} + i\hbar\gamma^{5}\frac{\partial}{\partial x^{5}} - mc\right)\Psi(x, x^{5}) = 0 , \qquad (24)$$

where the metric of 5-dimensional spacetime is of the signature (+, -, -, -, -) and $\gamma^5 = -\gamma_5 \equiv -i\gamma^0\gamma^1\gamma^2\gamma^3$. Obviously (23) can be treated as the massless 5-dimensional Dirac equation.

We can obtain the Lagrange density of a mass eigenstate of the free spin- $\frac{1}{2}$ particle from (22), that is

$$\mathcal{L}_{1m} = \bar{\psi}(x^{\mu}) \left(ic\hbar\gamma^{\mu} \frac{\partial}{\partial x^{\mu}} - m_{\psi}c^2 \right) \psi(x^{\mu}) , \quad \bar{\psi} \equiv \psi^{\dagger}\gamma^0 .$$
⁽²⁵⁾

Assume that a set of mass eigenstates $\psi_j(x^{\mu})$ given by (20) constitutes an *n*dimensional complete Hilbert space, where m_j is the corresponding rest mass. Then the nonmass eigenstate of spin- $\frac{1}{2}$ fields must be

$$\psi(x^{\mu}) = \sum_{j=1}^{n} a_j \psi_j(x^{\mu}) , \qquad \sum_{j=1}^{n} a_j a_j^* = 1 .$$
(26)

[†]In this letter \mathcal{L}_{1n} denotes the Lagrange density of one nonmass eigenstate and \mathcal{L}_{1m} denotes the Lagrange density of one mass eigenstate.

According to quantum mechanics, the rest mass of nonmass eigenstate $\psi(x^{\mu})$ given by (26) is therefore of the form

$$m = \sum_{j=1}^{n} a_j a_j^* m_j .$$
 (27)

In a word, we have given the equation for a nonmass eigenstate of spin- $\frac{1}{2}$ fields and the rest mass of a nonmass eigenstate.

The gauge field theories [2, 15, 16, 17, 18] tell us a good method to introduce the spin-1 fields in our framework. In quantum field theories, we are familiar with the electromagnetic field, which is a massless U(1) gauge field. From the Lagrange densities (22) and (25), we find that both of them admit the introduction of a U(1)gauge field. Since \mathcal{L}_{1m} can be treated as the Lagrangian of a free charged spin- $\frac{1}{2}$ particle in quantum field theory, introducing a U(1) gauge field into the Lagrange density (25) will directly give an electromagnetic field. The total Lagrangian

$$\mathcal{L}_{1mt} = \bar{\psi} \left(ic\hbar\gamma^{\mu} \frac{\partial}{\partial x^{\mu}} - e\gamma^{\mu}A_{\mu} - m_{\psi}c^{2} \right) \psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$$
(28)

is invariant under the following local U(1) transformation

$$\psi'(x^{\mu}) = e^{i\theta(x^{\mu})}\psi(x^{\mu}) , \qquad A'_{\mu} = A_{\mu} - \frac{\hbar c}{e} \frac{\partial \theta(x^{\mu})}{\partial x^{\mu}} , \qquad (29)$$

where $\theta(x^{\mu})$ is a function of x^{μ} , and A_{μ} is the electromagnetic field. The electromagnetic field strength tensor $F_{\mu\nu} = \frac{\partial A_{\nu}}{\partial x^{\mu}} - \frac{\partial A_{\mu}}{\partial x^{\nu}}$ is invariant under above transformation as well.

From now on we will pay more attention to introducing a U(1) gauge field into the Lagrange density \mathcal{L}_{1n} . To make the gauge invariance explicit, we formally define

$$x^4 = -x_4 = s , \quad \gamma^4 = -\gamma_4 = \mathbf{1} .$$
 (30)

Thus the Lagrange density of a nonmass eigenstate of free spin- $\frac{1}{2}$ fields (22) becomes

$$\mathcal{L}_{1n} = \bar{\psi}(x^{\mu}) \left(ic\hbar\gamma^{\alpha} \frac{\partial}{\partial x^{\alpha}} \right) \psi(x^{\mu}) , \quad \alpha = 0, 1, 2, 3, 4 .$$
(31)

Let us multiply the nonmass eigenstate $\psi(x^{\mu})$ by a local phase $e^{i\Theta(x^{\mu})}$, that is

$$\psi'(x^{\mu}) = e^{i\Theta(x^{\mu})}\psi(x^{\mu})$$
 (32)

We can reexpress the above equation as doing a U(1) transformation on $\psi(x^{\mu})$ because the phase factor $e^{i\Theta(x^{\mu})}$ is an element of U(1) group. So naturally

$$\bar{\psi}'(x^{\mu}) = e^{-i\Theta(x^{\mu})}\bar{\psi}(x^{\mu})$$
 (33)

The crucial result is that the total Lagrange density

$$\mathcal{L}_{1nt} = \bar{\psi}(x^{\mu}) \left(ic\hbar\gamma^{\alpha} \frac{\partial}{\partial x^{\alpha}} - g\gamma^{\alpha} \mathbf{A}_{\alpha}(x^{\mu}) \right) \psi(x^{\mu}) - \frac{1}{4} \mathbf{F}_{\alpha\beta}(x^{\mu}) \mathbf{F}^{\alpha\beta}(x^{\mu})$$
(34)

is invariant under a group of local gauge transformations, given by (32), (33) and

$$\mathbf{A}_{\alpha}'(x^{\mu}) = \mathbf{A}_{\alpha}(x^{\mu}) - \frac{\hbar c}{g} \frac{\partial \Theta(x^{\mu})}{\partial x^{\alpha}} , \qquad (35)$$

g in above equations is the coupling constant. The strength tensor of U(1) gauge field is of the form

$$\mathbf{F}_{\alpha\beta}(x^{\mu}) = \frac{\partial \mathbf{A}_{\beta}(x^{\mu})}{\partial x^{\alpha}} - \frac{\partial \mathbf{A}_{\alpha}(x^{\mu})}{\partial x^{\beta}} , \qquad (36)$$

which is invariant under the transformations of (32), (33) and (35) as well. Now we will prove that $\mathbf{A}_{\alpha}(x^{\mu})$ is a four dimensional covariant vector. To do this, we decompose $\mathbf{A}_{\alpha}(x^{\mu})$ into two parts

$$\mathbf{A}_{\alpha} \equiv \{\mathbf{A}_{\mu}, \mathbf{A}_{s}\} \ . \tag{37}$$

Considering (7) and (34) together, we get

$$\mathbf{A}_{s} = \frac{s}{ct}\mathbf{A}_{0} - \frac{s}{x}\mathbf{A}_{1} - \frac{s}{y}\mathbf{A}_{2} - \frac{s}{z}\mathbf{A}_{3} = \mathbf{n}^{\mu}\mathbf{A}_{\mu} .$$
(38)

Combining (35) with (38) gives

$$\mathbf{A}_{s}^{\prime} = \mathbf{n}^{\mu}\mathbf{A}_{\mu}^{\prime} = \mathbf{n}^{\mu}\left(\mathbf{A}_{\mu} - \frac{\hbar c}{g}\frac{\partial\Theta}{\partial x^{\mu}}\right) = \mathbf{A}_{s} - \frac{\hbar c}{g}\frac{\partial\Theta}{\partial s} .$$
(39)

Therefore \mathbf{A}_{α} is a four-vector and (38) is in accordance with (39).

Compare the Lagrange density (34) with (28), we find that the U(1) gauge field $\mathbf{A}_{\alpha}(x^{\mu})$ can be treated as a 5-dimensional Maxwell's electromagnetic field. Therefore

$$\frac{\partial \mathbf{F}_{\alpha\beta}(x^{\mu})}{\partial x^{\gamma}} + \frac{\partial \mathbf{F}_{\gamma\alpha}(x^{\mu})}{\partial x^{\beta}} + \frac{\partial \mathbf{F}_{\beta\gamma}(x^{\mu})}{\partial x^{\alpha}} = 0 , \qquad (40)$$

and

$$\frac{\partial \mathbf{F}_{\alpha\beta}(x^{\mu})}{\partial x_{\alpha}} = \mathbf{J}_{\beta}(x^{\mu}) , \qquad (41)$$

where $\mathbf{J}_{\beta}(x^{\mu})$ is the 5-dimensional current.

Substituting (36) into (41) we find that $\mathbf{A}_{\alpha}(x^{\mu})$ satisfies

$$\frac{\partial}{\partial x_{\alpha}} \frac{\partial \mathbf{A}_{\beta}(x^{\mu})}{\partial x^{\alpha}} - \frac{\partial}{\partial x^{\beta}} \frac{\partial \mathbf{A}_{\alpha}(x^{\mu})}{\partial x_{\alpha}} = \mathbf{J}_{\beta}(x^{\mu}) .$$
(42)

We may now make use of the freedom (35) and choose a particular $\Theta(x^{\mu})$ so that the transformed $\mathbf{A}_{\alpha}(x^{\mu})$ satisfies the following gauge condition:

$$\frac{\partial \mathbf{A}_{\alpha}}{\partial x_{\alpha}} = 0 \ . \tag{43}$$

In this special "choice of gauge", (42) becomes

$$\frac{\partial}{\partial x_{\alpha}} \frac{\partial \mathbf{A}_{\beta}(x^{\mu})}{\partial x^{\alpha}} = \mathbf{J}_{\beta}(x^{\mu}) .$$
(44)

In vacuo, there are no current, namely $\mathbf{J}_{\beta}(x^{\mu}) = 0$, (44) reduces to

$$\left(\frac{\partial^2}{c^2\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} - \frac{\partial^2}{\partial s^2}\right)\mathbf{A}_{\alpha} = 0.$$
(45)

This is the general equation of a nonmass eigenstate of free spin-1 fields. Under this special choice of gauge, let us consider a mass-squared eigenstate of \mathbf{A}_{μ} , which is given by

$$\hat{m}^2 \mathbf{A}_{\mu} = -\frac{\hbar^2}{c^2} \frac{\partial^2 \mathbf{A}_{\mu}}{\partial s^2} = m_{\mathbf{A}}^2 \mathbf{A}_{\mu} , \quad m_{\mathbf{A}} \ge 0 .$$
(46)

Then (45) reduces to

$$\left(\frac{\partial^2}{c^2\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2} + \frac{m_{\mathbf{A}}^2 c^2}{\hbar^2}\right)\mathbf{A}_{\mu} = 0.$$
(47)

Hence we have shown that the mass-squared eigenstates (or simply called "the mass eigenstates") of free vector field satisfies Proca equation (47). Therefore the gauge boson of U(1) gauge field that couples to the nonmass eigenstates can be massive.

Assume that a set of mass-squared eigenstates $[\mathbf{A}_{\mu}(x^{\mu})]_{j}$ given by (46) constitutes an *n*-dimensional complete Hilbert space, where m_{j} is the corresponding rest mass. Then the nonmass eigenstate of vector fields is expressed by

$$\mathbf{A}_{\mu}(x^{\mu}) = \sum_{j=1}^{n} a_j \left[\mathbf{A}_{\mu}(x^{\mu}) \right]_j , \qquad \sum_{j=1}^{n} a_j a_j^* = 1$$
(48)

It is proved that $[\mathbf{A}_{\alpha}(x, z)]_{j}$ are mass-squared eigenstates, one can calculate the square rest mass of the nonmass eigenstate defined by (48), namely

$$m^2 = \sum_{j=1}^n a_j a_j^* m_j^2 . (49)$$

Obviously the vector field has the same rest mass formula as that of the scalar field.

It is well-known that the mass term in Lagrangian of a charged particle must be invariant under the Lorentz transformations and the local gauge transformations, therefore, for a spin- $\frac{1}{2}$ nonmass eigenstate who couples with a U(1) gauge field, the rest mass operator must be invariant not only under the Lorentz transformations but also under the U(1) gauge transformations. Hence, the rest mass operator of $\psi(x^{\mu})$ in Lagrangian (34) should be

$$\hat{M} = -i\frac{\hbar}{c} \left(\frac{\partial}{\partial s} + i\frac{g}{\hbar c} \mathbf{A}_s \right) \ . \tag{50}$$

One can easily prove that $\bar{\psi}(x^{\mu})\hat{M}\psi(x^{\mu})$ is invariant under the Lorentz transformations and the local U(1) gauge transformations.

In the Standard Model, the Yukawa interactions of the quarks with the Higgs condensate cause the mismatch between the flavor eigenstates d'_k and the mass eigenstates d_l , namely $d'_k \equiv \sum_l V_{kl} d_l$, k, l = 1, 2, 3, and V is the Cabibbo-Kobayashi-Maskawa mixing matrix [19, 20], which is unitary $V^{\dagger}V = 1$. In our framework, the mass eigenstates of the quarks must be written as d_l , where m_l is the corresponding mass of the quark. Then the nonmass eigenstates of the quarks are of the form

$$d_k'' = \sum_{l=1}^3 V_{kl} d_l \ . \tag{51}$$

In the forthcoming papers [21, 22] we will prove that the nonmass eigenstates play an important role in constructing an electroweak model without Higgs mechanism.

By interpreting the proper time as the fifth coordinate, we define the operator of the rest mass and give the concepts of mass eigenstate and nonmass eigenstate. The general equations for nonmass eigenstates of free spin-0, spin- $\frac{1}{2}$ and spin-1 fields are obtained. It is found that there are two kinds of U(1) gauge fields: The U(1) gauge field of first kind merely couples to mass eigenstates, in which the gauge boson is massless. The second kind of U(1) gauge field couples to nonmass eigenstates, whose gauge boson may be massive.

Acknowledgement: I am grateful to Prof. Chao-Jun Feng and Prof. Dao-Jun Liu for their enlightening discussions. It is very important that Prof. Chao-Jun Feng indicated that the rest mass operator could be well defined by proper time without introducing an extra dimension.

References

- S. L. Glashow, Nucl. Phys. 22, 579 (1960); J. Goldstone, A. Salam and S. Weinberg, Phys. Rev. 127, 965 (1962); S. Weinberg, Phys. Rev. Lett. 19, 1264 (1967);
 S. L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D 2, 1285 (1970).
- [2] W. Greiner and B. Müller, *Gauge Theory of Weak Interactions*, (3rd. edition), (Springer-Verlag, 2000).
- [3] P. W. Higgs, Phys. Lett. 12, 132 (1964); P. W. Higgs, Phys. Rev. Lett. 13, 508 (1964); P. W. Higgs, Phys. Rev. 145, 1156 (1966); T. W. Kibble, Phys. Rev. 155, 1554 (1967); F. Englert and R. Brout, Phys. Rev. Lett. 13, 321 (1964).
- [4] Z. Z. Xing, Int. J. Mod. Phys. A **19**, 1 (2004).
- [5] O. M. Lecian and G. Montani, Int. J. Mod. Phys. D 15, 717 (2006).
- [6] K. S. Soh and P. Y. Pac, Phys. Rev. D 35, 544 (1987).
- [7] A. Macias and H. Dehnen, Class. Quantum Grav. 8, 203 (1991).

- [8] J. Polchinski, String Theory, Vols. I, II (Cambridge University Press, 1998).
- [9] L. Freidel, F. Girelli and E. R. Livine, Phys. Rev. D 75, 105016 (2007)[arXiv:hep-th/0701113].
- [10] D. J. Griffiths, Introduction to Quantum Mechanics, (2nd. edition), (Pearson Education, 2005).
- [11] S. Weinberg, *The Quantum Theory of Fields*, Vol. I, (Cambridge University Press, 1995).
- [12] N. Straumann, "Unitary Representations of the Inhomogeneous Lorentz Group and Their Significance in Quantum Physics", [arXiv:math-ph/0809.4942].
- [13] S. Ichinose, Phys. Rev. D 66, 104015 (2002)[arXiv:hep-th/0206187].
- [14] S.-Q. Wu, Phys. Rev. D 78, 064052 (2008) [arXiv:hep-th/0807.2114].
- [15] F. Scheck, *Electroweak and Strong Interactions*, (2nd. edition), (Springer-Verlag, 1996).
- [16] P. H. Frampton, Gauge Field Theories, (3rd. edition), (WILEY-VCH Verlag GmbH & Co. KGaA, 2008).
- [17] T.-P. Cheng and L.-F. Li, Gauge Theory of Elementary Particle Physics, (Clarendon Press, 1984).
- [18] L. H. Ryder, *Quantum Field Theory*, (2nd. edition), (Cambridge University Press, 1996).
- [19] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
- [20] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, 652 (1973).
- [21] X.-B. Huang, "Massive Gauge Bosons in Yang-Mills Theory without Higgs Mechanism", [arXiv:hep-ph/0906.2584].

[22] X.-B. Huang, "An Electroweak Model without Higgs Mechanism", in preparation.