

# Hadronic physics at KLOE

Roberto Versaci on behalf of the KLOE collaboration <sup>1</sup>

*Dipartimento di Energetica dell'Università "Sapienza", Roma, Italy  
Laboratori Nazionali di Frascati dell'INFN, Frascati, Italy*

**Abstract.** New KLOE results on scalar mesons,  $\gamma\gamma$  physics and  $\eta$  physics are presented.

**Keywords:** Scalar mesons,  $\gamma\gamma$  physics,  $\eta$  meson.

**PACS:** 13.25.Es, 13.25.Jx, 13.66.Bc, 14.40.Aq, 14.40.Cs

## 1. SCALAR PHYSICS

The structure of the scalars below 1 GeV needs to be further clarified. Several models have been proposed to describe them (e.g.  $q\bar{q}$ , four quarks,  $K\bar{K}$  molecules, etc.). The decay of the scalars into two pseudoscalars ( $S \rightarrow PP'$ ) can be used to investigate their nature because the branching ratios and the invariant mass of the two pseudoscalars are sensitive to the scalar structure.

$\phi \rightarrow a_0(980)\gamma \rightarrow \eta\pi^0\gamma$  decay [1]. For this measurement about 400 pb<sup>-1</sup> of KLOE collected data have been used. The analysis has been performed for two different  $\eta$  final states, i.e.  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ . A kinematic fit has been performed imposing the four momentum conservation, the photon velocity and the invariant masses of both  $\eta$  and  $\pi^0$ . The  $\eta\pi^0$  invariant mass distribution has been fitted with the “no-structure”[2] and the “kaon loop”[3] models after background subtraction. The results of the fit are shown in table 1. It is interesting to note that both models give a large coupling of the  $a_0(980)$  with the  $\phi$  meson, indicating a sizable strange quark content in the  $a_0(980)$ . The branching ratio obtained for the two different decay chains are in agreement:  $BR(\phi \rightarrow \eta\pi^0\gamma) = (7.01 \pm 0.10_{stat.} \pm 0.20_{syst.}) \times 10^{-5}$  for the  $\eta \rightarrow \gamma\gamma$  final state and  $BR(\phi \rightarrow \eta\pi^0\gamma) = (7.12 \pm 0.13_{stat.} \pm 0.22_{syst.}) \times 10^{-5}$  for the  $\eta \rightarrow \pi^+\pi^-\pi^0$  final state.

$\phi \rightarrow K^0\bar{K}^0\gamma$  decay [4]. This decay allegedly proceeds through the intermediate  $f_0(980)$  (I=0) and  $a_0(980)$  (I=1) scalar mesons:  $\phi \rightarrow (f_0 + a_0)\gamma \rightarrow K^0\bar{K}^0\gamma$ . The kaon pair is produced in a  $J^{PC} = 0^{++}$  state, so the two kaons are both  $K_S$  or  $K_L$ . We have searched for a final state with a  $K_S K_S$ , with both  $K_S$ ' decaying to  $\pi^+\pi^-$ . This request

---

<sup>1</sup> F. Ambrosino, A. Antonelli, M. Antonelli, F. Archilli, P. Beltrame, G. Bencivenni, S. Bertolucci, C. Bini, C. Bloise, S. Bocchetta, F. Bossi, P. Branchini, G. Capon, T. Capussela, F. Ceradini, P. Ciambrone, E. De Lucia, A. De Santis, P. De Simone, G. De Zorzi, A. Denig, A. Di Domenico, C. Di Donato, B. Di Micco, M. Dreucci, G. Felici, S. Fiore, P. Franzini, C. Gatti, P. Gauzzi, S. Giovannella, E. Graziani, G. Lanfranchi, J. Lee-Franzini, M. Martini, P. Massarotti, S. Meola, S. Miscetti, M. Moulson, S. Müller, F. Murtas, M. Napolitano, F. Nguyen, M. Palutan, E. Pasqualucci, A. Passeri, V. Patera, P. Santangelo, B. Sciascia, T. Spadaro, M. Testa, L. Tortora, P. Valente, G. Venanzoni, R. Versaci, G. Xu

**TABLE 1.** Output of the fit to the  $\eta\pi^0$  invariant mass with two different models: kaon loop and no structure.


	Kaon loop	No structure
$M_{a_0}$ [MeV]	$982.5 \pm 1.6 \pm 1.1$	982.5 (Fixed)
$g_{a_0 K^+ K^-}$ [GeV]	$2.15 \pm 0.06 \pm 0.06$	$2.01 \pm 0.07 \pm 0.28$
$g_{a_0 \eta \pi^0}$ [GeV]	$2.82 \pm 0.03 \pm 0.04$	$2.46 \pm 0.08 \pm 0.11$
$g_{\phi a_0 \gamma}$ [GeV $^{-1}$ ]	$1.58 \pm 0.10 \pm 0.16^*$	$1.83 \pm 0.03 \pm 0.08$
$\text{BR}(\phi \rightarrow \rho\pi \rightarrow \eta\pi\gamma)$	$(0.92 \pm 0.40 \pm 0.15) \cdot 10^{-6}$	$(0.05 \pm 4 \pm 0.07) \cdot 10^{-6}$
$\text{BR}(\eta \rightarrow \gamma\gamma)/\text{BR}(\eta \rightarrow \pi^+\pi^-\pi^0)$	$1.70 \pm 0.04 \pm 0.03$	$1.70 \pm 0.03 \pm 0.01$
$\chi^2$ probability	0.104	0.309

\* Not a free parameter of the fit in this model. Calculated from other fit outputs.

reduces the probability of observation to  $\sim 22\%$ , but selects a class of event with a clear signature: four tracks and a low energy photon coming from the interaction point. In this analysis the whole KLOE dataset,  $\sim 2.2 \text{ fb}^{-1}$ , has been used. At the end of which we have observed 5 events in the data, while we were expecting  $3.2 \pm 0.7$  background events from MC. A Cousins-Feldman approach has been used [5] and a 90% confidence level upper limit on the branching ratio has been obtained:  $\text{BR}(\phi \rightarrow K^0 \bar{K}^0 \gamma) < 1.9 \times 10^{-8}$ . This measurement excludes some of the theoretical predictions and is in agreement with expectations from other KLOE measurements (see figure 1 left, reference [4] and references therein).

## 2. $\gamma\gamma$ PHYSICS

KLOE has been making a pilot study for the search  $\gamma\gamma \rightarrow \sigma(600) \rightarrow \pi^0\pi^0$  using  $11 \text{ pb}^{-1}$  from the  $240 \text{ pb}^{-1}$  taken at  $\sqrt{s} = 1000 \text{ MeV}$  [6]. At this energy the background



**FIGURE 1.** **Left:** comparison between theoretical predictions for the  $\phi \rightarrow K^0 \bar{K}^0 \gamma$  branching ratio and KLOE upper limit. The grey area represents the expected range for the branching ratio using other KLOE results on scalar mesons. **Right:** search for  $\gamma\gamma \rightarrow \sigma(600) \rightarrow \pi^0\pi^0$ ; fit to the invariant mass of the four photons,  $M_{4\gamma}$ , with MC shapes of expected source of background. Dots: data; atched yellow: total background; red:  $\phi \rightarrow \eta\gamma \rightarrow \pi^0\pi^0\pi^0\gamma$ ; blue:  $e^+e^- \rightarrow \omega\pi^0 \rightarrow \pi^0\pi^0\gamma$ ; green:  $\phi \rightarrow K_S K_L$ ; cyan:  $\phi \rightarrow f_0\gamma$ ; magenta:  $e^+e^- \rightarrow \gamma\gamma$ .

**TABLE 2.** Output of the fit imposing or not the gluonium content to be zero.

	<b>Gluonium content forced to be zero</b>	<b>Gluonium content free</b>
$Z_G^2$	fixed 0	$0.115 \pm 0.036$
$\phi_P$	$(41.4 \pm 0.5)^\circ$	$(40.4 \pm 0.6)^\circ$
$Z_q$	$0.93 \pm 0.02$	$0.94 \pm 0.03$
$Z_s$	$0.82 \pm 0.05$	$0.83 \pm 0.05$
$\phi_V$	$(3.34 \pm 0.09)^\circ$	$(3.32 \pm 0.09)^\circ$
$m_s/\bar{m}$	$1.24 \pm 0.07$	$1.24 \pm 0.07$
$\chi^2 / \text{dof}$	14.7/4	4.6/3
$P(\chi^2)$	0.005	0.20


from  $\phi$  decays is very small. We have performed a fit to the four photons invariant mass ( $M_{4\gamma}$ ), using the shapes of the known sources, see figure 1, right. The result of the fit is very poor  $\chi^2/\text{dof} = 441/94$ , showing an excess of events in the expected  $\sigma(600)$  region, compared to what expected from MC, therefore pointing towards a search for the signal in the  $240 \text{ pb}^{-1}$ .

### 3. PSEUDOSCALAR PHYSICS

The  $\phi$  meson decays about 1.3% of times into  $\eta\gamma$ , this implies DAΦNE is an  $\eta$ -factory. KLOE has collected one of the largest sample of  $\eta$  mesons in the world, about  $10^8$ .

*$\eta - \eta'$  mixing and  $\eta'$  gluonium content [7].* The KLOE paper on  $\eta - \eta'$  mixing [7], suggesting for a  $3\sigma$  evidence of gluonium content in the  $\eta'$  meson, has triggered a large amount of discussion among theoreticians. Therefore we have decided to perform a new and more detailed study of this topic. We have considered  $\eta$  and  $\eta'$  in the quark mixing base as described in [8] ( $|\eta'\rangle = X_{\eta'}|q\bar{q}\rangle + Y_{\eta'}|s\bar{s}\rangle + Z_G|G\rangle$ ). The new fit we have performed has more constraints thus allowing an independent determination of more free parameters. We use the BR values from PDG 2008 [9] and the new KLOE results on the  $\omega$  meson [10]. The fit has been performed both imposing the gluonium content to be zero or allowing it free. The results are shown in table 2: gluonium content of the  $\eta'$  is confirmed at  $3\sigma$  level.

*$\eta$  decays into four charged particles [11].* KLOE has started to study the decays of the  $\eta$  into four charged particles, using  $1.7 \text{ fb}^{-1}$  of data. This decay is interesting because it allows us to probe the  $\eta$  internal structure exploiting the conversion of the virtual photon into a lepton pair [12]. It is also interesting because a non-CKM CP violating mechanism has been suggest to be present in this decay[13], and should manifest as an angular asymmetry  $A_\phi$ , between the pion and the electron decay planes in the  $\eta$  rest frame. After background rejection a fit of the sidebands of the four tracks invariant distribution has been performed to obtain the background scale factors. Most of the background is due to  $\phi$  decays, but there is still a non-negligible contribution from continuum events. Signal events have been counted in the  $\eta$  mass region, giving  $BR(\eta \rightarrow$



**FIGURE 2.** **Left and center:**  $\eta \rightarrow \pi^+\pi^-e^+e^-$  analysis;  $\pi^+\pi^-e^+e^-$  invariant mass and angular asymmetry distributions. Dots: data. The black histogram is the expected distribution, i.e. signal MC (dark grey),  $\phi$  background (light grey) and continuum background (white). **Right:**  $\eta \rightarrow e^+e^-e^+e^-$  analysis; fit of the four electron invariant mass,  $M_{eeee}$ .

$\pi\pi ee) = (26.8 \pm 0.9_{Stat.} \pm 0.7_{Syst.}) \times 10^{-5}$  and  $A_\phi = (-0.6 \pm 2.5_{Stat.} \pm 1.8_{Syst.}) \times 10^{-2}$  [11], see figure 2 left and center.

More recently KLOE has started studying the  $\eta \rightarrow e^+e^-e^+e^-$  decay. This decay, together with the  $\eta \rightarrow \mu^+\mu^-e^+e^-$ , is interesting for the  $\eta$  meson form factor because there are only leptons in the final state. The analysis strategy is similar to the  $\pi\pi ee$  one. Most of the background comes from continuum events and a small contribution is due to  $\phi$  decays. The latter is subtracted from data using the MC shape. The number of events is obtained fitting the data distribution of the 4 electron invariant mass,  $M_{eeee}$ , with signal and background shapes (figure 2, right). From the fit we obtain 413 events. This constitutes the first observation of this decay.

## REFERENCES

1. F. Ambrosino, et al. (2009), 0904.2539.
2. G. Isidori, L. Maiani, M. Nicolaci, and S. Pacetti, *JHEP* **05**, 049 (2006), hep-ph/0603241.
3. N. N. Achasov, and A. V. Kiselev, *Phys. Rev.* **D68**, 014006 (2003), hep-ph/0212153.
4. F. Ambrosino, et al. (2009), 0903.4115.
5. G. J. Feldman, and R. D. Cousins, *Phys. Rev.* **D57**, 3873–3889 (1998), physics/9711021.
6. F. Ambrosino, et al., *Nuovo Cim.* **31C**, 415 (2008).
7. F. Ambrosino, et al., *Phys. Lett.* **B648**, 267–273 (2007), hep-ex/0612029.
8. J. L. Rosner, *Phys. Rev.* **D27**, 1101 (1983).
9. C. Amsler, et al., *Phys. Lett.* **B667**, 1 (2008).
10. F. Ambrosino, et al., *Phys. Lett.* **B669**, 223–228 (2008), 0807.4909.
11. F. Ambrosino, et al., *Phys. Lett.* **B675**, 283–288 (2009), 0812.4830.
12. L. G. Landsberg, *Phys. Rept.* **128**, 301–376 (1985).
13. D.-N. Gao, *Mod. Phys. Lett.* **A17**, 1583–1588 (2002), hep-ph/0202002.