

Dark matter as a Ricci soliton.

Stuart Marongwe ¹ Stuart Kauffman ²

- 1 Physics Department, University of Botswana, 4775
Notwane Rd, Gaborone, Botswana
stuartmarongwe@gmail.com
- 2 University of Pennsylvania (emeritus)
stukauffman@gmail.com

Abstract: In this paper we consider clumped baryonic matter as a spherically symmetric barotropic fluid associated with a compact four-dimensional Einstein manifold with a four-radius that is determined by the fluid density. We further investigate the properties of these matter generated manifolds and show that because of their homogeneity and isotropy they exhibit FLRW metric properties and expand with cosmic time. At low densities such as those encountered in star clusters, galaxies and galactic clusters, these manifolds are large and show the same properties that are currently attributed to dark matter.

Submitted 28 March 2024

Essay written for the Gravity Research Foundation 2024 Awards for Essays on Gravitation

1. Introduction

In its vanilla form, the Mach Principle states that an inertial reference can be determined by measuring the motion of distance objects. Einstein was motivated by this principle in formulating the theory of General Relativity (GR). However, soon after its formulation it was found that GR does not follow the Mach's Principle. Although Einstein did not further pursue the idea of expressing Mach's Principle in mathematical form, there have been notable attempts [1-6] to formulate a theory of gravity based on this principle.

These alternate formulations of gravity have their merits and draw backs. Some formulations also attempt to explain the effects attributed to dark matter without the need of exotic non baryonic particles [7,8,9]. In this paper, we construct a theory of gravity based on two other flavors of the Mach Principle which are i) space-time cannot exist without matter ii) space-time is compact or a closed system. These flavors resonate well with Einstein's quote: "Time and space and gravitation have no separate existence from matter". Here, Einstein hints at matter generating a space-time manifold along with a gravitational field that appears when matter deforms Minkowski space. We pursue Einstein's hint using compact 4D Einstein manifolds that are generated by baryonic matter. This idea is motivated by the dualism that the Ricci tensor can be expressed as a compact energy-momentum tensor field $R_{\mu\nu} = \kappa \left(T_{\mu\nu} - \frac{1}{2} T g_{\mu\nu} \right) = k g_{\mu\nu}$ and that a compact Einstein manifold can be expressed as a stress-momentum tensor. $\kappa T_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = k g_{\mu\nu}$. Here κ is the Einstein constant and k is a constant proportional to the cosmological constant. We consider the compact energy-momentum tensor field $T_{\mu\nu} - \frac{1}{2} T g_{\mu\nu}$ as a clump of baryonic matter with a barotropic fluid like morphology. By being barotropic, it fulfills the condition of being homogeneous and isotropic within the volume of space it occupies.

This paper is arranged as follows: first we express Einstein's hint as a mathematical expression and then investigate the properties of the ensuing 4D manifolds generated

by the presence of matter and how they can offer an explanation to the enigmatic properties of dark matter. The paper is also an attempt to find a purely classical form of the quantum gravity version on the Nexus Paradigm of quantum gravity [10,11,12] postulated by the first author.

2. 4D compact Einstein manifolds from baryonic matter.

As suggested in the introduction, Einstein's field equations (EFE) need to be modified such that they include a matter generated 4-D compact Einstein manifold and thus expressing them as follows:

$$G_{\mu\nu} - kg_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \quad (1)$$

The first term on the left expresses the distortion of space-time by the presence of matter, the second term is a compact Einstein manifold associated with baryonic matter and the third term is a contribution by the cosmological constant. We consider a critical condition in which

$$G_{\mu\nu} - kg_{\mu\nu} = \kappa T_{\mu\nu} - \Lambda g_{\mu\nu} = 0 \quad (2)$$

Such a condition occurs when the density of baroynic matter is identical to that of DE. These conditions imply that space-time is covariantly flat and that the gravitional field is solely sourced from the Ricci soliton.

$$G_{\mu\nu} = kg_{\mu\nu} \quad (3)$$

We express the constant k as $k = \alpha^2 \Lambda = 3 \left(\frac{\alpha H_0}{c} \right)^2$ where αH_0 is the expansion rate of the Ricci soliton of reduced radius $r = \frac{c}{2\pi\alpha H_0}$. This implies a rotational velocity on the surface of the soliton of $v = H_0 r = \frac{c}{2\pi\alpha}$. Equation(3) must therefore satisfy a metric solution of the form

$$-c^2 \tau^2 = -c^2 t^2 + \alpha(t)^2 d\Sigma^2 \quad (4)$$

Where $d\Sigma^2 = \frac{dr^2}{1-kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2)$ in reduced-circumference polar coordinates.

The analytic solutions to equation (4) in which the energy momentum tensor is isotropic and homogeneous are the following equations:

$$\left(\frac{\dot{\alpha}}{\alpha} \right)^2 + \frac{\alpha^2 \Lambda c^2}{3\alpha^2} - \frac{\Lambda c^2}{3} = \frac{\kappa c^4}{3} \rho \rightarrow \left(\frac{\dot{\alpha}}{\alpha} \right)^2 = \frac{\kappa c^4}{3} \rho \quad (5)$$

$$2 \frac{\ddot{\alpha}}{\alpha} + \left(\frac{\dot{\alpha}}{\alpha} \right)^2 + \frac{\alpha^2 \Lambda c^2}{\alpha^2} - \Lambda c^2 = -\kappa c^2 p \rightarrow 2 \frac{\ddot{\alpha}}{\alpha} + \left(\frac{\dot{\alpha}}{\alpha} \right)^2 = -\kappa c^2 p \quad (6)$$

From the above equations the Ricci soliton expands exponentially when $p = -\frac{\rho c^2}{3}$ with a scale factor $\alpha(t) = \alpha_0 e^{H_0 t}$. This accelerated expansion occurs when the covariant flatness condition of equation (4) is satisfied. Under such conditions the centripetal acceleration on the surface of the soliton is given by the expression

$$\frac{v^2}{r} = \frac{(H_0 r)^2}{r} = H_0^2 r = \frac{H_0^2 c}{2\pi\alpha H_0} = \frac{H_0 c}{2\pi\alpha} \quad (7)$$

In the comoving reference frame $\alpha = 1$ which implies a scale invariant acceleration

$$a_0 = \frac{H_0 c}{2\pi} \quad (8)$$

Under such conditions the radius is computed from equation (7) as

$$r = \frac{2\pi v^2}{H_0 c} \quad (9)$$

Substituting r in the expression $\frac{GM(r)}{r} = v^2$ we obtain

$$v^4 = \frac{GM(r)H_0 c}{2\pi} \quad (10)$$

This is the baryonic Tully Fisher relation and $a_0 = \frac{H_0 c}{2\pi} \cong 1.1 \times 10^{-10} m/s^2$ is the empirically observed Milgrom's acceleration constant[13,14]. The theoretically computed value is in agreement with observations.

Since $r = r_0 e^{H_0 t}$ then the velocity evolves with the scale factor as follows:

$$v^4 = \frac{GM(r)H_0 c}{2\pi} e^{H_0 t} \quad (11)$$

We also obtain the following equations of galactic and galactic cluster evolution

$$r = \frac{1}{H_0} e^{(H_0 t)} (GM_B(r) \frac{H_0}{2\pi} c)^{\frac{1}{4}} = \frac{v_n}{H_0} \quad (12)$$

$$v = \frac{dr}{dt} = e^{(H_0 t)} (GM_B(r) \frac{H_0}{2\pi} c)^{\frac{1}{4}} = H_0 r \quad (13)$$

$$a = \frac{dv}{dt} = H_0 e^{(H_0 t)} (GM_B \frac{H_0}{2\pi} c)^{\frac{1}{4}} = H_0 v \quad (14)$$

The amplification of the radius of curvature with time explains the existence of ultra-diffuse galaxies and the spiral shapes of most galaxies. In the case of ultra-diffuse galaxies, most of their mass must reside at the core in the form of a dormant supermassive black hole for them to exhibit their properties. The increase in radial velocity with time explains why early-type galaxies composed of population II stars are fast rotators.

3. Summary

The nature of DM and DE are some of the most perplexing enigmas in modern physics. Here we have applied Mach's principle to explain the nature and properties of DM. We find that low matter density regions are associated with a large Ricci soliton which has the same properties attributed to DM. In the same manner, we can infer that the density of DE is the minimum baryonic mass density in the cosmos and is associated with a Ricci soliton of Hubble scale radius.

References

- 1 Brans, C., Dicke, R.H (1961), *Mach's Principle and a Relativistic Theory of Gravitation*, Physical Review **124** (Nov, 1961) 925-935
- 2 Hoyle,F.,Narlikar,J.V. (1964), *A New Theory of Gravitation*, Royal Society of London Proceedings Series A **282** (Nov.,1964) 191-207
- 3 Lichtenegger,H.,Mashhoon, B (2004), *Mach's Principle*, arxiv:physics/0407078v2
- 4 Sciama,D. (1953), *On the origin of inertia*, Mon. Not. Roy. Astron. Soc.' 1953, 113,34
- 5 Raine, D.J. (1975), *Mach's Principle in general relativity*, vol 171, p. 507-528
- 6 Das,S., *Mach's Principle and the origin of the quantum phenomenon*, arxiv:1206.0923 [physics.gen-ph]
- 7 Das,S. (2015),*Mach's Principle and a new theory of gravitation*, arive:1206.6755v2 [physics.gen-ph]
- 8 Kauffman, S., *Are dark matter, dark energy and inflation a construction of spacetime by matter?* OSF: <https://osf.io/5chn4>
- 9 Kauffman, S.,(2022) *On Quantum Gravity If Non-Locality Is Fundamental*, Entropy 2022 24(4),554; DOI: 10.3390/e24040554

- 10 Marongwe, S. (2021), *On the Electromagnetic Vacuum Origins of Dark Energy, Dark Matter, and Astrophysical Jets*, *Advances in High Energy Physics*, vol. 2021, Article ID 5588662, 8 pages
- 11 Marongwe, S.(2015) ,*The Schwarzschild solution to the nexus graviton field*, *Int. J. Geom. Methods Mod. Phys.* 12 1550042
- 12 Marongwe, S. (2024), *Energy momentum localization in quantum gravity*, *Phys. Scr.* 99 025306 27.
- 13 Milgrom, M. (1983), *A modification of the Newtonian dynamics — Implications for galaxies*, *Astrophys. J.* 270 371–389. 1983
- 14 Milgrom, M. (1986) , *Solutions for the modified Newtonian dynamics field equations*, *Astrophysical J.* 302 p617 DOI: 10.1086/164021