# Time & Fitness–Dependent Hamiltonian Biomechanics

Tijana T. Ivancevic

Society for Nonlinear Dynamics in Human Factors, Adelaide, Australia

and

CITECH Research IP Pty Ltd, Adelaide, Australia

e-mail: tijana.ivancevic@alumni.adelaide.edu.au

## Contents

| 1        | Introduction  | <b>2</b> |
|----------|---|----------|
| <b>2</b> | The Covariant Force Law                             | 2        |
| 3        | Autonomous Hamiltonian Biomechanics                 | 5        |
| <b>4</b> | Time–Dependent Hamiltonian Biomechanics             | 9        |
|          | 4.1 Biomechanical Bundle                            | 9        |
|          | 4.2 Biomechanical Jets                              | 10       |
|          | 4.3 Polysymplectic Dynamics                         | 12       |
|          | 4.4 Hamiltonian Connections                         | 13       |
|          | 4.5 Time–Dependent Dissipative Hamiltonian Dynamics | 13       |
|          | 4.6 Time & Fitness–Dependent Biomechanics           | 15       |
| 5        | Conclusion  | 16       |

#### Abstract

In this paper we propose the time & fitness-dependent Hamiltonian form of human biomechanics, in which *total mechanical + biochemical energy is not conserved*. Starting with the Covariant Force Law, we first develop autonomous Hamiltonian biomechanics. Then we extend it using a powerful geometrical machinery consisting of fibre bundles, jet manifolds, polysymplectic geometry and Hamiltonian connections. In this way we derive time-dependent dissipative Hamiltonian equations and the fitness evolution equation for the general time & fitnessdependent human biomechanical system.

Keywords: Human biomechanics, configuration bundle, Hamiltonian connections, jet manifolds, time & fitness-dependent dynamics

### 1 Introduction

Most of dynamics in both classical and quantum physics is based on *assumption of a total energy* conservation (see, e.g. [4]). Dynamics based on this assumption of time-independent energy, usually given by Hamiltonian (or Lagrangian) energy function, is called *autonomous*. This basic assumption is naturally inherited in human biomechanics, formally developed using Newton–Euler, Lagrangian or Hamiltonian formalisms (see [1, 2, 3, 5, 6, 8, 9, 10]).

And this works fine for most individual movement simulations and predictions, in which the total human energy dissipations are insignificant. However, if we analyze a 100 m-dash sprinting motion, which is in case of top athletes finished under 10 s, we can recognize a significant slow-down after about 70 m in *all* athletes – despite of their strong intention to finish and win the race, which is an obvious sign of the total energy dissipation. This can be seen, for example, in a current record-braking speeddistance curve of Usain Bolt, the world-record holder with 9.69 s, or in a former record-braking speeddistance curve of Carl Lewis, the former world-record holder (and 9 time Olympic gold medalist) with 9.86 s (see Figure 3.7 in [10]). In other words, the *total mechanical + biochemical energy* of a sprinter *cannot be conserved* even for 10 s. So, if we want to develop a realistic model of intensive human motion that is longer than 7–8 s (not to speak for instance of a 4 hour tennis match), we necessarily need to use the more advanced formalism of time-dependent mechanics.

In this paper, we will first develop the autonomous Hamiltonian biomechanics as a Hamiltonian representation of the covariant force law [see (1) in the next section] and the corresponding covariant force functor. After that we will extend the autonomous Hamiltonian biomechanics into the time-dependent one, in which *total mechanical + biochemical energy is not conserved*. for this, we will use the modern geometric formalism of jet manifolds and bundles.

## 2 The Covariant Force Law

Autonomous Hamiltonian biomechanics (as well as autonomous Lagrangian biomechanics), based on the postulate of conservation of the total mechanical energy, can be derived from the *covariant* force law [1, 2, 3, 4], which in 'plain English' states:

Force  $1-\text{form} = \text{Mass distribution} \times \text{Acceleration vector-field}$ ,

and formally reads (using Einstein's summation convention over repeated indices):

$$F_i = m_{ij} a^j. (1)$$

Here, the force 1-form  $F_i = F_i(t, q, p) = F'_i(t, q, \dot{q})$ , (i = 1, ..., n) denotes any type of torques and forces acting on a human skeleton, including excitation and contraction dynamics of muscularactuators [13, 12, 11] and rotational dynamics of hybrid robot actuators, as well as (nonlinear) dissipative joint torques and forces and external stochastic perturbation torques and forces [5].  $m_{ij}$  is the material (mass-inertia) metric tensor, which gives the total mass distribution of the human body, by including all segmental masses and their individual inertia tensors.  $a^j$  is the total acceleration vector-field, including all segmental vector-fields, defined as the absolute (Bianchi) derivative  $\dot{v}^i$  of all the segmental angular and linear velocities  $v^i = \dot{x}^i$ , (i = 1, ..., n), where n is the total number of active degrees of freedom (DOF) with local coordinates  $(x^i)$ . More formally, this *central Law of biomechanics* represents the *covariant force functor*  $\mathcal{F}_*$  defined by the commutative diagram:



Here,  $M \equiv M^n = \{x^i, (i = 1, ..., n)\}$  is the biomechanical configuration *n*-manifold, that is the set of all active DOF of the biomechanical system under consideration (in general, human skeleton), with local coordinates  $(x^i)$ .

The right-hand branch of the fundamental covariant force functor  $\mathcal{F}_* : TT^*M \to TTM$  depicted in (2) is Lagrangian dynamics with its Riemannian geometry. To each *n*-dimensional (*n*D) smooth manifold M there is associated its 2*n*D velocity phase-space manifold, denoted by TM and called the tangent bundle of M. The original configuration manifold M is called the base of TM. There is an onto map  $\pi : TM \to M$ , called the projection. Above each point  $x \in M$  there is a tangent space  $T_xM = \pi^{-1}(x)$  to M at x, which is called a fibre. The fibre  $T_xM \subset TM$  is the subset of TM, such that the total tangent bundle,  $TM = \bigsqcup_{m \in M} T_xM$ , is a disjoint union of

tangent spaces  $T_x M$  to M for all points  $x \in M$ . From dynamical perspective, the most important quantity in the tangent bundle concept is the smooth map  $v: M \to TM$ , which is an inverse to the projection  $\pi$ , i.e.,  $\pi \circ v = \mathrm{Id}_M$ ,  $\pi(v(x)) = x$ . It is called the *velocity vector-field*  $v^i = \dot{x}^i$ .<sup>1</sup> Its graph (x, v(x)) represents the cross-section of the tangent bundle TM. Velocity vector-fields are cross-sections of the tangent bundle. Biomechanical Lagrangian (that is, kinetic minus potential energy) is a natural energy function on the tangent bundle TM. The tangent bundle is itself a smooth manifold. It has its own tangent bundle, TTM. Cross-sections of the second tangent bundle TTM are the acceleration vector-fields.

The left-hand branch of the fundamental covariant force functor  $\mathcal{F}_*: TT^*M \to TTM$  depicted in (2) is Hamiltonian dynamics with its symplectic geometry. It takes place in the *cotangent bundle*  $T^*M_{rob}$ , defined as follows. A *dual* notion to the tangent space  $T_xM$  to a smooth manifold M at a point  $x = (x^i)$  with local is its cotangent space  $T_x^*M$  at the same point x. Similarly to the tangent bundle TM, for any smooth nD manifold M, there is associated its 2nD momentum phase-space manifold, denoted by  $T^*M$  and called the *cotangent bundle*.  $T^*M$  is the disjoint union of all its cotangent spaces  $T_x^*M$  at all points  $x \in M$ , i.e.,  $T^*M = \bigsqcup_{x \in M} T_x^*M$ . Therefore, the cotangent bundle of an n-manifold M is the vector bundle  $T^*M = (TM)^*$ , the (real) dual of the tangent bundle TM. Momentum 1-forms (or, covector-fields)  $p_i$  are cross-sections of the cotangent bundle. Biomechanical Hamiltonian (that is, kinetic plus potential energy) is a natural energy function on

<sup>&</sup>lt;sup>1</sup>This explains the dynamical term velocity phase-space, given to the tangent bundle TM of the manifold M.

the cotangent bundle. The cotangent bundle  $T^*M$  is itself a smooth manifold. It has its own tangent bundle,  $TT^*M$ . Cross-sections of the mixed-second bundle  $TT^*M$  are the force 1-forms  $F_i = \dot{p}_i$ .

There is a unique smooth map from the right-hand branch to the left-hand branch of the diagram (2):

$$TM \ni (x^i, v^i) \mapsto (x^i, p^i) \in T^*M.$$

It is called the *Legendre transformation*, or *fiber derivative* (for details see, e.g. [3, 4]).

The fundamental covariant force functor  $\mathcal{F}_*: TT^*M \to TTM$  states that the force 1-form  $F_i = \dot{p}_i$ , defined on the mixed tangent-cotangent bundle  $TT^*M$ , causes the acceleration vector-field  $a^i = \dot{v}^i$ , defined on the second tangent bundle TTM of the configuration manifold M. The corresponding *contravariant acceleration functor* is defined as its inverse map,  $\mathcal{F}^*: TTM \to TT^*M$ .

Representation of human motion is rigorously defined in terms of Euclidean SE(3)-groups<sup>2</sup> of full rigid-body motion in all main human joints [6]. The configuration manifold M for human musculo-skeletal dynamics is defined as a Cartesian product of all included constrained SE(3)groups,  $M = \prod_j SE(3)^j$  where j labels the active joints. The configuration manifold M is coordinated by local joint coordinates  $x^i(t)$ , i = 1, ..., n = total number of active DOF. The corresponding joint velocities  $\dot{x}^i(t)$  live in the velocity phase space TM, which is the tangent bundle of the configuration manifold M.

The velocity phase-space TM has the Riemannian geometry with the *local metric form*:

$$\langle g \rangle \equiv ds^2 = g_{ij} dx^i dx^j, \tag{3}$$

where  $g_{ij}(x)$  is the material metric tensor defined by the biomechanical system's mass-inertia matrix and  $dx^i$  are differentials of the local joint coordinates  $x^i$  on M. Besides giving the local distances between the points on the manifold M, the Riemannian metric form  $\langle g \rangle$  defines the system's kinetic energy:

$$T = \frac{1}{2}g_{ij}\dot{x}^i\dot{x}^j,$$

giving the Lagrangian equations of the conservative skeleton motion with kinetic-minus-potential energy Lagrangian L = T - V, with the corresponding geodesic form

$$\frac{d}{dt}L_{\dot{x}^i} - L_{x^i} = 0 \qquad \text{or} \qquad \ddot{x}^i + \Gamma^i_{jk}\dot{x}^j\dot{x}^k = 0, \tag{4}$$

<sup>2</sup>Briefly, the Euclidean SE(3)–group is defined as a semidirect (noncommutative) product (denoted by  $\triangleright$ ) of 3D rotations and 3D translations:  $SE(3) := SO(3) \triangleright \mathbb{R}^3$ . Its most important subgroups are the following:

| Subgroup   | Definition  |
|--|---|
| SO(3), group of rotations  | Set of all proper orthogonal                                  |
| in 3D (a spherical joint)  | $3 \times 3 - \text{rotational matrices}$                     |
|  | Set of all $3 \times 3$ – matrices:                           |
| SE(2), special Euclidean group   | $\begin{bmatrix} \cos\theta & \sin\theta & r_x \end{bmatrix}$ |
| in 2D (all planar motions)   | $-\sin\theta \cos\theta r_y$                                  |
|  |   |
| SO(2), group of rotations in 2D  | Set of all proper orthogonal                                  |
| subgroup of $SE(2)$ -group   | $2 \times 2 - $ rotational matrices                           |
| (a revolute joint)   | included in $SE(2) - \text{group}$                            |
| $\mathbb{R}^3$ , group of translations in 3D (all spatial displacements) | Euclidean 3D vector space                                     |

where subscripts denote partial derivatives, while  $\Gamma_{jk}^i$  are the Christoffel symbols of the affine Levi-Civita connection of the biomechanical manifold M.

The corresponding momentum phase-space  $P = T^*M$  provides a natural symplectic structure that can be defined as follows. As the biomechanical configuration space M is a smooth n-manifold, we can pick local coordinates  $\{dx^1, ..., dx^n\} \in M$ . Then  $\{dx^1, ..., dx^n\}$  defines a basis of the cotangent space  $T_x^*M$ , and by writing  $\theta \in T_x^*M$  as  $\theta = p_i dx^i$ , we get local coordinates  $\{x^1, ..., x^n, p_1, ..., p_n\}$  on  $T^*M$ . We can now define the canonical symplectic form  $\omega$  on  $P = T^*M$ as:

$$\omega = dp_i \wedge dx^i.$$

where ' $\wedge$ ' denotes the wedge or exterior product of exterior differential forms.<sup>3</sup> This 2-form  $\omega$  is obviously independent of the choice of coordinates  $\{x^1, ..., x^n\}$  and independent of the base point  $\{x^1, ..., x^n, p_1, ..., p_n\} \in T_x^*M$ . Therefore, it is locally constant, and so  $d\omega = 0.4$ 

If  $(P, \omega)$  is a 2nD symplectic manifold then about each point  $x \in P$  there are local coordinates  $\{x^1, ..., x^n, p_1, ..., p_n\}$  such that  $\omega = dp_i \wedge dx^i$ . These coordinates are called canonical or symplectic. By the Darboux theorem,  $\omega$  is constant in this local chart, i.e.,  $d\omega = 0$ .

## **3** Autonomous Hamiltonian Biomechanics

We develop autonomous Hamiltonian biomechanics on the configuration biomechanical manifold M in three steps, following the standard symplectic geometry prescription (see [1, 3, 4, 7]):

**Step A** Find a symplectic momentum phase-space  $(P, \omega)$ .

<sup>3</sup>Recall that an *exterior differential form*  $\alpha$  of order p (or, a p-form  $\alpha$ ) on a base manifold X is a section of the exterior product bundle  $\bigwedge^{p} T^*X \to X$ . It has the following expression in local coordinates on X

$$\alpha = \alpha_{\lambda_1 \dots \lambda_p} dx^{\lambda_1} \wedge \dots \wedge dx^{\lambda_p} \qquad (\text{such that } |\alpha| = p),$$

where summation is performed over all ordered collections  $(\lambda_1, ..., \lambda_p)$ .  $\Omega^p(X)$  is the vector space of p-forms on a biomechanical manifold X. In particular, the 1-forms are called the *Pfaffian forms*.

<sup>&</sup>lt;sup>4</sup>The canonical 1-form  $\theta$  on  $T^*M$  is the unique 1-form with the property that, for any 1-form  $\beta$  which is a section of  $T^*M$  we have  $\beta^*\theta = \theta$ .

Let  $f: M \to M$  be a diffeomorphism. Then  $T^*f$  preserves the canonical 1-form  $\theta$  on  $T^*M$ , i.e.,  $(T^*f)^*\theta = \theta$ . Thus  $T^*f$  is symplectic diffeomorphism.

Recall that a symplectic structure on a smooth manifold M is a nondegenerate closed<sup>5</sup> 2-form  $\omega$  on M, i.e., for each  $x \in M$ ,  $\omega(x)$  is nondegenerate, and  $d\omega = 0$ .

Let  $T_x^*M$  be a cotangent space to M at m. The cotangent bundle  $T^*M$  represents a union  $\cup_{m \in M} T_x^*M$ , together with the standard topology on  $T^*M$  and a natural smooth manifold structure, the dimension of which is twice the dimension of M. A 1-form  $\theta$  on M represents a section  $\theta: M \to T^*M$  of the cotangent bundle  $T^*M$ .

 $P = T^*M$  is our momentum phase-space. On P there is a nondegenerate symplectic 2-form  $\omega$  is defined in local joint coordinates  $x^i, p_i \in U$ , U open in P, as  $\omega = dx^i \wedge dp_i$ . In that case the coordinates  $x^i, p_i \in U$  are called canonical. In a usual procedure the canonical 1-form  $\theta$  is first defined as  $\theta = p_i dx^i$ , and then the canonical 2-form  $\omega$  is defined as  $\omega = -d\theta$ .

A symplectic phase-space manifold is a pair  $(P, \omega)$ .

#### **Step B** Find a Hamiltonian vector-field $X_H$ on $(P, \omega)$ .

Let  $(P, \omega)$  be a symplectic manifold. A vector-field  $X : P \to TP$  is called Hamiltonian if there is a smooth function  $F : P \to \mathbb{R}$  such that  $i_X \omega = dF$   $(i_X \omega$  denotes the *interior product* or *contraction* of the vector-field X and the 2-form  $\omega$ ). X is *locally Hamiltonian* if  $i_X \omega$  is closed.

Let the smooth real-valued Hamiltonian function  $H: P \to \mathbb{R}$ , representing the total biomechanical energy H(x,p) = T(p) + V(x) (T and V denote kinetic and potential energy of the system, respectively), be given in local canonical coordinates  $x^i, p_i \in U, U$  open in P. The Hamiltonian vector-field  $X_H$ , condition by  $i_{X_H}\omega = dH$ , is actually defined via symplectic matrix J, in a local chart U, as

$$X_H = J\nabla H = (\partial_{p_i} H, -\partial_{x^i} H), \qquad J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \tag{5}$$

where I denotes the  $n \times n$  identity matrix and  $\nabla$  is the gradient operator.

#### **Step C** Find a Hamiltonian phase-flow $\phi_t$ of $X_H$ .

Let  $(P, \omega)$  be a symplectic phase–space manifold and  $X_H = J\nabla H$  a Hamiltonian vector-field corresponding to a smooth real-valued Hamiltonian function  $H: P \to \mathbb{R}$ , on it. If a unique one– parameter group of diffeomorphisms  $\phi_t: P \to P$  exists so that  $\frac{d}{dt}|_{t=0} \phi_t x = J\nabla H(x)$ , it is called the Hamiltonian phase–flow.

$$d\beta = 0.$$

From this condition one can see that the closed form (the *kernel* of the exterior derivative operator d) is conserved quantity. Therefore, closed p-forms possess certain invariant properties, physically corresponding to the *conservation laws*.

Also, a p-form  $\beta$  that is an exterior derivative of some (p-1)-form  $\alpha$ ,

$$\beta = d\alpha,$$

is called *exact* (the *image* of the exterior derivative operator d). By *Poincaré lemma*, exact forms prove to be closed automatically,

$$d\beta = d(d\alpha) = 0.$$

Since  $d^2 = 0$ , every exact form is closed. The converse is only partially true, by Poincaré lemma: every closed form is locally exact.

Technically, this means that given a closed p-form  $\alpha \in \Omega^p(U)$ , defined on an open set U of a smooth manifold M any point  $m \in U$  has a neighborhood on which there exists a (p-1)-form  $\beta \in \Omega^{p-1}(U)$  such that  $d\beta = \alpha|_U$ . In particular, there is a Poincaré lemma for contractible manifolds: Any closed form on a smoothly contractible manifold is exact.

<sup>&</sup>lt;sup>5</sup>A *p*-form  $\beta$  on a smooth manifold *M* is called *closed* if its exterior derivative  $d = \partial_i dx^i$  is equal to zero,

A smooth curve  $t \mapsto (x^i(t), p_i(t))$  on  $(P, \omega)$  represents an *integral curve* of the Hamiltonian vector-field  $X_H = J\nabla H$ , if in the local canonical coordinates  $x^i, p_i \in U, U$  open in P, Hamiltonian canonical equations hold:

$$\dot{q}^i = \partial_{p_i} H, \qquad \dot{p}_i = -\partial_{q^i} H.$$
 (6)

An integral curve is said to be *maximal* if it is not a restriction of an integral curve defined on a larger interval of  $\mathbb{R}$ . It follows from the standard theorem on the *existence* and *uniqueness* of the solution of a system of ODEs with smooth r.h.s, that if the manifold  $(P, \omega)$  is Hausdorff, then for any point  $x = (x^i, p_i) \in U$ , U open in P, there exists a maximal integral curve of  $X_H = J\nabla H$ , passing for t = 0, through point x. In case  $X_H$  is complete, i.e.,  $X_H$  is  $C^p$  and  $(P, \omega)$  is compact, the maximal integral curve of  $X_H$  is the Hamiltonian phase-flow  $\phi_t : U \to U$ .

The phase–flow  $\phi_t$  is symplectic if  $\omega$  is constant along  $\phi_t,$  i.e.,  $\phi_t^*\omega=\omega$ 

 $(\phi_t^* \omega \text{ denotes the } pull-back^6 \text{ of } \omega \text{ by } \phi_t),$ 

iff  $\mathfrak{L}_{X_H}\omega = 0$ 

 $(\mathfrak{L}_{X_H}\omega \text{ denotes the } Lie \ derivative^7 \text{ of } \omega \text{ upon } X_H).$ 

Symplectic phase-flow  $\phi_t$  consists of canonical transformations on  $(P, \omega)$ , i.e., diffeomorphisms in canonical coordinates  $x^i, p_i \in U, U$  open on all  $(P, \omega)$  which leave  $\omega$  invariant. In this case the *Liouville theorem* is valid:  $\phi_t$  preserves the phase volume on  $(P, \omega)$ . Also, the system's total energy H is conserved along  $\phi_t$ , i.e.,  $H \circ \phi_t = \phi_t$ .

Recall that the Riemannian metrics  $g = \langle , \rangle$  on the configuration manifold M is a positive– definite quadratic form  $g: TM \to \mathbb{R}$ , in local coordinates  $x^i \in U$ , U open in M, given by (3) above. Given the metrics  $g_{ij}$ , the system's Hamiltonian function represents a momentum p-dependent quadratic form  $H: T^*M \to \mathbb{R}$  – the system's kinetic energy  $H(p) = T(p) = \frac{1}{2} \langle p, p \rangle$ , in local canonical coordinates  $x^i, p_i \in U_p, U_p$  open in  $T^*M$ , given by

$$H(p) = \frac{1}{2}g^{ij}(x,m) \, p_i p_j, \tag{7}$$

where  $g^{ij}(x,m) = g_{ij}^{-1}(x,m)$  denotes the *inverse* (contravariant) material *metric tensor* 

$$g^{ij}(x,m) = \sum_{\chi=1}^{n} m_{\chi} \delta_{rs} \frac{\partial x^{i}}{\partial x^{r}} \frac{\partial x^{j}}{\partial x^{s}}$$

$$f^*(\alpha \wedge \beta) = f^*\alpha \wedge f^*\beta, \qquad df^*\alpha = f^*(d\alpha),$$

for any two forms  $\alpha, \beta \in \mathbf{\Omega}^p(X)$ .

<sup>7</sup>The Lie derivative  $\mathfrak{L}_u \alpha$  of p-form  $\alpha$  along a vector-field u is defined by Cartan's 'magic' formula (see [3, 4]):

$$\mathfrak{L}_u \alpha = u | d\alpha + d(u | \alpha).$$

It satisfies the Leibnitz relation

$$\mathfrak{L}_u(\alpha \wedge \beta) = \mathfrak{L}_u \alpha \wedge \beta + \alpha \wedge \mathfrak{L}_u \beta.$$

Here, the contraction  $\rfloor$  of a vector-field  $u = u^{\mu} \partial_{\mu}$  and a p-form  $\alpha = \alpha_{\lambda_1...\lambda_p} dx^{\lambda_1} \wedge \cdots \wedge dx^{\lambda_p}$  on a biomechanical manifold X is given in local coordinates on X by

$$u \rfloor \alpha = u^{\mu} \alpha_{\mu \lambda_1 \dots \lambda_{p-1}} dx^{\lambda_1} \wedge \dots \wedge dx^{\lambda_{p-1}}.$$

It satisfies the following relation

$$u \rfloor (\alpha \land \beta) = u \rfloor \alpha \land \beta + (-1)^{|\alpha|} \alpha \land u \rfloor \beta$$

<sup>&</sup>lt;sup>6</sup>Given a map  $f: X \to X'$  between the two manifolds, the *pullback* on X of a form  $\alpha$  on X' by f is denoted by  $f^*\alpha$ . The pullback satisfies the relations

 $T^*M$  is an orientable manifold, admitting the standard volume form

$$\Omega_{\omega_H} = \frac{(-1)^{\frac{N(N+1)}{2}}}{N!} \omega_H^N$$

For Hamiltonian vector-field,  $X_H$  on M, there is a base integral curve  $\gamma_0(t) = (x^i(t), p_i(t))$  iff  $\gamma_0(t)$  is a geodesic, given by the one-form force equation

$$\dot{\bar{p}}_i \equiv \dot{p}_i + \Gamma^i_{jk} \, g^{jl} g^{km} \, p_l p_m = 0, \qquad \text{with} \qquad \dot{x}^k = g^{ki} p_i. \tag{8}$$

The l.h.s  $\dot{p}_i$  of the covariant momentum equation (8) represents the intrinsic or Bianchi covariant derivative of the momentum with respect to time t. Basic relation  $\dot{p}_i = 0$  defines the *parallel* transport on  $T^N$ , the simplest form of human-motion dynamics. In that case Hamiltonian vector-field  $X_H$  is called the *geodesic spray* and its phase-flow is called the *geodesic flow*.

For Earthly dynamics in the gravitational *potential* field  $V: M \to \mathbb{R}$ , the Hamiltonian  $H: T^*M \to \mathbb{R}$  (7) extends into potential form

$$H(p,x) = \frac{1}{2}g^{ij}p_ip_j + V(x),$$

with Hamiltonian vector-field  $X_H = J\nabla H$  still defined by canonical equations (6).

A general form of a *driven*, non-conservative Hamiltonian equations reads:

$$\dot{x}^{i} = \partial_{p_{i}}H, \qquad \dot{p}_{i} = F_{i} - \partial_{x^{i}}H, \tag{9}$$

where  $F_i = F_i(t, x, p)$  represent any kind of joint-driving *covariant torques*, including active neuromuscular-like controls, as functions of time, angles and momenta, as well as passive dissipative and elastic joint torques. In the covariant momentum formulation (8), the non-conservative Hamiltonian equations (9) become

$$\dot{\bar{p}}_i \equiv \dot{p}_i + \Gamma^i_{jk} g^{jl} g^{km} p_l p_m = F_i, \quad \text{with} \quad \dot{x}^k = g^{ki} p_i.$$

The general form of autonomous Hamiltonian biomechanics is given by dissipative, driven Hamiltonian equations on  $T^*M$ :

$$\dot{x}^{i} = \frac{\partial H}{\partial p_{i}} + \frac{\partial R}{\partial p_{i}},\tag{10}$$

$$\dot{p}_i = F_i - \frac{\partial H}{\partial x^i} + \frac{\partial R}{\partial x^i},$$
(11)

$$x^{i}(0) = x_{0}^{i}, \qquad p_{i}(0) = p_{i}^{0},$$
(12)

including contravariant equation (10) – the velocity vector-field, and covariant equation (11) – the force 1-form (field), together with initial joint angles and momenta (12). Here R = R(x, p) denotes the Raileigh nonlinear (biquadratic) dissipation function, and  $F_i = F_i(t, x, p)$  are covariant driving torques of equivalent muscular actuators, resembling muscular excitation and contraction dynamics in rotational form. The velocity vector-field (10) and the force 1-form (11) together define the generalized Hamiltonian vector-field  $X_H$ ; the Hamiltonian energy function H = H(x, p) is its generating function.

As a Lie group, the biomechanical configuration manifold  $M = \prod_j SE(3)^j$  is Hausdorff.<sup>8</sup> Therefore, for  $x = (x^i, p_i) \in U_p$ , where  $U_p$  is an open coordinate chart in  $T^*M$ , there exists a unique one-parameter group of diffeomorphisms  $\phi_t : T^*M \to T^*M$ , that is the *autonomous Hamiltonian* phase-flow:

$$\phi_t \quad : \quad T^*M \to T^*M : (p(0), x(0)) \mapsto (p(t), x(t)),$$

$$(\phi_t \quad \circ \quad \phi_s = \phi_{t+s}, \quad \phi_0 = \text{identity}),$$

$$(13)$$

given by (10-12) such that

$$\frac{d}{dt}\Big|_{t=0} \phi_t x = J\nabla H(x).$$

### 4 Time–Dependent Hamiltonian Biomechanics

In this section we develop time-dependent Hamiltonian biomechanics. For this, we first need to extend our autonomous Hamiltonian machinery, using the general concepts of bundles, jets and connections.

### 4.1 Biomechanical Bundle

While standard autonomous Lagrangian biomechanics is developed on the configuration manifold X, the *time-dependent biomechanics* necessarily includes also the real time axis  $\mathbb{R}$ , so we have an *extended configuration manifold*  $\mathbb{R} \times X$ . Slightly more generally, the fundamental geometrical structure is the so-called *configuration bundle*  $\pi : X \to \mathbb{R}$ . Time-dependent biomechanics is thus formally developed either on the *extended configuration manifold*  $\mathbb{R} \times X$ , or on the configuration bundle  $\pi : X \to \mathbb{R}$ , using the concept of *jets*, which are based on the idea of *higher-order tangency*, or higher-order contact, at some designated point (i.e., certain anatomical joint) on a biomechanical configuration manifold X.

In general, tangent and cotangent bundles, TM and  $T^*M$ , of a smooth manifold M, are special cases of a more general geometrical object called *fibre bundle*, denoted  $\pi: Y \to X$ , where the word *fiber* V of a map  $\pi: Y \to X$  is the *preimage*  $\pi^{-1}(x)$  of an element  $x \in X$ . It is a space which *locally* looks like a product of two spaces (similarly as a manifold locally looks like Euclidean space), but may possess a different *global* structure. To get a visual intuition behind this fundamental geometrical concept, we can say that a fibre bundle Y is a *homeomorphic generalization* of a *product space*  $X \times V$  (see Figure 1), where X and V are called the *base* and the *fibre*, respectively.  $\pi: Y \to X$  is called the *projection*,  $Y_x = \pi^{-1}(x)$  denotes a fibre over a point x of the base X, while the map  $f = \pi^{-1}: X \to Y$  defines the *cross-section*, producing the *graph* (x, f(x)) in the bundle Y (e.g., in case of a tangent bundle,  $f = \dot{x}$  represents a velocity vector-field).

More generally, a biomechanical configuration bundle,  $\pi : Y \to X$ , is a locally trivial fibred (or, projection) manifold over the base X. It is endowed with an atlas of fibred bundle coordinates  $(x^{\lambda}, y^{i})$ , where  $(x^{\lambda})$  are coordinates of X.

A linear connection  $\overline{\Gamma}$  on a biomechanical bundle  $Y \to X$  is given in local coordinates on Y by [14]

$$\bar{\Gamma} = dx^{\lambda} \otimes [\partial_{\lambda} - \Gamma^{i}{}_{j\lambda}(x)y^{j}\partial_{i}].$$
(14)

<sup>&</sup>lt;sup>8</sup>That is, for every pair of points  $x_1, x_2 \in M$ , there are disjoint open subsets (charts)  $U_1, U_2 \subset M$  such that  $x_1 \in U_1$  and  $x_2 \in U_2$ .



Figure 1: A sketch of a locally trivial fibre bundle  $Y \approx X \times V$  as a generalization of a product space  $X \times V$ ; left – main components; right – a few details (see text for explanation).

An affine connection  $\Gamma$  on a biomechanical bundle  $Y \to X$  is given in local coordinates on Y by

$$\Gamma = dx^{\lambda} \otimes [\partial_{\lambda} + (-\Gamma^{i}{}_{j\lambda}(x)y^{j} + \Gamma^{i}{}_{\lambda}(x))\partial_{i}].$$

Clearly, a linear connection  $\overline{\Gamma}$  is a special case of an affine connection  $\Gamma$ .

Every connection  $\Gamma$  on a biomechanical bundle  $Y \to X$  defines a system of first-order differential equations on Y, given by the local coordinate relations

$$y_{\lambda}^{i} = \Gamma^{i}(y). \tag{15}$$

Integral sections for  $\Gamma$  are local solutions of (15).

#### 4.2 Biomechanical Jets

A pair of smooth manifold maps,  $f_1, f_2 : M \to N$  (see Figure 2), are said to be k-tangent (or tangent of order k, or have a kth order contact) at a point x on a domain manifold M, denoted by  $f_1 \sim f_2$ , iff

$$f_1(x) = f_2(x) \quad \text{called} \quad 0-\text{tangent},$$
  

$$\partial_x f_1(x) = \partial_x f_2(x), \quad \text{called} \quad 1-\text{tangent},$$
  

$$\partial_{xx} f_1(x) = \partial_{xx} f_2(x), \quad \text{called} \quad 2-\text{tangent},$$
  
... etc. to the order k

In this way defined k-tangency is an equivalence relation.

A k-jet (or, a jet of order k), denoted by  $j_x^k f$ , of a smooth map  $f: M \to N$  at a point  $x \in M$  (see Figure 2), is defined as an equivalence class of k-tangent maps at x,

$$j_x^k f: M \to N = \{f': f' \text{ is } k - \text{tangent to } f \text{ at } x\}.$$

For example, consider a simple function  $f: X \to Y, x \mapsto y = f(x)$ , mapping the X-axis into the Y-axis in  $\mathbb{R}^2$ . At a chosen point  $x \in X$  we have: a 0-jet is a graph: (x, f(x));

a 1-jet is a triple: (x, f(x), f'(x));



Figure 2: An intuitive geometrical picture behind the k-jet concept, based on the idea of a higher-order tangency (or, higher-order contact).

a 2-jet is a quadruple: (x, f(x), f'(x), f''(x)), and so on, up to the order k (where  $f'(x) = \frac{df(x)}{dx}$ , etc). The set of all k-jets from  $j_x^k f : X \to Y$  is called the k-jet manifold  $J^k(X, Y)$ .

Formally, given a biomechanical bundle  $Y \to X$ , its first-order jet manifold  $J^1Y$  comprises the set of equivalence classes  $j_x^1 s, x \in X$ , of sections  $s : X \to Y$  so that sections s and s' belong to the same class iff

$$Ts \mid_{T_x X} = Ts' \mid_{T_x X}$$
.

Intuitively, sections  $s, s' \in j_x^1 s$  are identified by their values  $s^i(x) = {s'}^i(x)$  and the values of their partial derivatives  $\partial_\mu s^i(x) = \partial_\mu {s'}^i(x)$  at the point x of X. There are the natural fibrations [14]

$$\pi_1: J^1Y \ni j_x^1 s \mapsto x \in X, \qquad \pi_{01}: J^1Y \ni j_x^1 s \mapsto s(x) \in Y.$$

Given bundle coordinates  $(x^{\lambda}, y^{i})$  of Y, the associated jet manifold  $J^{1}Y$  is endowed with the adapted coordinates

$$(x^{\lambda}, y^{i}, y^{i}_{\lambda}), \qquad (y^{i}, y^{i}_{\lambda})(j^{1}_{x}s) = (s^{i}(x), \partial_{\lambda}s^{i}(x)), \qquad {y'}^{i}_{\lambda} = \frac{\partial x^{\mu}}{\partial {x'}^{\lambda}}(\partial_{\mu} + y^{j}_{\mu}\partial_{j})y'^{i}.$$

In particular, given the biomechanical configuration bundle  $M \to \mathbb{R}$  over the time axis  $\mathbb{R}$ , the 1-jet space  $J^1(\mathbb{R}, M)$  is the set of equivalence classes  $j_t^1 s$  of sections  $s^i : \mathbb{R} \to M$  of the configuration bundle  $M \to \mathbb{R}$ , which are identified by their values  $s^i(t)$ , as well as by the values of their partial derivatives  $\partial_t s^i = \partial_t s^i(t)$  at time points  $t \in \mathbb{R}$ . The 1-jet manifold  $J^1(\mathbb{R}, M)$  is coordinated by  $(t, x^i, \dot{x}^i)$ , that is by (time, coordinates and velocities) at every active human joint, so the 1-jets are local joint coordinate maps

$$j_t^1 s : \mathbb{R} \to M, \qquad t \mapsto (t, x^i, \dot{x}^i).$$

The repeated jet manifold  $J^1 J^1 Y$  is defined to be the jet manifold of the bundle  $J^1 Y \to X$ . It is endowed with the adapted coordinates  $(x^{\lambda}, y^i, y^i_{\lambda}, y^i_{(\mu)}, y^i_{\lambda\mu})$ .

The second-order jet manifold  $J^2Y$  of a bundle  $Y \to X$  is the subbundle of  $\widehat{J}^2Y \to J^1Y$ defined by the coordinate conditions  $y^i_{\lambda\mu} = y^i_{\mu\lambda}$ . It has the local coordinates  $(x^{\lambda}, y^i, y^i_{\lambda}, y^i_{\lambda\leq\mu})$  together with the transition functions [14]

$$y'^{i}_{\lambda\mu} = \frac{\partial x^{\alpha}}{\partial x'^{\mu}} (\partial_{\alpha} + y^{j}_{\alpha} \partial_{j} + y^{j}_{\nu\alpha} \partial^{\nu}_{j}) y'^{i}_{\lambda}.$$

The second–order jet manifold  $J^2Y$  of Y comprises the equivalence classes  $j_x^2s$  of sections s of  $Y \to X$  such that

$$y_{\lambda}^{i}(j_{x}^{2}s) = \partial_{\lambda}s^{i}(x), \qquad y_{\lambda\mu}^{i}(j_{x}^{2}s) = \partial_{\mu}\partial_{\lambda}s^{i}(x).$$

In other words, two sections  $s, s' \in j_x^2 s$  are identified by their values and the values of their first and second-order derivatives at the point  $x \in X$ .

In particular, given the biomechanical configuration bundle  $M \to \mathbb{R}$  over the time axis  $\mathbb{R}$ , the 2-jet space  $J^2(\mathbb{R}, M)$  is the set of equivalence classes  $j_t^2 s$  of sections  $s^i : \mathbb{R} \to M$  of the configuration bundle  $\pi : M \to \mathbb{R}$ , which are identified by their values  $s^i(t)$ , as well as the values of their first and second partial derivatives,  $\partial_t s^i = \partial_t s^i(t)$  and  $\partial_{tt} s^i = \partial_{tt} s^i(t)$ , respectively, at time points  $t \in \mathbb{R}$ . The 2-jet manifold  $J^2(\mathbb{R}, M)$  is coordinated by  $(t, x^i, \dot{x}^i, \ddot{x}^i)$ , that is by (time, coordinates, velocities and accelerations) at every active human joint, so the 2-jets are local joint coordinate maps<sup>9</sup>

$$j_t^2 s : \mathbb{R} \to M, \qquad t \mapsto (t, x^i, \dot{x}^i, \ddot{x}^i).$$

### 4.3 Polysymplectic Dynamics

Let  $Y \to X$  be a biomechanical bundle with local coordinates  $(x^{\lambda}, y^{i})$ . In jet terms, a *first-order* Lagrangian is defined (through the standard Lagrangian density  $\mathcal{L}$ ) to be a horizontal density  $L = \mathcal{L}\omega$  on the jet manifold  $J^{1}Y$ . The jet manifold  $J^{1}Y$  plays the role of the finite-dimensional configuration space of sections of the bundle  $Y \to X$ .

Lagrangian L yields the Legendre morphism  $\widehat{L}: J^1Y \to \Pi$ , from the 1-jet manifold  $J^1Y$  to the Legendre manifold, given by a product [14]

$$\Pi = V^* Y \wedge (\bigwedge^{n-1} T^* X) = V^* Y \wedge (\bigwedge^n T^* X) \otimes TX,$$
(16)

where V is called the *vertical bundle*. If plays the role of the finite-dimensional phase-space of sections of  $Y \to X$ . Given the bundle coordinates  $(x^{\lambda}, y^{i})$  on  $Y \to X$ , the Legendre bundle (16) has local coordinates  $(x^{\lambda}, y^{i}, p_{i}^{\lambda})$ , where  $p_{i}^{\lambda}$  are the holonomic coordinates with the transition functions

$${p'}_{i}^{\lambda} = \det(\frac{\partial x^{\varepsilon}}{\partial x'^{\nu}}) \frac{\partial y^{j}}{\partial y'^{i}} \frac{\partial x'^{\lambda}}{\partial x^{\mu}} p_{j}^{\mu}.$$
(17)

Relative to these coordinates, the Legendre morphism  $\widehat{L}$  reads

$$p_i^{\mu} \circ \widehat{L} = \pi_i^{\mu}. \tag{18}$$

The Legendre manifold  $\Pi$  in (16) is equipped with the generalized Liouville form

$$\Theta = -p_i^{\lambda} dy^i \wedge \omega \otimes \partial_{\lambda}, \tag{19}$$

<sup>&</sup>lt;sup>9</sup>For more technical details on jet spaces with their physical applications, see [14, 15]).

where  $\otimes$  denotes the standard tensor product. Furthermore, for any Pfaffian form  $\theta$  on X we have the relation

$$\Lambda |\theta = -d(\Theta |\theta).$$

This relation introduces the the *canonical polysymplectic form* on the Legendre manifold  $\Pi$ ,

$$\Lambda = dp_i^{\lambda} \wedge dy^i \wedge \omega \otimes \partial_{\lambda}, \tag{20}$$

whose coordinate expression (20) is maintained under holonomic coordinate transformations (17). It is a pullback-valued form [14].

### 4.4 Hamiltonian Connections

Let  $J^1\Pi$  be the jet manifold of the Legendre bundle  $\Pi \to X$ . It is endowed with the adapted coordinates  $(x^{\lambda}, y^i, p_i^{\lambda}, y_{\mu}^i, p_{i\mu}^{\lambda})$ . By analogy with the notion of an autonomous Hamiltonian vectorfield  $X_H$  in (5), a connection  $\gamma$  on the bundle  $\Pi \to X$ , given by

$$\gamma = dx^{\lambda} \otimes (\partial_{\lambda} + \gamma^{i}_{\lambda}\partial_{i} + \gamma^{\mu}_{i\lambda}\partial^{i}_{\mu}),$$

is said to be *locally Hamiltonian* if the exterior form  $\gamma \rfloor \Lambda$  is closed and *Hamiltonian* if the form  $\gamma \rfloor \Lambda$  is exact. A connection  $\gamma$  is locally Hamiltonian iff it obeys the conditions [14, 15]

$$\partial_{\lambda}^{i}\gamma_{\mu}^{j} - \partial_{\mu}^{j}\gamma_{\lambda}^{i} = 0, \quad \partial_{i}\gamma_{j\mu}^{\mu} - \partial_{j}\gamma_{i\mu}^{\mu} = 0, \quad \partial_{j}\gamma_{\lambda}^{i} + \partial_{\lambda}^{i}\gamma_{j\mu}^{\mu} = 0.$$
(21)

A p-form H on the Legendre bundle  $\Pi \to X$  is called a *general Hamiltonian form* if there exists a Hamiltonian connection such that

$$\gamma | \Lambda = dH.$$

A general, dissipative, time-dependent Hamiltonian form H on  $\Pi$  is said to be *Hamiltonian* if it has the splitting

$$H = H_{\Gamma} - \widetilde{H}_{\Gamma} = p_i^{\lambda} dy^i \wedge \omega_{\lambda} - (p_i^{\lambda} \Gamma_{\lambda}^i + \widetilde{\mathcal{H}}_{\Gamma})\omega = p_i^{\lambda} dy^i \wedge \omega_{\lambda} - \mathcal{H}\omega$$
(22)

modulo closed forms, where  $\Gamma$  is a connection on Y and  $H_{\Gamma}$  is a horizontal density. This splitting is preserved under the holonomic coordinate transformations (17).

Let a Hamiltonian connection  $\gamma$  associated with a Hamiltonian form H have an integral section s of  $\Pi \to X$ , that is,  $\gamma \circ s = J^1 s$ . Then s satisfies the system of first-order Hamiltonian equations on  $\Pi$ ,

$$y^i_{\lambda} = \partial^i_{\lambda} \mathcal{H}, \qquad p^{\lambda}_{i\lambda} = -\partial_i \mathcal{H}.$$

#### 4.5 Time–Dependent Dissipative Hamiltonian Dynamics

We can now formulate the time-dependent biomechanics as an n = 1-reduction of polysymplectic dynamics described above. The biomechanical phase space is the Legendre manifold  $\Pi$ , endowed with the holonomic coordinates  $(t, y^i, p_i)$  with the transition functions

$$p_i' = \frac{\partial y^j}{\partial {y'}^i} p_j.$$

 $\Pi$  admits the canonical form  $\Lambda$  (20), which now reads

$$\Lambda = dp_i \wedge dy^i \wedge dt \otimes \partial_t$$

As a particular case of the polysymplectic machinery of the previous section, we say that a connection

$$\gamma = dt \otimes (\partial_t + \gamma^i \partial_i + \gamma_i \partial^i)$$

on the bundle  $\Pi \to X$  is locally Hamiltonian if the exterior form  $\gamma \rfloor \Lambda$  is closed and Hamiltonian if the form  $\gamma \rfloor \Lambda$  is exact. A connection  $\gamma$  is locally Hamiltonian iff it obeys the conditions (21) which now take the form

$$\partial^i \gamma^j - \partial^j \gamma^i = 0, \quad \partial_i \gamma_j - \partial_j \gamma_i = 0, \quad \partial_j \gamma^i + \partial^i \gamma_j = 0.$$

Note that every connection  $\Gamma = dt \otimes (\partial_t + \Gamma^i \partial_i)$  on the bundle  $Y \to X$  gives rise to the Hamiltonian connection  $\widetilde{\Gamma}$  on  $\Pi \to X$ , given by

$$\widetilde{\Gamma} = dt \otimes (\partial_t + \Gamma^i \partial_i - \partial_j \Gamma^i p_i \partial^j).$$

The corresponding Hamiltonian form  $H_{\Gamma}$  is given by

$$H_{\Gamma} = p_i dy^i - p_i \Gamma^i dt.$$

Let H be a dissipative Hamiltonian form (22) on  $\Pi$ , which reads:

$$H = p_i dy^i - \mathcal{H} dt = p_i dy^i - p_i \Gamma^i dt - \mathcal{H}_{\Gamma} dt.$$
<sup>(23)</sup>

We call  $\mathcal{H}$  and  $\widetilde{\mathcal{H}}$  in the decomposition (23) the Hamiltonian and the Hamiltonian function respectively. Let  $\gamma$  be a Hamiltonian connection on  $\Pi \to X$  associated with the Hamiltonian form (23). It satisfies the relations [14, 15]

$$\gamma \rfloor \Lambda = dp_i \wedge dy^i + \gamma_i dy^i \wedge dt - \gamma^i dp_i \wedge dt = dH,$$
  

$$\gamma^i = \partial^i \mathcal{H}, \qquad \gamma_i = -\partial_i \mathcal{H}.$$
(24)

From equations (24) we see that, in the case of biomechanics, one and only one Hamiltonian connection is associated with a given Hamiltonian form.

Every connection  $\gamma$  on  $\Pi \to X$  yields the system of first-order differential equations (15), which now takes the explicit form:

$$\dot{y}^i = \gamma^i, \qquad \dot{p}_i = \gamma_i. \tag{25}$$

They are called the *evolution equations*. If  $\gamma$  is a Hamiltonian connection associated with the Hamiltonian form H (23), the evolution equations (25) become the *dissipative time-dependent* Hamiltonian equations:

$$\dot{y}^i = \partial^i \mathcal{H}, \qquad \dot{p}_i = -\partial_i \mathcal{H}.$$
 (26)

In addition, given any scalar function f on  $\Pi$ , we have the dissipative Hamiltonian evolution equation

$$d_H f = (\partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i) f, \qquad (27)$$

relative to the Hamiltonian  $\mathcal{H}$ . On solutions s of the Hamiltonian equations (26), the evolution equation (27) is equal to the total time derivative of the function f:

$$s^*d_H f = \frac{d}{dt}(f \circ s).$$

### 4.6 Time & Fitness–Dependent Biomechanics

The dissipative Hamiltonian system (26)–(27) is the basis for our time & fitness-dependent biomechanics. The scalar function f in (27) on the biomechanical Legendre phase-space manifold  $\Pi$  is now interpreted as an *individual neuro-muscular fitness function*. This fitness function is a 'determinant' for the performance of muscular drives for the driven, dissipative Hamiltonian biomechanics. These muscular drives, for all active DOF, are given by time & fitness-dependent Pfaffian form:  $F_i = F_i(t, y, p, f)$ . In this way, we obtain our final model for time & fitness-dependent Hamiltonian biomechanics:

$$\begin{aligned} \dot{y}^i &= \partial^i \mathcal{H}, \\ \dot{p}_i &= F_i - \partial_i \mathcal{H}, \\ d_H f &= (\partial_t + \partial^i \mathcal{H} \partial_i - \partial_i \mathcal{H} \partial^i) f. \end{aligned}$$

Physiologically, the active muscular drives  $F_i = F_i(t, y, p, f)$  consist of [1, 2]):

1. Synovial joint mechanics, giving the first stabilizing effect to the conservative skeleton dynamics, is described by the  $(y, \dot{y})$ -form of the Rayleigh-Van der Pol's dissipation function

$$R = \frac{1}{2} \sum_{i=1}^{n} (\dot{y}^{i})^{2} [\alpha_{i} + \beta_{i} (y^{i})^{2}],$$

where  $\alpha_i$  and  $\beta_i$  denote dissipation parameters. Its partial derivatives give rise to the viscousdamping torques and forces in the joints

$$\mathcal{F}_{i}^{joint} = \partial R / \partial \dot{y}^{i},$$

which are linear in  $\dot{y}^i$  and quadratic in  $y^i$ .

2. Muscular mechanics, giving the driving torques and forces  $\mathcal{F}_i^{musc} = \mathcal{F}_i^{musc}(t, y, \dot{y})$  with (i = 1, ..., n) for human biomechanics, describes the internal excitation and contraction dynamics of equivalent muscular actuators [11].

(a) The excitation dynamics can be described by an impulse force-time relation

$$F_i^{imp} = F_i^0(1 - e^{-t/\tau_i}) \quad \text{if stimulation} > 0$$
  
$$F_i^{imp} = F_i^0 e^{-t/\tau_i} \quad \text{if stimulation} = 0$$

where  $F_i^0$  denote the maximal isometric muscular torques and forces, while  $\tau_i$  denote the associated time characteristics of particular muscular actuators. This relation represents a solution of the Wilkie's muscular *active-state element* equation [12]

$$\dot{\mu} + \Gamma \mu = \Gamma S A, \quad \mu(0) = 0, \quad 0 < S < 1,$$

where  $\mu = \mu(t)$  represents the active state of the muscle,  $\Gamma$  denotes the element gain, A corresponds to the maximum tension the element can develop, and S = S(r) is the 'desired' active state as a function of the motor unit stimulus rate r. This is the basis for biomechanical force controller.

(b) The contraction dynamics has classically been described by the Hill's hyperbolic forcevelocity relation [13]

$$F_i^{Hill} = \frac{\left(F_i^0 b_i - \delta_{ij} a_i \dot{y}^j\right)}{\left(\delta_{ij} \dot{y}^j + b_i\right)},$$

where  $a_i$  and  $b_i$  denote the Hill's parameters, corresponding to the energy dissipated during the contraction and the phosphagenic energy conversion rate, respectively, while  $\delta_{ij}$  is the Kronecker's  $\delta$ -tensor.

In this way, human biomechanics describes the excitation/contraction dynamics for the ith equivalent muscle–joint actuator, using the simple impulse–hyperbolic product relation

$$\mathcal{F}_i^{musc}(t, y, \dot{y}) = F_i^{imp} \times F_i^{Hill}$$

Now, for the purpose of biomedical engineering and rehabilitation, human biomechanics has developed the so-called *hybrid rotational actuator*. It includes, along with muscular and viscous forces, the D.C. motor drives, as used in robotics

$$\mathcal{F}_k^{robo} = i_k(t) - J_k \ddot{y}_k(t) - B_k \dot{y}_k(t), \quad \text{with} \\ l_k i_k(t) + R_k i_k(t) + C_k \dot{y}_k(t) = u_k(t),$$

where k = 1, ..., n,  $i_k(t)$  and  $u_k(t)$  denote currents and voltages in the rotors of the drives,  $R_k, l_k$ and  $C_k$  are resistances, inductances and capacitances in the rotors, respectively, while  $J_k$  and  $B_k$ correspond to inertia moments and viscous dampings of the drives, respectively.

Finally, to make the model more realistic, we need to add some stochastic torques and forces:

$$\mathcal{F}_i^{stoch} = B_{ij}[y^i(t), t] \, dW^j(t),$$

where  $B_{ij}[y(t), t]$  represents continuous stochastic diffusion fluctuations, and  $W^{j}(t)$  is an N-variable Wiener process (i.e., generalized Brownian motion) [5], with

$$dW^j(t) = W^j(t+dt) - W^j(t),$$
 (for  $j = 1, \dots, n = \text{no. of active DOF}$ ).

### 5 Conclusion

We have presented the time-dependent generalization of an 'ordinary' autonomous Hamiltonian biomechanics, in which total mechanical + biochemical energy is not conserved. Starting with the Covariant Force Law, we have first developed autonomous Hamiltonian biomechanics. Then we have introduced a general framework for time-dependent Hamiltonian biomechanics in terms jets, Legendre manifolds and dissipative Hamiltonian connections associated with the extended musculo-skeletal configuration manifold, called the configuration bundle. In this way we formulated a general Hamiltonian model for time & fitness-dependent human biomechanics.

## References

- [1] Ivancevic, V., Ivancevic, T., Human–Like Biomechanics: A Unified Mathematical Approach to Human Biomechanics and Humanoid Robotics. Springer, Dordrecht, (2006)
- [2] Ivancevic, V., Ivancevic, T., Natural Biodynamics. World Scientific, Singapore (2006)
- [3] Ivancevic, V., Ivancevic, T., Geometrical Dynamics of Complex Systems: A Unified Modelling Approach to Physics, Control, Biomechanics, Neurodynamics and Psycho-Socio-Economical Dynamics. Springer, Dordrecht, (2006)
- [4] Ivancevic, V., Ivancevic, T., Applied Differential Geometry: A Modern Introduction. World Scientific, Singapore, (2007)
- [5] Ivancevic, V., Ivancevic, T., High–Dimensional Chaotic and Attractor Systems. Springer, Berlin, (2006)
- [6] Ivancevic, V., Ivancevic, T., Human versus humanoid robot biodynamcis. Int. J. Hum. Rob. 5(4), 699-713, (2008)
- [7] Ivancevic, V., Ivancevic, T., Complex Nonlinearity: Chaos, Phase Transitions, Topology Change and Path Integrals, Springer, Berlin, (008)
- [8] Ivancevic, T., Jovanovic, B., Djukic, M., Markovic, S., Djukic, N., Biomechanical Analysis of Shots and Ball Motion in Tennis and the Analogy with Handball Throws, J. Facta Universitatis, Series: Sport, 6(1), 51–66, (2008)
- [9] Ivancevic, T., Jain, L., Pattison, J., Hariz, A., Nonlinear Dynamics and Chaos Methods in Neurodynamics and Complex Data Analysis, Nonl. Dyn. (Springer), 56(1-2), 23–44, (2009)
- [10] Ivancevic, T., Jovanovic, B., Djukic, S., Djukic, M., Markovic, S., Complex Sports Biodynamics: With Practical Applications in Tennis, Springer, Berlin, (2009)
- [11] Hatze, H., A general myocybernetic control model of skeletal muscle. Biol. Cyber. 28, 143–157, (1978)
- [12] Wilkie, D.R., The mechanical properties of muscle. Brit. Med. Bull. 12, 177–182, (1956)
- [13] Hill, A.V., The heat of shortening and the dynamic constants of muscle. Proc. Roy. Soc. B76, 136–195, (1938)
- [14] Giachetta, G., Mangiarotti, L., Sardanashvily, G., New Lagrangian and Hamiltonian Methods in Field Theory, World Scientific, Singapore, (1997)
- [15] Sardanashvily, G.: Hamiltonian time-dependent mechanics. J. Math. Phys. 39, 2714, (1998)