.**Comment on the Paper "Stern-Gerlach Force on a** Precessing Magnetic Moment"by M. Conte et al¹⁾

(Bates 2008 report B/IR#08-01)

C. Tschalaer

1. **Introduction**

 The paper under discussion (ref. 1) is based on an earlier paper (ref. 2). The formula derived for the longitudinal Stern-Gerlach (SG) force in the laboratory frame in ref. 2 (eq. (22)) is not consistent with the total time derivative of the canonical momentum derived from the accepted Langrangian³⁾ which is (see ref. 4)

$$
L = -mc^2 / \gamma + e\vec{\beta} \cdot \vec{A} - e\phi + G'/\gamma
$$

where

$$
G' \equiv \vec{\mu} \cdot \vec{B}' / \gamma + (e/m)(1/(\gamma + 1))\vec{s} \cdot (\vec{\beta} \times \vec{E}' / c)
$$

is the total SG interaction and Thomas precession potential of a particle with spin \vec{s} , magnetic moment $\vec{\mu}$, mass m, and charge e; primed quantities refer to the rest frame of the particle.

The total time derivative of the canonical momentum *P* \rightarrow is thus

$$
cd\vec{P}/dt = d[mc^2\gamma\vec{\beta} + e\vec{A} + \delta(G'/\gamma)/\delta\vec{\beta}]/dt
$$

= $c\vec{\nabla}L = ec\vec{\nabla}(\vec{\beta}\cdot\vec{A}) - ec\vec{\nabla}\phi + (c/\gamma)\vec{\nabla}(G').$

For a particle traversing a localized field region F (all fields and their derivatives are zero outside For a particle traversing a localized field region Γ (
F), the resulting change $\Delta \vec{p}$ in the momentum \vec{p} is

$$
\Delta \vec{p} = \int_{F} dt \frac{d\vec{p}}{dt} = \int_{F} dt \frac{d\vec{P}}{dt} = mc\Delta(\gamma \vec{\beta})
$$

= $e \int_{F} dt \vec{\nabla} [(\vec{\beta} \cdot \vec{A}) - \phi] + (1/\gamma) \int_{F} (dt/\gamma) \vec{\nabla} (G')$.

The first integral is the electromagnetic interaction and the second the SG-Thomas interaction which contains no terms proportional to γ^2 and γ .

2. Paper-specific Comment

If, for the sake of discussion, we consider eq. (22) in ref. 2

$$
\vec{f}_{SG} = (1/\gamma)\hat{x}\delta(\vec{\mu}\cdot\vec{B}')/\delta x + (1/\gamma)\hat{y}\delta(\vec{\mu}\cdot\vec{B}')/\delta y + \hat{z}\delta(\vec{\mu}\cdot\vec{B}')/\delta z
$$
\n(22)

as the correct expression for the SG force and ignoring the Thomas precession, we can evaluate it using eq. (12) in ref. 2

$$
\delta/\delta z' = \gamma \delta/\delta z + (\gamma \beta/c) \delta/\delta t \tag{12}
$$

and by introducing the total time differential

$$
d/dt = \delta/\delta t + \beta c \delta/\delta z
$$

so that

$$
\delta/\delta z' = (\gamma \beta / c) d / dt - \gamma \beta^2 \delta / \delta z + \gamma \delta / \delta_z
$$

= $(\gamma \beta / c) d / dt + (1/\gamma) \delta / \delta z$.

Thus eq. (22) yields

$$
f_z = (\gamma \beta / c) d(\vec{\mu} \cdot \vec{B'}) / dt + (1/\gamma) \vec{\mu} \cdot \delta \vec{B'} / \delta z
$$

or

$$
dp_z/dt = f_z = mc \ d(\gamma \beta)/dt .
$$

The total longitudinal momentum change induced by SG forces from a localized field region F is

$$
\Delta p_z = \int\limits_F dt \cdot f_z = \int\limits_F dt (\gamma \beta / c) \ d(\vec{\mu} \cdot \vec{B}') / dt + \int\limits_F (dt / \gamma) \vec{\mu} \cdot \delta \vec{B}' / \delta z \ .
$$

Using partial integration and remembering that \vec{B}' \rightarrow is zero outside F we find

$$
\Delta p_z = -\int\limits_F dt [(\vec{\mu} \cdot \vec{B}') (mc^2)] d(\gamma \beta mc) / dt + \int\limits_F (dt / \gamma) \vec{\mu} \cdot \delta \vec{B}' / \delta z.
$$

Since

$$
\mu mc^2 = (g/2) 2.96 \cdot 10^{-11} (MeV)^2 / Tesla
$$

the SG interaction potential $\vec{\mu} \cdot \vec{B}'$ is much smaller than even the electron rest mass energy i.e.,

$$
(\vec{\mu} \cdot \vec{B}')/(mc^2) \approx 1.2 \cdot 10^{-10} B'/Tesla
$$

and much smaller for nucleons. Therefore we may approximate

$$
\Delta p_z + \int_F dt [\vec{\mu} \cdot \vec{B}' / (mc^2)] dp_z / dt \approx \Delta p_z [1 + (\vec{\mu} \cdot \vec{B}' / (mc^2))_{av}] \approx \Delta p_z
$$

or

$$
\Delta p_z = \int\limits_F (dt/\gamma) \vec{\mu} \cdot \delta \vec{B}' / \delta_z \times [1 + order(\vec{\mu} \cdot \vec{B}') (mc^2))]^{-1} ,
$$

i.e. there are no first-order γ^2 – terms in Δp_z . In ref. 1, this fact is acknowledged for the case of non-precessing spin. However, it is then claimed that, for a particular model with precessing spin, the γ^2 term survives. We now show that if the spin motion is accounted for properly, the γ^2 terms cancel again as in the case of fixed spin.

3. **Precessing spin in a T₀₁₁ field**

A particle traveling on axis through a rectangular RF cavity $(x = a/2)$; $y = b/2$) containing a T_{011} field as described in ref. 1 "sees" the following RF field \vec{B}' :

$$
B'_x = \gamma B_x + \gamma \beta E_y / c = 0
$$

\n
$$
B'_y = \gamma B_y - \gamma \beta E_x / c = -\gamma B_0 (b/d) [\cos(\pi \zeta) \cos \omega t - w \beta \sin(\pi \zeta) \sin \omega t]
$$

\n
$$
B'_z = B_z = 0
$$

where

$$
\zeta \equiv z/d
$$

\n
$$
\omega t = \pi w \zeta / \beta + \varphi
$$

\n
$$
w \equiv \sqrt{1 + d^2 / b^2} = \omega d / (\pi c) .
$$

Eqs. (12) and (22) then yield

Eqs. (12) and (22) then yield

$$
f_z = (\gamma \beta / c) \delta (\vec{\mu} \cdot \vec{B'}) / \delta t + \gamma \vec{\mu} \cdot \delta \vec{B'} / \delta z.
$$

Spin rotation in a magnetic field is described by 5)

$$
\delta \vec{\mu} / \delta t = \vec{\mu} \times \vec{B}' \cdot eg / (2m) .
$$

Therefore, $\vec{B}' \cdot \delta \vec{\mu}/\delta t$ is zero and

$$
f_z = (\gamma \beta / c) \vec{\mu} \cdot \delta \vec{B}' / \delta t + \gamma \vec{\mu} \cdot \delta \vec{B}' / \delta z
$$

$$
= \gamma \mu_y [\delta B'_y / \delta z + (\beta / c) \delta B'_y / \delta t .
$$

If we now apply a strong constant field $B'_x \gg B'_y$ as suggested in ref. 1 to precess the moment $\vec{\mu}$ which points in the z-direction on entering the cavity, the spin rotation equations yields

$$
d\mu_z / dt = (\mu_x B'_y - \mu_y B'_x)eg/(2m\gamma) \approx -\mu_y B'_x \text{ eg }/(2m\gamma) = -\Omega \mu_y
$$

$$
d\mu_y / dt = \mu_z B'_x \text{ eg }/(2m\gamma) = \Omega \mu_z
$$

where

$$
\Omega \equiv B'_x \, \, eg \, / (2m\gamma)
$$

which has the solutions

$$
\mu_{y} = \mu \sin[\Omega(t+t_0)], \quad \mu_{z} = \mu \cos[\Omega(t-t_0)].
$$

In ref. 1., B'_x was chosen to produce a 180° rotation of $\vec{\mu}$ in a cavity traversal i.e.

$$
\Omega(t-t_0) = \pi_{\zeta}
$$

and

$$
B'_x \, \, eg \, / (2m\gamma) = \pi \beta c \, / \, d
$$

However, spin rotation also results in a component μ_x produced by B'_y

$$
d\mu_x/dt = -\mu_z B'_y \, \,eg\,/(2m\gamma) = -\pi \beta c \mu B'_y / B'_x \cdot \cos(\pi \zeta) \,.
$$

After exiting the cavity, this moment μ_x

$$
\mu_x = -\pi \beta c \mu / B'_x \int_F dt \cos(\pi \zeta) B'_y
$$

will generate a SG force in the fringe field B'_x according to eq. (22):

$$
dp_z/dt = \gamma \mu_x \delta B_x / \delta z
$$

which produces a longitudinal momentum change Δp_z in traversing the B_x' fringe field

$$
\Delta p_2 = \mu_x \gamma / (bc) \cdot \int_{\text{fringe}} dz \, \delta B'_x / \delta z = -\mu_x B'_x \gamma / (\beta c)
$$

$$
= \gamma \mu \pi \int_{F} dt \cos(\pi \zeta) B'_y .
$$

The longitudinal momentum change Δp_1 produced by the SG force f_z in traversing the cavity is

$$
\Delta p_1 = \gamma \mu \int_F dt \, \sin(\pi \zeta) [\delta B_y' / \delta_z + (\beta/c) \delta B_y' / \delta t].
$$

The integrals for Δp_1 and Δp_2 can now be evaluated using

$$
dt=d\zeta\cdot d/(\beta c)\ .
$$

We find

$$
\delta B_y' / \delta z + (\beta/c)\delta B_y' / \delta t = \gamma \pi B_0 b / d^2 \cdot [(1 + w^2 \beta^2) \sin(\pi \zeta) \cos \omega t + 2w\beta \cos(\pi \zeta) \sin \omega t]
$$

and

$$
\Delta p_1 = \gamma^2 \pi \mu B_0 b / (\beta c d) \cdot \left[(1 + \beta^2 w^2) \right]_0^1 d\zeta \frac{1 - \cos(2\pi \zeta)}{2} \cos(\omega t) + 2w \beta \int_0^1 \frac{\sin(2\pi \zeta)}{2} \sin(\omega t) \right]
$$

$$
\Delta p_2 = -\gamma^2 \pi \mu B_0 b / (\beta c d) \left[\int_0^1 d\zeta \frac{1 + \cos(2\pi \zeta)}{2} \cos(\omega t) - w \beta \int_0^1 d\zeta \frac{\sin(2\pi \zeta)}{2} \sin(\omega t) \right].
$$

The total longitudinal momentum change is therefore

$$
\Delta p_1 + \Delta p_2 = \gamma^2 \mu B_0 b / (2 \beta c d) [\beta^2 w^2 I_0 - (2 + \beta^2 w^2) I_c + 3w \beta I_s]
$$

where

$$
I_0 \equiv \pi \int_0^1 d\zeta \cos(\pi w \zeta / \beta + \varphi) = (\beta / w) [\sin(\pi w / \beta + \varphi) - \sin \varphi]
$$

\n
$$
I_c \equiv \pi \int_0^1 d\zeta \cos(2\pi \zeta) \cos(\omega t) = (w / \beta) / (w^2 / \beta^2 - 4) \cdot [\sin(\pi w / \beta + \varphi) - \sin \varphi]
$$

\n
$$
I_s \equiv \pi \int_0^1 d\zeta \sin(2\pi \zeta) \sin(\omega t) = 2 / (w^2 / \beta^2 - 4) \cdot [\sin(\pi w / \beta + \varphi) - \sin \varphi].
$$

Thus

$$
\Delta p_1 + \Delta p_2 = [\gamma^2 \mu B_0 b / (2 \beta c d)] (2w/\beta) [(2\beta^2 - 1) / \gamma^2] [\sin(\pi w/\beta + \varphi) - \sin \varphi]
$$

= [\mu B_0 (b/d) w (2\beta^2 - 1) / (\beta^2 c)] [\sin(\pi w/\beta + \varphi) - \sin \varphi].

Therefore, the γ^2 terms of the longitudinal momentum change caused by the SG forces from the RF fields and by the fall-off of the precessing field B_x^{\dagger} cancel as predicted by the more general formalism.

The use of two equal cavities placed in series and containing RF fields of the same mode as was proposed in Ref. 1 simply extends the integrals I_0 , I_c , and I_s from 0 to 2 instead of from 0 to 1. Then the term sin $(\pi w/\beta + \varphi)$ in the last equation is replaced by sin $(2\pi w/\beta + \varphi)$, but this does not affect the cancellation of the γ^2 terms.

Conclusion

In ref. 1 it is claimed that the SG interaction results in a change of the longitudinal momentum proportional to γ^2 for a particle traversing a localized electromagnetic field and that this should be investigated as an effective way to polarize stored antiproton beams. However, this claim is inconsistent with predictions of refs. 3 and 4, from which it can be shown that such γ^2 -terms are absent.

In this paper, we have shown that the terms in γ^2 in ref. 1 are cancelled once the complete spin motion is included, thereby removing that contradiction with ref. 4.

It is worth noting that the Thomas precession included in ref. 4 does not change this picture but only modifies the strength of the non-cancelling terms.

References:

- 1) M. Conte et al., "Stern-Gerlach Force on a Precessing Magnetic Moment", Proceedings of PAC 07, (http://cern.ch/AccelConf/p07/PAPERS/THPAS105.pdf)
- 2) M. Conte et al., "The Stern-Gerlach Interaction between a Traveling Particle and a Time-Varying Magnetic Field"; INFN/TC-00/03, 22 March, 2000. (http://arxiv.org/PScache/physics/pdf/0003/0003069v1.pdf)
- 3) A.A. Pomeranskii and I. B. Khriplovich, Journal of Experimental and Theoretical Physics; Vol. **86**, No. 5, 1998
- 4) C. Tschalaer, 'The Relativistic Stern-Gerlach Force", Bates Report B/IR #06-02: (http://arxiv.org/ftp/arxiv/papers/0802/ 0802.0154.pdf)
- 5) J. D. Jackson, "Classical Electrodynamics", Wiley & Sons