

Dark Matter, Quasars, and Superstructures in the Universe (Also on the origin of supermassive physical singularities)

Xiaodong Huang

*Department of Mathematics, University of California, Los Angeles
Los Angeles, CA 90095, U.S.A.*

Wuliang Huang

*Institute of High Energy Physics, Chinese Academy of Sciences
P.O. Box 918(3), Beijing 100049, China*

Abstract: In this paper, from the observed results of the space distribution of quasars we deduced that neutrino mass, m_ν , is about $10^{-1} eV$. The fourth stable elementary particle δ with mass $m_\delta \sim 10^0 eV$ can help explain the energy resource mechanism in quasars, cosmic ultra-high energy particles, as well as the flatness of spiral galaxy rotation curves. The blue bump and IR bump in the quasar irradiation spectra, as well as the peaks of Extragalactic Background Light (EBL) around $10^0 eV$ and $10^{-1} eV$, are related to the annihilation of $\delta\bar{\delta}$ and $\nu\bar{\nu}$ respectively. Also, we explore how to use energy from dark matter. The search for δ -particle can be done using ground lab, or can be done at space station. Space electron spectrum ($> 10^0 TeV$) is related to cosmic ray spectrum (knee and ankle) at high energy regions. For all of these, the key is the super-strong gravitational/electromagnetic field from supermassive compact celestial bodies (SCCB) such as galactic nuclei, quasars etc. The first challenge is their origin. For the early era of the universe (EEU), this means: the density of the universe is ρ . If we suppose that $\rho/c^2 = const$ for $\rho > \rho_{cr}$, it will be an inflation stage, $\dot{R}^2 \propto R^2$, and there is a minimum radius r_{min} for all $\rho > \rho_{cr}$ black holes at EEU. This should be called Physical Singularity (PS) instead of black hole. Suppose $r_{min} = r_p$, then $\rho_{cr} \sim A^2 \cdot \rho_p$ and $A = m_{pl}/m_p$. During EEU, light speed c is not a constant. According to the mass scales sequence (MSS, see Appendix): M_n , $n=0,1\dots 8$; the SCCB are originated from this inflation-explosion stage ($n=5$ to 4) for one-time or two-times explosion. A small portion of the remaining explosion fragments should have scale $r_{min} = r_p$ and are composed of FTL matter as supermassive PSs connected to SCCB. The energy scale at $\rho_{cr}(M_4, R_4)$ is the critical energy E_{cr} . At E_{cr} , inner symmetries are degenerated; below E_{cr} , it can hold at most 4-fold SM. Appendix is the theoretical/phenomenal part related to MSS, inflation-explosion, spherical universe (M_4, R_4), BSM and new particles. The MSS, like a Mendeleev's periodic table, can be used to cosmology.

PACS numbers: 95.35.+d, 98.65.Dx, 98.54.-h, 26.65.+t, 89.20.-a

Keywords: dark matter, δ -particle, quasar, superstructure, new energy resource, MSS, inflation-explosion, spherical universe, beyond SM, new particles

Email address: huangwl39@yahoo.com, huangwl@ihep.ac.cn,
xhuang@ucla.edu

Supermassive compact celestial bodies (galactic nuclei, quasars...) originate from inflation-explosion stage at EEU. As noted in the Appendix, MSS^[1] from n=5 to n=4 is a fission process (inflation) with explosions. A small portion of the remaining explosion fragments should have scale r_{\min} and are composed of FTL matter, that are the PSs. Using the word “hermit” from “hermit crab”, the “hermit” galactic nucleus/quasar contains the PS, hiding in matter and dark matter density waves or clumps (for example the nucleus of Milky Way). We mainly discuss quasars in this paper.

Superstructure and neutrino mass

Superstructure in the universe connects with light dark matter. In Refs [1],[2] and [3], we mentioned that there exist dark matter particles ν ($\bar{\nu}$) with mass $m_\nu(m_{\bar{\nu}}) \sim 10^{-1} eV$, and may also exist dark matter particles δ ($\bar{\delta}$) with mass $m_\delta(m_{\bar{\delta}}) \sim m_\nu(m_{\bar{\nu}})$. The dark matter particles involve the superstructure with mass scale $M_F \sim 10^{19}$ solar mass and with length scale (at present time) $r_F \sim 10^3$ Mpc in the universe. Evidence pertaining to this topic has appeared: In Ref [4], from the quasar number count as a function of redshift, $N(z)$, there are several peaks between $z=0$ and $z=5$. There are strong peaks in $N(z)$ at $z \approx 0.24, 1.2, \text{ and } 1.8$.^[4] Using the results of the Fourier spectral analysis in that paper, we roughly adopted the redshift values of the peaks as follows:

i	1	2	3	4	5	6
z_i	0.24	0.5	0.8	1.2	1.8	3.0
d_i (Gpc)	0.86	1.55	2.15	2.75	3.40	4.22
Δd_i (Gpc)	0.86	0.69	0.60	0.60	0.65	0.82

Where d_i is the distance, $\Delta d_i = d_i - d_{i-1}$ and $d_0 = 0$. The average value of Δd_i is 0.7 Gpc ($\approx 10^3$ Mpc).^{**} This is the length scale of superstructure in the universe. Therefore, it can also be deduced that $m_\nu(m_{\bar{\nu}}) \sim 10^{-1} eV$ as in Refs [2] and [3], but it is not certain whether $m_\delta > m_\nu$ or $m_\delta < m_\nu$. In this paper, we are still using the order of magnitude estimation as in [2] and [3], and denote dark matter particles (ν and δ) as “ d ” and baryon particles as “ B ”.

S galaxies and quasars

In Ref [6], we discussed galaxies with two constituents: dark matter and baryon matter ($d + B, m_d \ll m_B$). In that paper, we introduced a parameter “ n ” which can be used to distinguish spiral (S) galaxy and elliptical (E) galaxy: $n > 0$ for S galaxies; $n < 0$ for E galaxies. From calculation, it is obvious that a galaxy contains more dark matter if $n > 0$. Therefore, S galaxies contain more dark matter than E galaxies. Comparing number count functions $N(z)$ of S and E galaxies^[7] with quasars,^[4] one can see that the $N(z)$ function of S galaxies is more similar to that of quasars. Thus, one of the quasar’s characteristics is that it contains abundant dark matter (and anti-dark matter).

^{**}Using the $N(z)$ data for quasi-stellar objects in Ref [5], we have a result of average $\Delta d_i \sim 0.6$ Gpc.

On the other hand, the fact that $N(z)$ of both S and E galaxies have several peaks^[7] means that both types of galaxy do not have a homogeneous space distribution in a superstructure. S and E galaxies may occupy outer region or central region respectively. In a superstructure, the light constituent d is enriched (with larger abundance rate^[8]) in outer regions and the heavy constituent B is enriched in the central region. This means that S galaxies and quasars are mainly distributed in the outer region of a superstructure.

δ particles and quasars

Since quasars contain abundant dark matter, the energy source of quasars can be related to dark matter. Thus, another energy resource mechanism is added to the standard black hole-accretion disk model of quasars.^{[9],[10]} Since there are super-strong gravitational field and super-strong electromagnetic field in quasars, dark matter and anti-dark matter are annihilated in super-strong gravitational / electromagnetic fields: $\delta + \bar{\delta} \rightarrow \text{graviton} + \text{graviton} / \nu + \bar{\nu} \rightarrow \gamma + \gamma$. The interaction between gravitons and the surrounding baryon matter produces blue light^[11], while the annihilation of $\nu\bar{\nu}$ produces IR photons. These two types of radiations are related to the “blue bump” and “IR bump”^{[12],[13],[14]} of quasars respectively.

The luminosity L_q of a quasar is about 10^{47} erg/sec. If the length scale of quasars is r_q (<1 pc), then the intensity flowing out ($\text{erg} / \text{sec} / \text{cm}^2$) of a quasar’s surface is $j_q \sim \frac{L_q}{r_q^2}$.

At the surface, the number density of d-particle (n_d) can be estimated as $(\frac{m_d c}{\hbar})^3$. During the annihilation process, the intensity entering the surface is $j_d \sim n_d (m_d c^2) \cdot c$. It follows that $j_d \sim \xi \cdot j_q$ and $\xi < 1$, so we have $m_d \sim (\frac{\xi \cdot L_q \hbar^3}{r_q^2 c^6})^{1/4} \sim 10^0 eV$, i.e. $m_\delta \sim 10^0 eV$. Therefore, $m_\delta > m_\nu$.

We discussed the mass of dark matter particles in Ref [6]. The massive neutrinos were used to explain the flatness of spiral galaxy rotation curves. From the calculation of 21 samples of S galaxies, the calculated values of neutrino mass are in the range of 5eV – 33eV.^[6] According to the analysis in this paper, the dark matter particles that cause the “flatness” are not the neutrinos, but rather the δ particles.

Cosmic blue light

The annihilation processes of $\delta\bar{\delta} / \nu\bar{\nu}$ may not only occur in quasars, but also in any places with super-strong gravitational field / electro-magnetic field in the universe, such as galactic nuclei, neutron stars, central regions of star, black holes, PSs etc. Gravitons, the product of $\delta\bar{\delta}$ annihilation, can travel a long way in space and interact with baryon matter anywhere to emit cosmic blue light. This means the 10^{14-15} Hz blue light and 10^{13-14} Hz IR radiation can occur at different z across the entire sky and make imprints on the EBL (Extragalactic Background Light). Thus, the peaks of EBL around $10^0 eV$ and $10^{-1} eV$ (a small peak) are related to the annihilation of $\delta\bar{\delta}$ and $\nu\bar{\nu}$ respectively.^{[15],[16]}

Solar neutrinos and energy resource

As discussed above, the annihilation processes of $\delta\bar{\delta}$ and $\nu\bar{\nu}$, especially the $\delta\bar{\delta}$ annihilation process ($m_\delta \gg m_\nu$), might provide a part of solar energy from LPS. This could be a reason for “missing” solar neutrinos. Since the “blue bump” and “IR bump” in the solar spectrum are not so clear, it is uncertain that the “missing solar neutrino” can be the result of this newly postulated additional energy source. We suggest another process: a portion of the solar neutrinos ν_e is annihilated with dark matter anti-neutrinos $\bar{\nu}_e$ at the central region of the sun and “missed”.

If we plan to examine the annihilation processes of $\delta\bar{\delta}$ and $\nu\bar{\nu}$ in a laboratory, the first task is to create equipment that can produce super-strong gravitational field, which may be similar to a mini National Ignition Facility (NIF). Since dark matter exists everywhere, it can be automatically accreted from the outside of the equipment, thus providing an unlimited source of fuel. The products (blue light, etc.) of this equipment can be directly transferred to electric power as solar energy equipment. So, this experiment equipment can become a new generation of artificial energy resource. Suppose the radius of reaction region is $r_c \sim 10^{-1}$ cm, and $m_\delta \sim 10^0$ eV, using the equation: $(m_\delta)^4 \sim \frac{\xi \cdot L_w \hbar^3}{r_c^2 c^6}$, the maximum output power (luminosity) $L_w \sim 10^4$ kw. This is a type of middle scale energy station, which can be used for individual living or spaceflight (if $m_\delta \sim 10^1$ eV, then $L_w \sim 10^8$ kw).

When accretion disks are consumed in quasars, it is possible that some point sources with “mono-color” gravitational wave / infrared emission can be observed across the sky. This will help us explore dark matter particles’ mass and the energy resource mechanism in quasars.

Search for δ particle

- (1) According to Dual SM/two-fold SM (see Appendix), δ particle is like neutrino but with baryon number. We may use high energy protons (>0.5 TeV) to collide carbon/beryllium target to produce δ particles; or reference equipment that was used to search for cosmic neutrinos.
- (2) For the ultra-high energy primary cosmic ray spectrum (UEPCRS), the “knee” may be related to the interaction between proton and $C\nu B$ ($p + \nu/\bar{\nu} \rightarrow n + e/\bar{e}$) and the “ankle” may be related to the interaction between proton and $C\delta B$ ($p + \delta/\bar{\delta} \rightarrow n + p/\bar{p}$). If space electron spectrum is correlated with UEPCRS, two abnormalities will appear around 10^0 TeV and 10^2 TeV. The latter is related to δ particle.
- (3) For cosmic ultra-high energy particles (such as ν_μ ^[17]), the key is the production of ultra-high energy neutrons, which may be created in the interaction process of δ and p .
- (4) Based on MSS (see Appendix) it is preferable to look for δ -particles in galaxy-clusters.
- (5) δ -particles are related to gravitons nearby PSs.

Summary:

For all of these above, the key is the super-strong gravitational/electromagnetic field from supermassive compact celestial bodies (SCCB) such as galactic nuclei, quasars etc. The first challenge is their origin. We'll discuss in Appendix for details, especially for cosmic evolution at EEU.

Appendix

Large number (A) and mass scales sequence (MSS)

The big universe described by MSS and represented by proton (m_p, r_p) has characteristic parameters: the speed of light c , the gravitation constant G , and the Planck constant \hbar . The fundamental block of mass is the most stable baryon proton with mass m_p and radius r_p ($r_p \sim \frac{\hbar}{m_p c}$).

From c, G, \hbar , we have

$$\text{Planck mass } m_{pl} \sim \sqrt{\frac{\hbar c}{G}} \sim 10^{19} GeV \quad (1)$$

$$\text{Planck length } r_{pl} \sim \sqrt{\frac{\hbar G}{c^3}} \sim \frac{G m_{pl}}{c^2} \quad (2)$$

then
$$\text{Large Number } A \sim \frac{m_{pl}}{m_p} \sim \frac{r_p}{r_{pl}} \sim 10^{19} \quad (3)$$

The mass scales sequence was suggested more than twenty years ago, now it can have the diagram (mass tree) as follows:

$$\begin{array}{ll} m_0 = M_0 & \\ m_1 & M_1 \\ m_2 & M_2 \\ m_3 & M_3 \\ m_4 & M_4 \\ m_5 & M_5 \\ m_6 & M_6 \\ m_7 & M_7 \\ m_8 & M_8 \end{array}$$

On the left side of the diagram are the mass scales of stable particles in micro-cosmos:

$$m_n = A^{\frac{n}{2}} m_{pl}, n = 0, 1, 2 \dots \quad (4)$$

(with length scale r_n)

On the right side are the corresponding mass scales of celestial bodies in macro-cosmos:

$$M_n = A^n m_{pl}, n = 0, 1, 2, 3 \dots \quad (5)$$

(with length scale R_n and density ρ_n)

For n=0, $m_0 = M_0 = m_{pl}$

For n=1, $m_1 = A^{-0.5} m_{pl}$

$M_1 = A \cdot m_{pl}$ (lightest black hole - LBH at EEU, or directly, lightest physical singularity - LPS)

For n=2, $m_2 = A^{-1} m_{pl} = m_p$ (proton)

$M_2 = A^2 m_{pl} = A^3 m_p = M_{star}$ (star / black hole)

For n=3, $m_3 = A^{-1.5} m_{pl} = A^{-0.5} m_p = m_\nu \sim 10^{-1} eV$ (neutrino)

$M_3 = A^3 m_{pl} = A \cdot M_{star}$ (superstructure)

For n=4, $m_4 = A^{-2} m_{pl}$

$M_4 = A^4 m_{pl} = A^3 M_1$ (the end of inflation-explosion, critical energy E_{cr} and critical density ρ_{cr})

For n=5, $m_5 = A^{-2.5} m_{pl}$

$M_5 = A^5 m_{pl} = A^3 \cdot M_{star} = M_u$ (the beginning of inflation-explosion, u-particle)

⋮

For n=8, $m_8 = A^{-4} m_{pl} = A^{-3} m_p = m_A$ (A-particle, the most elementary particle)

$M_8 = A^8 m_{pl} = A^3 \cdot M_u$ (heaviest celestial body for big universe)

We can insert $M' \sim M_{starcluster}$ and $M'' \sim M_{galaxycluster}$ between the mass scale M_2 and M_3 .

The corresponding particle masses are $m' = m_e$ (electron) and $m'' = m_\delta$ (δ particle) as well as $m_2 = m_p$ and $m_3 = m_\nu$. We have a total of four stable elementary particles. The δ particle is the fourth stable elementary particle, which is fermion, and the mass is about $10^0 - 10^1 eV$. It is similar to a neutrino of $10^{-1} eV$ mass, but with a baryon number.

Critical density ρ_{cr} and lightest physical singularity (LPS)

The MSS, like a Mendeleev's periodic table, can summarize and speculate on some important physical results, as listed above. If the density of the universe is ρ and suppose

that $\frac{\rho}{c^2} = \text{const}$ for $\rho > \rho_{cr}$, our universe will naturally have an inflation stage ($\dot{R}^2 \propto R^2$) and there is a minimum radius r_{\min} for all $\rho > \rho_{cr}$ celestial bodies as a physical singularity (PS). For example, $R_8 \sim R_7 \sim R_6 \sim R_5 \sim r_{\min}$. We supposed $r_{\min} = r_p$. When $\rho = \rho_{cr}$, PS is the back hole indeed. We can name it as lightest physical singularity (LPS) or lightest back hole (LBH). As $r_{\min} \sim c/\sqrt{G \cdot \rho_{cr}}$, the mass of LPS $M_{LPS} \sim r_{\min} \cdot c^2/G$, so $M_{LPS} \sim A^2 \cdot m_p \sim A \cdot m_{pl} \sim M_1$, $R_1 = r_p$ and the critical density $\rho_{cr} \sim A^2 \cdot \rho_p$, $\rho_p \sim m_p/r_p^3$.

Critical energy, inflation-explosion, and TOE

Imagining the inverse process of evolution of the universe, when ρ arrive at ρ_{cr} (n=4) from present CMB density (ρ_{CMB}): $M_4 \sim A^4 m_{pl} \sim A^3 M_1$, $R_4 = R_{star} \sim A \cdot R_1$ and

$\rho_4 \sim \frac{M_4}{R_4^3} = \frac{M_1}{R_1^3} \sim \rho_1 = \rho_{cr}$. It means that, at this time, our universe (M_4, R_4) includes

A^3 LPSs. Since then, the LPSs were merged and collapsed to a “u particle”, which has mass scale $M_u = M_5 \sim A^5 \cdot m_{pl}$ and length scale $R_5 \sim r_p$.

It is obvious that the positive process of evolution from M_5 to M_4 is exactly the inflation-explosion process in a bounded space-time. It also means that a small portion of remaining explosion fragments should have scale r_{\min} and are composed of FTL matter, for example the nucleus of Milky Way. Using the word “hermit” from “hermit crab”, the “hermit” galactic nucleus/quasar contains PS and hides in matter and dark matter density waves or clumps.

Inflation means “fission” step by step to lightest physical singularities (LPSs), which expand to form CMB, and CMB may have a fine grain structure.

The energy scale at R_4 is the critical energy E_{cr} . From ρ_{cr} we have

$E_{cr} \sim (m_p^2 c^9 \hbar G^{-1})^{1/4}$. At E_{cr} , for elementary particles there are symmetries degenerated.

At this time LSPs are produced, the boundaries of elementary particles for SM or BSM disappeared. According to MSS we will confront five back ground particles/fields:

m_4, m_5, m_6, m_7, m_8 in the big universe. This is the TOE in a new manner.

BSM, dark matter particles with low mass

The diagram of mass scales sequence is like a “mass tree”. The diagram of different SM of particle physics is like “pods” (with symmetry) on the tree.

We suggest Dual Standard Model diagram as follows:

u	c	t	γ	u_l	c_l	t_l	G
d	s	b	g	d_l	s_l	b_l	g_l
ν_e	ν_μ	ν_τ	Z^0	δ	δ'	δ''	Z'
e	μ	τ	W^\pm	p	p'	p''	W'

Where $u_l, c_l, t_l, d_l, s_l, b_l$ are lept-quarks, g_l is lept-gluon, G is graviton. Z', W' are the gauge bosons about a new type of interaction (relax interaction) related to δ particles. We can look for $G, p'...$ etc. The speed of photon γ is c . The speed of graviton G is c' . There is a CGB similar to the CMB. We also suggest Two-fold Standard Model:

u	c	t	γ	u'	c'	t'	G
d	s	b	g	d'	s'	b'	g'
ν_e	ν_μ	ν_τ	Z^0	δ	δ'	δ''	Z'
e	μ	τ	W^\pm	p	p'	p''	W'

In these models with weakened symmetry, new quarks may be related to new particles ($\pi_{0l}/\pi'_0; K_{0l}/K'_0; e_{0l}/e'_+$...) with mass $10^2 GeV - 10^1 TeV$. During the cooling process of LPS at R_4 (and also of the collision fire ball in laboratory), if a lot of electrons are created before protons, the Dual SM is preferred.

In order to search for new particles, a possible candidate is π_{0l}/π'_0 .

Discussion

(1) CMB cold spot and spherical universe.

The report about CMB cold spot^{[18],[19]} has existed for a long time. One of the possible explanations is that our universe is a spherical universe ($M_4; R_4$) and the earth is not at the center. If it is so, we can see the CMB anisotropy spectrums of cold semi-sphere is different to that of hot semi-sphere (when we make observation facing towards/away from the CMB cold spot, the first peak of CMB anisotropy spectrum will drop/rise while the sixth peak will rise/drop). The peaks of CMB anisotropy spectrum may reflect the mass spectrum of ν and δ particles, and the CMB anisotropy spectrum itself is "anisotropic".

(2) The cosmic microwave background (CMB) is from the atomic level, and correspondingly, the cosmic gravitational background (CGB) is from the nuclear level. If the collisions of electrons and positrons produce photons, then the collisions of protons and anti-protons (or deuterons and anti-deuterons) produce gravitons.

(3) For MSS, c is not a constant; it has a mutation (phase transition).

For M_n ($n \geq 4$), $c_n = c \cdot A^{\frac{n-4}{2}}$. The FTL matter cannot be explained by the theory of special relativity.

(4) From two basic supposes: " $\rho/c^2 = const$ for $\rho > \rho_{cr}$ "; " $r_{min} = r_p$ ", we have: (i) SCCB is originated from PS.

- (ii) Below E_{cr} , it can hold at most 4-fold SM.
- (iii) At E_{cr} , inner symmetries are degenerated.
- (iv) At EEU, celestial bodies have r_p scale.

References:

- [1] Wuliang Huang, et al., arXiv: astro-ph/9909321.
- [2] Huang Wu-Liang (Wuliang Huang), et al., Commun. Theor. Phys. 51 (2009) 575.
- [3] Wuliang Huang, et al., arXiv: 0804.2680.
- [4] J.G. Hartnett, arXiv: 0712.3833v2.
- [5] G. Burbidge and W.M. Napier, Astrophys. J. 121 (2001) 21.
- [6] Huang Wu-Liang (Wuliang Huang), et al., Commun. Theor. Phys. 2 (1983) 1453.
- [7] S.P. Driver, et al., Astrophys. J. 496 (1998) L93.
- [8] K. Cohen, the Theory of Isotope Separation, McGraw-Hill Book Company, 1951.
- [9] N.I. Shakura & R.A. Sunyaev, A&A 24 (1973) 337.
- [10] C.W.Morgan, et al., arXiv: 0707.0305v1.
- [11] M. Kishimoto, et al., Nature (London) 454 (2008) 492.
- [12] M. Elvis, et al., Astrophys. J. Suppl. 95 (1994) 1.
- [13] Z. Shang, et al., Astrophys. J. 619 (2005) 41.
- [14] D.B. Sanders, et al., Astrophys. J. 347 (1989) 29s.
- [15] A.Coaray, et al., arXiv: 0902.2372v1.
- [16] J.D. Finke, et al., arXiv: 0905.1115v1.
- [17] The 34th ICRI Proceedings (2015);
The Astronomer's Telegram: ATel #7856 (2015).
- [18] M. Cruz et al, arXiv: astro-ph/0405341.
- [19] Collaboration, Plank, arXiv: 1303.50