



v-matter



s-matter

## Cosmological Models Without Singularity Based on RW Metric or A New Metric and their Explanation for Evolution of the Universe

Shi-Hao Chen

<sup>1</sup>Department of Physics, Jilin University; <sup>2</sup>Institute of Theoretical Physics, Northeast Normal University, Changchun 130024, China. shchen@nenu.edu.cn

(Date textdate; Received textdate; Revised textdate; Accepted textdate; Published textdate)

#### Abstract

A new conjecture is proposed that there are two sorts of matter called s-matter and v-matter which are symmetric and whose gravitational masses are opposite to each other, although both masses are positive. Therefore, there are two sorts of symmetry breaking called V-breaking and Sbreaking. In the S-breaking, s-particles get their masses and form s-galaxies etc., while v-fermions and v-gauge bosons are still massless and form v-SU(5) singlets. There is no interaction among the v-SU(5) color-single states except gravitation so that they distribute loosely in space, cannot be observed and cause space to expand with an acceleration. When temperature increases to the critical temperature because space contracts, masses of all particles are zero so that s-particles and v-particles transform from one to another and the gravitational mass density becomes negative. Consequently, space stop to contract and inflation must occur. After reheating, space first expands with a deceleration and then expands with an acceleration up to now. There is no space-time singularity. There are the critical temperature, the highest temperature and the least scale in the universe. A formula describing distance-redshift is obtained. A huge void is not empty, and is equivalent to a huge concave lens. The densities of hydrogen in the huge voids must be much less than that predicted by the conventional theory. The gravitation between two galaxies with distance long enough will be less than that predicted by the conventional theory. A black hole with its mass and density big enough will transform into a white hole. Primordial nucleosynthesis and CMB are explained. It is possible that the universe is composed of infinite cosmic islands. The problem of energy conservation is discussed.

Keywords: Cosmology of theories beyond the SM; Cosmic singularity; Physics of the early universe; Inflation.

PACS: 98.80.Cq; 98.80.Bp; 98.80.Es; 95.35.+d; 98.80.-k

\*Supported by tau-charm physics research KJ(x2-yw-N29); National Natural Science Foundation of China (No. 11075064).

#### Contents

# I The cosmological model without singularity based on the RW metric

I. Introduction to the model based on the RW metric	7
II. Conjectures, action, energy-momentum tensor and field equations	10
A. Conjectures	10
B. Action	13
C. Energy-momentum tensor and field equations	16
III. Spontaneous symmetry breaking	18
IV. Evolution equations of space in RW metric	20
V. Temperature effect	21
A. Effective potentials	22
B. Critical temperatures $T_{cr}$	24
VI. Contraction of space, the highest temperature, inflation of space and	
there is no singularity	26
A. The contracting process of space and the highest temperature	26
B. A theorem related to singularity	29
C. Expansion and inflation of space	31
D. The result above is not contradictory to the singularity theorems	31
VII. Evolving process of space after inflation	33
A. The reheating process	33
B. The process of space inflation	35
C. To determine $a(t)$	36
D. The relationship between luminosity distance and its redshift	37

VIII. Existing forms and	l distribution	forms of s	s - SU(5)	color single states	38
--------------------------	----------------	------------	-----------	---------------------	----

IX. New predictions and an inference	39
A. New predictions	39
1. It is possible that huge voids are not empty and are equivalent to huge	
concave lenses. The density of hydrogen inside the huge voids is more less	
than that predicted by the conventional theory.	39
2. The gravitation between two galaxies with distant long enough will be less	S
than that predicted by the conventional theory.	40
3. A black hole with its mass and density big enough will transform into a	
white hole	41
B. An inference : $\lambda_{eff} = \lambda = 0$ , although $\rho_{vac} \neq 0$	41
	40
A. Discussion and conclusions	42
A. Discussion	42
B. Conclusions of the model based on the RW metric	43

## II An explanation for some phenomena based on the first model

All inforduction to explanation for some phenomena	40
A. The $F - W$ dark matter model	46
B. The cold-dark matter model	48
XII. Primordial nucleosynthesis	49
A. Explanation for primordial nucleosynthesis the by the Friedmann model	49
B. Explanation for primordial nucleosynthesis by this cosmological model and	
the F-W dark matter model or the cold dark matter model	50
XIII. Cosmic microwave background radiation	52
A. The recombination temperature $T_{rec}$	52
B. The temperature of matter-radiation equality	52
C. Decoupling temperature	54
D. Space is open	56

### XIV. Dynamics of structure formation and the distributive form of the

SU(5) singlets	58
A. The equations of structure formation	58
B. Three predictions.	60
XV. Conclusions of the explanation for some phenomena	62

# III A cosmological model without singularity based on a new metric

XVI.	Introduction to the model based on a new metric	62
XVII.	Conjectures	64
	A. Conjectures	64
	B. The new metric is applicable to incomplete hypersurfaces of simultaneity	64
XVIII.	Evolution equations of space	65
	A. The evolution equations in a new metric	66
	B. The asymptotic solution of the equations $(18.24) - (18.27)$	69
	1. The asymptotic solution of $K(t, r)$	69
	2. Discussion of $K$	70
XIX.	Contraction of space, the highest temperature and inflation of space	75
	A. Contraction of space	75
	B. There is non-singularity in the second model, the highest temperature and	
	inflation of space	76
XX.	Evolving process of space after inflation	78
	A. The three stage of the universe evolution	79
	B. Determination of $a(t)$ in the second model	80
XXI.	Transformation of repulsive potential energy chiefly into the kinetic	
	energy of $SU(5)$ singlets	81
XXII.	New predictions	82

	A. The second model can obtain the three predictions of the first model and	
	explain primordial, $CMBR$ and the large scale structure.	82
	B. The universe is composed of infinite s-cosmic islands and v-cosmic islands	83
	C. Mass redshifts	85
XXIII.	Conclusions of the model based on a new metric	85
XXIV.	Summation	87
	Acknowledgments	87
	References	88

Part I

# The cosmological model without singularity based on the RW metric

The cosmological model without singularity can be constructed based on the RW metric and a new metric. The curvature factor K in the RW metric can be -1, 0 or 1. The difference between the new metric and the RW metric is that K in the new metric is a function of the gravitational mass density in comoving coordinates. In addition to the major inferences of the first model (this model based on the RW metric), the second model (this model based on the new metric) has new inferences. For example, the universe is composed of infinite s-cosmic islands and v-cosmic islands according to the second model.

The present paper is composed of three parts. The first part describes the first model, the second part explains primordial nucleosynthesis CMBR and the large scale structure, the third part describes the second model.

#### I. INTRODUCTION TO THE MODEL BASED ON THE RW METRIC

It is impossible to solve the space-time singularity issue and the cosmological constant issue in the frame of the conventional theory, hence new conjectures are necessary. Based on a new essential conjecture, we have constructed a cosmological model and solve the two issues, well explain the evolution of the universe, primordial nucleosynthesis, cosmic microwave background radiation (CMBR) and give new predictions.

As is now well known, there is space-time singularity under certain conditions<sup>[1]</sup>. These conditions fall into three categories. First, there is the requirement that gravity shall be attractive. Secondly, there is the requirement that there is enough matter present in some region to prevent anything escaping from that region. The third requirement is that there should be no causality violations. Because of the theorems, there must be space-time singularity in the conventional theory. On the other hand, there should be no space-time singularity in physics. Hence this problem must be solved.

Recent astronomical observations show that the universe expanded with a deceleration earlier while is expanding with an acceleration now. This implies that there is dark energy. 73 percent of the total energy density of the universe is dark energy density<sup>[2]</sup>. What is dark energy? Many possible answers have been given. One possible interpretation is in terms of the effective cosmological constant  $\lambda_{eff} = \lambda + \rho_{gvac}$ , here  $\lambda$  and  $\rho_{gvac}$  are the Einstein cosmological constant and the gravitational mass density of the vacuum state, respectively. According to the equivalence principle,  $\rho_{gvac} = \rho_{vac}$ ,  $\rho_{vac}$  is the energy density of the vacuum state, hence  $\lambda_{eff}$  may be written as  $\lambda + \rho_{vac}$ .  $\lambda_{eff}$  cannot be derived from basic theories<sup>[3]</sup>. and  $\rho_{vac} \gg \lambda_{eff}$ . Hence the interpretation is unsatisfactory. Alternatively, dark energy is associated with the dynamics of scalar field  $\phi(t)$  that is uniform in space<sup>[4]</sup>. This is a seesaw cosmology<sup>[5]</sup>. Thus, discussion about the universe expansion with an acceleration is still open to the public.

 $\rho_{gvac} = \rho_{vac} \gg \lambda_{eff}$  originates from the conventional quantum field theory and the equivalent principle.  $\rho_{vac} \gg \lambda_{eff}$  and the singularity issue imply that the conventional theory is not self-consistent.  $\rho_{vac} = 0$  is a necessary result of our quantum field theory without divergence<sup>[6]</sup>. In this theory, there is no divergence of loop corrections as well, and dark matter which can form dark galaxies is predicted<sup>[7]</sup>. It is different from the supersymmetric quantum field theory in which  $\rho_{vac} = 0$  can be obtained in only some models but is not necessary. Thus, issue of the cosmological constant is open as well, because  $\lambda_{eff}$  is not still determined.

Huge voids in the cosmos have been observed<sup>[8]</sup>. Such a model in which hot dark matter (e.g. neutrinos) is dominant can explain the phenomenon. However, it cannot explain the structure with middle and small scales. Why are there the huge voids? This is an open problem as well.

We consider that all important existing forms of matter (including dark matter and dark energy) have appeared. Hence the basic problems should be solved. As mentioned above, we have constructed a quantum field theory without divergence which predicts that there must be dark matter. We construct a cosmological model without singularity which can solve the space-time singularity and cosmological constant issues and explain well the evolution of the universe in the present paper.

The bases of the present model are the general relativity, the conventional quantum field theory and grand unified theory (GUT).

We consider the following condition to be necessary in order to solve the space time singularity and the cosmological constant problems on the basis of the classical cosmology and in the frame of the conventional quantum field theory.

Condition: There are two sorts of matter with positive masses which are symmetric, whose gravitational masses are opposite to each other, although whose masses are all positive.

The two sorts of matter are called s-matter and v-matter, respectively. The condition implies that if  $\rho_s = \rho_v$ , then  $\rho_{gs} = -\rho_{gv}$ , here  $\rho_g$  denotes a gravitational mass density,  $\rho_s > 0$ and  $\rho_v > 0$  and there is no negative energy or negative probability at all. The conditions cannot be realized in the conventional theory, but can be realized in the present model. In order to uniformly solve the above four problems consistently on one basis, we present a new conjectures equivalent to the condition and construct two cosmological model, i.e. [9] and this model in the present paper.

The basic idea of the present model is only conjecture 1. The present model has the following results:

**1**. There is no space-time singularity. A theorem related to singularity is presented.

2. The evolution of the universe and the relation between the optical distance and the redshift predicted by the present model are consistent with the observations up to now.

There are two sorts of spontaneous symmetry breaking in the present model because of

conjecture 1, and they are called S – breaking and V – breaking.

According to the present model, the evolving process of space is as follows.

In the S – breaking, space contracts so that temperature T rises. When T arrives the critical temperature  $T_{cr}$ , the universe is in the most symmetric state with  $s - SU(5) \times v - SU(5)$  symmetry. When space continues to contract so that T arrives the highest temperature  $T_{\text{max}}$ , space expands and then inflates. After inflation, the state with the highest symmetry transits to the state with the V – breaking. After reheating, space expands with a deceleration, then expands with an acceleration up to now.

**3**. There are the critical temperature  $T_{cr}$ , the highest temperature  $T_{\max} \ge T_{cr}$ , the least scale  $R_{\min} > 0$  and the largest energy density  $\rho_{\max}$  in the universe.  $R_{\min}$  and  $T_{cr}$  are two new important constants,  $T_{\max}$  and  $\rho_{\max}$  are finite and determined by  $R(T_{cr})$ .

4. We generalize equations governing nonrelativistic fluid motion to present model. The equations of structure formation have been derived out. According to the equations, galaxies can form earlier than that in the conventional theory.

5. Three predictions are given.

6. Primordial nucleosynthesis and cosmic microwave background radiation are explained.

7. Dark energy is explained as s - matter in the V - breaking. In contrast with the dark energy,  $\rho_{sg} = -\rho_s < 0$ .

8.  $\rho_{gvac} = \rho_{s,vac} - \rho_{v,vac} = 0$  is proven, although  $\rho_{vac} = \rho_{s,vac} + \rho_{v,vac}$  is still very large. Consequently,  $\lambda_{eff} = \lambda = 0$ .

Problems 4 and 6 will be discussed in the following paper.

In section 2, conjectures are presented; In section 3, action, energy-momentum tensor and field equations are presented; In section 4, spontaneous symmetry breaking is given, and evolution equations of space are derived; In section 5, temperature effects are considered; In section 6, contraction of space, the highest temperature and inflation of space are discussed; In section 7, evolving process of space after inflation is discussed; In section 8, expansion of space after inflation is discussed; In section 9, new predictions and an inference are given; Section 10 is discussion; Section 11 is the conclusions.

# II. CONJECTURES, ACTION, ENERGY-MOMENTUM TENSOR AND FIELD EQUATIONS

#### A. Conjectures

In order to solve the above problems, we propose the following conjectures:

**Conjecture 1** There are two sorts of matter with positive masses which are called solid – matter (s - matter) and void – matter (v - matter), respectively. Both are symmetric and their contributions to the Einstein tensor are opposite each other. There is no other interaction between both except the interaction (2.10) of s – Higgs fields and v – Higgs fields.

**Conjecture 2** When SU(5) symmetry holds and temperature is low, all particles must exist in SU(5) color single states.

This conjecture is obviously a direct generalization of SU(3) color single states.

Other premises of the present model are the cosmological principle, the RW metric (k = -1 in the present model) and the conventional SU(5) grand unified theory (GUT). But it is easily seen that the present model does not rely on the special GUT. Provided conjecture 1 and such a coupling as (2.10) are kept, the GUT can be accepted.

All the following inferences hold when  $S \rightleftharpoons V$  and  $s \leftrightarrows v$  due to the conjecture 1.

The gravitational properties of matter and the mode of symmetry breaking determine the features of space-time. We consider that there are only two possibilities.

**A**. The first possibility can be described by the conventional theory. There is only one sort of matter so that the equivalence principle strictly holds. This theory based on the conjecture is simple, but there must be essential difficulties. For example, there must be the singularity and cosmological constant issues which cannot be solved in the frame of this theory because of the Hawking theorems etc.

**B**. The basis of the second possibility is conjecture 1.

We explain it in detail as follows:

1. It must be emphasized that there is no negative mass or negative probability in the present model at all. Conjecture 1 implies that  $\rho_{sg} = -\rho_{vg}$  when  $\rho_s = \rho_v$ . In the *S*-breaking,  $\rho_{sg} = \rho_s \ge 0$  and  $\rho_{vg} = -\rho_v \le 0$ . Here  $\rho_g$  denotes a gravitational mass density. From this

we can regard the gravitation charges of s - matter and v - matter to be opposite, i.e.  $\alpha = -\beta = 1$ , here  $\alpha$  and  $\beta$  are the gravitation charges. The energy-momentum tensor should be independent of the gravitation charges. Hence it is necessary to eliminate  $\alpha$  and  $\beta$ from the definition of  $T_{\mu\nu}$  by the operator  $(\partial/\alpha + \partial/\beta)$ . Consequently, both the s - energyand the v - energy must be positive (see (2.20) – (2.21) in detail).

2. The observation basis of conjecture 1 is that space expands with an acceleration. One of the two sorts of matter must loosely distribute in whole space, can cause space to expand with an acceleration and cannot be observed as so-called dark energy (see 7 below).

**3**. Because of conjecture 1, there must be two sorts of symmetry breaking.

Because of conjecture 1, s - Higgs fields and v - Higgs fields must be symmetric as well. If this symmetry is not broken, s - matter and v - matter will exist in the same form at arbitrary time and place. This implies that nature is simply duplicate. This is impossible because nature does like duplicate. Of course, this contradicts experiments and observations. Consequently the symmetry must be broken, i.e.  $\langle \omega_s \rangle_0 \neq 0$  and  $\langle \omega_v \rangle_0 = 0$ or  $\langle \omega_v \rangle_0 \neq 0$  and  $\langle \omega_s \rangle_0 = 0$ , here  $\omega$  denotes an arbitrary Higgs field. Thus the coupling constant  $\Lambda$  etc. in (2.10) must be positive so that there must be the two sorts of breaking.

The existing probability of the S-breaking and the V-breaking must be equal because the s-Higgs fields and the v-Higgs fields are symmetric. This equality can be realized by two sorts of modes.

(1) The universe is composed of infinite s - cosmic islands with the S - breaking and v - cosmic islands with the V - breaking; This possibility will be discussed in the third part (see also Ref. [9])

(2) The whole universe is in the same breaking (e.g. the S - breaking). But one sort of breaking can transform to another as space contracts to the least scale  $R_{\min}$  (see later). We discuss the case in the present paper. The RW metric is applicable to the case.

4. S - matter and v - matter are no longer symmetric after the symmetry breaking and there is no interaction except the gravitation among v - SU(5) color-single states.

In the S-breaking,  $\langle \omega_s \rangle_0 \neq 0$  and  $\langle \omega_v \rangle_0 = 0$ . Consequently, s - SU(5) is broken finally to  $s - SU(3) \times U(1)$  and v - SU(5) still strictly holds. Thus, s-particles get their masses and form s-atoms, s-observers and s-galaxies etc.; while all v-fermions and v-gauge bosons are massless and all v-particles must form v - SU(5) color-single states after reheating because of conjecture 2.

There is no interaction (e.g. electroweak) except the gravitation among v - SU(5) colorsingle states, because SU(5) is a simple group. Consequently the v - SU(5) color-single states cannot form v - atoms etc., and must distribute loosely in space as the so-called dark energy.

Thus, in the S-breaking, s-matter is identified with the conventional matter, while v-matter is similar to dark energy, because of the following reasons 5 and 6. In contrast with the dark energy, the gravitational masses of v-matter is negative.

5. The equivalence principle still strictly holds for the s - particles  $(m_{sg} = m_s)$ , but is violated by the v - particles  $(m_{vg} = -m_v)$  in the S - breaking. Although it is such, the motion equations of all s - particles and all v - particles are still independent of their masses.

In the S-breaking, there are only the s-observers and the s-galaxies, and there is no v-observer and v-galaxy. Hence the gravitational masses of s-particles must be positive, i.e.  $m_{sg} = m_s > 0$ , while the gravitational masses of v-matter must be negative relatively to s-matter, i.e.  $m_{vg} = -m_v < 0$ , because of conjecture 1. Thus, a s-photon falling in a gravitational field must have purple shift, but a v-particle (there is no v-photon and there are only v - SU(5) color single states) falling in the same gravitational field will have 'redshift'. Although the equivalence principle is violated by v-matter in the S-breaking, this result does not contradict any observation or experiment, because v-SU(5) color-single states cannot be observed by an s-observer (see 6 and 7 below).

6. There is only the repulsion between s - matter and v - matter so that any bound state is composed of only s - particles or only v - particles.

Because of conjecture 1, there is the repulsion between s - matter and v - matter and the coupling constant of the repulsion is the same as that of the gravitation.

The interaction (2.10) is repulsive as well. The interaction (2.10) can be neglected after reheating, because the masses of the Higgs particles  $\Omega_s$  and  $\Omega_v$  are very large after reheating. Thus, there is no the transformation of s - particles and v - particles from one into another in low temperatures. The interaction (2.10) is important when temperature is high enough  $(T \sim T_{cr})$ .

7. The v-SU(5) color-single states cannot be observed by an s-observer in fact, because of the following reasons.

As mentioned above, the v - SU(5) color-single states cannot form any atom or any celestial body with a large mass and must distribute loosely in space because there is no interaction except the gravitation among them. Hence the repulsion between a s - body and the v - SU(5) color-single states must be too small for observation.

On the other hand,  $\rho_v$  must be very small when  $\rho_s$  is large because there is the repulsion and there is only the repulsion determined by conjecture 1 between s-matter and v-matterafter reheating.

After reheating, the interaction (2.10) is too small to observation as well.

In fact, in the S – *breaking*, only the cosmological effects of v – *matter* are important and are consistent with the observed data up to now.

It is seen from 5-7 that although the equivalence principle is violated by the SU(5) singlets, but there is no contradiction with experiments and observations up to now.

8.  $\rho_s$  and  $\rho_v$  can transform from one into another by (2.10) when  $T \sim T_{cr}$ , because the expectation values of all Higgs fields and the masses of all particles are zero in this case. Consequently,  $\rho_s$  and  $\rho_v$  can transform from one into another by (10) so that  $\rho_s = \rho_v$ ,  $T_s = T_v \sim T_{cr}$  and the symmetry of  $v - SU(5) \times s - SU(5)$  holds in this case. Thus space cannot contract to infinite small and inflation must occur.

9. From (2.16) – (2.18) and (2.22) we see that  $T^{\mu\nu}_{g;\nu} = 0$  can be derived from the field equation, but  $T^{\mu\nu}_{;\nu} = 0$  cannot be derived. This implies that there is no restriction for  $T^{\mu\nu}$ . Thus, it is possible that the differential law of energy-momentum conservation  $T^{\mu\nu}_{,\nu} = 0$  holds, where  $T^{\mu\nu}$  may contain the contribution of gravitational field as well.

In summary, in the S – breaking, the v – SU(5) color single states cannot be observed and have only the cosmological effects. Conjecture 1 does not contradict any experiment and observation up to now.

We will see in the following that the evolution of the universe can be well explained and the singularity and cosmological constant issues can be solved.

#### B. Action

It is impossible that there are simultaneously the S – breaking and the V – breaking in the same region because of (2.10). There are only s – observers and only  $I_S$  is applicable in the S – breaking, and there are only v – observers and only  $I_V$  is applicable in the V-breaking. Hence in any case, the action is unique. But the universe in the S-breaking can transform to the universe in the the V-breaking, hence both  $I_S$  and  $I_V$  are necessary. Thus, the actions should be written as two sorts of form,  $I_S$  in the S-breaking and  $I_V$  in the V-breaking. Because of conjecture 1, the structures of  $I_S$  and  $I_V$  are the same, i.e.  $I_S \rightleftharpoons I_V$  when  $S \rightleftharpoons V$  and  $s \rightleftharpoons v$ . Thus, at the zero-temperature we have

$$I_V = I_g + I_{VM} = I_S \left( s \leftrightarrows v, S \longrightarrow V \right), \quad I_S = I_g + I_{SM}, \tag{2.1}$$

$$I_g = \frac{1}{16\pi G} \left( \int_{\Sigma} R\sqrt{-g} d^4 x + 2 \int_{\partial \Sigma} K\sqrt{\pm h} d^3 x \right), \qquad (2.2)$$

$$I_{SM} = \int d^4x \sqrt{-g} \mathcal{L}_{SM},$$
  
$$\mathcal{L}_{SM} = \alpha \left( \mathcal{L}_{Ss} + V_0 \right) + \beta \mathcal{L}_{Sv} + \frac{1}{2} \left( \alpha + \beta \right) V_{sv},$$
 (2.3)

$$I_{VM} = \int d^4x \sqrt{-g} \mathcal{L}_{VM},$$
  
$$\mathcal{L}_{VM} = \alpha \left( \mathcal{L}_{Vv} + V_0 \right) + \beta \mathcal{L}_{Vs} + \frac{1}{2} \left( \alpha + \beta \right) V_{vs},$$
 (2.4)

The physics quantities with the subscript 'S' represent which have meaning in only the S - breaking, and have unmeaning in the V - breaking! It is the same for 'V' as that for 'S', because of conjecture 1. For simplicity, the subscripts 'S' and 'V' are elided in the following when there is not confusion.

$$\mathcal{L}_{s} = \mathcal{L}_{sM} \left( \Psi_{s}, g_{\mu\nu}, g_{\mu\nu}, \lambda \right) + V_{s} \left( \omega_{s} \right), \qquad (2.5)$$

$$\mathcal{L}_{v} = \mathcal{L}_{vM}\left(\Psi_{v}, g_{\mu\nu}, g_{\mu\nu}, \lambda\right) + V_{v}\left(\omega_{v}\right), \qquad (2.6)$$

$$V_{sv}(\omega_s, \omega_v) = V_{vs}(\omega_s, \omega_v); \qquad (2.7)$$

$$\omega_s \equiv \Omega_s, \ \Phi_s, \ \chi_s; \quad \omega_v \equiv \Omega_v, \ \Phi_v, \ \chi_v,$$

where the meanings of the symbols are as follows:  $g = \det(g_{\mu\nu})$ , and  $g_{\mu\nu} = diag(-1, 1, 1, 1)$ in flat space. R is the scalar curvature. Here  $\alpha$  and  $\beta$  are two parameters and we finally take  $\alpha = -\beta = 1$ .  $V_0$  is a parameter which is so taken that  $V_{s\min}(\varpi_s) + V_0 = 0$  in the S - breaking at the zero - temperature, here  $\varpi = \langle \omega \rangle$ .  $\mathcal{L}_{sM}(\mathcal{L}_{vM})$  is the Lagrangian density of all s - fields (v - fields) and their couplings of the SU(5) GUT except the Higgs potentials  $V_s$ ,  $V_v$  and  $V_{sv}$ .  $\Psi_s$  and  $\Psi_v$  represents all s - fields and all v - fields, respectively.  $\mathcal{L}_s$  and  $\mathcal{L}_s$  do not contain the contribution of the gravitation energy and the repulsion energy. It may be seen that the set of equation (2.1) - (2.7) is unchanged when the subscripts  $s \rightleftharpoons v$  and  $S \rightleftharpoons V$ . This shows the symmetry between s - matter and v - matter.

Gibbons and Hawking pointed out that in order to get the Einstein field equations<sup>[10]</sup>, it is necessary that

$$I'_g = \frac{1}{16\pi G} \int_{\Sigma} R\sqrt{-g} d^4 x \longrightarrow I_g$$
$$= \frac{1}{16\pi G} \left( \int_{\Sigma} R\sqrt{-g} d^4 x + 2 \int_{\partial \Sigma} K\sqrt{\pm h} d^3 x \right).$$

This is because it is not necessary that  $\delta\Gamma^{\alpha}_{\mu\nu} = 0$  on the boundary  $\partial\Sigma$ . Hence  $I'_g$  is replaced by  $I_g$  in (2.2).  $\Sigma$  is a manifold with four dimensions.  $\partial\Sigma$  is the boundary of  $\Sigma$ .  $K = trK^i_j$ .  $K_{ij} = -\nabla_i n_j$  is the outer curvature on  $\partial\Sigma$ .  $n_j$  is the vertical vector on  $\partial\Sigma$ .  $h = |h_{ij}|$ , and  $h_{ij}$ is the induced outer metric on  $\partial\Sigma$ . When  $\partial\Sigma$  is space-like,  $\sqrt{\pm h}$  takes positive sign. When  $\partial\Sigma$  is time-like,  $\sqrt{\pm h}$  takes negative sign.

The Higgs potentials in (2.5) - (2.7) is the following:

$$V_{s} = -\frac{1}{2}\mu^{2}\Omega_{s}^{2} + \frac{1}{4}\lambda\Omega_{s}^{4}$$
  
$$-\frac{1}{2}w\Omega_{s}^{2}Tr\Phi_{s}^{2} + \frac{1}{4}a\left(Tr\Phi_{s}^{2}\right)^{2} + \frac{1}{2}bTr\left(\Phi_{s}^{4}\right)$$
  
$$-\frac{1}{2}\varsigma\Omega_{s}^{2}\chi_{s}^{+}\chi_{s} + \frac{1}{4}\xi\left(\chi_{s}^{+}\chi_{s}\right)^{2},$$
 (2.8)

$$V_{v} = -\frac{1}{2}\mu^{2}\Omega_{v}^{2} + \frac{1}{4}\lambda\Omega_{v}^{4}$$
  
$$-\frac{1}{2}w\Omega_{v}^{2}Tr\Phi_{v}^{2} + \frac{1}{4}a\left(Tr\Phi_{v}^{2}\right)^{2} + \frac{1}{2}bTr\left(\Phi_{v}^{4}\right)$$
  
$$-\frac{1}{2}\varsigma\Omega_{v}^{2}\chi_{v}^{+}\chi_{v} + \frac{1}{4}\xi\left(\chi_{v}^{+}\chi_{v}\right)^{2},$$
 (2.9)

$$V_{sv} = \frac{1}{2}\Lambda\Omega_{s}^{2}\Omega_{v}^{2} + \frac{1}{2}\alpha\Omega_{s}^{2}Tr\Phi_{v}^{2} + \frac{1}{2}\beta\Omega_{s}^{2}\chi_{v}^{+}\chi_{v} + \frac{1}{2}\alpha\Omega_{v}^{2}Tr\Phi_{s}^{2} + \frac{1}{2}\beta\Omega_{v}^{2}\chi_{s}^{+}\chi_{s}, \qquad (2.10)$$

where  $\Omega_a$ ,  $\Phi_a = \sum_{i=1}^{24} (T_i/\sqrt{2}) \varphi_{ai}$  and  $\chi_a$  are respectively <u>1</u>, <u>24</u> and <u>5</u> dimensional representations of the SU(5) group, a = s, v, here the couplings of  $\Phi_a$  and  $\chi_a$  are ignored for short<sup>[11]</sup>. The meaning of  $\alpha$  and  $\beta$  in (2.10) is different from that in (2.3) – (2.4). Here  $\alpha$  and  $\beta$  are coupling constants. The coupling constants in (2.8) – (2.10) are all positive, especially, as mentioned before,  $\Lambda$ ,  $\alpha$  and  $\beta$  in (2.10) must be positive.

We do not consider the terms coupling to curvature scalar, e.g.  $\xi R\Omega^2$ , for a time. In fact,  $\xi R(\Omega_s^2 - \Omega_v^2) \sim 0$  when temperature T is high enough due to the symmetry between s - matter and v - matter.

#### C. Energy-momentum tensor and field equations

By the conventional method, from (2.2) we can get

$$\delta I_g = \frac{1}{16\pi G} \int_{\Sigma} \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) \delta g^{\mu\nu} \sqrt{-g} d^4 x.$$
(2.11)

Considering  $\alpha = -\beta = 1$ , from (2.3) - [2.4] we have

$$\delta I_{SM} = \int \frac{1}{\sqrt{-g}} \left[ \frac{\partial \mathcal{L}_{SM} \sqrt{-g}}{\partial g^{\mu\nu}} - \left( \frac{\partial \mathcal{L}_{SM} \sqrt{-g}}{\partial g^{\mu\nu}_{,\sigma}} \right)_{,\sigma} \right] \delta g^{\mu\nu} \sqrt{-g} d^4 x$$
$$= \int \frac{1}{2} \left[ T_{s\mu\nu} - g_{\mu\nu} V_0 - T_{\nu\mu\nu} \right] \delta g^{\mu\nu} \sqrt{-g} d^4 x, \qquad (2.12)$$

$$\delta I_{VM} = \int \frac{1}{2} \left[ T_{\nu\mu\nu} - g_{\mu\nu} V_0 - T_{s\mu\nu} \right] \delta g^{\mu\nu} \sqrt{-g} d^4 x, \qquad (2.13)$$

$$T_{a\mu\nu} = T_{aM\mu\nu} - g_{\mu\nu}V_a, \quad a = s \text{ or } v,$$
 (2.14)

$$T_{aM\mu\nu} = \frac{2}{\sqrt{-g}} \left[ \frac{\partial \left( \sqrt{-g} \mathcal{L}_{aM} \right)}{\partial g^{\mu\nu}} - \left( \frac{\partial \left( \sqrt{-g} \mathcal{L}_{aM} \right)}{\partial g^{\mu\nu}, \sigma} \right), \sigma \right].$$
(2.15)

From (2.11) - (2.13) we obtain

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -8\pi G T_{Ag\mu\nu}, \quad A = S \quad or \quad V.$$
 (2.16)

In the S-breaking,

$$T_{Sg\mu\nu} \equiv T_{s\mu\nu} - g_{\mu\nu}V_0 - T_{v\mu\nu} = T_{SMg\mu\nu} - g_{\mu\nu}V_{Sg}$$
$$T_{SMg\mu\nu} \equiv T_{sM\mu\nu} - T_{vM\mu\nu}, \quad V_{Sg} = V_s + V_0 - V_v.$$
(2.17)

In the V - breaking,

$$T_{Vg\mu\nu} \equiv T_{\nu\mu\nu} - g_{\mu\nu}V_0 - T_{s\mu\nu} = T_{VMg\mu\nu} - g_{\mu\nu}V_{Vg}$$
$$T_{VMg\mu\nu} \equiv T_{\nu M\mu\nu} - T_{sM\mu\nu}, \quad V_{Vg} = V_v + V_0 - V_s.$$
(2.18)

It is seen from (2.17)-(2.18) that  $V_{Ag}$  is independent of  $V_{sv}$ . This implies that the potential energy  $V_{sv}$  is different from other energies in essence. There is no contribution of  $V_{sv}$  to  $R_{\mu\nu}$ , i.e., there is no gravitation and repulsion of the potential energy  $V_{sv}$ . This does not satisfy the equivalence principle. But this does not cause any contradiction with all given experiments and astronomical observations because  $V_{sv} = 0$  in either of breaking modes.

We will see that, in fact,  $V_{v\min}(\varpi_v) = 0$  because  $\langle \omega_v \rangle = 0$  in the *S* – breaking, and  $V_{s\min}(\varpi_s) = 0$  because  $\langle \omega_s \rangle = 0$  in the *V* – breaking. Hence

$$V_{Ag\min}\left(\varpi_s, \varpi_v\right) = V_{a\min} + V_0. \tag{2.19}$$

 $T_{Ag\mu\nu}$ ,  $T_{AMg\mu\nu}$  and  $V_{Ag}$  are the gravitational energy-momentum tensor density, the gravitational energy-momentum tensor density without the Higgs potential and the gravitational potential density of the Higgs fields in the A - breaking, respectively.

From (2.1) the energy-momentum tensor density which does not contain the energymomentum tensor of gravitational and repulsive interactions can be defined as

$$T_{A\mu\nu} = \frac{2}{\sqrt{-g}} \left( \frac{\partial}{\partial \alpha} + \frac{\partial}{\partial \beta} \right) \\ \cdot \left[ \frac{\partial \left( \sqrt{-g} \mathcal{L}_A \right)}{\partial g^{\mu\nu}} - \left( \frac{\partial \left( \sqrt{-g} \mathcal{L}_A \right)}{\partial g^{\mu\nu}, \sigma} \right), \sigma \right] \\ \equiv T_{As\mu\nu} + T_{Av\mu\nu} - g_{\mu\nu} \left( V_{sv} + V_0 \right) \\ = T_{AM\mu\nu} - g_{\mu\nu} V_A \equiv T_{\mu\nu}, \qquad (2.20)$$

$$T_{AM\mu\nu} = T_{sM\mu\nu} + T_{vM\mu\nu} \equiv T_{M\mu\nu},$$
  
$$V_A = V_s + V_v + V_{sv} + V_0 \equiv V.$$
 (2.21)

Both  $\alpha$  and  $\beta$  in (2.3) – (2.4) may be regarded as the gravitation charges (the gravitation charges of  $T_{s\mu\nu}$ ,  $T_{v\mu\nu}$  and  $g_{\mu\nu}V_{s\nu}$  are regarded as 1, -1 and 0 in the *S* – *breaking*, respectively.). The energy-momentum tensor should be independent of the gravitation charges. Hence it is necessary to eliminate  $\alpha$  and  $\beta$  from the definition of  $T_{\mu\nu}$  by the operator  $(\partial/\alpha + \partial/\beta)$  which is the only difference between the definition of  $T_{\mu\nu}$  in the present model and that in the conventional theory. This definition does not contradict any basic principle and it is completely consistent with the conventional theory (the conventional theory corresponds to one matter type so that  $\beta = 0$ ). Both the s - energy and the v - energy must be positive because of the definition (2.20) – (2.21).

It should be pointed out that only (2.16) and (2.17) is applicable in the S – breaking, and only (2.16) and (2.18) applicable in the V – breaking.

It is proved that the necessary and sufficient condition of  $T^{\mu\nu}_{;\nu} = 0$  is  $I_M$  to be a scalar quantity<sup>[12]</sup>.  $I_S$  and  $I_V$  are all scalar quantities, and  $T^{\mu\nu}_{;\nu}$  in the conventional theory corresponds to  $T^{\mu\nu}_{g;\nu}$ . Hence we have

$$T_{Sg;\nu}^{\mu\nu} = T_{Vg;\nu}^{\mu\nu} = 0.$$
 (2.22)

Of course, (2.22) can been derived from (2.16).

#### III. SPONTANEOUS SYMMETRY BREAKING

Ignoring the couplings of  $\Phi_s$  and  $\chi_s$  and suitably choosing the parameters of the Higgs potential, analogously to Ref. [11], we can prove from (2.8) - (2.10) that there are the following vacuum expectation values at the zero-temperature and under the tree-level approximation,

$$\langle 0 | \omega_v | 0 \rangle \equiv \overline{\omega}_{v0} = 0, \quad \langle 0 | \omega_s | 0 \rangle \equiv \overline{\omega}_{s0} \neq 0,$$
 (3.1)

$$\langle 0 \left| \Omega_s \right| 0 \rangle = \upsilon_{\Omega 0}, \tag{3.2}$$

$$\langle 0 | \Phi_s | 0 \rangle = Diagonal\left(1, 1, 1, -\frac{3}{2}, -\frac{3}{2}\right) \upsilon_{\varphi 0}, \qquad (3.3)$$

$$\langle 0 | \chi_s | 0 \rangle^+ = \frac{v_{\chi 0}}{\sqrt{2}} (0, 0, 0, 0, 1),$$
 (3.4)

The breaking satisfied (3.1) is called the S-breaking. Ignoring the contributions of  $\Phi_s$  and  $\chi_s$  to  $\langle 0 | \Omega_s | 0 \rangle$ , at the zero-temperature we get

$$v_{\Omega 0}^2 = \frac{\mu^2}{f}, \quad f \equiv \lambda - \frac{15w^2}{(15a+7b)} - \frac{\varsigma^2}{\xi}.$$
 (3.5)

$$v_{\varphi 0}^2 = \frac{2w}{15a + 7b} v_{\Omega 0}^2, \tag{3.6}$$

$$v_{\chi 0}^2 = \frac{2\zeta}{\xi} v_{\Omega 0}^2. \tag{3.7}$$

We take  $\Lambda > \lambda > 15w^2/(15a + 7b) + \zeta^2/\xi$ . From (2.9)-(2.10) and (3.1)-(3.7) it can be proved that all v - Higgs bosons can get their big enough masses. The masses of the Higgs particles exclusive of the  $\Phi_s - particles$  and the  $\chi_s - particles$  in the S - breaking are respectively

$$m^2\left(\Omega_s\right) = 2\mu^2,\tag{3.8}$$

$$m^2\left(\Omega_v\right) = \Lambda v_{\Omega 0}^2 - \mu^2,\tag{3.9}$$

$$m^{2}(\Phi_{v}) = \frac{1}{2}\alpha v_{\Omega 0}^{2}, \qquad (3.10)$$

$$m^2\left(\chi_v\right) = \beta v_{\Omega 0}^2. \tag{3.11}$$

We can choose such parameters that

$$m(\Omega_s) \simeq m(\Omega_v)$$
  
$$\gg m(\varphi_v) \sim m(\varphi_s) \gg m(\chi_v) \sim m(\chi_s), \qquad (3.12)$$

e.g.,  $m(\Omega_s) \sim 10^{16} Gev$ ,  $m(\varphi_s) \sim 10^{14} Gev$  and  $m(\chi_s) \sim 10^2 Gev$ . It is easily seen from (3.8) - (3.11) that all real components of  $\Phi_v$  have the same mass  $m(\Phi_v)$ , and all real components of  $\chi_v$  have the same mass  $m(\chi_v)$  in the S – breaking.

The *S*-breaking and the *V*-breaking are symmetric because *s*-matter and *v*-matter are symmetric. Hence when  $s \rightleftharpoons v$  and  $S \rightleftharpoons V$  in (3.1) – (3.12), the formulas are still kept.

#### IV. EVOLUTION EQUATIONS OF SPACE IN RW METRIC

Based on the RW metric metric,

$$(ds)^{2} = -(dt)^{2} + R^{2}(t) \left\{ \frac{(dr)^{2}}{1 - kr^{2}} + (rd\theta)^{2} + (r\sin\theta d\varphi)^{2} \right\}.$$
(4.1)

In the present model, we take k = -1.

Matter in the universe may approximately be regarded as ideal gas distributed evenly in space. Considering the potential energy densities in (2.14), we can write  $T_{a\mu\nu}$  as

$$T_{a\mu\nu} = \left[\widetilde{\rho}_a + \widetilde{p}_a\right] U_{a\mu} U_{a\nu} + \widetilde{p}_a g_{\mu\nu}, \qquad (4.2)$$

$$\widetilde{\rho}_a = \rho_a + V_a\left(\varpi_a\right), \quad \widetilde{p}_a = p_a - V_a(\varpi_a), \tag{4.3}$$

where  $U_{a\mu}$  is a 4-velocity,  $U_{a\mu} = \delta^0_{\mu} = U_{\mu}$ , and a = s or v.  $(-g_{\mu\nu}V_0)$  can be written as

$$-g_{\mu\nu}V_{0} = (\tilde{\rho}(V_{0}) + \tilde{p}(V_{0}))U_{\mu}U_{\nu} + g_{\mu\nu}\tilde{p}(V_{0}), \qquad (4.4)$$
$$\tilde{\rho}(V_{0}) = V_{0}, \quad \tilde{p}(V_{0}) = -V_{0}.$$

Considering  $U_{\mu} = \delta^{0}_{\mu}$ , substituting (4.2) - (4.4) and the RW metric in (4.1) into (2.16), we get the evolution equations

$$\stackrel{\cdot}{R}^{2} + k = \eta \left[ \rho_{g} + V_{g} \right] R^{2}, \quad \eta \equiv 8\pi G/3, \tag{4.5}$$

$$\ddot{R} = -\frac{1}{2}\eta \left[ (\rho_g + 3p_g) - 2V_g \right] R.$$
(4.6)

In the S-breaking,

$$\rho_g = \rho_s - \rho_v, \quad p_g = p_s - p_v, \quad V_g = V_s + V_0 - V_v. \tag{4.7}$$

In the V-breaking,

$$\rho_g = \rho_v - \rho_s, \quad p_g = p_v - p_s, \quad V_g = V_v + V_0 - V_s. \tag{4.8}$$

Comparing (4.5) - (4.6) with the Friedmann equations, we see that provided  $\rho$ , p and V in the Friedmann equations are replaced by  $\rho_g$ ,  $p_g$  and  $V_g$ , (4.5) - (4.6) are obtained.

In contrast with the conventional theory,  $\rho_g > 0$ , = 0 and  $\rho_g < 0$  are all possible in the present model.

From (4.5) - (4.6) we have

$$\dot{\rho}_g + 3\left(\rho_g + p_g\right)\frac{R}{R} = -\dot{V}_g.$$
(4.9)

Pressure density is a function of masses of particles and temperature. In order to determine the pressure at a given temperature, we divide the particles into three sorts according to their masses. The first sort is composed of such particles whose masses are larger than  $m_M \gtrsim 1 MeV$ . The second sort is composed of such particles with their masses  $m_l$ , and  $1 MeV \gtrsim m_l \gtrsim 1 eV$ . We suppose  $p_l = \kappa \rho_l$ . It is obvious  $1/3 \gtrsim \kappa \gtrsim 0$ . The third sort is composed of photon-like particles whose masses are less than  $m_l$  and may be regarded as zero. Thus, we have

$$\rho_g = \rho_{Mg} + \rho_{\lg} + \rho_{\gamma g}, \tag{4.10}$$

and  $p_{\gamma g} = \rho_{\gamma g}/3$ . In the V – breaking,  $\rho_{Mg} = \rho_{vMg} - \rho_{sMg}$ ,  $\rho_{lg} = \rho_{v \, lg} - \rho_{s \, lg}$ . Considering all s – particles must be in s - SU(5) color single states whose masses are not zero so that  $p_{s\gamma} = 0$ , we have  $\rho_{\gamma g} = \rho_{v\gamma g} - \rho_{s\gamma g} = \rho_{v\gamma g}$ . Thus, when T is so large that all masses may be neglected, from (4.9) – (4.10) we have

$$p_g = \rho_g/3, \ \frac{d(\rho_g R^4)}{dt} = -\dot{V}_g R^4.$$
 (4.11)

When  $m_M \gtrsim T \gtrsim m_l$ ,  $\rho_M \gg p_M \sim 0$ . Letting  $p_l = \kappa \rho_l$ , we have

$$\frac{d\left(\rho_{Mg}R^{3}\right)}{dt} + \frac{d\left(\rho_{\lg}R^{3(1+\kappa)}\right)}{R^{3\kappa}dt} + \frac{d\left(\rho_{\gamma g}R^{4}\right)}{Rdt} = -\dot{V}_{g}R^{3}.$$
(4.12)

When  $T < m_l \sim 1 eV$ ,  $\rho_M \gg p_M \sim 0$  and  $\rho_l \gg p_l \sim 0$  so that

$$\frac{d\left(\rho_{mg}R^{3}\right)}{dt} + \frac{d\left(\rho_{\gamma g}R^{4}\right)}{Rdt} = -\dot{V}_{g}R^{3}.$$
(4.13)

where  $\rho_{mg} = \rho_{Mg} + \rho_{lg}$ .

#### V. TEMPERATURE EFFECT

The thermal equilibrium between the v - particles and the s - particles can be realized by only (2.10). The Higgs bosons  $\Omega_s$  and  $\Omega_v$  are hardly produced because their masses are all very big after reheating. Consequently, the interaction between the v - particlesand the s - particles may be ignored so that there is no thermal equilibrium between the v - particles and the s - particles. Thus, when temperature is low, we should use two sorts of temperature  $T_v$  and  $T_s$  to describe the thermal equilibrium of v - matter and the thermal equilibrium of s - matter, respectively. Generally speaking,  $T_v \neq T_s$ . We will see when  $T_s \gtrsim T_{cr}$ ,  $\langle \Omega_s \rangle = \langle \Omega_v \rangle = 0$  so that  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  because of (2.8) – (2.10). Thus, the masses of all particles are zero and  $\rho_s$  and  $\rho_v$  can transform from one to an other by (2.10). In this case, if  $\dot{R} = \ddot{R} = 0$ , there must be  $T_v = T_s$  and  $\rho_v = \rho_s$ ; If  $\dot{R} < 0$  and  $\ddot{R} > 0$ , there must be the least scale  $R_{\min} > 0$ . When space contracts to  $R_{\min}$ ,  $T_s = T_{\max}$  and then inflation occurs. There is no such a case  $\dot{R} < 0$  and  $\ddot{R} \leq 0$  in the present model.

#### A. Effective potentials

The influence of finite temperature on the Higgs potential in the present model are consistent with the conventional theory. For short, we consider only  $\Omega_a$  and  $\varphi_a$ , a = s or v. When  $\chi_a$  is considered as well, the following inferences are still qualitatively valid. For

$$V_{s\Omega} = -\frac{\mu^2}{2}\Omega_s^2 + \frac{\lambda}{4}\Omega_s^4,\tag{5.1}$$

to ignore the terms proportional to  $\lambda^n (n > 1)$ , the finite-temperature effective potential approximate to 1-loop in flat space is<sup>[13][14]</sup>

$$V_{s\Omega,eff}^{(1)T} = -\frac{1}{2} \left( \mu^2 - \frac{\lambda}{4} T_s^2 \right) \overline{\Omega}_s^2 + \frac{\lambda}{4} \overline{\Omega}_s^4 - \frac{\pi^2}{90} \left( kT_s \right)^4 + \frac{\mu^2}{24} T_s^2.$$
(5.2)

Considering the influence of the expectation values  $v_{\Omega v}(T_s, T_v)$ ,  $v_{\varphi s}(T_s, T_v)$  and  $v_{\varphi v}(T_s, T_v)$ , and ignoring the terms irrelevant to  $\Omega_s$ , we have

$$V_{s\Omega,eff}^{(1)T} = -\frac{1}{2}\mu_s^2 \left(T_s, T_v\right)\overline{\Omega}_s^2 + \frac{\lambda}{4}\overline{\Omega}_s^4,\tag{5.3}$$

$$\mu_s^2 \equiv \mu^2 - \frac{\lambda}{4} T_s^2 - \Lambda v_{\Omega v}^2 - \frac{15}{2} \left( \alpha v_{\varphi v}^2 - w v_{\varphi s}^2 \right).$$
(5.4)

Similarly (5.1) - (5.4), we have

$$V_{v\Omega,eff}^{(1)T} = -\frac{1}{2}\mu_v^2 \left(T_s, T_v\right)\overline{\Omega}_v^2 + \frac{\lambda}{4}\overline{\Omega}_v^4,\tag{5.5}$$

$$\mu_v^2 \equiv \mu^2 - \frac{\lambda}{4} T_v^2 - \Lambda v_{\Omega s}^2 - \frac{15}{2} \left[ \alpha v_{\varphi s}^2 - w v_{\varphi v}^2 \right].$$
(5.6)

In the *S*-breaking, we can prove  $v_{\Omega v} (T_s, T_v) = v_{\varphi v} (T_s, T_v) = 0$ . This is because  $v_{\Omega v} = v_{\varphi v} = 0$  in low temperatures, and  $T_v \sim T_s$  holds when  $T_s$  is high enough so that  $v_{\Omega v} = v_{\varphi v} = 0$  still holds. Thus

$$\mu_s^2(T_s, T_v) = \mu^2 - \frac{\lambda}{4}T_s^2 + \frac{15}{2}wv_{\varphi s}^2(T_s, T_v) .$$
$$v_{\Omega s}^2(T_s, T_v) = \mu_s^2(T_s, T_v) / \lambda, \text{ when } \mu_s^2(T_s, T_v) > 0$$
$$v_{\Omega s}(T_s, T_v) = 0, \text{ when } \mu_s^2(T_s, T_v) \le 0.$$

For

$$V_{s\varphi} = \frac{1}{2} \left( \alpha \Omega_v^2 - w \Omega_s^2 \right) Tr \Phi_s^2 + \frac{1}{4} a (Tr \Phi_s^2)^2 + \frac{1}{2} b Tr \Phi_s^4, \tag{5.7}$$

ignoring the contributions of the Higgs fields and the fermion fields to one loop correction, and only considering the contribution of the gauge fields, when  $\overline{\varphi}_s \ll kT$ , here k is the Boltzmann constant, we get the effective potential at finite-temperature approximate to 1-loop in flat space<sup>[13][14]</sup>

$$V_{s\varphi,eff}^{(1)T} = V(\overline{\varphi}_s) + B\overline{\varphi}_s^4 \left( \ln \frac{\overline{\varphi}_s^2}{\sigma^2} - \frac{25}{6} \right) + CT_s^2 \overline{\varphi}_s^2 - \frac{\pi^2}{15} \left( kT_s \right)^4, \tag{5.8}$$

where  $B = (5625/1024\pi^2) g^4$ , and  $\overline{\Phi}_s = Diagonal (1, 1, 1, -3/2, -3/2) \overline{\varphi}_s$ . In general,  $w = \alpha < \lambda \sim g^4 < C = (75/16) (kg)^2$ . We take  $w = \alpha$  for simplicity. Here  $\sigma$  is a parameter at which the renormalization coupling-constant  $\lambda$  is defined as below,

$$\frac{d^4 V_{s\varphi,eff}^{(1)T}}{d\overline{\varphi}_s} \mid_{\overline{\varphi}_s=\sigma} = \lambda.$$
(5.9)

Only considering the contribution of the expectation values of  $\Omega_s$  and  $\Omega_v$  to  $V_{s\varphi,eff}^{(1)T}$ , taking (15/16)(15a+7b) = (11/3)B, and ignoring the term  $(kT_s)^4$  unconnected with  $\overline{\varphi}_v$  (it may be added to in  $\rho_{rg}$ ), from (2.8) and (5.7) – (5.8) we have

$$V_{s\varphi,eff}^{(1)T} = A_s^2 \overline{\varphi}_s^2 + B \overline{\varphi}_s^4 \left( \ln \frac{\overline{\varphi}_s^2}{\sigma^2} - \frac{1}{2} \right), \qquad (5.10)$$

$$A_{s}^{2} \equiv \frac{15}{4} w \left( v_{\Omega v}^{2} \left( T_{v} \right) - v_{\Omega s}^{2} \left( T_{s} \right) \right) + C T_{s}^{2}.$$
(5.11)

Similarly, from (2.9) we have

$$V_{v\varphi,eff}^{(1)T} = A_v^2 \overline{\varphi}_v^2 + B \overline{\varphi}_v^4 \left( \ln \frac{\overline{\varphi}_v^2}{\sigma^2} - \frac{1}{2} \right), \qquad (5.12)$$

$$A_{v}^{2} \equiv \frac{15}{4} w \left( v_{\Omega s}^{2} \left( T_{s} \right) - v_{\Omega v}^{2} \left( T_{v} \right) \right) + C T_{v}^{2}.$$
(5.13)

#### **B.** Critical temperatures $T_{cr}$

In the following  $V_{eff} = V_{eff}^{(1)T}$ .  $T_s$  and  $T_v$  must rise as space contracts. From (5.4), (5.6), (5.11) and (5.13) we see that there must be critical temperatures  $T_{s,\varphi cr}$  and  $T_{cr}$  so that  $\langle \varphi_s \rangle = \langle \varphi_v \rangle = 0$  when  $T_s > T_{s,\varphi cr}$  and  $\langle \Omega_s \rangle = \langle \Omega_v \rangle = 0$  when  $T_s > T_{cr}$ . By suitably choosing the parameters  $\mu^2$ ,  $\Lambda$ ,  $\lambda$ , C and w, we can get  $T_{cr} > T_{s,\varphi cr}$ . Consequently,  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  when  $T_s > T_{cr}$ .

We discuss  $T_{s,\varphi cr}$  and  $T_{cr}$  in detains as follows.  $V_{eff} = V_{eff}^{(1)T}$  in the following.

1. The critical temperature  $T_{s,\varphi cr}$  in the S – breaking.

Let  $T_v = T_{v,\varphi cr}$  when  $T_s = T_{s,\varphi cr}$  in the S-breaking.  $T_{s,\varphi cr}$  is the critical temperature at which the minima are degenerate, i.e.  $V_{eff\min}(\overline{\varphi}_s, T_{s,\varphi cr}, T_{v,\varphi cr}) = V_{eff}(v_{s,\varphi cr}, T_{s,\varphi cr}, T_{v,\varphi cr}) =$  $V_{eff}(0, T_{s,\varphi cr}, T_{v,\varphi cr})$ . In other words,  $\langle \varphi_s \rangle = v_{\varphi s} \neq 0$  when  $T_s < T_{s,\varphi cr}$ , and  $v_{\varphi s} = 0$  when  $T_s > T_{s,\varphi cr}$ .  $T_{s,\varphi cr}$  and  $A_v^2(\overline{\varphi}_s, T_{s,\varphi cr}, T_{v,\varphi cr})$  can be determined from (5.12) – (5.13) by that when  $\overline{\varphi}_s = v_{\varphi scr}$ ,

$$V_{eff}\left(\overline{\varphi}_{s}, T_{s,\varphi cr}, T_{v,\varphi cr}\right) = V_{eff}\left(0, T_{s,\varphi cr}, T_{v,\varphi cr}\right), \quad \frac{\partial}{\partial \overline{\varphi}_{s}} V_{eff}\left(\overline{\varphi}_{s}, T_{s,\varphi cr}, T_{v,\varphi cr}\right) = 0.$$

When  $A_{\varphi}^2 = A_{\varphi}^2(\overline{\varphi}_s, T_{s,\varphi cr}, T_{v,\varphi cr}) \equiv A_{cr}^2 = B\sigma^2 e^{-1/2}$ ,  $V_{s,eff\min}$  is degenerate, here  $\overline{\varphi}_{scr}^2 \equiv \sigma^2 e^{-1/2}$ , hence we have

$$T_{s,\varphi cr}^{2} = \frac{A_{cr}^{2} + (15/4) \left(w\mu^{2}/\lambda\right)}{C + 15w/16},$$
(5.14)

Here  $v_{\Omega v} = v_{\varphi s} = v_{\varphi v} = 0$  when  $T_s > T_{s\varphi cr}$  is considered. When  $T_s < T_{s\varphi cr}$ , i.e.,  $A_s^2 < A_{cr}^2$ ,  $V_{s\varphi,eff\min} < V_{s\varphi,eff}(0)$  and  $v_{\varphi s} \neq 0$ . When  $T_s > T_{s\varphi cr}$ , i.e.,  $A_s^2 > A_{cr}^2$ ,  $v_{\varphi s} = 0$  and  $V_{s\varphi,eff\min} = V_{s\varphi,eff}(0)$ . Let  $\mu_a^2 = \mu_{a\varphi cr}^2$  and  $A_v^2 = A_{v\varphi cr}^2$  when  $T_s = T_{s\varphi cr}$ , then when  $T_s \ge T_{s\varphi cr}$ , from (5.6) and (5.13) we have

$$\mu_s^2 \equiv \mu^2 - \frac{\lambda}{4} T_s^2, \tag{5.15}$$

$$\mu_{s\varphi cr}^2 = \frac{\mu^2 - (\lambda/4C) A_{cr}^2}{1 + 15w/16C},\tag{5.16}$$

$$\mu_v^2 = \left(1 - \frac{\Lambda}{\lambda}\right)\mu^2 - \frac{\lambda}{4}\left(T_v^2 - \frac{\Lambda}{\lambda}T_s^2\right),\tag{5.17}$$

$$\mu_{v\varphi cr}^2 = \left(1 - \frac{\Lambda}{\lambda}\right)\mu^2 - \frac{\lambda}{4}\left(T_{v\varphi cr}^2 - \frac{\Lambda}{\lambda}T_{s\varphi cr}^2\right),\tag{5.18}$$

$$A_v^2 = \frac{15}{4} \frac{w}{\lambda} \mu^2 + CT_v^2 - \frac{15}{16} wT_s^2, \qquad (5.19)$$

 $\lambda \left( -2 \Lambda -2 \right)$ 

$$A_{v\varphi cr}^{2} = A_{v\varphi}^{2} \left( T_{s\varphi cr} \right) = \frac{\left( 15w/4\lambda \right) \mu^{2} - \left( 15w/16C \right) A_{cr}^{2}}{1 + 15w/16C} + CT_{v\varphi cr}^{2}$$
(5.20)

It is obvious that  $v_{a\Omega}(T_s, T_v) = 0$  when  $\mu_a^2 \leq 0$ . Considering  $\Lambda \gg \lambda$ ,  $C > \lambda > w$ , taking  $A_{cr}^2 < \mu^2$ , from (5.17) – (5.20) and (5.14) we see that provided  $T_v \sim T_s$ , there are

$$\mu_v^2 < 0, \quad A_{v\varphi cr}^2 > A_{cr}^2,$$
(5.21)

$$v_{v\varphi} = v_{v\Omega} = 0, \quad \text{when } T_s \ge T_{s\varphi cr},$$

$$(5.22)$$

$$T_{s\varphi cr} < \frac{2\mu}{\sqrt{\lambda}}.\tag{5.23}$$

In fact,  $T_v \gtrsim T_s$  when  $T_s > T_{s\varphi cr}$ . Hence (5.21) – (5.23) must hold when  $T_s > T_{s\varphi cr}$ . The masses of particles originating from the couplings of particles with  $\langle \omega_s \rangle$  and  $\langle \omega_v \rangle$  will be small, because  $v_{s\varphi} = v_{v\varphi} = v_{v\Omega} = 0$  and  $v_{s\Omega}$  becomes small when  $T_s > T_{s\varphi cr}$ . Thus, photon-like particles are dominative and the transformation of  $\rho_s$  to  $\rho_v$  is striking due to  $V_{g,eff} > 0$  (see section 6.1 in detail). Consequently,

$$T_v \gtrsim T_s, \ p_g \sim \rho_g/3, \ \text{ when } T_s > T_{s\varphi cr}.$$
 (5.24)

On the other hand, the transformation of  $\rho_s$  and  $\rho_v$  from one to another may be neglected in low temperatures. Thus, it is easily proven that  $v_{v\varphi} = v_{v\Omega} = 0$  when  $T_s < T_{s\varphi cr}$  in the S - breaking. Hence,  $v_{\Omega v} = v_{\varphi v} = 0$  holds always in the S - breaking.

2. Critical temperatures  $T_{cr}$ 

(5.15) holds when  $T_s > T_{s\varphi cr}$  due to  $v_{\varphi s} = 0$  and (5.22). Thus, from (5.3), (5.17), (5.19), (5.5), (5.10) and (5.12) we have

$$v_{s\Omega} = 0 = v_{s\varphi} = v_{v\varphi} = v_{v\Omega}, \tag{5.25}$$

$$m_s = m_v = 0$$
, when  $T_s \ge T_{cr} \equiv \frac{2\mu}{\sqrt{\lambda}}$ , (5.26)

where  $m_s$  and  $m_v$  denote the masses of all s-particles and the masses of all v-particles originating from the couplings of particles with  $v_{s\Omega}$ ,  $v_{v\Omega}$ ,  $v_{\varphi s}$  and  $v_{\varphi v}$ , respectively.

The symmetry  $s - SU(5) \times v - SU(5)$  holds because of (5.25). The state with this symmetry is called the most symmetric state.  $\rho_s$  and  $\rho_v$  can transform from one to another because (5.26) and (2.10). Consequently, if space is static and  $T_s \geq T_{cr}$ , we have

$$T_s = T_v = T, \ \rho_g(T) = 0, \ p_g = 0$$
 (5.27)

$$\rho_s = \rho_v, \quad p_s = \rho_s/3, \quad p_v = \rho_v/3,$$
(5.28)

$$V_{v,eff} = V_{s,eff} = 0, \ V_{g,eff} = V_0.$$
(5.29)

Here  $V_{a,eff} = V_{a,eff} \left(\overline{\varphi}_a, \overline{\Omega}_a\right)$  which does not contain the terms containing  $(kT_a)^4$  and  $\mu^2 T_a^2$ . The terms are added to  $\rho_a$  (see (6.6)). In fact, space cannot be static, and the contracting process is not a thermal equilibrium process because  $\ddot{R} > 0$  and  $\ddot{R} < 0$ . Hence (5.27) – (5.28) does not hold in the contracting process, but (5.29) possibly holds because  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$ .

### VI. CONTRACTION OF SPACE, THE HIGHEST TEMPERATURE, INFLATION OF SPACE AND THERE IS NO SINGULARITY

On the basis of the cosmological principle, if there is the space-time singularity, it is a result of space contraction. Thus, we discuss the contracting process. From the contracting process we will see that there is no space-time singularity in present model.

We do not consider the couplings of the Higgs fields with the Ricci scalar R for this time. We will see in the following paper that the following conclusions still hold when such couplings as  $\xi R\Omega_s^2$  are considered. In fact,  $\xi R\left(\overline{\Omega}_s^2 - \overline{\Omega}_v^2\right) = 0$  because there is the strict symmetry between s - matter and v - matter when  $T \gtrsim T_{cr}$ .

We chiefly discuss change of  $\langle \Omega_a(T_a) \rangle$  and  $\langle \varphi_a(T_a) \rangle$  as temperature in the contracting process of space for short. When  $\langle \chi_a(T_a) \rangle$  is considered as well, the inferences are still valid qualitatively.

#### A. The contracting process of space and the highest temperature

#### 1. Space contracts in the S - breakng.

In the S - breaking, we consider the space-contraction process in which R < 0. In low temperatures  $T_s \sim T_v \gtrsim 0$ ,  $\langle \omega_s \rangle = \langle \omega_s \rangle_0$  and  $\langle \omega_v \rangle = 0$ . Consequently  $V_{s,eff} = -V_0$ ,  $V_{v,eff} = 0$ 

$$V_{g,eff} = V_{s,eff} - V_{v,eff} + V_0 = 0, (6.1)$$

and  $\rho_v$  and  $\rho_s$  cannot transform from one into other and  $V_{g,eff} \sim 0$ , because the masses of the Higgs particles  $\Omega_s$  and  $\Omega_v$  are very large. There must be  $\rho_g = \rho_s - \rho_v > 0$  because of (6.1), (4.5) - (4.7) and  $\dot{R} < 0$ . In the stage, space will contract faster and faster, i.e.  $\ddot{R} < 0$ due to (4.6).

2.  $V_{g,eff} > 0$  and  $\rho_q < 0$ .

The temperature must rise monotonously because space contracts, because the non-zero momentum of a free particle  $p \propto 1/R(t)$ ,  $\Delta p \Delta x \gtrsim 1$  and  $\rho_m \propto 1/R^3(t)$ . In the contracting process, as mentioned section 5.2,  $\langle \omega_v \rangle = 0$  holds always. Consequently,  $V_{v,eff} = V_{v,eff} = 0$  holds always. In contrast with  $V_{v,eff}$ ,  $V_{s,eff}$  will rise from  $V_{s,eff} = -V_0$  (when  $T_s \sim 0$ ) to  $V_{s,eff}(T_{cr}) = 0$ .  $\langle \varphi_s \rangle \sim 0$  when  $T_s \gtrsim T_{s\varphi cr}$  and  $\langle \Omega_s \rangle \sim 0$  when  $T_s \sim T_{cr}$ . It is seen that  $\langle \omega_s \rangle$  can strikingly change in the period  $T_{s\varphi cr} \sim T_s \lesssim T_{cr}$ . Hence  $V_{s,eff}$  must strikingly change and  $V_{s,eff} > 0$ . On the other hand, the transformation of  $\rho_s$  into  $\rho_v$  is striking in the period  $T_{s\varphi cr} \sim T_s \lesssim T_{cr}$ . This is because in the period, in addition  $\omega_v \rangle = 0$ ,  $\langle \varphi_s \rangle \sim 0$  or  $\langle \Omega_s \rangle \sim 0$  so that  $m_s \sim m_v \sim 0$ . Thus, from (4.9) we have

$$\dot{V}_{g,eff} = \dot{V}_{s,eff} - \dot{V}_{v,eff} = \dot{V}_{s,eff} > 0,$$
 (6.2)

$$\rho_g = \rho_s - \rho_v < 0. \tag{6.3}$$

When  $T_s = T_{cr}$ ,  $V_{s,eff} = V_{v,eff} = 0$ , because  $\langle \omega_v \rangle = \langle \omega_v \rangle = 0$ . Thus,  $V_{g,eff} = V_0$  and (4.5) - (4.7) is reduced to

$$\left(\frac{\dot{R}}{R}\right)^2 = -\frac{k}{R^2} + \eta \left[\rho_g + V_0\right],\tag{6.4}$$

$$\frac{R}{R} = \eta \left[ V_0 - \rho_g \right]. \tag{6.5}$$

$$\rho_a = \frac{\pi^2}{30} \left( g^* - \frac{7}{3} k^4 \right) T_a^4 + \frac{\mu^2}{24} T_a^2, \quad a = s, v,$$
(6.6)

$$\rho_g = \rho_s - \rho_v, \quad p_g = \rho_g/3. \tag{6.7}$$

Here (5.2), (5.8), (5.28) and  $m_s = m_v = 0$  are considered.  $g_a^* = g_{aB} + 7g_{aF}/8$ , a = s, v, is the total number of the spin states, and  $g_{aB}(g_{aF})$  is the total number of the spin states of a-bosons (a-fermions). Considering s-matter and v-matter are symmetric, we have

$$g_s^* = g_{sB} + 7g_{sF}/8 = g_v^* = g_{vB} + 7g_{vF}/8 \equiv g^*.$$

As a consequence of  $\rho_g < 0$ ,  $\rho_g(T_{cr}) < 0$  can be obtained.  $\rho_g(T_{cr}) < 0$  can be realized, in addition to (6.2) - (6.3), because R < 0 and R > 0 in the contracting process so that it is not a thermal equilibrium process.

3. To realize  $\rho_g(T_{cr}) < 0$ 

When  $T_{s\varphi cr} \sim T_s \lesssim T_{cr}$ ,  $v_{s\varphi} = v_{v\varphi} = v_{v\Omega} = 0$  and (5.4) and (5.6) are reduced to

$$\mu_s^2 \equiv \mu^2 - \frac{\lambda}{4} T_s^2, \quad m_{s\Omega}^2 = 2\mu_s^2 = 2\lambda v_{\Omega s}^2, \tag{6.8}$$

$$\mu_v^2 \equiv \mu^2 - \frac{\lambda}{4} T_v^2 - \Lambda v_{\Omega s}^2, \quad m_{v\Omega}^2 = -2\mu_v^2$$
(6.9)

It is seem from (6.8) – (6.9) when  $\mu_v \sim 0$ ,  $m_{s\Omega} \gtrsim 0$ . Hence

$$m_{s\Omega} \gtrsim m_{v\Omega}, \text{ when } m_{v\Omega} \sim 0,$$
 (6.10)

Thus,  $\rho_s$  can transform to  $\rho_v$ , i.e.  $\dot{\rho}_g < 0$ , when  $T_{s\varphi cr} \sim T_s \lesssim T_{cr}$  due to (6.10) and (2.10), i.e.

$$\Omega_s + \Omega_s \rightleftharpoons \Omega_v + \Omega_v, \quad \Omega_s + \Omega_s \rightleftharpoons \varphi_v + \varphi_v,$$
  
$$\varphi_s + \varphi_s \rightleftharpoons \Omega_v + \Omega_v. \tag{6.11}$$

The transformation of  $\rho_s$  to  $\rho_v$  is very fast because the coupling constant  $\Lambda$  is very large and  $m_{s\Omega}$  and  $m_{v\Omega}$  are very small in the case. Consequently,  $\rho_s = \rho_v$  can be realized when  $T_s < T_{cr}$  or  $t \equiv t_{eq} < t_{cr}$ . There must be  $\dot{\rho}_g(t_{eq}) < 0$ , because  $V_{g,eff}(t_{eq}) < 0$  when  $T_s(t_{eq}) < T_{cr}$ . Thus  $\dot{V}_{g,eff}(t_{eq}) > 0$  and  $\dot{\rho}_g(t_{eq}) = -\dot{V}_{g,eff}(t_{eq}) < 0$  because of (4.9) and  $\rho_g(t_{eq}) = 0$ . Consequently,

$$\rho_g(t_{cr}) = \rho_g(t_{eq}) + \dot{\rho}_g(t_{eq})(t_{cr} - t_{eq}) < 0.$$
(6.12)

#### 4. The boundary condition

Only when the boundary condition is correct, the solution of a partial differential equation can be correct. we consider the correct physics boundary condition of the equations (6.4) - (6.5) is

$$R = 0$$
 when  $R > 0.$  (6.13)

The condition

R < 0 when R = 0

is not correct, because it is impossible that R < 0 when R = 0.

The condition (6.12) requires  $\rho_g < 0$  in the contracting process. The condition (6.12) can be realized provided  $\rho_g(T_{cr}) < 0$  (see the following theorem), because  $\rho_g(t) \sim R^{-4}$  so that there must be  $-k/R^2 + \eta (-C_g/R^4 + V_0) = 0$  at  $R = R_{\min} > 0$ . Here  $C_g > 0$  is a constant.

If  $V_{g,eff} > 0$  (i.e.,  $\rho_g < 0$ ) when  $t \ge t_{cr}$ , there are such solutions satisfying  $\rho_g(t_{cr}) = 0$ (i.e.  $T_s = T_v$ ), (6.14) and (6.4) - (6.5). We do not discuss the case for a time.

#### B. A theorem related to singularity

**Theorem 3** Let  $V_g = 0$  and  $p_g = \kappa \rho_g$ ,  $1/3 \ge \kappa \ge 0$ , for a contracting process in which R < 0 at the initial time  $t_1$ , then in the case  $\rho_g < 0$ , all solutions of (4.5) - (4.6) satisfy the boundary condition  $R(t_2) = 0$  and  $R(t_2) > 0$ , here  $t_2 > t_1$ ; In the case k > 0 and  $\rho_g > 0$ , when  $\rho_g(t_2) + 3p_g(t_2) - 2V_g \le 0$ , the solutions of (4.5) - (4.6) satisfy the boundary condition; When k > 0,  $\rho_g = 0$  and  $V_g > 0$ , the solutions of (4.5) - (4.6) satisfy the boundary condition.

It is necessary due to  $V_g = 0$  and (4.5) that  $V_g = V_{g0} \ge 0$ . This is because if  $V_{g0} < 0$ , there will be  $\left(\frac{R}{R}\right)^2 \longrightarrow \eta V_{g0} < 0$  when  $R \longrightarrow \infty$ . This is impossible. Thus we discuss only the case  $V_{g0} \ge 0$ .

When  $k \ge 0$ ,  $V_g = 0$  and  $\rho_g < 0$ , there is no solution of (4.5) - (4.6). In the case  $\rho_g > 0$ , when  $k \le 0$  or k > 0 but  $\rho_g(t_2) + 3p_g(t_2) - 2V_g > 0$ , or in the case  $k \le 0$ ,  $\rho_g = 0$  and  $V_g > 0$ , the solutions of (4.5) - (4.6) do not satisfy the boundary condition.

**Proof.** If there is such a time  $t_2$  satisfying R/R = 0, there must be  $d\left(\frac{R}{R}\right)/dt > 0$  in the period  $t_1 < t \le t_2$ , because R < 0 at the initial time  $t_1$ . From (4.5) - (4.6) we have

$$\frac{d}{dt}\left(\frac{\dot{R}}{R}\right) = \frac{k}{R^2} - \frac{3\eta}{2}\left(\rho_g + p_g\right).$$
(6.14)

When  $R(t_2)/R(t_2) = 0$ , (6.14) is reduced to

$$\frac{d}{dt}\left(\frac{\dot{R}}{R}\right) = \frac{\ddot{R}}{R} = -\frac{\eta}{2}\left(\rho_g + 3p_g - 2V_g\right).$$
(6.15)

because of (4.5). In the case with  $k \leq 0$ , only when  $\rho_g < 0$ ,  $d\left(\frac{\dot{R}}{R}\right)/dt > 0$  is possible because of (6.14) and  $p_g = \kappa \rho_g$  ( $\kappa \geq 0$ ). On the other hand, when  $\rho_g < 0$ ,  $d\left(\frac{\dot{R}}{R}\right)/dt > 0$ must be satisfied due to (6.15), and the contracting process ( $\dot{R}(t_1) < 0$ ) can come into being due to (4.5) provided k < 0 or k = 0 but  $V_{g0} > 0$ .

In the case with  $k \leq 0$  and  $\rho_g > 0$  or  $V_g > 0$ , there are the solutions of (4.5) – (4.6) and the contracting process. But the solutions cannot satisfy the boundary condition because (6.14) cannot be satisfied.

In the case with k > 0 and  $\rho_g < 0$ , It is obvious that (6.14) must be satisfied. But there

is no solution of (4.5) - (4.6) when  $V_{g0} = 0$  in the case. When  $V_g > 0$ , there are the solutions of (4.5) - (4.6), and the solutions satisfy the boundary condition.

In the case with k > 0 and  $\rho_g > 0$  or  $V_g > 0$ , there are the solutions of (4.5) - (4.6)and the contracting process. But only when there is such a time  $t_2$  satisfying  $-k/R(t_2) + \eta(\rho_g(t_2) + V_{g0}) = 0$  and  $\rho_g(t_2) + 3p_g(t_2) - 2V_{g0} \le 0$ , the solutions satisfy the boundary condition due (4.5) and (6.15). Otherwise, the solutions cannot satisfy the boundary condition, e.g. it is obvious due to (6.15) that the boundary condition cannot be satisfied when k > 0,  $\rho_g > 0$  and  $V_{g0} = 0$ .

In the discussion above, (4.12) is considered. Without loss of generality, for simplicity, from (4.12) we write  $\rho_g$  as

$$\rho_g = \frac{C_g}{R^{3(1+\kappa)}}, \quad 0 \le \kappa \le 1/3$$

It is possible that  $C_g > 0$ , = 0 or  $C_g < 0$ . Thus,  $|\rho_g|$  rises monotonously and faster as R decrease than  $|k/R^2|$ . Proof ends.

The boundary condition (13) is just the condition without singularity. From the discussion above, we see  $\rho_g < 0$  to be necessary in order to eliminate the singularity.

 $R_{\min} > 0$  so that there is no singularity of space-time

Space contracts with a deceleration R > 0, because  $\rho_g(t_{cr}) < 0$  and (6.5). As mentioned before,  $V_{g,eff} = V_0$  and  $\dot{V}_{g,eff} = 0$  due  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  when  $t > t_{cr}$ . Thus  $\rho_g(t) = -C_g/R^4$ . Substituting  $\rho_g(t) = -C_g/R^4$  into (6.4), taking k = -1, we see that there must be such a  $R_{\min}$  satisfying

$$\left(\frac{\dot{R}_{\min}}{R_{\min}}\right)^2 = \frac{1}{R_{\min}^2} + \eta \left[-\frac{C_g}{R_{\min}^4} + V_0\right] = 0, \qquad (6.16)$$

$$\frac{\ddot{R}_{\min}}{R_{\min}} = \eta \left[ V_0 + \frac{C_g}{R_{\min}^4} \right] > 0, \ R_{\min} > 0.$$
(6.17)

It is seen that there is the least scale  $R_{\min}$  in the present model.

Let  $R = R_{\min}$  when  $t = t_{FI}$ , then  $t_{FI}$  is the final time of the s - world and the initial time of the v - world. It is obvious that  $T_v(t_{FI})$  is the highest temperature  $T_{v\max}$ ,  $T_s(t_{FI}) = T_{s\max} < T_{v\max}$ ,  $\rho_s(t_{FI}) = \rho_{s\max}$ ,  $\rho_v(t_{FI}) = \rho_{v\max} > \rho_{s\max}$ ,  $\rho_s(t_{FI}) + \rho_v(t_{FI}) = \rho_{\max}$  is the largest energy density  $\rho_{\max}$ . All the physics quantities  $R_{\min}$ ,  $\rho_{s\max}$ ,  $\rho_{v\max}$ ,  $T_{s\max}$  and  $T_{v\max}$  must be finite. Because of the cosmological principle, all  $\rho_s$ ,  $\rho_v$ ,  $\rho_g$ ,  $V_s$ ,  $V_v$ ,  $V_g$  and  $p_g \leq \rho_g/3$  are finite when  $t < t_{FI}$ . Consequently  $T_{Ss\mu\nu}$ ,  $T_{Sv\mu\nu}$  and  $T_{Sg\mu\nu}$  must be finite. On the other hand, because of the cosmological principle, it is obvious that when space is not in contracting process or does not contract to  $R_{\min}$ , the physical quantities must be finite as well. Substituting the finite  $T_{Sg\mu\nu}$  or  $T_{Vg\mu\nu}$  into the Einstein field equation, we see that  $R_{\mu\nu}$  and  $g_{\mu\nu}$  must be finite.

Consequently, there is no singularity of space-time in the present model.

#### C. Expansion and inflation of space

It is seen from (6.16) – (6.17) that space will expands with an acceleration, because  $\ddot{R}_{\min} = 0$  and  $\ddot{R}_{\min} > 0$ . The expanding process is different from the contracting process in essence. (6.14) is no longer the boundary condition for the expanding process. (6.16) – (6.17) or (6.14) is the initial condition. In the initial stage,  $\langle \omega_v \rangle = \langle \omega_v \rangle = m_s = m_v = 0$  so that  $\rho_s$  and  $\rho_v$  can transform from one to another. Hence there must be such a time  $\tilde{t}_{cr}$  at which

$$\rho_g = \rho_v - \rho_s = 0, \quad T_s = T_v,$$

$$V_{g,eff} = V_{v,eff} + V_0 - V_{s,eff} = V_0.$$
(6.18)

The transformation of  $\rho_v$  and  $\rho_s$  from one to another is very fast because the coupling constant  $\Lambda$  is very large and  $m_s = m_v = 0$ . The universe is in the most symmetric state when  $t \sim \tilde{t}_{cr}$ . Thus, (4.5) and (4.6) are reduced to

$$R^{2} = 1 + \eta V_0 R^2 = 0, (6.19)$$

$$R = \eta V_0 R > 0. \tag{6.20}$$

From (6.19) – (6.20) we see that space inflation must occur when  $t > \tilde{t}_{cr}$ .

$$R(t) = \sqrt{\frac{1}{\eta V_0}} \sinh H \left( t - \tilde{t}_{cr} + \tau \right), \quad H \equiv \sqrt{\eta V_0}, \tag{6.21}$$

$$\sim \frac{1}{2} \exp H\left(t - \widetilde{t}_{cr} + \tau\right), \text{ when } H\left(t - \widetilde{t}_{cr} + \tau\right) \gg 1, \tag{6.22}$$

$$R\left(\tilde{t}_{cr}\right) = \sqrt{\frac{1}{\eta V_0}} \sinh H\tau.$$
(6.23)

#### D. The result above is not contradictory to the singularity theorems

We first intuitively explain the reasons that there is no space-time singularity. It has been proved that there is space-time singularity under certain conditions<sup>[1]</sup>. These conditions

fall into three categories. First, there is the requirement that gravity shall be attractive. Secondly, there is the requirement that there is enough matter present in some region to prevent anything escaping from that region. The third requirement is that there should be no causality violations.

Hawking considers it is a reasonable conjecture that  $\rho_g = \rho > 0$  and  $p_g \ge 0^{[1]}$ . But this conjecture is not valid in the present model, because  $\rho_g = \rho_s - \rho_v > 0$ , = 0 or < 0 are all possible.

As mentioned above, there must be  $\rho_g \leq 0$  when  $T \gtrsim T_{cr}$ . It is seen that  $\rho_g$  does not only stop increasing, but also decreases from  $\rho_g > 0$  (when  $t < t_{cr}$ ) to  $\rho_g(t_{cr}) = 0$  and  $\rho_g(t) < 0$ when  $t > t_{cr}$ . Hence the second condition of the singularity theorem is violated.

The key of non-singularity is  $\rho_{sg} = -\rho_{vg}$  when  $\rho_s = \rho_v$  and  $\rho_s$  and  $\rho_v$  can transform from one to another when  $T \gtrsim T_{cr}$ 

We explain the reasons that there is no space-time singularity from the Hawking theorem as follows. S.W. Hawking has proven the following theorem<sup>[1]</sup>.

The following three conditions cannot all hold:

- (a) every inextendible non-spacelike geodesic contains a pair of conjugate point;
- (b) the chronology condition holds on  $\mu$ ;
- (c) there is an achronal set  $\mathfrak{T}$  such that  $E^+(\mathfrak{T})$  or  $E^-(\mathfrak{T})$  is compact.

The alternative version of the theorem can obtained by the following two propositions. Proposition  $1^{[1]}$ :

If  $R_{ab}V^aV^b \geq 0$  and if at some point  $p = \gamma(s_1)$  the tidal force  $R_{abcd}V^cV^d$  is non-zero, there will be values  $s_0$  and  $s_2$  such that  $q = \gamma(s_0)$  and  $r = \gamma(s_2)$  will be conjugate along  $\gamma(s)$ , providing that  $\gamma(s)$  can be extended to these values.

Proposition  $2^{[1]}$ :

If  $R_{ab}V^{a}V^{b} \geq 0$  everywhere and if at  $p = \gamma(v_{1})$ ,  $K^{a}K^{b}K_{[a}R_{b]cd[e}K_{f]}$  is non-zero, there will be  $v_{0}$  and  $v_{2}$  such that  $q = \gamma(v_{0})$  and  $r = \gamma(v_{2})$  will be conjugate along  $\gamma(v)$  provided that  $\gamma(v)$  can be extended to these values.

An alternative version of the above theorem is as following.

Space-time  $(\mu, g)$  is not timelike and null geodesically complete if:

(1)  $R^{ab}K_aK_b \ge 0$  for every non-spacelike vector **K**.

(2) The generic condition is satisfied, i.e. every non-spacelike geodesic contains a point at which  $K_{[a}R_{b]cd[e}K_{f]}K^{c}K^{d} \neq 0$ , where **K** is the tangent vector to the geodesic. (3) The chronology condition holds on  $\mu$  (i.e. there are no closed timelike curves).

(4) There exists at least one of the following:

(A) a compact achronal set without edge,

(B) a closed trapped surface,

(C) a point p such that on every past (or every future) null geodesic from p the divergence  $\hat{\vartheta}$  of the null geodesics from p becomes negative (i.e. the null geodesics from p are focussed by the matter or curvature and start to reconverge).

In fact,  $R_{ab}$  is determined by the gravitational energy-momentum tensor  $T_{gab}$ . According to the conventional theory,  $T_{gab} = T_{ab}$  so that the above theorem holds.

In contrast with the conventional theory, according to conjecture 1,

$$S_{g\mu\nu} \equiv T_{g\mu\nu} - \frac{1}{2}g_{\mu\nu}T_g = (T_{\nu\mu\nu} - T_{s\mu\nu}) - \frac{1}{2}g_{\mu\nu}(T_v - T_s), \qquad (6.24)$$

 $S_{g00} > 0$ , = 0 and < 0 are all possible. Thus, although the strong energy condition still holds, i.e.

$$\left[ \left( T_s^{ab} + T_v^{ab} \right) - \frac{1}{2} g^{ab} \left( T_s + T_v \right) \right] K_a K_b \ge 0, \tag{6.25}$$

the conditions of propositions 1 and 2 and condition (1) no longer hold, because the gravitational mass density  $\rho_g$  determines  $R_{\mu\nu}$  and  $\rho_g = \rho_v - \rho_s \neq \rho_v + \rho_s = \rho$ . Hence (a) and (c) do not hold, but (b) still holds, and  $\mu$  is timelike and null geodesically complete.

#### VII. EVOLVING PROCESS OF SPACE AFTER INFLATION

#### A. The reheating process

After inflation, the temperature must sharply descend. In this case, it is easily seen that the most symmetric state with  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  is no longer stable and must decay into such a state with  $V_{\min}$ . This is the reheating process. Either of the *S*-breaking and the *V*-breaking can come into being, because *s*-matter and *v*-matter are completely symmetric at  $T \sim T_{cr}$ . Letting the *V*-breaking comes into being, then  $v-SU(5) \longrightarrow v-SU(3) \times U(1)$  and s-SU(5)symmetry is still kept. After the phase transition, we have

$$\varpi_v = \varpi_{v0}, \ \varpi_s = 0, \ V_v = -V_0, \ V_s = V_g = 0.$$
(7.1)
After transition,  $V_v(T_{cr}) - V_v(T_v \sim 0) = V_0$  must first transform into v - particles by the v - Higgs fields. On the other hand, because of the coupling (2.10), the v - Higgs bosons can transform into s - particles by the s - Higgs fields as well. Letting  $\alpha V_0$  transform the v - energy, then  $(1 - \alpha)V_0$  transforms the s - energy. It is necessary  $\alpha > (1 - \alpha)$ . There must be  $\rho'_v = \rho'_s$  before the transition because of (6.17). Thus, after transition, we have

$$\rho_v = \rho'_v + \alpha V_0 > \rho_s = \rho'_s + (1 - \alpha) V_0.$$
(7.2)

After reheating process, all particles must exist in the form of plasma in the initial stage. After temperature descends further, v - particles will exist in the forms of nucleons, leptons and photons, and s - particles will form s - SU(5) color single states whose masses are all non-zero so that  $\rho_{s\gamma} = 0$ . Consequently,  $\rho_v = \rho_{vm} + \rho_{v\gamma}$  and  $\rho_s = \rho_{sm}$ , here  $\rho_{vm} \equiv \rho_{vM} + \rho_{vl}$ and  $\rho_{sm} \equiv \rho_{sM} + \rho_{sl}$ . Thus, letting  $\rho_{sm} > \rho_{vm}$  and the reheating process ends at  $t_2$ , from (7.2) we have

$$\rho_{vm}\left(t\right) + \rho_{v\gamma}\left(t\right) > \rho_{sm}\left(t\right) > \rho_{vm}\left(t\right), \text{ when } t > t_2.$$

$$(7.3)$$

After reheating,  $dV_g/dt \sim 0$  and the temperature effects may be neglected. Ignoring  $p_{sm}$ and  $p_{vm}$ , considering  $p_{v\gamma} = \rho_{v\gamma}/3$  and  $\rho_{s\gamma} = 0$  in the V – breaking, from (4.13) and (7.3) we have

$$\rho_g R^3 = (\rho_{vm} - \rho_{sm}) R^3 = -C_{mg} = \rho_{g0} R_0^3 < 0, \tag{7.4}$$

$$\rho_{s\gamma g}R^4 = \rho_{s\gamma}R^4 = C_{\gamma g} = \rho_{s\gamma 0}R_0^4 > 0, \tag{7.5}$$

where both  $C_{mg}$  and  $C_{\gamma g}$  are constants. From (4.8) and (7.4) - (7.5), (4.5) - (4.6) is reduced to

$$\overset{.}{R}^{2} = 1 + \eta \left( -\frac{C_{mg}}{R} + \frac{C_{\gamma g}}{R^{2}} \right),$$
 (7.6)

$$\overset{\cdots}{R} = \frac{\eta}{2} \left( \frac{C_{mg}}{R^2} - 2\frac{C_{\gamma g}}{R^3} \right).$$
(7.7)

We discuss (7.6) - (7.7) as follows.  $1 - \eta C_{mg}^2/4C_{\gamma g}$  is important for the function R of  $R^{-1}$ .

1. If  $1 - \eta C_{mg}^2/4C_{\gamma g} > 0$ , there is not R = 0. When  $R < R_1 \equiv 2C_{\gamma g}/C_{mg}$ ,  $\ddot{R} < 0$ and R > 0, i.e. space expands with a deceleration; when  $R = R_1$ ,  $\ddot{R} = 0$  and  $\ddot{R} = 0$   $\dot{R}_{\min} = (1 - \eta C_{mg}^2 / 4C_{\gamma g})^{1/2} > 0$ ; when  $R > R_1$ ,  $\ddot{R} > 0$  and  $\ddot{R} > 0$  i.e. space expands with an acceleration. In the process,  $\ddot{R}$  increases from  $\ddot{R} = 0$  to  $\ddot{R}_{\max} = \eta C_{mg}^3 / 54D_{\gamma g}^2$  when  $R = 3D_{\gamma g}/C_{mg}$ , then  $\ddot{R}$  decreases from  $\ddot{R}_{\max}$  to  $\ddot{R}_0 > 0$ .

2. If  $1 - \eta C_{mg}^2/4C_{\gamma g} = 0$ , when  $R < R_1 \equiv 2C_{\gamma g}/C_{mg}$ ,  $\ddot{R} < 0$  and  $\ddot{R} > 0$ ; when  $R = R_1$ ,  $\ddot{R} = \dot{R} = 0$ ; In the case, space can be static.

3. If  $1 - \eta C_{mg}^2/4C_{\gamma g} < 0$ , when  $R < R_2 \equiv (\eta C_{mg}/2) \left[1 - \sqrt{1 - 4C_{\gamma g}/\eta C_{mg}^2}\right]$ ,  $\ddot{R} < 0$  and  $\dot{R} > 0$ ; when  $R = R_2$ ,  $\ddot{R} < 0$  and  $\ddot{R} = 0$ . In the case, space will begin to contract.

The first case is consistent with observations. A computation in detail is the same as that of Ref. [9].

Even  $\chi_s$  and  $\chi_v$  are considered, the above conclusions still hold qualitatively.

#### B. The process of space inflation

Supposing  $\lambda \sim g^4$  and  $g^2 \sim 4\pi/45$  for SU(5), and considering  $m(\Omega_s) = \sqrt{2}\mu$ , from (5.17) we can estimate  $T_{cr}$ ,

$$T_{cr} = \frac{2\mu}{\sqrt{\lambda}} \sim \frac{2\mu}{g^2} \sim \frac{\sqrt{2}m(\Omega_s)}{4\pi/45} = 5m(\Omega_s). \tag{7.8}$$

The temperature will strikingly decrease in the process of inflation, but the potential energy  $V(\varpi_s \sim \varpi_v \sim 0) \sim V_0$  cannot decrease to  $V_{\min}(T_v)$  at once, because this is a super-cooling process.

We can get the expecting results by suitably choosing the parameters in (2.8) - (2.10). In order to estimate  $H = \sqrt{\eta V_0}$ , taking  $V_0 \sim \mu^4/4\lambda$ , from (7.8) we have

$$H = aT_{cr}^2, \quad a \equiv \sqrt{\eta\lambda}/8 \sim g^2 \sqrt{\eta}/8.$$
(7.9)

We can take  $T_{cr}$  to be the temperature corresponding to GUT because the SU(5) symmetry strictly holds at  $T_{cr}$ .

Taking  $T_{cr} \sim 5m(\Omega_s) \sim 5 \times 10^{15} Gev$  and  $\sqrt{\lambda}/8 \sim g^2 \sim 0.035$ , we have  $H^{-1} = 10^{-35}s$ . If the duration of the super-cooling state is  $10^{-33}s \sim (10^8 Gev)^{-1}$ ,  $R_{cr}$  will increase  $e^{100} \sim 10^{43}$ times. The result is consistent with the Guth's inflation model<sup>[15]</sup>.

If there is no v - matter, because of contraction by gravitation, the world would become a thermal-equilibrating singular point, i.e., the world would be in the hot death state. As seen, it is necessary that there are both s - matter and v - matter and both the S - breakingand the V - breaking.

### C. To determine a(t)

Letting  $a = R/R_0$ ,  $\dot{a}_0^2 = H_0^2 \equiv \eta \rho_{gc}$ ,  $\Omega_{gm0} = (\rho_{sm0} - \rho_{vm0})/\rho_{gc}$ ,  $\Omega_{v\gamma0} = \rho_{v\gamma0}/\rho_{gc}$  and  $\Omega_{g0} = \Omega_{gm0} - \Omega_{v\gamma0}$ , and considering

$$\rho_{sm} - \rho_{vm} - \rho_{v\gamma} = \rho_{gc} \left[ \Omega_{gm0} / a^3 - \Omega_{v\gamma0} / a^4 \right], \qquad (7.10)$$

$$H_0^2 (1 + \Omega g_0) = 1/R_0^2, \ k = -1, \tag{7.11}$$

we rewrite (7.6) as

$$\dot{a}^{2} = H_{0}^{2} \left(1 + \Omega_{g0}\right) \\ \cdot \left[1 - \frac{1}{\left(1 + \Omega_{g0}\right)} \left(\frac{\Omega_{gm0}}{a} - \frac{\Omega_{v\gamma0}}{a^{2}}\right)\right].$$
(7.12)

From (7.12) we have

$$t = t_0 - \frac{1}{H_0\sqrt{1 + \Omega_{g0}}} \{\sqrt{1 - M + \Gamma} - \sqrt{a^2 - Ma + \Gamma} + \frac{M}{2} \ln \frac{2 - M + 2\sqrt{1 - M + \Gamma}}{2a - M + 2\sqrt{a^2 - Ma + \Gamma}} \},$$
(7.13)

If  $t_0$  is taken as

$$t_{0} = \frac{1}{H_{0}\sqrt{1+\Omega_{g0}}} \{1 - M + \Gamma + \frac{M}{2} \ln\left[1 - \frac{2}{M} - \frac{2\sqrt{1-M+\Gamma}}{M}\right]\},$$
(7.14)

then

$$t = \frac{1}{H_0\sqrt{1+\Omega_{g0}}} \{\sqrt{a^2 - Ma + \Gamma} + \frac{M}{2} \ln\left[1 - \frac{2a}{M} - \frac{2\sqrt{a^2 - Ma + \Gamma}}{M}\right]\},$$
(7.15)

where  $M = \Omega_{gm0} / (1 + \Omega_{g0})$ , and  $\Gamma = \Omega_{v\gamma0} / (1 + \Omega_{g0})$ .

Taking  $\Omega_{v\gamma0} = 0.001$ ,  $\Omega_{gm0} = 0.3\Omega_{v\gamma0} + 2\sqrt{\Omega_{v\gamma0}}$ ,  $H_0^{-1} = 9.7776 \times 10^9 h^{-1} yr^{[8]}$ . and h = 0.8, we get  $a(t) \cdot a(t)$  is shown by the curve B in the figure 1 and describes evolution of the universe from  $14 \times 10^9 yr$  ago to now. Taking  $\Omega_{v\gamma0} = 0.05$ ,  $\Omega_{gm0} = 2\sqrt{\Omega_{v\gamma0}}$ , we get the a(t) which is shown by the curve A in the figure 1 and describes evolution of the cosmos from  $13.7 \times 10^9 yr$  ago to now. Provided  $2\left(\Omega_{v\gamma0} + \sqrt{\Omega_{v\gamma0}}\right) > \Omega_{gm0}$  which is equivalent to  $1 - \eta C_{mg}^2/4C_{\gamma g} > 0$ , we can get a curve of a(t) which describes evolution of the cosmological scale.

Figure 1

From the two curves we see that the cosmos must undergo a period in which space expands with a deceleration in the past, and undergo the present period in which space expands with an acceleration.

It should be noted that  $\rho_{g0} = \rho_{v0} - \rho_{s0}$  in the V - breaking, but here  $\Omega_{gm0} = (\rho_{s0} - \rho_{v0})/\rho_c = -\rho_{g0}/\rho_c$ . Ignoring  $\Omega_{v\gamma0}$ , taking k = -1 and

$$\sqrt{1-M} = \frac{1}{\sqrt{1+\Omega_{mg}}}, \quad M = \frac{\Omega_{mg}}{1+\Omega_{mg}}, \quad (7.16)$$

$$\sqrt{a^2 - Ma} = \frac{\sqrt{1 - z\Omega_{mg}}}{(1 + z)\sqrt{1 + \Omega_{mg}}}, \ a = \frac{1}{1 + z},$$
(7.17)

and taking

$$t_0 = \frac{1}{H_0 \left(1 + \Omega_{mg}\right)} \left[ 1 + \frac{\Omega_{mg}}{2 \left(1 + \Omega_{mg}\right)^{1/2}} \ln \frac{2 + \Omega_{mg} + 2\sqrt{(1 + \Omega_{mg})}}{(-\Omega_{mg})} \right],$$
(7.18)

we can reduce (7.15) to

$$t = \frac{1}{H_0 (1 + \Omega_{mg})} \{ \frac{\sqrt{1 - z\Omega_{mg}}}{(1 + z)} + \frac{\Omega_{mg}}{2\sqrt{1 + \Omega_{mg}}} \cdot \frac{1}{2\sqrt{1 + \Omega_{mg}}} \cdot \frac{1}{1 + z} \frac{1}{(1 + z)(1 - z\Omega_{mg})} (1 - z\Omega_{mg})}{(1 + z)(-\Omega_{mg})} \}.$$
(7.19)

Replacing  $\Omega_{gm0}$  by  $(-\Omega_{gm0})$  because  $\Omega_{gm0} = (\rho_{s0} - \rho_{v0}) / \rho_c = -\rho_{g0} / \rho_c$ , we see (7.19) to be the same as (3.44) in Ref. [8].

### D. The relationship between luminosity distance and its redshift

From (7.12) and the RW metric we have

$$\int_{a}^{1} \frac{cda}{R\dot{a}} = -\int_{r}^{0} \frac{dr}{\sqrt{1+r^{2}}},$$
(7.20)

$$H_{0}d_{L} = H_{0}R_{0}r(1+z) = \frac{2c}{\left(\Omega_{gm0} - 2\Omega_{v\gamma0}\right)^{2} - 4\Omega_{v\gamma0}} \cdot \left\{2\left(1+\Omega_{g0}\right) - \left(1+z\right)\Omega_{gm0} - \left[2\left(1+\Omega_{g0}\right) - \Omega_{gm0}\right] \cdot \sqrt{1 - \left(\Omega_{gm0} - 2\Omega_{v\gamma0}\right)z + \Omega_{v\gamma0}^{2}z^{2}}\right\},$$
(7.21)

where z = (1/a) - 1 is the redshift caused by R increasing.

Considering  $\Omega_{gm0}$  in (7.21) corresponds to  $-\Omega_{m0}$  in (3.81) in Ref. [8] and  $\Omega_{g0} = \Omega_{gm0} - \Omega_{v\gamma0}$ , we see that (7.21) is consistent with (3.81).

Ignoring  $\Omega_{v\gamma 0}$ , taking  $\Omega_{gm 0} \longrightarrow -\Omega_{gm 0}$ , we reduce (7.21) to

$$H_0 d_L = \frac{2c}{\Omega_{gm0}^2} \cdot \left\{ 2 + \Omega_{gm0} \left( 1 - z \right) - \left[ 2 + \Omega_{gm0} \right] \sqrt{1 - \Omega_{gm0} z} \right\},$$
(7.22)

which is consistent with (3.78) in Ref. [8]. Approximating to  $\Omega_{m0}^1$  and  $z^2$ , we obtain

$$H_0 d_L = z + \frac{1}{2} z^2 \left( 1 + \frac{1}{2} \Omega_{gm0} \right).$$
(7.23)

Taking  $\Omega_{v\gamma0} = 0.001$ ,  $\Omega_{gm0} = 0.3\Omega_{v\gamma0} + 2\sqrt{\Omega_{v\gamma0}}$  and  $H_0^{-1} = 9.7776 \times 10^9 h^{-1} yr^{[8]}$  and h = 0.8, from (7.22) we get the  $d_L - z$  relation which is shown by the curve A in the figure 2; Taking  $\Omega_{v\gamma0} = 0.05$ ,  $\Omega_{gm0} = 2\sqrt{\Omega_{v\gamma0}}$  we get the  $d_L - z$  relation which is shown by the curve B in the figure 2.

Figure 2

# VIII. EXISTING FORMS AND DISTRIBUTION FORMS OF s - SU(5) COLOR SINGLE STATES

In the V – breaking, all s – gauge particles and s – fermions are massless. When the temperature  $T_s$  is high enough, all s – particles must exist in plasma form. When  $T_s$  is low, all s – particles will exist in s – SU(5) color-single state form (conjecture 2). Let A, B, C, D, E denote the 5 sorts of colors. A component of <u>10</u> representation carries color  $\alpha\beta$ ,  $\alpha$ ,  $\beta = A, B, C, D, E, \alpha \neq \beta$ . A component of <u>5</u> representation carries color  $\alpha$ . A gauge boson carries colors  $\alpha\beta^*$ . There are the following sorts of the s – SU(5) color-single states.

2-fermion states:  $\underline{\alpha\alpha}^*$  or  $(\underline{\alpha\beta})(\underline{\alpha\beta})^*$ ,  $\alpha \neq \beta$ . 3-fermion states:  $(\underline{AB})(\underline{CD})\underline{E}$ or  $(\underline{AB})\underline{A}^*\underline{B}^*$ . 4-fermion states:  $(\underline{AB})\underline{CDE}$ . 5-fermion states:  $\underline{ABCDE}$  or  $(\underline{AB}) (\underline{BC}) (\underline{CD}) (\underline{DE}) (\underline{EA}). \text{ Gauge boson single-states: } (\underline{\alpha\beta^*}) (\underline{\alpha^*\beta}) \text{ or } (\underline{\alpha\beta^*}) (\underline{\beta\gamma^*}) (\underline{\gamma\alpha^*}) \text{ etc.}, \alpha \neq \beta \neq \gamma. \text{ Fermion-gauge boson single-states: } \underline{\alpha^*} (\underline{\alpha\beta^*}) \underline{\beta}, \underline{\alpha^*} (\underline{\alpha\gamma^*}) (\underline{\gamma\beta^*}) \underline{\beta} \text{ etc.}.$ 

The masses of all color single states are non-zero, hence  $\rho_{s\gamma} = 0$ . The fermions with the spin s = 1/2 and the least mass and the bosons with the spin s = 0 and the least mass are stable, because there is no the electroweak interaction among SU(5) color single states. Of course, there are the s – antiparticles corresponding to the s – colour single states above as well.

There are interaction among the SU(5) color single states by exchanging the s - SU(5)color single states. The interaction radius must be very small because the masses of all s - SU(5) color single states are non-zero. Thus we can approximately regard the s - SU(5)color single states as ideal gas without collision. Consequently the s - SU(5) color single states cannot form clusters and must distribute loosely in space, but it is possible they form s - superclusterings as the neutrinos.

# IX. NEW PREDICTIONS AND AN INFERENCE

## A. New predictions

1. It is possible that huge voids are not empty and are equivalent to huge concave lenses. The density of hydrogen inside the huge voids is more less than that predicted by the conventional theory.

There is no interaction except the gravitation among the s - SU(5) singlets (the interaction by exchanging SU(5) singlets between two SU(5) singlets may be neglected), because SU(5) is a simple group. Hence the s - SU(5) singlets cannot form massive S - objects, the s - SU(5) color single states can be regarded as ideal gas without collision, the decoupling temperature of the s - SU(5) singlets must be very high, and the velocities of the s - SU(5)singlets must be very large and invariant.

The ideal gas has the effect of free flux damping for clustering, i.e. the s - SU(5) singlets with very high velocities prevents clustering to form. Thus the s - SU(5) singlets must loosely distribute in space or form s - superclusters (similar to neutrino superclusters) which are the huge v - voids for v - observers.

Consequently, huge v - voids in the V - breaking are, in fact, superclusters of s - SU(5)

singlets. The huge v - voids are not empty. There must be s - matter inside them, and  $\rho'_s \gg \rho'_v$ ,  $\rho'_s > \rho_s$ , and  $\rho'_v < \rho_v$  because of the repulsion between v - matter and s - matter. Here  $\rho'_s$  and  $\rho'_v$  denote the densities inside the huge v - voids, and  $\rho_s$  and  $\rho_v$  denote the densities outside the huge v - voids. Because there is the repulsion between s - matter and v - matter and there is the gravitation among s - particles, a huge v - void can form. The characters of such a huge v - void are as follows:

A. A v - void must be huge, because there is no other interaction among the s - SU(5) color single states except the gravitation and the masses of the s - SU(5) singlets are very small.

B. When v - photons pass through such a huge v - void, the v - photons must suffer repulsion from  $\rho'_s$  and are scattered by the v - void as they pass through a huge concave lens. Consequently, the galaxies behind the huge v - void seem to be darker and more remote. Hence the huge voids are equivalent to huge concave lenses.

C. Both density of matter and density of dark matter in huge voids must be more lower than those predicted by the conventional theory. Consequently, the density of hydrogen and the density of helium inside the huge voids must be more less than that predicted by the conventional theory.

The predict can be confirmed or negated by the observation of hydrogen distribution.

This is a decisive prediction which distinguishes the present model from other models.

2. The gravitation between two galaxies with distant long enough will be less than that predicted by the conventional theory.

There must be s-matter between two v-galaxies with distance long enough, hence the gravitation between the two v-galaxies must be less than that predicted by the conventional theory due to the repulsion between s-matter and v-matter. When the distance between two v-galaxies is small, the gravitation is not influenced by s-matter, because  $\rho_s$  must be small when  $\rho_v$  is big.

#### 3. A black hole with its mass and density big enough will transform into a white hole

Letting there be a v-black hole with its mass and density to be so big that its temperature can arrive at  $T_v \gtrsim T_{cr} = 2\mu/\sqrt{\lambda}$  since the black hole contracts by its self-gravitation, then the expectation values of the Higgs fields inside the v-black hole will change from  $\varpi_v = \varpi_{v0}$ and  $\varpi_s = 0$  into  $\varpi_v = \varpi_s = 0$ . Consequently, inflation must occur. After inflation, the most symmetric state will transit into the V-breaking. Thus, the energy of the black hole must transform into both v - energy and s - energy. Thus, a v - observer will find that the black hole disappears and a white hole appears.

In the process, part of v - energy transforms into s - energy. A v - observer will consider the energy not to be conservational because he cannot detect s - matter except by repulsion. The transformation of black holes is different from the Hawking radiation. This is the transformation of the vacuum expectation values of the Higgs fields. There is no contradiction between the transformation and the Hawking radiation or another quantum effect, because both describe different processes and based on different conditions. According to the present model, there still are the Hawking radiation or other quantum effects of black holes. In fact, the universe is just a huge black hold. The universe can transform from the S - breaking into the V - breaking because of its contraction. This transformation is not quantum effects.

# **B.** An inference : $\lambda_{eff} = \lambda = 0$ , although $\rho_{vac} \neq 0$

The effective cosmological constant  $\lambda_{eff} = \lambda + \rho_{g,vac}$ . The conventional theory can explain evolution with a small  $\lambda_{eff}$ , but  $\rho_{g,vac} = \rho_{vac} \gg \lambda_{eff}$ , Consequently, the issue of the cosmological constant appears.

 $\rho_{vac} = 0$  can be obtained by some supersymmetric model, but it is not a necessary result. On the other hand, the particles predicted by the supersymmetric theory have not been found, although their masses are not large.

 $\rho_{vac} = 0$  is a necessary result of our quantum field theory without divergence<sup>[6]</sup>. In this theory,  $\rho_{vac} = 0$  is naturally obtained without normal order of operators, there is no divergence of loop corrections, all known results are obtained, and dark matter which can form dark galaxies is predicted<sup>[7]</sup>. As mention above, this model can explain evolution of the universe without  $\lambda_{eff}$ , hence

$$\lambda_{eff} = 0. \tag{9.1}$$

Applying the conventional quantum field theory to the present model, we have  $\rho_{vac} = \rho_{s,vac} + \rho_{v,vac}$ , here  $\rho_{vac}$  is the energy density of the vacuum state. According to the conjecture 1, s-particles and v-particles are symmetric. Hence both ground states must be symmetric as well. Hence

$$\rho_{s,vac} = \rho_{v,vac} = \rho_{vac}/2. \tag{9.2}$$

According to conjecture 1,  $\rho_{gs} = -\rho_{gv}$  when  $\rho_s = \rho_v$ . Consequently, although

$$\rho_{vac} = \rho_{s,vac} + \rho_{v,vac} = 2\rho_{s,vac} \gg 0 \tag{9.3}$$

we have still

$$\rho_{g,vac} = \rho_{sg,vac} + \rho_{vg,vac} = \rho_{s,vac} - \rho_{v,vac} = 0, \qquad (9.4)$$

$$\lambda_{eff} = \lambda + \rho_{g,vac} = \lambda = 0. \tag{9.5}$$

Here  $\lambda$  is the Einstein cosmological constant. This is a direct inference of the present model, and independent of a quantum field theory.

Because of (9.4), for the vacuum state in the S – breaking or the V – breaking, the Einstein field equation is reduced to

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -8\pi G \left(T_{s,vac,\mu\nu} - T_{v,vac,\mu\nu}\right)$$
  
=  $-8\pi G \left(T_{v,vac,\mu\nu} - T_{s,vac,\mu\nu}\right) = 0.$  (9.6)

This is a reasonable result.

## X. DISCUSSION AND CONCLUSIONS

#### A. Discussion

The problem of total energy conservation in the general relativity is unsolved up to now, because tensors at different points cannot summed up. Based on the same reason, the problem of total gravitation mass conservation in the general relativity is unsolved as well. Hence I can only discuss  $T_{g;\nu}^{\mu\nu} = 0$  and  $T_{,\nu}^{\mu\nu}$ . From (2.22) and (4.9) we see that there are the following corresponding relations between the conventional theory and the present model.

$$T^{\mu\nu}_{;\nu} = 0 \longrightarrow T^{\mu\nu}_{g;\nu} = 0, \qquad (10.1)$$

$$\dot{\rho} + 3(\rho + p)\frac{\dot{R}}{R} = -\dot{V}$$
$$\longrightarrow \dot{\rho}_g + 3(\rho_g + p_g)\frac{\dot{R}}{R} = -\dot{V}_g.$$
(10.2)

(10.2) is consistent with (10.1). For  $T_g^{\mu\nu} = T_s^{\mu\nu} - T_v^{\mu\nu}$ , there is no such a condition similar to the dominant energy condition. Hence, there is no such a theorem similar to the positive energy theorem for  $T_g^{\mu\nu}$ . All  $\rho_g > 0$ , = 0 or  $\rho_g < 0$  are possible.

If  $V_g = 0$ ,  $\rho_s$  cannot transform into  $\rho_v$ . For example, when temperature is low or so high that  $\langle \omega_s \rangle = \langle \omega_v \rangle = V_s = V_v = 0$  and  $V_g = V_0$ , the transformation of  $\rho_s$  and  $\rho_v$  from one to another may be neglected. In the case,  $\rho_g$  is the same as  $\rho$  on the left of (10.2) and (10.1) and (10.2) obviously hold. It i.e. seen that in general, (10.1) and (10.2) can hold.

When  $V_g \neq 0$ ,  $\rho_s$  and  $\rho_v$  can transform from one to another. Although  $\rho_s$  transforms to  $\rho_v$  so that  $\dot{\rho}_g < 0$ , (10.1) and (10.2) still hold. For example, all  $V_s$ ,  $\rho_s$  and  $\rho_v$  will increase when space contracts in the S – *breaking*, but (10.1) and (10.2) still hold. Such a process is not different from a equilibrium or stable process.

It is possible that the differential law of energy-momentum conservation holds,

$$T_{S,\nu}^{\mu\nu} = T_{V,\nu}^{\mu\nu} = T_{,\nu}^{\mu\nu} = 0, \qquad (10.3)$$

where  $T^{\mu\nu}$  may contain the contribution of gravitational field as well. This is because there is no restriction for  $T^{\mu\nu}$ .

### B. Conclusions of the model based on the RW metric

A new conjecture is proposed that there are s - matter and v - matter which are symmetric, whose gravitational masses are opposite to each other, although whose masses are all positive. Both can transform from one to another when temperature  $T \gtrsim T_{cr}$ . Consequently there is no singularity in the model. The cosmological constant  $\lambda_{eff} = \lambda = 0$  is determined although the energy density of the vacuum state is still very large. A theorem related to singularity is presented.

The conjecture are not in contradiction with all given experiments and astronomical observations up to now, although the conjecture corrects the equivalence principle. This is because SU(5) singlets which violate the equivalence principle cannot be observed and can only loosely distribute in space as so-called dark energy.

There are two sorts of breaking modes, i.e. the S - breaking and the V - breaking. In the V - breaking v - SU(5) is broken into  $v - SU(3) \times U(1)$  and s - SU(5) is still kept. Consequently, v - particles get their masses and form v - atoms, v - observers and v - galaxies etc., while s - gauge bosons and s - fermions are still massless and must form s - SU(5) color-single states after reheating. There is no interaction among the s - SU(5)color-single states except the gravitation, because SU(5) group is a simple group. Hence they must distribute loosely in space, cannot be observed and can cause space to expand with an acceleration. Thus, v - matter is identified with conventional matter (include dark matter) and s - matter is similar to the dark energy. But in contrast with the dark energy, the gravitational mass of s - matter is negative in the V - breaking.

Based on the present model, the space evolving process is as follows. Firstly, in the S - breaking,  $\rho_g = \rho_s - \rho_v > 0$ , hence space contracts and  $T_s$  rises. When  $T_s$  arrives the critical temperature  $T_{cr}$ ,  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$ , the  $[v - SU(5)] \times [s - SU(5)]$  symmetry is realized, and the masses of all particles originating from the couplings with the Higgs fields are zero so that  $\rho_s$  and  $\rho_v$  can transform from one into another. As a consequence,  $\rho_g = \rho_s - \rho_v < 0$ , and  $V_g = V_0$ . When space contracts further, R arrives the least scale  $R_{\min} > 0$  and  $T_v$  arrives the highest temperature  $T_{\max}$ . Then space expands and inflation must occur. After the inflation, the phase transition of the vacuum (the reheating process) occurs. After the reheating process, this state with the highest symmetry transits to the state with the V - breaking. Space in the V - breaking have two evolving stages. Space firstly expands with a deceleration because  $\rho_g = \rho_v - \rho_s > 0$ ; then expands with an acceleration up to now because  $\rho_g < 0$  and k = -1. The results above is still valid when  $V \rightleftharpoons S$  and  $v \rightleftharpoons s$ . It is seen that the world in the S - breaking and the world in the V - breaking can transform from one into another.

There are the critical temperature  $T_{cr}$ , the highest temperature  $T_{max}$ , the least scale  $R_{min}$ and the largest energy density  $\rho_{max}$  in the universe.  $V_0$  and  $T_{cr}$  are two new important constants.

A formula is derived which well describes the relation between a luminosity distance and

the redshift corresponding to it.

Three new predicts have been given.

Huge v - voids in the V - breaking are not empty, but are superclusters of s - particles. The huge voids are equivalent to huge concave lens. The density of hydrogen and the density of helium in the huge voids predicted by the present model must be much less than that predicted by the conventional theory.

The gravitation between two galaxies with distant long enough will be less than that predicted by the conventional theory.

It is possible that a v - black hole with its big enough mass and density can transform into a huge white hole by its self-gravitation.

It is possible that the differential law of energy-momentum conservation holds,  $T^{\mu\nu}_{,\nu} = 0$ , where  $T^{\mu\nu}$  may contain the contribution of gravitational field as well. This is because there is no restriction for  $T^{\mu\nu}$ .

# Part II

# An explanation for some phenomena based on the first model

# XI. INTRODUCTION TO EXPLANATION FOR SOME PHENOMENA

The evolution of the universe has been explained and the cosmological constant issue has been solved based on the first model. There are two ways to explain the primordial nucleosynthesis and the cosmic microwave background radiation (CMBR) by suitably choosing parameters in this model. The first way is based on this model and the F - W dark matter model<sup>[7]</sup> or mirror dark matter model<sup>[16]</sup>; The second way is based on this model and the cold-dark matter model. Generalizing the equations governing nonrelativistic fluid motion<sup>[8]</sup> to the present model, we have derived the equations of structure formation. According to the equations, galaxies can form earlier than that according to the Friedmann model; Galaxies distribution is not uniform in radial direction. There are two sphere layers in which the density of galaxies is relatively larger. SU(5) color single states must loosely distribute in space or form superclusters which are equivalent to huge voids for observers.

Only the expanding process after reheating is considered in the present part. According to the model, after reheating, the density of the gravitational potential energy  $V_g = V_v + V_0 - V_s \sim 0$  and  $\dot{V}_g \sim 0$ . Thus the evolution equations (4.5) – (4.6) is reduced to

$$R^{2} + k = \eta \rho_g R^2, \ k = -1, \ \eta \equiv 8\pi G/3,$$
 (11.1)

$$\overset{\cdots}{R} = -\frac{1}{2}\eta \left(\rho_g + 3p_g\right) R.$$
(11.2)

$$\rho_g = \rho_v - \rho_s, \ p_g = p_v - p_s, \ \text{in the } V - breaking.$$
(11.3)

# A. The F - W dark matter model

The F - W dark matter model<sup>[7]</sup> is a necessary inference of the quantum field theory without divergence<sup>[6]</sup>. The F - W dark matter model is similar with the mirror dark matter model.

According to the mirror dark matter model, it is impossible that the density of matter is equal to that of mirror matter in order to explain the primordial nucleosynthesis and CMBR. This is too difficult to understanding, because matter and mirror matter are symmetric and both can transform from one into another when temperature is high enough.

In contrast with the mirror dark matter model, according to the F-W model, F-matterand W-matter are completely symmetric in both properties and density, there is no interaction except the gravitation and Higgs couplings between F-matter and W-matter. Thus, W-matter is dark matter for a F-observer, and vice versa. Both F-matterand W-matter belong to v-matter. We regard F-matter as conventional matter, and W-matter as dark matter or mirror matter.

In order to analyze the primordial nucleosynthesis, we divide F - matter into three sorts. The first sort is called F - massive particles whose masses are denoted by  $M_{vF}$ , and  $M_{vF} > 1 MeV$ . It is composed of F - nucleons and unknown F - particles. The unknown F - particles are invisible for a time so that they belong to cold dark matter. The second sort is called F - light particles whose masses are denoted by  $m_{vFl}$ ,  $1 MeV > m_{vFl} \gg 1eV$ , e.g. F - electrons. The third sort is called F - photon - like particles whose masses are zero, e.g. F - photons and F - neutrinos whose masses are approximately regarded as zero.

Let the densities of the three sorts of particles are  $\rho_{vFM}$ ,  $\rho_{vFl}$ , and  $\rho_{vF\gamma}$ , respectively, then

$$\rho_{vF} = \rho_{vFM} + \rho_{vFl} + \rho_{vF\gamma}. \tag{11.4}$$

Considering F - matter and W - matter to be completely symmetric, we have

$$\rho_{vFM} = \rho_{vWM}, \ \rho_{vFl} = \rho_{vWl}, \ \rho_{vF\gamma} = \rho_{vW\gamma}, \tag{11.5}$$

$$\rho_{vW} = \rho_{vWM} + \rho_{vWl} + \rho_{vW\gamma} = \rho_{vF}. \tag{11.6}$$

$$\rho_v = \rho_{vF} + \rho_{vW} = 2\rho_{vF} = \rho_{vM} + \rho_{vl} + \rho_{v\gamma}.$$
(11.7)

$$\rho_{vM} = 2\rho_{vFM}, \ \rho_{vl} = 2\rho_{vFl}, \ \rho_{v\gamma} = 2\rho_{vF\gamma}.$$
(11.8)

$$\rho_{vFM} = \rho_{vFB} + \rho_{vUF}, \ \rho_{vUF} = \rho_{vUW}, \ \rho_{vFB} = \rho_{vWB}, \tag{11.9}$$

where  $\rho_{vFB}$  is the density of F-baryons and  $\rho_{vUF}$  is the density of unknown F-particles. Both W - matter and unknown F - particles are invisible for a F - observer, hence the density  $\rho_{dm}$  of dark matter is

$$\rho_{dm} = \rho_{vW} + \rho_{vUF} \equiv \rho_{cd} + \rho_{hd}, \qquad (11.10)$$

$$\rho_{cd} \equiv \rho_{vWB} + \rho_{vUW} + \rho_{vUF}, \qquad (11.11)$$

$$\rho_{hd} \equiv \rho_{vWl} + \rho_{vW\gamma},\tag{11.12}$$

where  $\rho_{cd}$  and  $\rho_{hd}$  are the density of cold-dark matter and the density of hot-dark matter, respectively. Observations show  $\rho_{hd0} \ll \rho_{cd0}$ . It is seen from (11.6) and (11.9) – (11.10) that dark matter is composed of three parts. The first part is the W – baryons which can form galaxies (dark galaxies) as the F – baryons. The second part is unknown F – matter which is invisible for a time. The third part is composed of unknown W – matter, W – photons and light W – particles which are possibly always invisible and have only cosmological effects.

The density of visible matter is

$$\rho_{vis} = \rho_{vFB} + \rho_{vFl} + \rho_{vF\gamma}$$

 $\rho_{vF}$  and  $\rho_{vW}$  can transform from one into another when temperature is high enough by the couplings of F - Higgs and W - Higgs fields.

In V – breaking, all s – particles must form s - SU(5) color single states whose masses are not zero, i.e. there is no s – photon so that  $\rho_{s\gamma} = 0$ . Thus, s-particles may be divided into two sorts. The first sort is composed of massive s-particles whose masses  $M_s > 1 MeV$ and density is  $\rho_{sM}$ ; The second sort is composed of light s-particles with their masses  $m_{sl}$ and density  $\rho_{sl}$ ,  $1 MeV > m_{sl} \gg 1 eV$ . Thus,

$$\rho_{sm} = \rho_{sM} + \rho_{sl}.\tag{11.13}$$

 $\rho_{sl}/\rho_{sM}$  is a undetermined parameter.

When temperature T < 1 MeV,  $p_{sM} \ll \rho_{sM}$  and  $p_{vM} \ll \rho_{vM}$  so that  $p_{sM}$  and  $p_{vM}$  may be neglected. Considering  $V_g = \dot{V}_g = 0$  after particles decouple, and  $p_{v\gamma} = \rho_{v\gamma}/3$ , letting  $p_{sl} = \kappa \rho_{sl}$  and  $p_{vl} = \kappa \rho_{vl}$ , from (11.1) – (11.2) we have

$$\frac{d\left(\rho_{Mg}R^{3}\right)}{dt} + \frac{d\left(\rho_{lg}R^{3(1+\kappa)}\right)}{R^{3\kappa}dt} + \frac{d\left(\rho_{v\gamma g}R^{4}\right)}{Rdt} = 0,$$
  

$$\rho_{Mg}R^{3} = \left(\rho_{vM} - \rho_{sM}\right)R^{3} = \rho_{vM}\left(1 - S_{M}\right)R^{3} \sim C_{Mg},$$
  

$$\rho_{lg}R^{3(1+\kappa)} = \left(\rho_{vl} - \rho_{sl}\right)R^{3(1+\kappa)} \sim C_{lg}, \quad \rho_{v\gamma g}R^{4} \sim C_{v\gamma g},$$
(11.14)

where all  $C_{mg}$ ,  $C_{lg}$  and  $C_{v\gamma g}$  are parameters, and  $S_M \equiv \rho_{sM}/\rho_{vM}$  is a undetermined parameter. When temperature is high enough, e.g.  $T \gtrsim T_{cr}$ , all masses may be regarded photon-like particles.  $\kappa$  is a function of temperature,

$$\kappa \lesssim 1/3$$
, when  $T \gtrsim 1 MeV$ ,  
 $0 < \kappa < 1/3$  when  $1 MeV > T \gtrsim 1 eV$   
 $0 \lesssim \kappa$ , when  $T \lesssim 1 eV$ . (11.15)

#### B. The cold-dark matter model

Both matter and dark matter belong to v-matter according to this cosmological model. Cold-dark matter is composed of such particles which have large masses and very weak interaction. The present observations show that there are dark energy and dark matter. Visible matter is ordinary baryon matter. Let the density of dark energy, the density of visible matter, the density of cold-dark matter, the density of hot-dark matter be  $\rho_{de}$ ,  $\rho_B$ ,  $\rho_{cd}$ , and  $\rho_{hd}$ , respectively, then  $\rho_{dm} = \rho_{cd} + \rho_{hd}$  and the total density is  $\rho_t = \rho_{de} + \rho_{dm} + \rho_B + \rho_l + \rho_{\gamma}$ . Observations show

$$\frac{\rho_{de0}}{\rho_{t0}} = 0.73, \quad \frac{\rho_{dm0}}{\rho_{t0}} = 0.23, \quad \frac{\rho_{B0}}{\rho_{t0}} = 0.04,$$
  

$$\rho_{dm0}/\rho_{B0} \sim 6, \quad \rho_{cd0} \gg \rho_{hd0} \sim \rho_{\gamma0}, \quad \rho_{dm0} \sim \rho_{cd0}.$$
(11.16)

After nucleons decouple,  $\rho_{dm} \sim R^{-3}$  and  $\rho_{cd} \sim R^{-3}$ . From (11.16) we have

$$\rho_{dm} \sim \rho_{cd} \sim 6\rho_B. \tag{11.17}$$

According to this cosmological model, there is no dark energy, but there are s-matter and v-matter. In the V-breaking, s-matter corresponds the dark energy, but  $\rho_{sg} = -\rho_s$ . For primordial nucleosynthesis and CMBR, (11.1) - (11.3) is applicable. The hot-dark matter is composed of neutrinos mainly and  $\rho_{hd} \sim \rho_{\gamma}$ . For simplicity, we replace  $\rho_{hd} + \rho_{\gamma}$  by  $\rho_{\gamma}$ . Combining this cosmological model and the the cold-dark matter model, we have

$$\rho_v = \rho_B + \rho_{cd} + \rho_l + \rho_\gamma, \quad \rho_{hd} + \rho_\gamma \longrightarrow \rho_\gamma \tag{11.18}$$

Thus (11.1) can be write as

$$H^{2} = \left(\frac{\dot{R}}{R}\right)^{2} = -k/R^{2} + \eta \{(\rho_{B} + \rho_{cd} + \rho_{l} + \rho_{\gamma}) - (\rho_{sM} + \rho_{sl})\}.$$
(11.19)

In section 12, primordial nucleosynthesis is explained; In section 13, CMBR is explained; In section 14, dynamics of structure formation and the distributive form of the SU(5) color single states; Section 15 is the conclusion of the part.

# XII. PRIMORDIAL NUCLEOSYNTHESIS

#### A. Explanation for primordial nucleosynthesis the by the Friedmann model

The primordial helium abundance  $Y_4$  is determined by  $n_n/n_p^{[17]}$ ,

$$Y_4 = 2/ \left[ 1 + (n_n/n_p)^{-1} \right],$$

$$n_n/n_p = \exp\left(-\bigtriangleup m/kT_d\right), \ \bigtriangleup m = m_n - m_p,$$
(12.1)

where  $n_n/n_p$  is the neutron-proton ratio in the unit comoving volume at the freeze-out temperature  $T_d$ .  $T_d$  is determined by  $\Gamma = \Gamma(T)$  and H(T) = R/R, e.g.

$$\Gamma(T) = H(T), \qquad (12.2)$$

here  $\Gamma$  is the interaction rate experienced by a nucleon. According to the conventional theory

$$\Gamma \sim G_F^2 T^5, \tag{12.3}$$

where  $G_F$  is the weak interaction Fermi constant.

Considering the cold-dark matter model and the Friedmann model with k = 0 and the effective cosmological constant  $\lambda_{eff}$ , we have

$$\widetilde{H}^2 = \eta \left(\rho_B + \rho_{cd} + \rho_l + \rho_\gamma + \lambda_{eff}\right)$$
(12.4)

The freezing-out temperature is  $T_d \sim 0.8 MeV$  at which  $\Gamma(Td) \sim H(T_d)$ . Consequently  $\rho_B$  may be neglected. The reasons are as follows.

$$\rho_B = \rho_{B0} \left(\frac{R_0}{R}\right)^3, \quad \rho_\gamma = \rho_{\gamma 0} \left(\frac{R_0}{R}\right)^4,$$
$$\frac{\rho_B}{\rho_\gamma} = \frac{\rho_{B0}}{\rho_{\gamma 0}} \frac{R}{R_0}, \quad \frac{R}{R_0} \sim \frac{T_{\gamma 0}}{T_\gamma}.$$
(12.5)

 $T_{\gamma}$  is the effective temperature of photons.  $T_{\gamma} = T$  before photons decouple.  $T_{\gamma 0} = 0.235 \times 10^{-4} eV$ . Thus  $T_d/T_0 \sim 10^{10}$ . Substituting  $T_d/T_0$  and  $\rho_{B0}/\rho_{\gamma 0} \sim 6000^{[5]}$  into (12.4), we have  $\rho_B(T_d) \ll \rho_{\gamma}(T_d)$  so that  $\rho_B(T_d)$  may be neglected. According to the cold-dark matter model,  $\rho_{dm} \sim \rho_{cd} \sim 6\rho_B \sim R^{-3}$  when  $T \sim T_d$ , and  $\lambda_{eff}$  is very small and is invariant. Hence  $\rho_{cd}$  and  $\lambda_{eff}$  may be neglected when  $T \sim T_d$  (see (11.14), (11.17) – (11.18)).

When  $T \sim T_d$ , electrons and neutrinos may be regarded as photon-like particles. Considering three generations of neutrinos and electrons and their antiparticles, we have

$$g^* = 2 + (7/8) \times (2 \times 2 + 3 \times 2) = 43/4$$

Thus, from the Friedmann model we have

$$\widetilde{H}^{2}(T_{d}) \sim \eta \left(\rho_{l} + \rho_{\gamma}\right) = \eta \frac{\pi^{2}}{30} g^{*} T_{d}^{4}.$$
 (12.6)

Taking (12.2) as a rough approximation, from (12.3) and (12.6) we get  $T_d \sim 0.8 MeV$ . Substituting  $T_d \sim 0.8 MeV$  into (12.1), we get

$$n_n/n_p \sim 1/7, \quad Y_4 \sim 1/4.$$
 (12.7)

# B. Explanation for primordial nucleosynthesis by this cosmological model and the F-W dark matter model or the cold dark matter model

As mentioned before, F-matter is regarded as the conventional matter. Thus  $\rho_{vFB} = \rho_B$ ,  $\rho_{vFl} = \rho_l, \ \rho_{vF\gamma} = \rho_{\gamma} \text{ and } T_v = T.$  Thus, (12.3) still holds, provided T is replaced by  $T_v$ . This is because both W-particlesand s-particles do not influence F-nuclear interactions. Thus, provided (12.6) is derived from the present model, this model can explain primordial nucleosynthesis.

Observations show  $\rho_{dm}/\rho_B \sim 6$ . As mentioned above, when  $T \sim T_d$ ,  $\rho_B \ll \rho_{\gamma}$ , i.e.  $\rho_{vFB} \ll \rho_{vF\gamma}$ . Consequently,  $\rho_{vFB}$  and  $\rho_{cd}$  may be neglected when  $T \sim T_d$  due to  $\rho_{cd} \lesssim \rho_{dm}$ . Thus  $\rho_{vM}$  may be neglected due to (11.8) – (11.9).

Before photons decouple,  $T_{\gamma} = T$ . For the stage in which photon-like particles is dominative,

$$\frac{T_s}{T_{s0}} = \frac{R_0}{R} = \frac{T_v}{T_{v0}}$$
(12.8)

we have  $T_s \sim T_v$ . Thus, because of (11.15) – (11.17) and  $\rho_{sM} \sim \rho_{vM} \ll \rho_{vF\gamma}$ ,  $\rho_{sM}$  may be neglected when  $T_v \sim T_d$ . On the other hand, when  $T_v \sim T_d$ ,

$$k/\eta R^2 \ll \rho_{vF\gamma} \sim R^{-4},\tag{12.9}$$

so that  $k/\eta R^2$  may be neglected when  $T_v \sim T_d$ .

Taking

$$\rho_{vWl}(T) + \rho_{vW\gamma}(T) - \rho_{sl}(T) \sim 0, \text{ when } T \gtrsim T_d, \qquad (12.10)$$

neglecting  $\rho_{vFM}$  and  $\rho_{sM}$ , when  $T_v \sim T_d$ , we can reduce (11.1) as

$$H^{2} = \eta \{ 2 \left( \rho_{vFM} + \rho_{vFl} + \rho_{vF\gamma} \right) - \left( \rho_{sM} + \rho_{sl} \right) \} \sim \eta \left( \rho_{vFl} + \rho_{vF\gamma} \right) = \eta \frac{\pi^{2}}{30} g^{*} T_{d}^{4},$$
(12.11)

where  $g^* = 43/4$ . Thus, (12.11) is the same as (12.6). It is seen that this model and the F - W model can still explain the primordial nucleosynthesis and  $Y_4$ , although  $\rho_{vF} = \rho_{vW}$ . This is different from the mirror dark matter model.

It is easily seen from (11.10) - (11.12) that the cold dark matter model is equivalent to  $\rho_{vWl} = \rho_{vW\gamma} = 0$ . From (12.10) we see that provided  $\rho_{sl} = 0$  is taken, when  $T \sim T_d$ , we can still explain the primordial nucleosynthesis based this model and the cold dark matter model.

For the cold-dark matter model,  $(-k/R^2)$ ,  $\rho_B$ ,  $\rho_{cd}$  and  $\rho_{sM}$  may be neglected because of the same reasons as above. From (11.18) – (11.19) we see that provided  $\rho_{sl} \sim 0$  is taken when  $T \sim T_d$ , we can still explain the primordial nucleosynthesis based this cosmological model and the cold dark matter model.

#### XIII. COSMIC MICROWAVE BACKGROUND RADIATION

## A. The recombination temperature $T_{rec}$

It is the same as the conventional theory that there are the inflation and big bang processes in the present model. Hence there must be CMBR.

As mentioned before, v - F - matter is regarded as the conventional matter. Thus,  $T = T_v$ . The recombination mechanism and temperature of this model is the same as those of the conventional theory, because there is no interaction between the v - F - particlesand the v - W - particles except the gravitation and there is no interaction between the v - F - particles and the s - particles except the repulsion as well. From the Saha formulas<sup>[17]</sup> we can determine the recombination temperature  $T_{vrec}$ ,

$$\frac{1-\chi}{\chi^2} = 1.1 \times 10^{-8} \xi T_v^{3/2} \exp \frac{13.6}{T_v} 
= 3.96 \times 10^{-14} \xi A^{3/2} \exp \frac{57872}{A}, \qquad (13.1) 
\chi = n_p / (n_p + n_H), 
A \equiv T_{\gamma v} / T_{v \gamma 0} = T_v / T_{v \gamma 0},$$

where  $n_p = n_e$  and  $n_H$  are the number densities of protons and hydrogens, respectively,  $T_{v\gamma 0} = 2.35 \times 10^{-4} ev$ , and 13.6 ev is the ionization potential energy of hydrogen. Before v - F - photons decouple,  $T_{\gamma v} = T_v$ .

Taking  $\xi \sim 5 \times 10^{-10}$  and  $\chi = 0.1$ , we obtain<sup>[5]</sup>.

$$T_{vrec} = 0.295ev = T_{rec}, \ (1 + z_{rec}) = T_{vrec}/T_{v\gamma 0} = 1255.$$
 (13.2)

 $T_{vrec}$  is just the result of the conventional theory.

## B. The temperature of matter-radiation equality

Let the temperature of matter-radiation equality be  $T_{eq}$ , then  $T_{eq} \sim T_{vrec} \ll m_e \sim m_{vFl}$ . Thus  $p_{vFl}$  may be neglected so that  $\rho_{vFl} \propto R^{-3}$ . When  $T_v \leq T_{eq}$ , because of (11.8) and (11.14), the density of v - matter can be written as

$$\rho_{vm} = 2\rho_{vFm} = 2\left(\rho_{vFM} + \rho_{vFl}\right) = \rho_{vm0}R_0^3/R^3.$$
(13.3)

In contrast with the conventional theory, according to the F - W model, not only there are F - photon (ordinary photons), but also W - photons (dark photons), and  $\rho_{vF\gamma} = \rho_{vW\gamma}$ . From this we can estimate  $T_{eq}$  as follows.

Because (11.14) and (11.8) or (11.5), the density of photon-like particles is

$$\rho_{v\gamma} \equiv 2\rho_{vF\gamma} = 2\rho_{vF\gamma0}R_0^4/R^4 = 2 \times \frac{\pi^2}{30}g_{\gamma}^*T_{v\gamma}^4, \qquad (13.4)$$

$$g_{\gamma}^* = 2 + \frac{7}{8} \times 6 \times \left(\frac{4}{11}\right)^{4/3} = 3.36.$$
 (13.5)

Here the photons and the three species of neutrinos are considered [18].

Letting  $\rho_{vm}(T_{eq}) = \rho_{v\gamma}(T_{eq})$ , Considering  $(R_0/R) = (T_{v\gamma}/T_{v\gamma 0})$ , from (13.3) – (13.4) we have

$$\frac{\rho_{vm0}}{2\rho_{vF\gamma0}} = \frac{1}{2} \cdot \frac{\rho_{m0}}{\rho_{\gamma0}} = \frac{R_0}{R} = \frac{T_{v\gamma}}{T_{v\gamma0}}$$
(13.6)

Considering  $T_{vF\gamma 0} = T_0 = 2.35 \times 10^{-4} ev$ , from (13.4) - (13.5) we get

$$\rho_{v\gamma0} = 2\rho_{vF\gamma0} = 6.7425 \times 10^{-51} Gev^4.$$
(13.7)

Observations show that the total density of matter and dark matter is  $\rho_0 = \Omega_0 \rho_c = 0.27 \rho_c$ ,  $H_0^2 \equiv \eta \rho_c$ . According to the F - W model, this implies

$$\rho_{0} = \rho_{v0} = 2 \left( \rho_{vFM0} + \rho_{vFl0} + \rho_{vF\gamma0} \right)$$
  

$$\simeq 2 \left( \rho_{vFM0} + \rho_{vFl0} \right) = \rho_{vm0} = \Omega_{0}\rho_{c}$$
  

$$= 1.8789 \times 10^{-26} h^{2}\Omega_{0} \cdot kg \cdot m^{-3}$$
  

$$= 9.238 \times 10^{-48} Gev^{4}, \text{ when } h = 0.65.$$
(13.8)

where h = 0.5 - 0.8. Here  $\rho_{\gamma 0}$  is neglected due to  $\rho_{B0} / \rho_{\gamma 0} \sim 6000^{[5]}$ . From (13.6) – (13.8) we obtain

$$T_{veq} = T_{v\gamma eq} = 0.32ev$$
, when  $h = 0.65$ . (13.9)

 $\rho_{\gamma 0} = \rho_{v\gamma 0}/2$  and  $\rho_{m0} = \rho_{vm0}$  in (13.6) are the densities of radiation and matter, respectively, in the conventional theory. Let  $T'_{eq}$  be is the temperature of matter-radiation equality in the conventional theory, from (13.6) we see

$$T'_{eq} = 2T_{veq} = 0.64ev = 2T_{veq}.$$
(13.10)

#### C. Decoupling temperature

The decoupling temperature of photons  $T_{dec}$  is determined by  $\Gamma(T_{dec})$  and  $H(T_{dec})$ . According to the conventional theory<sup>[17]</sup>,

$$\Gamma = n_e \sigma_{Th}, \ n_e = n_p \equiv \chi \xi n_{\gamma}$$

$$(n_p + n_H) = \xi n_{\gamma} = \xi \left(\frac{2.4}{\pi^2} T^3\right),$$

$$\xi \sim 5 \times 10^{-10}, \ \sigma_{Th} = 1.71 \times 10^3 GeV^{-2}.$$

$$\Gamma = 5.4 \times !0^{-36} \chi \xi A^3 GeV,$$
(13.11)

where  $A \equiv T_{\gamma}/T_{\gamma 0} = T_v \gamma/T_{v \gamma 0}$ .

From the present model and the F - W model [7] we have the same result as (13.11), because F-matter is regarded as conventional matter. In order to determine the decoupling temperature  $T_{dec}$ , only it is necessary to determine  $H(T_v)$ .

When  $T_v = T_{vdec}$ ,  $\rho_{vm}$  is the same as (13.3) because  $T_{dec} \sim T_{veq} < 1ev$ . Let when  $T_v = T_{veq}$ ,  $T_s = T_{sq}$  and  $\rho_{sm} = \rho_{sM} + \rho_{sl} \equiv s_{mq}\rho_{vm} (T_{veq})$ . Considering

$$\frac{T_{sq}}{T_s} = \frac{R}{R_{eq}} = \frac{T_{veq}}{T_v} \tag{13.12}$$

and  $\rho_{v\gamma,eq} = \rho_{vm,eq}$ , we can rewrite  $\rho_{vm}$ ,  $\rho_{v\gamma}$  and  $\rho_{sm}$  as follows.

$$\rho_{vm} = \frac{\rho_{vm}}{\rho_{vm0}} \frac{\rho_{vm0}}{\rho_c} \rho_c = \frac{R_0^3}{R^3} \Omega_{m0} \rho_c$$
$$= A^3 \Omega_{m0} \rho_c, \ A \equiv \frac{T_{v\gamma}}{T_{v\gamma0}} = \frac{R_0}{R},$$
(13.13)

$$\rho_{v\gamma} = \frac{\rho_{v\gamma}}{\rho_{v\gamma,eq}} \frac{\rho_{v\gamma,eq}}{\rho_{vm,eq}} \frac{\rho_{vm,eq}}{\rho_{vm0}} \frac{\rho_{vm0}}{\rho_c} \rho_c$$

$$= \left(\frac{T_{v\gamma0}}{T_{v\gamma,eq}}\right) A^4 \Omega_{m0} \rho_c,$$
(13.14)

$$\rho_{sm} = \frac{\rho_{sm}}{\rho_{sm,eq}} \frac{\rho_{sm,eq}}{\rho_{vm,eq}} \frac{\rho_{vm,eq}}{\rho_{vm0}} \frac{\rho_{vm0}}{\rho_c} \rho_c$$

$$=s_{mq}A^3\Omega_{m0}\rho_c,\tag{13.15}$$

$$s_{mq} \equiv \frac{\rho_{sm,eq}}{\rho_{vm,eq}} = \frac{\rho_{sm0}}{\rho_{vm0}}, \text{ e.g. } s_{mq} = 1.5.$$
 (13.16)

(11.14) is considered in (13.16). k = -1 may be neglected because  $R_0/R_{vdec} \sim T_{vdec}/T_0 \sim$ 

 $10^3$  and

$$(-k/\eta R_0^2) (R_0/R_{vdec})^2 \ll \rho_{g0} (R_0/R_{vdec})^3,$$

$$R_{vdec} = R_0 T_{v\gamma 0}/T_{vdec}, \ T_{vdec} = T_{v\gamma dec}.$$
(13.17)

Considering  $H_0 = \sqrt{\eta \rho_c} = 65 km \cdot (s \cdot Mpc)^{-1} = 1.4 \times 10^{-42} Gev$ ,  $\Omega_{vm0} = 0.27$ , and  $T_{v0}/Tveq = (2.35 \times 10^{-4})/0.32 = 0.734 \times 10^{-3}$ , from (13.13) - (13.16) we get

$$H^{2} = \eta \rho_{g} = \eta \left( \rho_{vm} + \rho_{v\gamma} - \rho_{sm} \right)$$
  
= 0.27 × 1.4<sup>2</sup> × 10<sup>-82</sup> A<sup>3</sup>.  
 $\left( 1 - s_{mq} + 0.734 \times 10^{-3} A \right) (Gev)^{2}$ . (13.18)

This is the only difference between this model and the Friedmann model in order to determine the decoupling temperature.

Taking the crude approximation

$$\Gamma\left(T_{v}\right) = H\left(T_{v}\right),\tag{13.19}$$

from (13.11) and (13.18) – (13.19) we can represent  $\chi$  and  $\chi A^{3/2}$  by A. Thereby from (13.1), (13.11) and (13.18) – (13.19) we can determine A and  $\chi$ .

$$\chi A^{3/2} = \xi^{-1} 1.347 \times 10^{-7} \left( 1 - t_{mq} + 0.734 \times 10^{-3} A \right)^{1/2}$$
(13.20)

$$A^{3/2} = \xi^{-1} 1.347 \times 10^{-7} \left( 1 - t_{mq} + 0.734 \times 10^{-3} A \right)^{1/2} + \xi^{-1} 7.185 \times 10^{-28} \times \left( 1 - t_{mq} + 0.734 \times 10^{-3} A \right) \times \exp \frac{57872}{A}$$
(13.21)

Taking  $s_{mq} = 1.5$  and  $\xi = 5 \times 10^{-10}$ , we have

$$A = 1 + z_{dec} = 1096.5, \tag{13.22}$$

$$T_{vdec} = AT_{\gamma 0} = 0.257 eV, \quad \chi = 0.004.$$
 (13.23)

This result is consistent with that of the Friedmann model  $T_{dec} = 0.25 eV$  and  $\chi = 0.004^{[17]}$ . It can be proved that  $z_{dec}$  is not susceptible for change of  $s_{mq}$  in the scope 1.1 - 1.7. For the cold-dark matter model, (13.11) still holds. Thus, provided  $\rho_{sl} = 0$  and  $s_{mq} = 1.1$  in (13.18) and (13.21), considering  $\rho_{vm} = \rho_B + \rho_{cd}$ ,  $\rho_{\gamma} = \rho_{v\gamma}$  and  $\rho_{sm} = \rho_{sM}$ , from (11.19) we can still obtain the result (13.22) – (13.23). In fact, in the case, there is only one sort of photon-like particles, i.e. ordinary photons and neutrinos. Hence  $Tveq \rightarrow T'veq = 2Tveq = 0.64eV$  so that  $0.734 \times 10^{-3} = (T_{v0}/Tveq) = (2.35 \times 10^{-4})/0.32$  becomes  $(T_{v0}/T'veq) = (2.35 \times 10^{-4})/0.64$ . In order to get the result (13.22) – (13.23), provided  $s_{mq} = 1.5 \rightarrow 1.1$ . Thus, we can still obtain the results above.

Sum up, we see that the primordial nucleosynthesis and CMBR can be explained based on this model and the F - W dark matter model or the cold dark matter model.

### D. Space is open

From (13.2) we see  $T_{vrec} = T_{rec}$ . On the other hand, as mentioned before, F - matter is same as the conventional matter and there are the same reheating process so that  $T_{vreh} = T_{reh}$ .

The interaction of s - Higgs fields and v - Higgs fields (2.10) may be neglected, because  $m(\Omega_s)$  and  $m(\Omega_v)$  are very large after reheating. Thus, there is no interaction except the repulsion or the gravitation among F - matter, W - matter and s - matter. The speed in F - matter is not influenced by W - matter and s - matter. On the other hand, F - matter and W - matter are symmetric and F - matter is identified as conventional matter. Hence we have

$$c_{vFs} = c_{vWs} \equiv c_v = c_s, \tag{13.24}$$

$$T_{vrec} = T_{rec}, \quad T_{vreh} = T_{reh} \tag{13.25}$$

Here  $c_{vFs} = \partial p_{vF} / \partial \rho_{vF}$  and  $c_{vWs} = \partial p_{vW} / \partial \rho_{vW}$  are the sound speeds in the F – matter or in the W – matter, respectively.

After reheating, temperature is so high that there are the F - plasma, the W - plasmaand the S - plasma. The F - plasma and the W - plasma are the same as the conventional plasma, because F - matter and W - matter are symmetric.

Let  $\Delta t_{vhc}$  be the duration in which  $T_{vreh}$  descends into  $T_{vrec}$  according to this model and  $\Delta t_{hc}$  be that according to the Friedmann model, there must be

$$\Delta t_{vhc} > \Delta t_{hc}. \tag{13.26}$$

The reasons are as follows.

 $\Delta t_{hc}$  is determined by the Friedmann model and the cold-dark matter model in the conventional theory. As mentioned before, according to the Friedmann model and the cold-dark matter model,  $\lambda_{eff} \ll \rho_B \ll \rho_\gamma$  when  $T \gtrsim T_d$ . Thus  $\lambda_{eff}$ ,  $\rho_B$  and  $\rho_{cd} \sim 6\rho_B$  may be neglected. Thus, (12.4) is reduced to

$$\widetilde{H}^2 \sim \eta \left(\rho_l + \rho_\gamma\right) = \eta \frac{\pi^2}{30} g^* T^4 \quad \text{when} T \gtrsim T_d, \tag{13.27}$$

$$\widetilde{H}^2 = \eta \left(\rho_B + \rho_{cd} + \rho_l + \rho_\gamma\right) \text{ when } T_d > T \gtrsim T_{rec}.$$
(13.28)

Thus  $\Delta t_{hc}$  is determined by (13.27) – (13.28). When  $T_d > T \gtrsim T_{rec}$ ,  $\rho_B$ ,  $\rho_{cd}$  and  $\rho_{sM}$  cannot be neglected because  $\rho_B$  and  $\rho_{sM} \propto R^{-3}$  and  $\rho_{\gamma} \propto R^{-4}$ 

 $\Delta t_{vhc}$  is determined by (11.1) and (12.10). As mentioned before, k may be neglected when  $T_v \gtrsim T_{rec} \sim 1 eV$ . Thus we can rewrite (11.1) as

$$H^{2}(T_{v}) = \eta [2(\rho_{vFM} + \rho_{vFl} + \rho_{vF\gamma}) - (\rho_{sM} + \rho_{sl})].$$
(13.29)

Considering  $\rho_B = \rho_{vFB}$ ,  $\rho_l = \rho_{vFl}$ ,  $\rho_{\gamma} = \rho_{vF\gamma}$  (12.10) and (11.11), neglecting  $\rho_{sM}$  and  $\rho_{vFM}$ , we again get

$$H^{2} = \eta \left( \rho_{vFl} + \rho_{vF\gamma} \right) = \eta \frac{\pi^{2}}{30} g^{*} T_{v}^{4} \text{ when } T_{v} \gtrsim T_{d}.$$
(13.30)

Considering  $T_v = T$ , we have

$$H = \tilde{H}$$
 when  $T \gtrsim T_d$ . (13.31)

From (11.8) - (11.9), (11.11) and (13.16) we have

$$2\rho_{vFM} = \rho_B + \rho_{cd}, \ \rho_{vFl} + \rho_{vF\gamma} = \rho_l + \rho_{\gamma}.$$
  
$$\rho_{sM} \sim 1.5 \times 2\rho_{vFM}.$$
 (13.32)

When  $T_d > T \gtrsim T_{rec}$ , from (11.14) – (11.15) and (12.10) we have

$$\rho_{vWl}(T) + \rho_{vW\gamma}(T) - \rho_{sl}(T) \lesssim 0.$$
(13.33)

Considering (13.32) - (13.33) and comparing (13.28) and (13.29), we find

$$H < H$$
 when  $T_d > T \gtrsim T_{rec}$ . (13.34)

Consequently, it is necessary due to (13.31) and (13.34) that  $T_{reh} = T_{vreh}$  can descend faster to  $T_{rec} = T_{vrec}$  according to the Friedmann model than that according to the present model and the F - W model. Thus, (13.26) must holds. Considering  $c_{vs} = c_s$ , we have

$$c_{vs} \bigtriangleup t_{vhc} = c_s \bigtriangleup t_{vhc} > c_s \bigtriangleup t_{hc}. \tag{13.35}$$

Based on  $\Delta t_{hc} = 3.8 \times 10^5 yr$ , space is regarded to be flat according to the conventional theory. But based on  $\Delta t_{vhc} > \Delta t_{hc}$  and  $c_{vs} = c_s$ , space is open (k < 0) according to the this model. This shows this model to be self consistent.

Combining the cold dark matter model and this model, we can still get the result similar to (13.35). The cold dark matter model is equivalent to  $\rho_{vm} = \rho_B + \rho_{cd}$ ,  $\rho_{vl} + \rho_{v\gamma} = \rho_l + \rho_{\gamma}$ ,  $\rho_{sM} = \rho_{sm}$ ,  $\rho_{vWl} + \rho_{vW\gamma} = 0$  so that  $\rho_{sl} = 0$ , and  $s_{sq} = \rho_{sm}/\rho_{vm} = 1.1$ . When  $T \gtrsim T_d$ , (13.30) still holds because  $\rho_{sm}$ ,  $\rho_B + \rho_{cd}$  may be neglected. When  $T_d > T \gtrsim T_{rec}$ , we have

$$H^{2} = \eta \left(\rho_{B} + \rho_{cd} + \rho_{vl} + \rho_{v\gamma} - \rho_{sm}\right).$$
(13.36)

It is seen from (13.28) and (13.36) that (13.34) still holds because  $\rho_{sm} \sim 1.1 \rho_{vm}$ .

# XIV. DYNAMICS OF STRUCTURE FORMATION AND THE DISTRIBUTIVE FORM OF THE SU(5) SINGLETS

#### A. The equations of structure formation

Generalizing the equations governing nonrelativistic fluid motion<sup>[8]</sup> to the model without singularity, in the V – *breaking*, considering (1) – (3), we have

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_v \cdot \nabla\right) \mathbf{v}_v = -\frac{\nabla p_v}{\rho_v} - \nabla \Phi, \qquad (14.1)$$

$$\frac{\partial}{\partial t}\rho_v + \nabla \cdot (\rho_v \mathbf{v}_v) = 0 \tag{14.2}$$

$$\nabla^2 \Phi = 4\pi G \left( \rho_v - \rho_s \right), \tag{14.3}$$

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_s \cdot \nabla\right) \mathbf{v}_s = -\frac{\nabla p_s}{\rho_s} + \nabla \Phi, \qquad (14.4)$$

$$\frac{\partial}{\partial t}\rho_s + \nabla \cdot (\rho_s \mathbf{v}_s) = 0, \qquad (14.5)$$

where  $\partial/\partial t + \mathbf{v}_v \cdot \nabla$  is call the convective derivative<sup>[8]</sup>. We can produce the linearized equations of motion by collecting terms of first order in perturbations about a homogeneous background  $\rho_v = \rho_{v0} + \delta \rho_v$  etc.. Letting

$$\mathbf{v}_v = \mathbf{v}_{v0} + \delta \mathbf{v}_v, \qquad \mathbf{v}_s = \mathbf{v}_{s0} + \delta \mathbf{v}_s \tag{14.6}$$

$$\rho_v = \rho_{v0} + \delta \rho_v, \qquad \rho_s = \rho_{s0} + \delta \rho_s, \tag{14.7}$$

$$\rho_v = \rho_{v0} + \delta\rho_v, \qquad \rho_s = \rho_{s0} + \delta\rho_s, \qquad (14.7)$$
$$\delta_v = \frac{\delta\rho_v}{\rho_{v0}}, \qquad \delta_s = \frac{\delta\rho_s}{\rho_{s0}}, \qquad (14.8)$$

where  $\mathbf{v}_0 = H\mathbf{x} = \left(\frac{\dot{a}}{a}\right)\mathbf{x}$  is the Hubble expansion. Now, for sufficiently small perturbations, we get

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_{v0} \cdot \nabla\right) \delta \mathbf{v}_v = -\frac{\nabla \delta p_v}{\rho_{v0}} - \nabla \delta \Phi - \left(\delta \mathbf{v}_v \cdot \nabla\right) \mathbf{v}_{v0},\tag{14.9}$$

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_{v0} \cdot \nabla\right) \delta_v = -\nabla \cdot \delta \mathbf{v}_v, \qquad (14.10)$$

$$\nabla^2 \Phi = 4\pi G \left( \rho_{v0} \delta_v - \rho_{s0} \delta_s \right), \qquad (14.11)$$

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_0 \cdot \nabla\right) \delta \mathbf{v}_s = -\frac{\nabla \delta p_s}{\rho_{s0}} + \nabla \delta \Phi - \left(\delta \mathbf{v}_s \cdot \nabla\right) \mathbf{v}_0, \tag{14.12}$$

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_0 \cdot \nabla\right) \delta_s = -\nabla \cdot \delta \mathbf{v}_s. \tag{14.13}$$

Defining the comoving spatial coordinates

$$\mathbf{x}(t) = a(t)\mathbf{r}(t), \quad \delta \mathbf{v}(t) = a(t)\mathbf{u}(t), \quad \nabla_x = \frac{\nabla_r}{a}.$$
(14.14)

(14.9) - (14.10) and (14.12) - (14.13) can be rewritten as

$$\dot{\mathbf{u}}_{v} + 2\frac{\dot{a}}{a}\mathbf{u}_{v} = \frac{\nabla\delta\Phi}{a^{2}} - \frac{\nabla\delta p_{v}}{\rho_{v0}}$$
(14.15)

$$\dot{\delta}_v = -\nabla \cdot \delta \mathbf{u}_v, \tag{14.16}$$

$$\dot{\mathbf{u}}_{s} + 2\frac{\dot{a}}{a}\mathbf{u}_{s} = -\frac{\nabla\delta\Phi}{a^{2}} - \frac{\nabla\delta p_{s}}{\rho_{s0}}$$
(14.17)

$$\dot{\delta}_s = -\nabla \cdot \delta \mathbf{u}_s, \tag{14.18}$$

Letting

$$\delta_{vk} = \delta_{vk} \left( t \right) \exp\left( -i\mathbf{k}_v \cdot \mathbf{r} \right), \quad c_v^2 = \frac{\partial p_v}{\partial \rho_v}, \tag{14.19}$$

$$\delta_{sk} = \delta_{sk} \left( t \right) \exp\left( -i\mathbf{k}_s \cdot \mathbf{r} \right), \quad c_s^2 = \frac{\partial p_s}{\partial \rho_s}, \tag{14.20}$$

from (14.15) - (14.20) and (14.11) we can get

$$\ddot{\delta}_{vk} + 2\frac{\dot{a}}{a}\dot{\delta}_{vk} = 4\pi G \left(\rho_{v0}\delta_{vk} - \rho_{s0}\delta_{sk}\right) - \frac{c_v^2 k_v^2}{a^2}\delta_{vk}, \qquad (14.21)$$

$$\ddot{\delta}_{sk} + 2\frac{\dot{a}}{a}\dot{\delta}_{sk} = 4\pi G \left(\rho_{s0}\delta_{sk} - \rho_{v0}\delta_{vk}\right) - \frac{c_s^2 k_s^2}{a^2}\delta_{sk},$$
(14.22)

Where dots stand for  $d/dt = (\partial/\partial t + \mathbf{v}_0 \cdot \nabla_r)$ . It is necessary that  $\delta_{sk} < 0$  when  $\delta_{vk} > 0$ and that  $\delta_{sk} > 0$  when  $\delta_{vk} < 0$ , because there is only repulsion between s – matter and v – matter. Consequently,

$$\rho_{v0}\delta_{vk} - \rho_{s0}\delta_{sk} = \rho_{v0}\delta_{vk} + \rho_{s0} \mid \delta_{sk} \mid,$$
(14.23)

$$\rho_{s0}\delta_{sk} - \rho_{v0}\delta_{vk} = \rho_{s0}\delta_{sk} + \rho_{v0} \mid \delta_{vk} \mid .$$
(14.24)

#### B. Three predictions.

1. Galaxies can form faster and earlier according to the present model than that according to the Friedmann model.

From (14.21) and (14.23) we see that  $\delta_{vk}(t)$  can grow faster than that determined by the Friedmann model because  $\rho_{v0}\delta_{vk} + \rho_{s0} | \delta_{sk} | > \rho_{v0}\delta_{vk}$ . The origin is the repulsion between s - matter and v - matter.

2. Galaxies distribution is not uniform in radial direction. There are two sphere layers in which the density of galaxies is relatively larger.

According to the present model, there is  $a_{\min}$ . If  $a_{\min}$  is so small that it may be neglected,  $v_0$  may be neglected as well. For simplicity,  $\rho_{s0}\delta_{sk}$  in (14.21) is neglected, because  $\rho_{v0}\delta_{vk}+\rho_{s0}$  |  $\delta_{sk}$  |>  $\rho_{v0}\delta_{vk}$ . Thus, in the period in which  $a_{\min}$  is may be neglected, for long-wavelength, i.e.  $(4\pi G\rho_{v0} - c_v^2 k_v'^2) > 0$ , from (14.21) and (14.23) we see that  $\delta_{vk}(t)$  can grow exponentially,

$$\overset{\cdot\cdot}{\delta_{vk}}(t) = \left(4\pi G\rho_{v0} - c_v^2 k_v'^2\right) \delta_{vk}, \qquad \delta_{vk}(t) = \exp(t/\tau), \qquad (14.25)$$

where  $\tau = 1/\sqrt{4\pi G \rho_{v0} - c_v^2 k_v'^2}$ ,  $k'_v = k_v/a$ . For observation, this implies that the density of galaxies and the brightness will be relatively larger in the sphere layer corresponding to  $\dot{a}_{\min} \sim 0$  than those in near other sphere layer. Of course, the density of galaxies on the sphere-surface is still uniform.

If  $(|\delta_{sk}|/\delta_{vk})$  in (14.23) is regarded as invariant approximately, we have

$$\tau = \left[4\pi G \left(\rho_{v0} + \rho_{s0} \mid \delta_{sk} \mid /\delta_{vk}\right) - c_v^2 k_v^{\prime 2}\right]^{-1/2}.$$
(14.26)

In the V-breaking, v-particles can form particles with large masses as v-atoms and v-molecules by the electroweak interaction. The density  $\rho_{v\gamma}$  of photon-like particles may be neglected and matter is dominative in low temperatures. In this stage, the velocities  $u'_v s$  of the particles with large masses may be neglected so that  $(\dot{a}/a) \cdot \mathbf{u}_v$  in (14.15) may be neglected as well. Thus, in the period in which  $\rho_{vm}$  is relatively larger and  $(\dot{a}/a) \cdot \mathbf{u}_v$  may be neglected, for long-wavelength, from (14.15) and (14.21) we get again the result similar (14.25).

Thus, galaxies distribution is not uniform in radial direction. There are two sphere layers in which the density of galaxies is relatively larger.

3. The s - SU(5) color single states must loosely distribute in space or form s - superclustering, i.e. huge v - voids.

There is no interaction except the gravitation among the s - SU(5) color single states, because SU(5) is a simple group and SU(5) symmetry holds in the V-breaking. There is no interaction except the repulsion between v - matter and s - matter. Consequently, the s - SU(5) color single states have decoupled after reheating, and can be regarded as ideal gas without collision. The ideal gas has the effect of free flux damping for clustering, i.e. the s - SU(5) color single states with very high velocities prevents  $\delta_{sk}(t)$  to increase. The velocities  $u_s$  of the s - SU(5) color single states must be very large and invariant, because their decoupling temperature, i.e. the temperature after the reheating, is very high, and there is no interaction except the repulsion or gravitation. Thus  $\delta_s = -\nabla \cdot \mathbf{u}_s$  cannot be small. Hence when a/a cannot be neglected,  $\left(a\delta_{sk}/a\right)$  in (14.22) cannot be neglected as well. The damping term  $\left(a\delta_{sk}/a\right)$  reduces  $\delta_{sk}(t)$  to grow. When  $\left(a/a\right)$  may be neglected, the perturbation  $\delta_{sk}$  will slowly grow in power rules. Thus, the s - SU(5) color single states must loosely distribute in space or form s - superclusters which are the huge v - voids for v - observers. In the huge v - voids,  $\rho_s \gg \rho_v$ , because of the repulsion between v - matterand s - matter. The features of the huge v-voids have been discussed before.

## XV. CONCLUSIONS OF THE EXPLANATION FOR SOME PHENOMENA

Based on the first model without singularity and the F - W dark matter model or cold dark matter model, the primordial nucleosynthesis and CMBR are explained. Based on the analysis of the baryon-elementary wave which began at reheating and ended at recombination, we see that space is open. This shows this model to be self-consistent. Generalizing equations governing nonrelativistic fluid motion to the present model, we have derived the equations of structure formation. Based on the equations, the following three predictions are obtained. Galaxies can form faster and earlier according to the present model than according to the Friedmann model; Galaxies distribution is not uniform in radial direction. There are two sphere layers in which the density of galaxies is relatively larger. SU(5) singlets must loosely distribute in space or form superclusters which are equivalent to huge voids for observers.

# Part III

# A cosmological model without singularity based on a new metric

# XVI. INTRODUCTION TO THE MODEL BASED ON A NEW METRIC

As mentioned in section 2, the existing probability of the S – breaking and the V – breaking must be equal, because the s – Higgs fields and the v – Higgs fields are symmetric. This equality can be realized by the following two sorts of modes.

(1) The whole universe is in the same breaking (e.g. the S - breaking), but this sort of breaking (the S - breaking) can transform to the other (the V - breaking) as space contracts to the least scale  $R_{\min}$ .

In this case, a hypersurface of simultaneity must be complete as the same as the Friedmann model. The 3-dimensional volume of a complete hypersphere is finite, and the 3-dimensional volume of a complete hyperplane or hyper-hyperboloid is infinite. Consequently, if the density of matter is finite and the cosmological principle holds, k cannot be changeable, because a complete hypersphere cannot transform into a complete hyperplane or hyper-hyperboloid in the case. This mode has been discussed in the first part.

(2) There are the two sorts of breaking in the universe simultaneously. Consequently, the universe must be composed of infinite s-cosmic islands with the S-breaking and v-cosmic islands with V-breaking. Every cosmic island must be finite. Thus, the supersurface of a cosmic island must be incomplete and the radial coordinate of a cosmic island is finite. For example, we may take the maximum of the radial coordinate to be  $r_{max} \leq 1$  as the same as this case k = 1 in the Friedmann model. This implies that not only is the maximum of the radial coordinate of a supersphere finite, but also the maximum of the radial coordinate of a superplane or a hyper-hyperboloid is finite and may be taken as 1. We discuss the first sort of modes in the present part.

In this case, not that we describe the evolution of the whole universe, but that we describe approximately the evolution of a cosmic island. In fact, in contrast with the conventional theory, the universe as a whole does not change in essence according to the present model, although every cosmic island is changing (expands, contracts or inflates).

We will see that the probability that particles get out of a cosmic island is very little. Consequently a cosmic island seems to be the whole cosmos for an observer inside it.

In this case, the cosmological principle does no longer hold strictly, the RW metric is no longer applicable, and a new metric is necessary for the incomplete hypersurfaces.

The inferences of the first model can still be obtained by the second model. The differences between the two model are follows.

1. In addition to conjectures 1 and 2 in the first model, the second model increases a new conjecture about incomplete supersurfaces.

2. There are new inferences.

A. The universe must be composed of infinite s - cosmic islands, v - cosmic islands and transitional regions.

B. Some huge redshifts (e.g. the big redshifts of quasi-stellar objects) possibly are 'the mass redshifts' which is caused by less mass  $m_{eT}$  of an electron than given  $m_e$ .

C. It is possible that a v-black hole with its big enough mass and density can transform into a huge white hole in the V-breaking or the S-breaking.

In section 17, the conjecture about incomplete supersurfaces are presented; in section 18, in section 6, contraction of space, the highest temperature and inflation of space are

discussed; in section 7, some new predictions are given; section 8 is the conclusion of the part.

## XVII. CONJECTURES

#### A. Conjectures

In addition to conjectures 1 and 2, the following conjecture is necessary for the present model. In order to describe the conjecture, we first define incomplete hypersurfaces of simultaneity.

**Definition 4** Such hypersurfaces of simultaneity are called incomplete, if a hypersphere has one hole, and a hyperplane or a hyper-hyperboloid is finite.

**Conjecture 5** The metric describing incomplete hypersurfaces can be obtained by the replacement of the curvature factor k in the RW metric by  $K = K\left(\underline{\rho}_g\right)$  which is a monotone and finite function of  $\underline{\rho}_g$  and  $dK/d\underline{\rho}_g > 0$ . Here  $\underline{\rho}_g$  is the gravitational mass density in the comoving coordinates.

In this case, K does no longer describe the quality of an incomplete supersurface in whole and only describe differential quality of the incomplete supersurface. K corresponding to a cosmic island can change from K > 0 to K = 0 or K < 0, because a cosmic island is finite and its gravitational mass density  $\rho_g$  can change from  $\rho_g > 0$  to  $\rho_g = 0$  or < 0.

#### B. The new metric is applicable to incomplete hypersurfaces of simultaneity

The only difference between the new metric and the RW metric is that K is changeable in the new metric.

A few explanations for the conjecture as follows:

(1) The basic promise of a hypersurface of simultaneity to be complete is that the universe evolves as a whole. If the universe is composed of infinite s - cosmic islands and v - cosmic islands which evolve individually, the hypersurface of every cosmic island must be incomplete.

k in the RW metric cannot change from k = 1 to 0 and -1 for complete hypersurfaces because of the relative topological theorems. But the topological theorems are not applicable to the incomplete supersurfaces. In fact, there is no the same breaking, no unified evolution, and no the same features for the cosmic islands. Hence the metrices of the cosmic islands must be different from each other.

k can describe the topology feature for complete hypersurfaces. But  $K\left(\underline{\rho}_{g}\right)$  describes only the differential feature for the incomplete hypersurfaces.

The radial coordinate r in the spherical coordinates is finite for an incomplete hypersurface of simultaneity and  $r_{\text{max}}$  satisfies  $r_{\text{max}}^2 \leq 1/K_{\text{max}}$  ( $K_{\text{max}} > 0$  in this model), because a cosmic island must be finite.

(2) In the Friedmann model the equation of R is regarded as a first integral of the equation of R because of the matter conservation equation  $\dot{\rho} = -3(\rho + p)R/R$ . In this model, the equation of  $\dot{\rho}$  becomes the equation of  $\dot{\rho}_g$  (18.50) or (18.30) which is more complex, because  $K \neq 0$ . The equation (18.50) or (18.30) does not contradict any known experiment and observation, because  $\rho_g$  is the gravitational mass density, rather than mass density. The gravitational mass may be not conservational, although energy is still conservational. In other words,  $K \neq 0$  and energy conservation are compossible. On the other hand, from (18.49), (18.44) and (18.41) we see that the the conservation equation of  $\rho_g$  holds approximately when  $K \sim \partial K/\partial r \sim 0$ .

(3) Let  $h_{ab}(t, x^i)$  be the induced metric of the space-time metric  $g_{ab}$  on a homogeneous hypersurface. Only when the uniqueness of the homogeneous hypersurface clan holds, the variables t and  $x^i$  of  $h_{ab}(t, x^i)$  can be separated, i.e.  $h_{ab}(t, x^i) = a^2(t) \hat{h}_{ab}(x^i)$ , so that the metric can be written as the RW metric<sup>[19]</sup>. When  $\rho_g = 0$  or a hypersurfaces is incomplete, the uniqueness of the homogeneous hypersurface clan cannot hold. Thus, t and  $x^i$  in  $h_{ab}(t, x^i)$  cannot be separated completely and the new metric  $(K = K(t, x^i))$  is necessary.

In fact, only when  $\rho_g \sim 0$ ,  $K \neq 0$ . When  $\underline{\rho}_g \neq 0$  so that  $K \sim 0$ , the evolution equations of the present model are consistent with those of the Friedmann model in form.

# XVIII. EVOLUTION EQUATIONS OF SPACE

The only difference between the two models is the curvature factors in them to be different from each other. Hence (2.1) - (3.12) are still applicable to the second model, because they

are independent of the curvature factor.

# A. The evolution equations in a new metric

Based on the new metric described by conjecture 3, we have

$$(ds)^{2} = -(dt)^{2} + R^{2}(t) \left\{ \frac{(dr)^{2}}{1 - K(\underline{\rho}_{g})r^{2}} + (rd\theta)^{2} + (r\sin\theta d\varphi)^{2} \right\}.$$
 (18.1)

(18.1) implies  $g_{00} = -1$ ,  $g_{11} = R^2(t) / (1 - K(\underline{\rho}_g)r^2)$ ,  $g_{22} = R^2(t)r^2$ ,  $g_{33} = R^2(t)r^2\sin^2\theta$ and the others to be zero. Form this we can determine  $\Gamma^{\lambda}_{\mu\nu}$  and  $R_{\mu\nu}$ ,

$$R_{00} = g_{00} \left\{ \frac{3\ddot{R}}{R} + \frac{r^2}{2\left(1 - Kr^2\right)} \left[ \frac{2\ddot{R}\ddot{K}}{R} + \ddot{K} + \frac{3r^2\ddot{K}}{2\left(1 - Kr^2\right)} \right] \right\},$$
(18.2)  
$$R_{11} = g_{11} \left\{ \frac{2\ddot{R}}{R^2} + \frac{2K}{R^2} + \frac{\ddot{R}}{R} \right\}$$

$$R_{22} = g_{22} \left[ \frac{2R}{R^2} + \frac{2K}{R^2} + \frac{R}{R} + \frac{r^2}{2(1 - Kr^2)} \frac{RK}{R} + \frac{r\partial K}{2R^2\partial r} \right],$$
 (18.4)

$$R_{33} = g_{33} \left[ \frac{2R}{R^2} + \frac{2K}{R^2} + \frac{R}{R} + \frac{r^2}{2(1 - Kr^2)} \frac{RK}{R} + \frac{r\partial K}{2R^2\partial r} \right],$$
(18.5)

$$R_{01} = R_{10} = \frac{r\dot{K}}{(1 - Kr^2)}$$
, the others are zero. (18.6)

Matter in the universe may be regarded approximately as ideal gas distributed evenly in the whole space. The energy-momentum tensor densities of the ideal gas are

$$T_{aM\mu\nu} = (\rho_a + p_a) U_{a\mu} U_{a\nu} + p_a g_{\mu\nu}, \ a = s, \ v.$$
(18.7)

Considering the potential energy densities in (2.14) - (2.15), we can rewrite (18.7) as

$$\widetilde{T}_{a\mu\nu} = \left[\widetilde{\rho}_a + \widetilde{p}_a\right] U_{\mu} U_{\nu} + \widetilde{p}_a g_{\mu\nu}, \qquad (18.8)$$

$$\widetilde{\rho}_a = \rho_a + \widetilde{V}_a(\varpi_a), \quad \widetilde{p}_a = p_a - \widetilde{V}_a(\varpi_a).$$
(18.9)

From (2.17) - (2.18) we have

$$\widetilde{V}_{s}(\varpi_{s}) = V_{s}(\varpi_{s}) + V_{0}, \quad \widetilde{V}_{v}(\varpi_{v}) = V_{v}(\varpi_{v}), \quad (18.10)$$

$$T_{Sg\mu\nu} = \widetilde{T}_{s\mu\nu} - \widetilde{T}_{v\mu\nu}.$$
(18.11)

in the S-breaking, and

$$\widetilde{V}_{s}(\varpi_{s}) = V_{s}(\varpi_{s}), \quad \widetilde{V}_{v}(\varpi_{v}) = V_{v}(\varpi_{v}) + V_{0}, \quad (18.11)$$

$$T_{Vg\mu\nu} = \widetilde{T}_{\nu\mu\nu} - \widetilde{T}_{s\mu\nu}, \qquad (18.13)$$

in the V - breaking. In fact,  $V_v = 0$  because  $\langle \omega_v \rangle = 0$  in the S - breaking, hence  $\tilde{\rho}_v = \rho_v$ and  $\tilde{p}_v = p_v$ . Similarly,  $V_s = 0$ ,  $\tilde{\rho}_s = \rho_s$  and  $\tilde{p}_s = p_s$  in the V - breaking. In the present model, the 4-velocities are

$$U^{\mu} = \frac{dx^{\mu}}{d\tau}, \ x^{1} = r, \ x^{2} = \theta, \ x^{3} = \varphi,$$
(18.14)

$$U^{1} = \frac{dr}{d\tau} = \frac{q}{\sqrt{1 - v^{2}}}, \ U^{2} = U^{3} = 0,$$
(18.15)

$$q = \frac{dr}{dt}, \quad v = q\sqrt{\frac{R^2}{1 - Kr^2}},$$
 (18.16)

$$U_0 = g_{0\nu}U^{\nu} = g_{00}U^0 = -U^0 = \frac{-1}{\sqrt{1 - v^2}},$$
(18.17)

$$U_1 = g_{1\nu}U^{\nu} = \sqrt{\frac{R^2}{1 - Kr^2}} \cdot \frac{v}{\sqrt{1 - v^2}}.$$
(18.18)

Let

$$S_{g\mu\nu} \equiv T_{g\mu\nu} - \frac{1}{2}g_{\mu\nu}T_g, \ T_g = T_{g\lambda}^{\lambda} = -\widetilde{\rho}_g + 3\widetilde{p}_g,$$
(18.19)

$$S_{g00} = \frac{1}{2} \left( \tilde{\rho}_g + 3\tilde{p}_g \right) + \frac{v^2}{1 - v^2} \left( \tilde{\rho}_g + \tilde{p}_g \right),$$
(18.20)

$$S_{g11} = g_{11} \left[ \frac{1}{2} \left( \tilde{\rho}_g - \tilde{p}_g \right) + \frac{v^2}{1 - v^2} \left( \tilde{\rho}_g + \tilde{p}_g \right) \right],$$
(18.21)

$$S_{g22} = g_{22} \frac{1}{2} \left( \tilde{\rho}_g - \tilde{p}_g \right), \quad S_{33} = g_{33} \frac{1}{2} \left( \tilde{\rho}_g - \tilde{p}_g \right), \tag{18.22}$$

$$S_{g01} = S_{g10} = \sqrt{\frac{R^2}{1 - Kr^2} \cdot \frac{-v}{1 - v^2}} \left(\tilde{\rho}_g + \tilde{p}_g\right).$$
(18.23)

the others are all zero. Substituting (18.2) - (18.23) into (2.16), we obtain

$$\dot{R}^{2} + K + \frac{r^{2}}{3(1 - Kr^{2})}RRK + \frac{r\partial K}{3\partial r}$$

$$= \eta \left[\rho_{g} + V_{g} + \frac{v^{2}}{1 - v^{2}}(\rho_{g} + p_{g})\right]R^{2}, \ \eta \equiv 8\pi G/3,$$
(18.24)

$$\ddot{R} - \frac{r^2}{6(1 - Kr^2)} \dot{R} \dot{K} - \frac{r\partial K}{6R\partial r} = -\frac{\eta}{2} \left[ (\rho_g + 3p_g) - 2V_g + \frac{4v^2}{1 - v^2} (\rho_g + p_g) \right] R.$$
(18.25)

$$\frac{R^2 r^2}{(1 - Kr^2)} \left[ \frac{3RK}{R} + \ddot{K} + \frac{3r^2 K^2}{2(1 - Kr^2)} \right] + \frac{r\partial K}{\partial r} \\
= 6\eta \frac{v^2}{1 - v^2} \left(\rho_g + p_g\right) R^2,$$
(18.26)

$$\frac{rK}{1 - Kr^2} = 3\eta \frac{(-v)}{1 - v^2} \sqrt{\frac{R^2}{1 - Kr^2}} \left(\rho_g + p_g\right), \qquad (18.27)$$

In the S-breaking,

$$\rho_g = \rho_s - \rho_v, \quad V_g = V_s + V_0 - V_v, \quad p_g \equiv p_s - p_v. \tag{18.28}$$

In the V-breaking,

$$\rho_g = \rho_v - \rho_s, \quad V_g = V_v + V_0 - V_s, \quad p_g \equiv p_v - p_s.$$
(18.29)

From (18.24) - (18.27) we get

$$\dot{K}\left\{1 + \frac{r^{2}}{1 - Kr^{2}} \cdot \left[\frac{K}{3} + \frac{r}{6}\frac{\partial K}{\partial r} - \frac{\eta}{2}R^{2}\left(1 + \frac{2v^{2}}{1 - v^{2}}\right)(\rho_{g} + p_{g})\right]\right\} = \eta\left\{R^{2}\frac{d}{dt}\left[\rho_{g} + V_{g} + \frac{v^{2}}{1 - v^{2}}(\rho_{g} + p_{g})\right] + \left(3 + \frac{4v^{2}}{(1 - v^{2})}\right)(\rho_{g} + p_{g})R\dot{R}\right\}$$
(18.30)

#### **B.** The asymptotic solution of the equations (18.24) - (18.27)

### 1. The asymptotic solution of K(t,r)

As mentioned in the section above, the radial coordinate r must be finite and  $r_{\text{max}}^2 < 1/K_{\text{max}}$ . K(t,r) in (18.24) – (18.27) has such an asymptotic solution in the following form,

$$K = \frac{D - D_0}{1 + S(r, D)},\tag{18.31}$$

$$S(r,D) \equiv (r^{2}-1) \left[1 - \exp(-Ar^{2})\right], \ A \equiv \frac{1-D^{2}}{D^{2}(t) + \delta^{2}}$$
(18.32)

$$D\left(\underline{\rho}_{g}\right) \equiv \frac{2}{\pi} \arctan Q\left(\underline{\rho}_{g}\right), \ \underline{\rho}_{g} = V_{g} + \frac{\rho_{g}R^{3}}{R_{1}^{3}}, \tag{18.33}$$

where  $\underline{\rho}_g$  is the gravitational mass density in the comoving coordinates,  $Q\left(\underline{\rho}_g\right)$  is a monotonously increasing function of  $\underline{\rho}_g$ , and  $0 \leq D_0 < 1$ ,  $\delta$  is a small positive parameter.  $Q\left(\underline{\rho}_g\right)$  can be so chosen that

$$Q(0) = 0, \quad dQ/d\underline{\rho}_g > 0,$$

$$Q\left(\underline{\rho}_g\right) \longrightarrow \infty \text{ when } \underline{\rho}_g \longrightarrow \infty,$$

$$Q\left(\underline{\rho}_g\right) \longrightarrow -\infty \text{ when } \underline{\rho}_g \longrightarrow -\underline{\rho}_{g0}/A_{\rho},$$
(18.34)

e.g.

$$Q\left(\underline{\rho}_{g}\right) = \ln \frac{A_{\rho}\underline{\rho}_{g} + \underline{\rho}_{g0}}{\underline{\rho}_{g0}},\tag{18.35}$$

where  $\underline{\rho}_{gc}$  and  $A_{\rho}$  are all positive parameters. In the present model,  $\underline{\rho}_{g}$  must be finite in any case. We will see  $V_0 \geq \underline{\rho}_g > -\underline{\rho}_{g0}/A_{\rho}$ .

In general,  $\underline{\rho}_g = \underline{\rho}_g(r, t)$  so that D = D(r, t). In fact, we will see that  $\underline{\rho}_g \simeq \underline{\rho}_g(t)$  in many cases, because the cosmological principle holds approximately. Thus, we take  $\underline{\rho}_g = \underline{\rho}_g(t)$  and D = D(t) as an approximate solution in the following.

 $D\left(\underline{\rho}_{g}\right)$  is a monotone and finite function of  $\underline{\rho}_{g}$  due to (18.31) - (18.35),

$$D \sim 0$$
 provided  $\underline{\rho}_g \sim 0$ ,  
 $K \sim -\frac{D_0}{r^2}$  only when  $r$  is large and  $\underline{\rho}_g \sim 0$ , (18.36)

$$K \sim -D_0$$
, when  $r \sim 0$  and  $\underline{\rho}_g \sim 0$  (18.37)

$$K \longrightarrow \pm 1 - D_0, \text{ when } Q\left(\underline{\rho}_g\right) \longrightarrow \pm \infty.$$
 (18.38)
It is considered in (18.36) – (18.37) that  $V_g = 0$  when  $\underline{\rho}_q \sim 0$ , because  $V_g \sim 0$  after reheating.

It is seen from (18.36) - (18.38) that K(r,t) is dependent on r only when r is very large and  $\underline{\rho}_g \sim 0$ . In this case,  $K \sim (r\partial K/\partial r) \sim -D_0/r^2 \sim 0$  and  $(rK) \sim D/r \sim 0$  may still be neglected, because r is very large. Thus (18.24) - (18.27) is still approximately independent of r.

#### 2. Discussion of K

1.  $K \sim 0$  when  $Q^2 \longrightarrow \infty$ .

For monotonously expanding space,  $\underline{\rho}_g(t)$  is a monotone decreasing function of t, hence  $\underline{\dot{\rho}}_g \lesssim 0$ . Considering  $D\left(\underline{\rho}_g\right)$  and K are all monotonously increasing functions of  $\underline{\rho}_g$  since  $dQ/d\underline{\rho}_g = (2/\pi) a/\left(\underline{\rho}_g + \underline{\rho}_{g0}/A_{\rho}\right) > 0$ , from (18.31) – (18.35) we have

$$\overset{\cdot}{D} = \frac{2}{\pi} \frac{dQ/d\underline{\rho}_g}{1+Q^2} \overset{\cdot}{\underline{\rho}_g} \lesssim 0, \qquad (18.39)$$

$$\dot{K} = \frac{D}{1+S} \left[ 1 + r^2 \left( r^2 - 1 \right) \frac{K}{1+S} \frac{2D \left( 1 + \delta^2 \right)}{\left( D^2 + \delta^2 \right)^2} \exp \left( -Ar^2 \right) \right],$$
  
$$\dot{K} \sim \frac{\dot{D}}{1+S}, \text{ when } D, K, \text{ or } r \sim 0, \text{ or } Ar^2 \text{ is large}$$
(18.40)

Considering K, S and D are all finite, from (18.31) - (18.32) we have

$$\ddot{K} \sim 0$$
, provided  $D \sim 0$  or  $S$  is large, (18.41)  
 $\ddot{K} \sim 0, \ \ddot{D} \sim 0, \text{ provided } \dot{\underline{\rho}}_g \sim 0$   
and  $\ddot{\underline{\rho}}_g \sim 0 \text{ or } Q^2 \longrightarrow \infty.$  (18.42)

D, D, K and K must be finite.

2.  $r\partial K/\partial r \sim 0$ , provided  $A \sim 0$ , or  $r \sim 0$ , or  $K \sim 0$ .

From (18.31) - (18.32) we have

$$\frac{r\partial K}{\partial r} = -\frac{2Kr^2}{(1+S)} \{1 - \left[1 + A\left(1 - r^2\right)\right] \exp\left(-Ar^2\right)\}.$$
(18.43)

From (18.43), and (18.26) - (18.27) we

$$\frac{r\partial K}{\partial r} \sim 0, \text{ provided } A \sim 0, \text{ or } r \sim 0, \text{ or } K \sim 0,$$
  
or  $\ddot{K} \sim \ddot{K} \sim 0.$  (18.44)

When r is very large and  $\underline{\rho}_g \sim 0$ 

$$\frac{r\partial K}{\partial r} \sim -\frac{2D_0}{r^2}.\tag{18.45}$$

In the case,  $r\partial K/\partial r$  may still be neglected, when  $D_0/r^2 \ll 1$ . It is seen that (18.31) is indeed an asymptotic solution of K(t, r).

Sum up, it is seen that we may take  $r\partial K/\partial r \sim 0$  so that  $K(r,t) \sim K(t)$  approximately.

3. Reduction of (18.24) - (18.27) and (18.30) when  $Q^2$  is very large.

As mentioned above,  $K \sim \pm 1 - D_0$  and  $r \partial K / \partial r \sim K \sim 0$  provided  $Q^2 \longrightarrow \infty$ , and  $K \sim D(t) - D_0$  when  $r \sim 0$ . In the two cases, , from (18.26) - (18.27) we get

$$\frac{(-v)}{1-v^2}\left(\rho+p\right) \sim 0, \ \frac{r\partial K}{\partial r} \sim 0.$$
(18.46)

and (18.24) - (18.25) is reduced to

$$\dot{R}^2 + K = \eta \left[ \rho_g + V_g \right] R^2, \tag{18.47}$$

$$\ddot{R} = -\frac{\eta}{2} \left[ (\rho_g + 3p_g) - 2V_g \right] R.$$
(18.48)

When  $Q^2 \longrightarrow \infty$ , (18.30) is reduced to

$$R^{2} \frac{d}{dt} \left[ \rho_{g} + V_{g} \right] + 3 \left( \rho_{g} + p_{g} \right) R\dot{R} = 0.$$
(18.49)

The equations are independent of r.

When  $r \sim 0$ , (18.30) is reduced to

$$\dot{D} \sim \dot{K} = \eta \{ R^2 \frac{d}{dt} \left( \rho_g + V_g \right) + 3 \left( \rho_g + p_g \right) R R \} < 0.$$
(18.50)

When  $\underline{\rho}_g \sim 0$  and  $r \sim 0$ , considering  $V_g = 0$  after reheating and  $p_g = \rho_{\gamma g}/3 > 0$  when  $\rho_g = 0$ , from (18.24) - (18.25) and (18.31) - (18.33) we have

$$\overset{\cdot}{R}^{2} = D_{0} > 0, \, \ddot{R}/R = -(3\eta/2) \, p_{g} < 0.$$
 (18.51)

4. Reduction of (18.24) - (18.27) and (18.30) when  $\underline{\rho}_{q} \sim 0$ .

The results (18.46) - (18.49) still hold for very large r, provided  $Q^2$  is large enough. But the results do no long hold for very large r and  $\underline{\rho}_g \sim 0$  due to Q(0) = 0. When  $\underline{\rho}_g \sim 0$ and r is very large,  $D\left(\underline{\rho}_g\right) \sim 0$  and  $K \sim -D_0/r^2$  due to (18.37). From (18.39) we have  $\dot{D} = (2/\pi) \left(A_{\rho}/\rho_{g0}\right) \dot{\underline{\rho}}$  and  $\dot{K} = \dot{D}/r^2$  when  $\underline{\rho}_g \sim 0$ . Thus,  $r\partial K/\partial r \sim K \sim -D_0/r^2 \sim 0$  and  $(rK) \sim D/r \sim 0$  may be neglected when  $\underline{\rho}_g \sim 0$  and r is very large. Considering  $V_g = 0$  after reheating, (18.24) - (18.27) is reduced to

$$\left(\frac{\dot{R}}{R}\right)^2 \simeq -\frac{\dot{D}}{3\left(1+D_0\right)}\frac{\dot{R}}{R} + \eta\rho_g,\tag{18.52}$$

$$\frac{\ddot{R}}{R} \simeq \frac{D\dot{R}}{6(1+D_0)R} - \frac{3\eta}{2}p_g,$$
(18.53)

$$3\frac{\dot{R}\dot{D}}{R} + \ddot{D} + \frac{3\dot{D}}{2(1+D_0)} \simeq 0, \qquad (18.54)$$

$$0 \simeq \frac{(-v)}{1 - v^2} p_g,\tag{18.55}$$

 $\dot{R} > 0$  because space is expanding. Hence

$$D < 0, \ \underline{\dot{\rho}} < 0, \ \ddot{R} < 0 \text{ when } \underline{\rho}_g \sim 0.$$
 (18.56)

due to (18.52) - (18.53).

When  $\underline{\rho}_g \sim 0$  and  $r \sim 0$ , we have  $D\left(\underline{\rho}_g\right) \sim 0$ ,  $K \sim -D_0$ ,  $K = D = (2/\pi) \left(\frac{A_{\rho}}{\rho_{g0}}\right) \underline{\dot{\rho}}$ , and  $r\partial K/\partial r \sim \left(r\dot{K}\right) \sim 0$  may be neglected. Thus (18.24) – (18.25) is reduced to

$$\dot{R}^2 \simeq D_0 + \eta \rho_g > 0, \quad \ddot{R} \simeq -\frac{3\eta}{2} p_g R \simeq -\frac{\eta}{2} \rho_{\gamma g} R < 0.$$
 (18.57)

In this case, (18.56) still holds. When  $\underline{\rho}_{q} \sim 0$  and r is large,

$$\dot{\rho}_g = -\frac{3\dot{R}}{2R}p_g = \frac{D}{2\left(1+D_0\right)}p_g < 0.$$

When  $\underline{\rho}_g \sim 0$  and  $r \sim 0$ ,

$$\dot{\rho}_g = -3p_g R/R + D/\eta R^2 < 0.$$

5. It is possible that  $\ddot{R} > 0$  when R = 0.

As mentioned above, when  $\underline{\rho}_g \sim 0$ , we have  $\ddot{R} < 0$  and  $\dot{R} > 0$  so that  $\dot{\rho}_g < 0$ . Hence space will continue to expand with a deceleration and  $\underline{\rho}_g$  will become negative.

Let  $t = t_2$  when  $\underline{\rho}_g(t_2) = 0$ . Because of (18.56),  $\underline{\dot{\rho}}_g < 0$  so that  $\underline{\rho}_g(t) = \underline{\rho}_g(t_2) + \underline{\dot{\rho}}_g(t-t_2) < 0$  when  $t > t_2$ . Thus R(t) = 0 is possible when  $t = t_1 > t_2$ , because R < 0 when  $t = t_2$  and  $\underline{\rho}_g(t) < 0$ .

When  $R(t_1) = 0$ , from (18.24) - (18.25) we have

$$\ddot{R} = -\frac{K}{2R} - \frac{3\eta}{2} \left[ p_g + \frac{v^2}{1 - v^2} \left( \rho_g + p_g \right) \right] R.$$
(18.58)

From (18.31) – (18.35), (18.40) and (18.43) we see that when  $A_{\rho}\underline{\rho}_g/\underline{\rho}_{g0} \sim -1$ ,  $K \sim -1 - D_0$ ,  $Q^2 \longrightarrow \infty$ ,  $\dot{K} \sim 0$  and  $r\partial K/\partial r \sim 0$ . Thus,  $r\dot{K}$  and  $r\partial K/\partial r$  may be neglected when  $\left(-A_{\rho}\underline{\rho}_g/\underline{\rho}_{g0}\right)$  is large. Consequently, (18.58) may be reduced to

$$\ddot{R} = -\frac{K}{2R} - \frac{3\eta}{2} p_g R \simeq -\frac{K}{2R} - \frac{\eta}{2} \rho_{\gamma g} R.$$
(18.59)

It i.e. seen that all  $\ddot{R} > 0$ , = 0 and  $\ddot{R} < 0$  are possible when  $\dot{R}(t_1) = 0$ , because  $K \sim -1 - D_0 < 0$ . If  $\ddot{R}(t_1) < 0$ , space will contract with a deceleration when  $t > t_1$ . If  $\ddot{R}(t_1) = 0$ , space will be static when  $t > t_1$ . If  $\ddot{R}(t_1) > 0$ , space will expand with an acceleration when  $t > t_1$ . This is because  $\underline{\rho}_g$  will decrease as space expands (see (18.65)). We will discuss the case  $\ddot{R}(t_1) > 0$  in the present paper.

6. A conservation equation.

It is impossible that the s-particles transforms into the v-particles when temperature is low. When temperature is high enough  $(T \gtrsim T_{cr})$ , see the following), it is possible that  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  so that the masses of all particles are zero. Thus, the s-particles and the v-particles can transform from one to another by (2.10). We discuss (18.49) as follows.

As mentioned above,  $K \sim 0$  when  $Q^2 \longrightarrow \infty$ . In the case, (18.49) is a conservation equation. Let  $\rho_g = \rho_{mg} + \rho_{lg} + \rho_{\gamma g}$ ,  $p_{mg} \sim 0$  and  $p_{lg} = \kappa \rho_{lg}$  ( $0 < \kappa < 1/3$ ) for a given temperature. For photon-like particles,  $p_{\gamma g} = \rho_{\gamma g}/3$ . Here  $\rho_m$  is the mass density of the particles with so large masses that  $\rho_m \gg p_m$ ,  $\rho_l$  is the mass density of the particles whose masses are small, but cannot be neglected for the given temperature. From (18.49) we have

$$-R^{2}\frac{d}{dt}V_{g} = \frac{1}{R}\frac{d}{dt}\left(\rho_{mg}R^{3}\right) + \frac{1}{R^{1+3\kappa}}\frac{d}{dt}\left(\rho_{\lg}R^{3(1+\kappa)}\right) + \frac{1}{R^{2}}\frac{d}{dt}\left(\rho_{\gamma g}R^{4}\right).$$
(18.60)

 $V_g = dV_g/dt = 0$  after reheating. (18.60) may be regarded as the conservation equation of the gravitation mass. The transformation of the s - particles and the v - particles may be neglected when temperature is low. After particles decouple, from (18.60) we have

$$\rho_{sm}R^{3} = C_{sm}, \ \rho_{vm}R^{3} = C_{vm}, \ \rho_{sl}R^{3(1+\kappa)} = C_{sl},$$
  
$$\rho_{vl}R^{3(1+\kappa)} = C_{vl}, \ \rho_{s\gamma}R^{4} = C_{s\gamma}, \ \rho_{v\gamma}R^{4} = C_{v\gamma},$$
(18.61)

 $C_{sm}$  etc. are constants. It is seen that the individual conservation of s-energy and v-energy holds only approximately in low temperatures. In the S - breaking,  $\rho_{v\gamma} = 0$  because the masses of all v - SU(5) color-single states are not zero. Similarly, in the V - breaking,  $\rho_{s\gamma} = 0$ .

When K cannot be neglected, the conservation equation (18.49) is no long held. From (18.50) we see that there must be  $\rho_g \neq 0$  when  $K \neq 0$ . This does not contradict any known experiment and observation, because  $\rho_g = \rho_v - \rho_s$  or  $= \rho_s - \rho_v$ , rather than energy density  $\rho = \rho_v + \rho_s$ . Although the gravitational mass is not conservational, the total energy, whose density is  $\rho$ , can still be conservational. Consequently  $K \neq 0$  and energy conservation are compossible.

7. K cannot be neglected in the period from K = 0 to  $\rho_g = 0$ .

Let  $D\left(\underline{\rho}'_g\right) - D_0 \sim 0$ , then  $K \sim 0$  and  $\underline{\rho}_g > 0$  due to (18.31) and (18.33), but  $K \neq 0$ so that v = v(r,t) cannot be neglected due to (18.27). In the case, (18.24) - (18.27) is dependent on r. (18.24) - (18.27) is independent of r when  $\rho_g \sim 0$  due to (18.52) - (18.57). When  $D\left(\underline{\rho}_g\right) > D_0$  due to  $\underline{\rho}_g > \underline{\rho}'_g$ , K > 0 and Q is large. In the case, as mentioned above, K and  $r\partial K/\partial r$  may be neglected and (18.24) - (18.27) is reduced to (18.46) - (18.49).

8. The definition of  $\underline{\rho}_{a}$ .

Letting  $(\rho_{mg1} + \rho_{\lg 1} + \rho_{\gamma g1}) \neq 0$   $(\rho_{gm1} \equiv \rho_{gm}(t_1)$  etc.), considering the Higgs potential, we define  $\underline{\rho}_g$  as

$$\underline{\rho}_{g} = V_{g} + (\rho_{mg} + \rho_{\lg} + \rho_{\gamma g}) R^{3} / R_{1}^{3}, \qquad (18.62)$$

where  $\rho_{bg} = \rho_{sb} - \rho_{vb}$ , b = m, l or  $\gamma$ ,  $\rho_{v\gamma g} = 0$  and  $V_g = V_s + V_0 - V_v$  in the S - breaking. When  $T \gtrsim T_{cr}$ , then  $\rho_g = \rho_s - \rho_v = 0$  and  $V_g = V_0$ . Thus

$$\underline{\rho}_a = V_0 > 0 \tag{18.63}$$

so that  $K \sim 0$ .

 $V_g = 0$  after inflation. Thus, we have

$$\underline{\rho}_g = \left(\rho_{mg} + \rho_{\lg} + \rho_{\gamma g}\right) R^3 / R_1^3. \tag{18.64}$$

When temperature is so low that  $\rho_l \gg p_l$ , we may take  $\rho_l \sim 0$ . Thus, considering  $\rho_{gm}R^3 \simeq \rho_{gm1}R_1^3$  and  $\rho_{g\gamma}R^4 = \rho_{g\gamma1}R_1^4$ , we can reduce (18.64) to

$$\underline{\rho}_{g} = \rho_{vm1} - \rho_{sm1} + \rho_{g\gamma1} R_1 / R.$$
(18.65)

When  $R_1/R \ll 1$ , we have  $K \sim \frac{\dot{\rho}}{\rho_g} \sim 0$ .

### XIX. CONTRACTION OF SPACE, THE HIGHEST TEMPERATURE AND IN-FLATION OF SPACE

The temperature effects in the second model are the same as those in the first model. Analogously to the first model, there are  $T_{cr}$ ,  $T_{max}$ ,  $R_{min}$  and inflation, and there is no singularity in the second model as well.

#### A. Contraction of space

The contracting process in the second model is similar to that in the first model. The difference is that the boundary condition (6.13) is more obviously satisfied in the second model than in the first model.

In the S - breakng, we consider the space contraction process. In low temperatures,  $V_g = 0, \rho_{sm} \gg \rho_{s\gamma}, \underline{\rho}_g = \underline{\rho}_{sm} - \underline{\rho}_{vm} \gtrsim \underline{\rho}_{g0}$ . In this case,  $K \sim 0$  and K > 0 could be regarded as a constant and space will contract because of (18.47) - (18.48). When  $T_s \ll T_{cr}$ , the transformation  $\rho_s$  into  $\rho_v$  may be ignored. In this case, space will monotonously contract faster and faster.

 $T_s$  and  $T_v$  must go up high as R decreases. When  $T_s \sim T_{cr}$ , the masses all particles tend to zero. Consequently the thermal equilibrium between s - matter and v - matter comes into being due to (2.10). After thermal equilibrium, the number and the energy of every sort of particles will satisfy statistical distribution determined by their spins. Considering (5.26) and the symmetry of s - particles and v - particles, when  $T_s \gtrsim T_{cr}$ , we have

$$T_{s} = T_{v} = T, \quad \rho_{s} (T_{s}) = \rho_{v} (T_{v}) \equiv \rho (T) = \frac{\pi^{2}}{30} g^{*} T^{4}, \quad (19.1)$$
$$g_{s}^{*} = g_{sB} + 7g_{sF}/8 = g_{v}^{*} = g_{vB} + 7g_{vF}/8 \equiv g^{*}.$$

where  $g_{aB}(g_{aF})$  is the total number of the spin states of a - bosons(a - fermions), and

$$\rho_a(T_a) R^4(T_a) = D_a, \quad \rho_a(T_a) = \frac{\pi^2}{30} g_a^* T_a^4,$$
(19.2)

$$TR(T) = T_{cr}R(T_{cr}).$$
(19.3)

 $V_s = V_v = 0$  when  $T_s \ge T_{cr}$  due to  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$ . Considering (5.26), (18.63), (18.47) -

(18.48) and (19.1), when  $T_s \ge T_{cr}$ , we have

$$\rho_g + V_g = \rho_s \left( T_s \right) - \rho_v \left( T_v \right) + V_s + V_0 - V_v = V_0, \tag{19.4}$$

$$K = K(V_0) \equiv K_{cr} = 1 - D_0 > 0, \qquad (19.5)$$

$$\dot{R}^2 = -K_{cr} + \eta V_0 R^2, \tag{19.6}$$

$$\ddot{R} = \eta V_0 R > 0.$$
 (19.7)

It is seen that when  $T \ge T_{cr}$ , s - matter and v - matter are completely symmetric, and both s - SU(5) and v - SU(5) hold strictly. (19.4) and (19.6) - (19.7) are the certain results of space contraction and conjecture 1. This is different from the conventional theory in essence. If there is only one sort of matter as the conventional theory or  $\rho_s$  and  $\rho_v$  cannot transform from one into other, space will continue to contract and  $T_s$  and  $T_v$  will continue to rise provided  $\rho_g R^2 - K > 0$ . In fact, in this case, space will contract to a singular point and  $T_s$  and  $T_v$  tend to infinite.

 $\langle \omega_v \rangle = \langle \omega_s \rangle = 0$  is the sufficient but is not the necessary condition for (19.4). In fact, provided the following conditions are realized, (19.4) can come into being when  $T_s < T_{cr}$  as well.

A.  $m(\omega_a, T_v, T_s)$ ,  $m(f_a, T_v, T_s)$  and  $m(g_a, T_v, T_s)$  will decrease because  $T_v$  and  $T_s$  arise. Here  $f_a$  and  $g_a$  denote fermions and gauge bosons, respectively. When  $m(\omega_a, T_v, T_s) \sim m(f_a, T_v, T_s)$  or  $m(g_a, T_v, T_s)$ ,  $\Omega_a$  or  $\varphi_a$  and  $f_a$  or  $g_a$  can transform from one into other by the SU(5) couplings.

B. When  $m(\omega_s, T_v, T_s) \sim m(\omega_v, T_v, T_s)$ ,  $\omega_s$  and  $\omega_v$  can transform from one into other by (2.10).

The two conditions can be realized as well when  $T_s \sim T_v \lesssim T_{cr}^{[9]}$ .

Even when  $\chi_s$  and  $\chi_v$  are considered, the above conclusions still hold qualitatively.

# B. There is non-singularity in the second model, the highest temperature and inflation of space

It is seen from (19.6) - (19.7) there no singularity of space-time in the second model.

If space does not contract, it is necessary that there is no space-time singularity because of the cosmological principle. (19.6) - (19.7) is consistent with the Lemaitre model without singularity in which  $\rho_g = 0$ , K = 1 and the cosmological constant  $\lambda_{eff} > 0^{[20]}$ . It is seen that there is no singularity in the present model. This can be explained in detail as follows.

Let 
$$R_{cr} = R(T_{cr})$$
. If  
 $R_{cr}^{2} = -K_{cr} + \eta V_0 R_{cr}^{2} \ge 0$ , i.e.  $R_{cr} \ge \sqrt{K_{cr}/\eta V_0}$ , (19.8)

R can continue to decrease with a deceleration or stop contracting. Hence there must be the least scale  $R_{\min} \leq R_{cr}$ , the critical temperature  $T_{cr}$ , the highest temperature  $T_{\max}$  and the largest energy density  $\rho_{\max}$ :

$$0 < R_{\min} = \sqrt{K_{cr}/\eta V_0} \le R_{cr}, \quad T_{cr} \equiv 2\mu/\sqrt{\lambda}, \tag{19.9}$$

$$T_{\max} = T(R_{\min}) = T_{cr}R_{cr}/R_{\min} \ge T_{cr},$$
 (19.10)

$$\rho_{\max} = \rho_{s\max} + \rho_{v\max} = 2\frac{\pi^2}{30}g^*T_{\max}^4.$$
(19.11)

(19.2) - (19.3) is considered in (19.9) - (19.11).

It is seen from (19.9) that the boundary condition (6.13) must be satisfied. Consequently, there must be no singularity in the second model as well, because of the theorem related to singularity and the discussion in section 6.

It is seen from (19.6) - (19.7) that when R decreases to  $R_{\min}$ , space inflation must occur.

$$R = \sqrt{\frac{K_{cr}}{\eta V_0}} \cosh \sqrt{\eta V_0} (t - t_{FI})$$
$$= \sqrt{\frac{K_{cr}}{\eta V_0}} \cosh H (t - t_{FI}), \quad \sqrt{\eta V_0} \equiv H,$$
(19.12)

$$=\sqrt{\frac{K_{cr}}{\eta V_0}} = R_{\min} > 0 \text{ when } t = t_{FI}$$
(19.13)

$$\sim \frac{1}{2} \sqrt{\frac{K_{cr}}{\eta V_0}} \exp H \left( t - t_{FI} \right) \text{ when } H \left( t - t_{FI} \right) >> 1,$$
 (19.14)

 $t_{FI}$  is the final of the s-world and the beginning moment.  $(K_{cr}/R_{\min}^2)$  or  $V_0$  and  $T_{cr}$  are two new important constants, and  $T_{\max}$  and  $\rho_{\max}$  are determined by  $R_{\min}$ ,  $R_{cr}$  and  $T_{cr}$ . It is seen from (19.9) – (19.11) that all R, T and  $\rho$  must be finite in this case. The meanings of the parameters are that when  $T = T_{cr}$ ,  $\langle \omega_s \rangle = \langle \omega_v \rangle = 0$  and  $R = R_{cr}$ , and when  $R = R_{\min}$ ,  $T = T_{\max}$  or  $\rho = \rho_{\max}$  and  $\dot{R} = 0$ .

We know that the duration of inflation  $\tau$  must be finite, because T will descend extremely fast due to inflation and the inflation process is a supercooled process. When  $T < T_{cr}$ ,  $V = V_0$  is the maximum so that the phase transition (i.e. reheating process) must occur. After  $\tau$ , R has a large enough increase.

 $R_{cr} \geq \sqrt{K_{cr}/\eta V_0}$  in (19.8) is the condition of space inflation. By (19.1) – (19.3) and (19.9) – (19.11), considering  $V_0 = \mu^4/4\lambda$  and  $K_{cr} > 0$ , we can rewrite  $R_{cr} \geq \sqrt{K_{cr}/\eta V_0}$  as

$$(TR)^{2} = (T_{cr}R_{cr})^{2} \ge \frac{K_{cr}}{\eta} \left(\frac{4}{\mu}\right)^{2}$$
 or  
 $D_{s} = \rho_{cr}R_{cr}^{4} \ge g^{*}\frac{(\pi K_{cr})^{2}}{30\eta^{2}} \left(\frac{4}{\mu}\right)^{4} \equiv D_{cr}.$  (19.15)

If  $R(T_{cr}) < \sqrt{K_{cr}/\eta V_0}$  or  $D_s < D_{cr}$ , this implies that  $\dot{R} = 0$  already occurs before R contracts to  $R_{cr}$  or  $T_s$  rises to  $T_{cr}$ , i.e.,  $R_{\min} > R_{cr}$  and  $T(R_{\min}) = T_{\max} < T_{cr}$ . Consequently  $T_{cr}$  and  $R_{cr}$  cannot be reached. there are still  $\langle \omega_s(T_s) \rangle \neq 0$  and  $\langle \omega_v \rangle = 0$ . In this case, there are still  $\langle \omega_s(T_s) \rangle \neq 0$  and  $\langle \omega_v \rangle = 0$  and  $\langle \omega_v \rangle = 0$ , and all  $R_{\min}$ ,  $T_{\max}$ ,  $\rho_g$ ,  $\rho_s$  and  $\rho_v$  must still be finite, i.e. there is no space-time singularity. In this case, it is necessary that

$$R = 0, \quad R > 0, \quad \text{when } R = R_{\min} > R_{cr},$$
 (19.16)

because  $R_{\min}$  is the end of contracting R. In this case, space will expand still in the S – *breaking*, but space inflation cannot occur.

We see from mentioned above that in any case of the contracting process, there must be  $R_{\min} > 0$  and the finite  $T_{\max}$ . Because of the cosmological principle, all  $\rho_s$ ,  $\rho_v$ ,  $\tilde{\rho}_{Sg} = \tilde{\rho}_s - \tilde{\rho}_v$ ,  $\tilde{\rho}_{Sg} = \tilde{p}_s - \tilde{p}_v$  and  $p_a \leq \rho_a/3$  are finite. Hence all  $T_{s\mu\nu}$ ,  $T_{v\mu\nu}$  and  $T_{g\mu\nu} = T_{s\mu\nu} - T_{v\mu\nu}$  are finite. Substituting the finite  $T_{g\mu\nu}$  into the Einstein field equation (2.13), we see that  $R_{\mu\nu}$  and  $g_{\mu\nu}$  must be finite.

According to the present model, the universe is composed of infinite s-cosmic islands and v-cosmic islands. The cosmic islands are different from each other. Hence every possibility can be realized. There must be no singularity in any case.

Thus, we have proved that there is no singularity in present model.

The process of space inflation is the same as that in the first model (see section 7.2).

#### XX. EVOLVING PROCESS OF SPACE AFTER INFLATION

We discuss the expanding process after reheating as follows. After reheating process, expansion of space can be divide into three stages. The three stages are the early stage, the stage from  $\rho_g \sim 0$  to  $\dot{R} \sim 0$  and the final stage in which space expands with an acceleration.

#### A. The three stage of the universe evolution

1. The reheating process and the early stage in which space expands with a deceleration

As the same as the first model, there is the reheating process after inflation. (7.1) - (7.3) is still applicable for the second model.

After the reheating,  $V_g = 0$ . K, K and  $r\partial K/\partial r$  may be neglected because  $\underline{\rho}_g = \underline{\rho}_v - \underline{\rho}_s \gg \underline{\rho}_g/A_{\rho}$  in the early stage. Thus, (18.24) - (18.25) and (18.30) are reduced to (18.47) - (18.49) in which  $V_g = 0$ . Thus, space must expand with a deceleration in the early stage after reheating.

2. The stage from  $\rho_g(t_2) \sim 0$  to  $\overset{\cdot}{R}(t_1) \sim 0$ .

After temperature descends sharply,  $\rho_g$  will decrease to  $\rho_g(t_2) = 0$  and finally  $\rho_g < 0$ , because  $\rho_{mg} = \rho_{vm} - \rho_{sm} \sim R^{-3}$ ,  $\rho_{lg} = \rho_{vl} - \rho_{sl} \sim R^{-3(1+\kappa)}$  and  $\rho_{\gamma g} = \rho_{v\gamma} \sim R^{-4}$ . When temperature is so low that  $p_l \ll \rho_l$ , we may rewrite  $\rho_v = \rho_{vm} + \rho_{v\gamma}$  and  $\rho_s = \rho_{sm}$ . The stage from  $\rho_g(t_2) \sim 0$  to  $R(t_1) \sim 0$   $(t_1 > t_2)$  has been discussed in section 18.2 (see (18.52) - (18.59)). In contrast with the conventional theory, both  $\rho_g(t_1) < 0$  and  $R(t_1) > 0$ are possible when  $R(t_1) = 0$  so that space will expand with an acceleration up to now.

3. The final stage in which space expands with an acceleration

Space will continue to expand because  $\ddot{R}(t_1) > 0$  after  $R(t_1) = 0$ . Thus,  $\rho_g(t)$  will continue to decrease from  $\rho_g(t_1) < 0$  as R increases. Temperature will decrease as Rincreases. When temperature is so low that  $p_\alpha \ll \rho_\alpha$ , we may rewrite  $\rho_v = \rho_{vm} + \rho_{v\gamma}$  and  $\rho_s = \rho_{sm}$ . Considering  $\rho_{mg} \sim R^{-3}$  and  $\rho_{\gamma g} \sim R^{-4}$ , we obtain  $\underline{\rho}_g$  determined by (18.65). Thus  $\underline{\rho}_g \simeq \underline{\rho}_{mg1}$  when  $R_1/R \ll 1$  due to (18.65) and  $p_g \sim \rho_{\gamma g}/3$  may be neglected. Consequently,  $K \sim 0, \ K \sim 0, \ v(\rho_g + p_g) \sim 0$  and  $r\partial K/\partial r \sim 0$ , and (18.24) – (18.25) and (18.30) are reduced to

$$\dot{R}^2 = -K + \eta \rho_g R^2, \ \ddot{R} = -\frac{\eta}{2} \rho_g R > 0, \ K \sim -1 - D_0$$
 (20.1)

$$\dot{\rho}_g + 3\rho_g R/R = 0.$$
 (20.2)

Thus, space will expand with an acceleration when  $t > t_1$ .

Sum up, we see that after reheating, space first expands with a deceleration, then comes to static (in fact, it is  $R/R = (R/R)_{\min} \ge 0$ ), and finally expands with an acceleration up to now.

#### **B.** Determination of a(t) in the second model

The exact solutions of K(t,r), v(t,r), R(t,r),  $\rho_g(t,r)$  and  $p_g(t,r)$  are hardly determined. Hence we have to discuss the asymptotic solutions. The asymptotic solution of K(t,r) had been discussed in section 18.2. We discuss the asymptotic solutions of R(t,r) and  $\rho_g(t,r)$ as follows.

The evolution equations have been derived in section 18.2. From (18.38) we see that  $K \sim 1 - D_0 > 0$  when  $A_{\rho}\underline{\rho}_g \gg \underline{\rho}_{g0}$  and  $K \sim -1 - D_0 < 0$  when  $A_{\rho}\underline{\rho}_g \longrightarrow -\underline{\rho}_{g0}$ . In the two cases, (18.24) - (18.27) is reduced to (18.46) - (18.49). When  $K \sim 0$ , K cannot be neglected and (18.24) - (18.27) is dependent on r. When  $\underline{\rho}_g \sim 0$  and  $r \sim 0$ ,  $K \sim -D_0$  and (18.24) - (18.25) are reduced to (18.57). When  $\underline{\rho}_g \sim 0$  and r is large,  $K \sim -D_0/r^2$  and (18.24) - (18.25) is reduced to (18.52) - (18.53). When  $R \sim 0$ ,  $0 > K > -1 - D_0$  and (18.25) is reduced to (18.59). The equations (18.52) - (18.53), (18.57) and (18.59) are all independent of r.

After reheating,  $\underline{\rho}_g$  and K decrease as space expands. The process is as follows.  $K \sim 1 - D_0, \underline{\rho}_g \gg \underline{\rho}_{g0}/A_\rho$  and  $\ddot{R} < 0 \longrightarrow K = 0, \underline{\rho}_g = \underline{\rho}'_g > 0$  and  $\ddot{R} < 0 \longrightarrow K \sim -D_0, \underline{\rho}_g = 0$  and  $\ddot{R} < 0 \longrightarrow K \gtrsim -1 - D_0, \dot{R} = 0$  and  $\ddot{R} > 0 \longrightarrow K \sim -1 - D_0, \dot{R} > 0$  and  $\ddot{R} > 0$ .

In the early stage after reheating,  $\rho_g$  is so large that  $\eta \rho_g R^2 \gg K$ . Hence, in this case, K cannot influence evolution of space markedly so that K may be taken as -1 for convenience. The two stages from  $K \sim 0$  to  $\rho_g \sim 0$  and from  $\rho_g \sim 0$  to  $R \sim 0$  are short. As mentioned above,  $K \leq 0$  in the two stages. As a rough approximation for the whole evolving process, we take K = -1 in the two stages. When  $A_{\rho}\rho_g + \rho_{g0} \gtrsim 0$ ,  $K \sim -1 - D_0$ , we may take K = -1 because  $K/R^2$  describe the curvature of space and is measurable. From the approximation we can see the main evolving features of the universe. Based on the approximation,  $rK = r\partial K/\partial r = 0$  and (18.24) - (18.27) is reduced to

$$\dot{R}^2 + K = \eta \rho_g R^2, \quad K = -1,$$
(20.3)

$$\ddot{R} = -\frac{\eta}{2} \left( \rho_g + 3p_g \right) R, \quad , \tag{20.4}$$

$$R^{2}\frac{d}{dt}\rho_{g} + 3\left(\rho_{g} + p_{g}\right)R\dot{R} = 0.$$
(20.5)

(20.3) - (20.5) are the same as (4.5) - (4.6) and (4.9) in which  $V_g = V_g = 0$ . Hence the inferences (7.10) - (7.23) can be derived from (20.3) - (20.5).

## XXI. TRANSFORMATION OF REPULSIVE POTENTIAL ENERGY CHIEFLY INTO THE KINETIC ENERGY OF SU(5) SINGLETS

In the section we consider the repulsion between s-matter and v-matter in the flat spacetime and the Newtonian mechanics.

The repulsion potential energy between v - matter and s - matter is determined by the distributing mode of s - matter and v - matter. In V - breaking, v - particles with their masses can form v - celestial bodies, but s - SU(5) color single states cannot form any dumpling and must distribute loosely in space. Consequently, the huge repulsion potential energy must chiefly transform into the kinetic energy of s - SU(5) color single states when the v - celestial bodies form or space expands. In fact, when flat space expands N times, i.e.,  $R \longrightarrow NR$ , the repulsive-potential energy density  $V_r$  becomes  $V_r/N$  and

$$\Delta V_r = (1 - 1/N) V_r.$$
(21.1)

Consider a system in flat space which is composed of a v - body with its mass M and a s - colour single state with its mass m. It is easy to get the rate  $\Delta E_m / \Delta E_M$  for static M and m at the initial moment.

$$\frac{\triangle E_m}{\triangle E_M} = \frac{2M + \triangle V_r}{2m + \triangle V_r}.$$
(21.2)

Because  $M \gg m$ ,  $\triangle E_m > \triangle E_M$ .

Space expansion is not the necessary condition to transform repulsive potential energy into kinetic energy. Supposing R = 0 and some v - matter gathers to a region and forms a galaxy, s - color single states, which are initially in the region, must be repulsed away from the region by the celestial bodies in the galaxy. Consequently, the repulsive potential energy chiefly transforms into the kinetic energy of the s - matter.

There are four of causes for  $\dot{\underline{\rho}}_{a} \neq 0$ :

1. It is seen from (18.50) that  $K \neq 0$  can cause  $\underline{\rho}_g \neq 0$ . Of course,  $\underline{\rho}_g \neq 0$  is the cause of  $K \neq 0$ .

2. The change of  $\underline{\rho}_g$  can be caused by the transformation of s-particles and v-particles from one to another when temperature is high enough;

3. The expansion or contraction of space causes the change of  $\underline{\rho}_g$ , because of  $\rho_{sm}$  and that  $\rho_{sm} \sim R^{-3}$  and  $\rho_{v\gamma} \sim R^{-4}$ .

4. The repulsion potential chiefly transform to kinetic energy of the SU(5) singlets when  $\dot{R} \sim 0$  and galaxies form.

One effect of the repulsion between s - matter and v - matter is that it can prevent v - galaxies are teared by expansion of space with an acceleration. Any galaxy must undergo a pressure coming from s-matter surrounding the galaxy. The pressure can prevent the v - galaxy are teared by expansion of space with an acceleration.

#### XXII. NEW PREDICTIONS

A. The second model can obtain the three predictions of the first model and explain primordial, CMBR and the large scale structure.

The three predictions of the first model can be obtained by the second model as well. For the third prediction, in contrast with the first model, after a black hole with its mass and density big enough inflate, the black hole can transit into either of the S – breaking and the V – breaking.

Letting there be a v - black hole with its mass and density to be so huge that its temperature can arrive to  $T_v \gtrsim T_{cr} = 2\mu/\sqrt{\lambda}$  when the black hole contracts by its selfgravitation, then the expectation values of the Higgs fields inside the v - black hole will change from  $\varpi_v = \varpi_{v0}$  and  $\varpi_s = 0$  into  $\varpi_v = \varpi_s = 0$ . Consequently, inflation must occur. After inflation, the most symmetric state will transit into the S - breaking or the V - breaking. No matter which breaking appears, the energy of the black hole must transform into both v - energy and s - energy. Thus, a v - observer will find that the black hole disappears and a white hole appears.

As the same as the first model, the second model can explain primordial nucleosynthesis, CMBR and the large scale structure of space-time. As mentioned in section 20.2, after inflation and reheating,  $V_g = \dot{V}_g = 0$ , the evolution equation can approximately be reduced to (20.3) - (20.5) in which  $K \sim -1$ . In the case, (20.3) - (20.5) are the same as (4.5) - (4.6)and (4.9) of the first model. Hence the second model, as the same as the first model, can explain primordial nucleosynthesis, CMBR and the large scale structure as well.

#### B. The universe is composed of infinite s-cosmic islands and v-cosmic islands

If there is one sort of spontaneous symmetry breaking, the hypersurfaces of the universe must be complete. Hence K cannot change and there is only one sort of evolving mode in the whole universe. In contrast with the conventional theory, there are two sorts of matter and the two sorts of breaking. The S – breaking and the V – breaking are different in essence and symmetric. Hence it is possible that some regions of the universe can exist in the S – breaking, and the others can exist in the the -V – breaking. Consequently the universe must be composed of s – cosmic islands with the S – breaking and v – cosmic islands with the V – breaking. Of course, a transitional region (T – region) must exist between a s – island and a v – island.

Every cosmic island is only one part of the universe, must be finite and evolves individually. Consequently its hypersurface of simultaneity must be incomplete and its metric must be different with the RW-metric which describes a complete hypersurface.

In the s - islands,  $\langle \omega_s \rangle = \langle \omega_s \rangle_0$  and  $\langle \omega_v \rangle = 0$ , s - particles forms s - galaxies and v - particles form v - SU(5) singlets which corresponds to so-called dark energy. The same results hold for v - islands because of conjecture 1.

Through the transitional regions,  $\langle \omega_v \rangle = \langle \omega_v \rangle_0$  and  $\langle \omega_s \rangle = 0$  will transit to  $\langle \omega_s \rangle = \langle \omega_s \rangle_0$ and  $\langle \omega_v \rangle = 0$ . Hence all  $\langle \omega_s \rangle$ ,  $\langle \omega_v \rangle$ ,  $V_s$  and  $V_v$  are not equal to zero and the expectation values  $\langle \omega_s \rangle_T$  and  $\langle \omega_v \rangle_T$  inside the T – region must satisfy

$$0 < |\langle \omega_s \rangle_T | < |\langle \omega_s \rangle_0 |, \quad 0 < |\langle \omega_v \rangle_T | < |\langle \omega_v \rangle_0 |.$$

$$(22.1)$$

There must be only v - islands neighboring a s - island. This is because that if two s - cosmic islands are neighboring, they must form one new larger s - cosmic island in order that  $V = V_{min}$ .

Based on the following reasons, the probability is very little that a v – observer receives messages from a s – island:

1. The probability must be very small that a s - particle (a quark, a lepton or a photon) in a s - island comes into a v - island, because a s - particle in the s - island is non-color single state. If a s - particle comes into the v - island, it would still be non-color single state. This is impossible due to color confinement. Such a bound state as  $(u\overline{u} \mp d\overline{d})/\sqrt{2}$ is a SU(5) color single state in both V - breaking and S - breaking. It seems that it can comes into the v - cosmic island. In fact, it still hardly comes into the v - island, because both masses are very different from each other. On the other hand, such states must be unstable and will decay very fast in s - island.

2. The probability must be very small that a v - particle in the s - island come into the v - island as well. The v - particle (a fermion or a gauge boson) in the s - cosmic island must be massless. If the v - particle comes into the v - island, its mass will change from  $m_0 = 0$  to  $m_0 > 0$ . Thus it must suffer a strong-repulsive interaction, and hence it hardly comes into the v - cosmic island.

3. It is important that  $\langle \omega_s \rangle_T \neq 0$  and  $\langle \omega_v \rangle_T \neq 0$ . Hence it is necessary that  $m_s \neq m_T$  and  $m_v \neq m_T$ , where  $m_s$ ,  $m_v$  and  $m_T$  are the masses of a particle when it is in a s – *island*, a v – *island* or a T – *region*, respectively. Any particle coming into a T – *region* must emit or absorb Higgs particles in order to change its mass from  $m_s$  or  $m_v$  to  $m_T$ , i.e.

$$f \longrightarrow f_T + h, \quad h \longrightarrow g_T + \overline{g}_T, \quad f + h \longrightarrow f_T,$$
 (22.2)

where f is a s- or v-particle coming into a T-region,  $f_T$  is the particle f with its mass  $m_T$  in the T-region, h is a s- or v-Higgs particle, and  $g_T$  and  $\overline{g}_T$  are any particle and its anti-particle. The process hardly occur, because annihilation and creation of h will alter the Higgs potential.

4. Higgs particles in the s - cosmic island must decay very fast into fermions or gauge bosons, and hence they cannot come to the v - cosmic island.

In summary, any particle hardly go to another cosmic island. Particles coming to a T - region will be reflected by the T - region. Thus a galaxy inside a cosmic island is similar to an electron static in a metal sphere shell.

A v-cosmic island and a s-cosmic island can influence each other by the Higgs potential in the T-region between both.

An observer in the cosmic island can regard the cosmic island as the whole cosmos. It is possible that some cosmic islands are forming or expanding, and the other cosmic islands are contracting.

Thus, according to the present model the cosmos as a whole is infinite and its properties are always unchanging, and there is no starting point or end of time.

The sketch of the universe according to the second model is shown in figure 3. Figure 3

#### C. Mass redshifts

Hydrogen spectrum is

$$\omega_{nk} = (E_n - E_k)/\hbar = -\frac{\mu e^4}{2\hbar^3} (\frac{1}{n^2} - \frac{1}{k^2}), \qquad (22.3)$$

$$\mu = \frac{mM}{m+M},\tag{22.4}$$

where m is the mass of an electron, and M is the mass of a proton. According the unified model,  $m \propto v_e$ , the mass of a quark  $m_q \propto v_q$ , and  $M \propto m_q$ , where  $v_e$  and  $v_q$  are the expectation values of the Higgs fields coupling with the electron and the quark, respectively.

If there are some galaxies inside a T - region, from (22.1) we see that the mass  $m_T$  of an electron and the mass  $M_T$  of a proton inside the T - region must be

$$m_T < m, \quad M_T < M. \tag{22.5}$$

Thus we have

$$\mu_T < \mu, \tag{22.6}$$

$$\Delta \omega_{nk} = \omega_{nk} - \omega_{nkT} = -\frac{(\mu - \mu_T)e^4}{2\hbar^3} (\frac{1}{n^2} - \frac{1}{k^2}) < 0.$$
(22.7)

This sort of red-shifts is called mass redshift. The mass redshift is essentially different from the cosmological red-shift mentioned before. Thus, the photons coming from the star in a T - region must have larger red-shift than that determined by the Hubble formula at the same distance. Thereby we guess that some quasars are just the galaxies in the T - region and they have the mass redshifts. The fine-structure constant is considered to be changeable based on the redshifts of some quasars. In contrast with the guess, we consider that the phenomenon is possibly because the mass of electrons changes, but is not because the fine-structure constant changes.

An ordinary s - galaxy and a s - quasar can be neighboring, because a T - region must be neighboring to an ordinary region.

#### XXIII. CONCLUSIONS OF THE MODEL BASED ON A NEW METRIC

Based on the two conjectures in the first model and a new metric which describes the incomplete hypersurface, the second sort of the cosmological models without singularity is constructed. The inferences of the first model can be derived from the present model as well. There is no singularity in the model and the cosmological constant  $\lambda = \lambda_{eff} = 0$  is determined although  $\rho_0$  is still very large and there is no the fine tuning problem, even if  $\lambda_{eff} \neq 0$ .

Dark energy is explained as SU(5) singlets. This is because the SU(5) singlets can only distribute loosely in space or form huge superclusters, cannot be observed and can cause space to expand with an acceleration. In contrast with the dark energy, the gravitational masses of the SU(5) singlets is negative, although their masses are all positive.

Space in the V – *breaking* has three evolving stages: Space first expands with a deceleration because  $\rho_g = \rho_v - \rho_s > 0$  and K > 0; then comes to static; and finally expands with an acceleration up to now because  $\rho_g < 0$  and K < 0.

There are the critical temperature  $T_{cr}$ , the highest temperature  $T_{\max}$ , the least scale  $R_{\min}$ and the largest energy density  $\rho_{\max}$  in the universe.  $V_0$  or  $(K_{cr}/R_{\min}^2)$  and  $T_{cr}$  are two new important constants, and  $T_{\max}$  and  $\rho_{\max}$  are determined by  $R(T_{cr})$ .

The three new predicts of the first model have been obtained as well. Besides, there are two new predictions in the present model.

The universe is composed of infinite s - cosmic islands and v - cosmic islands. Every cosmic island is finite and evolves individually. The hypersurface of every cosmic island must be incomplete. Hence the K - changeable metric is presented. An observer in the cosmic island can regard the cosmic island as the whole cosmos. It is possible that some cosmic islands are forming or expanding, and the other cosmic islands are contracting. Thus, according to the present model the cosmos as a whole is infinite and its properties are always unchanging, and there is no starting point or end of time.

There is mass redshift, because  $m_{eT} < m_{e0}$ , here  $m_{eT}$  is the mass of an electron in a transitional region and  $m_{e0}$  is the given mass of an electron. We guess that some quasars are just the galaxies in a transitional region and they have the mass redshifts. The fine-structure constant is considered to change based on the redshifts of some quasars according to the conventional theory. In contrast with the guess, we consider that the phenomenon is possibly because the mass of electrons changes, but is not because the fine-structure constant changes.

An ordinary s - galaxy and a s - quasar can be neighboring, because a T - region must be neighboring to an ordinary region. There is no physical restriction for  $T_{\mu\nu}$  originating from the field equation. Hence we can suppose  $T^{\mu\nu}_{,\nu} = 0$ , where  $T^{\mu\nu}$  may contain contribution gravitational field.

#### XXIV. SUMMATION

Essential new conjecture 1 is presented. Although conjecture 1 corrects the equivalence principle (the principle is violated by the SU(5) singlets), but it does not contradict experiments and observations up to now.

Based on conjecture 1, two cosmological models without singularity are constructed. There is no singularity in the two models. Based on either of the two models, the cosmological constant issue is solved, i.e.  $\lambda = \lambda_{eff} = 0$ ; Dark energy is explained as SU(5) singlets; The evolution of the universe is explained well and the distance-redshift relationship is derived out which is consistent with observations up to now; The primordial nucleosynthesis, CMBRand the large scale structure of space-time are explained; The five predictions are given (see parts 1 and 2). According to either of the two models, there are the critical temperature  $T_{cr}$ , the highest temperature  $T_{max}$ , the least scale  $R_{min}$  and the largest energy density  $\rho_{max}$ in the universe, and  $V_0$  or  $(K_{cr}/R_{min}^2)$  and  $T_{cr}$  are two new important constants in the two models.

In contrast with the first model, according to the second model, the universe is composed of infinite s-cosmic islands and v-cosmic islands; Every cosmic island is finite and evolves individually; The cosmos as a whole is infinite and its properties are always unchanging, and there is no starting point or end of time. There is mass redshift which is different from the cosmological redshift.

#### Acknowledgments

I am very grateful to professor Zhao Zhan-yue, professor Wu Zhao-yan, professor Zheng Zhi-peng and professor Zhao zheng-guo for their helpful discussions and best support. I am very grateful to professor Liu Yun-zuo, professor Lu Jingbin, doctor Yang Dong and doctor Ma Keyan for their helpful discussions and help in the manuscript.

- Hawking S. W. and Ellis G. F. R., 1999, The Large Scale Structure of Space-Time, Cambidge University Press, p7, 98, 101, 137, 256-298.
- [2] Caldwell R.R. 2004, Phys. World 17, 37; Padmanabhan, T. 2003, Phys. Rep. 380 325; Peebles
   P.J.E. and Ratra B. 2003, Rev. Mod. Phys. 75 559.
- [3] Weinberg S. 1987, Phys. Rev. Lett., 59 2607; Martel H., Shapiro P.R. and Weinberg S. 1998, Astrophys. J 492 29.
- [4] Peebles P.J.E. and Ratra B. 1988, Astrophys. J. **325** L17; Ratra B. and Peebles, P.J.E. 1988,
   Phys. Rev. D **37** 3406; Peebles P.J.E. and Ratra B. 2003, Rev. Mod. Phys. **75** 559.
- [5] Hall L.J., Nomura Y. and Oliver S.J., Phys. 2005, Rev. Lett. 95, 141302.
- Chen S-H , Quantum Field Theory Without Divergence A 2002, hep-th/0203220; 2005, Quantum Field Theory: New Research, Kovras O. Editor, Nova Science Publishers, Inc. 103-170.
- [7] Chen S-H. 2001, A Possible Candidate for Dark Matter, hep-th/0103234; 2005, Progress in Dark Matter Research, Editor: J. Val Blain, Nova Science Publishers, 65-72. Inc. arXiv: 1001.4221.
- [8] Peacock J. A. 1999, Cosmological Physics, Cambridge University Press, 579, 458, 78, 90, 89, 460-464, 664, 296.
- [9] Chen S-H. 2009, Discussion of a Possible Universal Model without Singularity, arXiv. 0908.1495v2; 2006, A Possible Universal Model without Singularity and its Explanation for Evolution of the Universe, hep-ph/0611283.
- [10] Gibbons G. W. and Hawking, S. W. 1977, Phys. Rev D, 15, 2752.
- [11] Chaichian M. and Nelipa N.F. 1984, Introduction to Gauge Field Theories, Springer-Verlag Berlin Heidelberg, 269; Ross G. G. 1984, Grand Unified Theories, The Benjamin/Cummings Publishing Company, INC, 177-183.
- [12] Weinberg S. 1972, Gravitation and Cosmology, New York, Wiley Chanter 12 section 3.
- [13] Coleman S. and Weiberg E.J. 1973, Phys. Rev. D7, 1888; Brandenberg R.H. 1985, Rev. of Mod. Phys., 57. 1.
- [14] Liu L. Jiang Y. and Qian Z. 1989, Progress in Physics, V.9, No. 2, 159 [Use the language of Chinese].

- [15] Guth A. H., 1981, Phys. Rev. D 23, 347.
- [16] Ignatiev A. Y. and Volkas R. R., Phys. Rev. (2003) D 68, 023518; Berezhiani Z., Comelli D. and Villante F.L., Phys. Lett. (2001) B 503, 362.
- [17] Yu Yunqiang., Lectures in Cosmological Physics (Chinese), Peking University Press, (2002), 151, 170-172.
- [18] Liddle A. R. and Lyth D. H., 2000, Cosmological Inflation and Large-Scale Structure, Cambridge University Press, 20, 248.
- [19] Liang C.B and Zhou B, 2006, The Course of Differential Geometry and General Relativity (Chinese), Second Edition, (China, Science Press) p 360-367.
- [20] Ohanian H.C. and Ruffini R., (1994), Gravitation and Spacetime (2nd ed.), W.W. Norton and Company, Inc., Section 9.9.

# Cosmological Models Without Singularity Based on RW Metric or A New Metric and their Explanation for Evolution of the Universe

Shi-Hao Chen

<sup>1</sup>Department of Physics, Jilin University; <sup>2</sup>Institute of Theoretical

(Date textdate; Received textdate; Revised textdate; Accepted textdate; Published textdate)

Abstract

Caption

#### Contents

Caption

Figure 1 The curve A describes evolution of a(t) from  $14 \times 10^9 yr$  ago to now; The curve

B describes evolution of a(t) from  $13.7 \times 10^9 yr$  ago to now.

Figure 2 The curve A describes the  $d_L - z$  relation when  $\Omega_{v\gamma 0} = 0.001$  and  $\Omega_{m0} = 0.3\Omega_{v\gamma 0} + 2\sqrt{\Omega_{v\gamma 0}}$ ; The curve B describes the  $d_L - z$  relation when  $\Omega_{v\gamma 0} = 0.05$  and  $\Omega_{m0} = 2\sqrt{\Omega_{v\gamma 0}}$ .

Figure 3  $A_v$ ,  $B_v$  and  $C_v$  denote the v - cosmic islands with K = 1, -1 and 0, respectively;

 $A_s$ ,  $B_s$  and  $C_s$  denote the s - cosmic islands with K = 1, -1 and 0, respectively. Every cosmic island is finite so that its supersurface is incomplete.