

# CAN AN INFINITE LEFT-PRODUCT OF NONNEGATIVE MATRICES BE EXPRESSED IN TERMS OF INFINITE LEFT-PRODUCTS OF STOCHASTIC ONES?

ALAIN THOMAS

ABSTRACT. If a left-product  $M_n \dots M_1$  of square complex matrices converges to a nonnull limit when  $n \rightarrow \infty$  and if the  $M_n$  belong to a finite set, it is clear that there exists an integer  $n_0$  such that the  $M_n$ ,  $n \geq n_0$ , have a common right-eigenvector  $V$  for the eigenvalue 1. Now suppose that the  $M_n$  are nonnegative and that  $V$  has positive entries. Denoting by  $\Delta$  the diagonal matrix whose diagonal entries are the entries of  $V$ , the stochastic matrices  $S_n = \Delta^{-1}M_n\Delta$  satisfy  $M_n \dots M_{n_0} = \Delta S_n \dots S_{n_0} \Delta^{-1}$ , so the problem of the convergence of  $M_n \dots M_1$  reduces to the one of  $S_n \dots S_{n_0}$ . In this paper we still suppose that the  $M_n$  are nonnegative but we do not suppose that  $V$  has positive entries. The first section details the case of the  $2 \times 2$  matrices, and the last gives a first approach in the case of  $d \times d$  matrices.

2000 Mathematics Subject Classification: 15A48.

## INTRODUCTION

The problem of the convergence of  $M_1 \dots M_n$  or  $M_n \dots M_1$  for all the sequences  $(M_n)$  with terms in a finite set of complex matrices, is studied for instance in [1], [2] and [3]. The same problem in case of stochastic matrices is also classical, see for instance [9, chapter 4]. On the other side there exist much results about the distribution of the product matrix  $M_1 \dots M_n$  where the  $M_i$  are taken in a set of stochastic matrices endowed with some probability measure. In [6] and [7], Mukherjea, Nakassis and Ratti give conditions for the limit distribution of  $M_1 \dots M_n$  to be discrete or continuous singular; this contains for instance the case of the Erdős measure [4]. It is well known that in much cases the normalized product  $M_1 \dots M_n$ , if the  $M_i$  are taken in a finite set of nonnegative matrices endowed with some positive probability  $P$ , converges  $P$ -almost everywhere to a rank one matrix; but this general result should be more consistent if one can to specify the  $P$ -negligeable set of divergence.

---

*Key words and phrases.* LCP sets, RCP sets, products of nonnegative matrices, products of stochastic matrices.

Notice that the Erdős measure is studied more in detail in [8] by an other method, using a finite set of matrices and the asymptotic properties of the columns in the products of matrices taken in this set.

In the present paper we first consider (§1) the left-products and the right-products of  $2 \times 2$  matrices (resp.  $2 \times 2$  stochastic matrices). For the left-products of stochastic matrices, the hypothesis that the matrices belong to a finite set is not necessary. We recover the known results by giving several formulations of the necessary and sufficient conditions of convergence.

In §2 we associate to any sequence  $(M_n)$  of nonnegative  $d \times d$  matrices, some sequences of stochastic ones, let  $(S_n^{(i)})$  for  $1 \leq i \leq t$ . The convergence of  $M_n \dots M_1$  is equivalent to the one of the  $S_n^{(i)} \dots S_1^{(i)}$  and some additional condition.

## 1. PRODUCTS OF $2 \times 2$ MATRICES

**1.1. Convergent left-products.** Suppose that the left-product  $M_n \dots M_1$  of some non-negative matrices  $M_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$  converges to a nonnull limit, let  $Q$ , and that the set  $\{M ; \exists n, M = M_n\}$  is finite. It is clear that from a rank  $n_0$ , the  $M_n$  belong to the set

$$\{M ; \text{ there exists infinitely many } n \text{ such that } M = M_n\}$$

and consequently, for any  $M$  in this set,  $MQ = M$ . In other words the nonnull columns of  $Q$  are eigenvectors – for the eigenvalue 1 – of any  $M_n$ ,  $n \geq n_0$ . If for instance this eigenvector is  $\begin{pmatrix} v_1 \\ 0 \end{pmatrix}$ , then  $M_n = \begin{pmatrix} 1 & b_n \\ 0 & d_n \end{pmatrix}$  and  $M_n \dots M_{n_0} = \begin{pmatrix} 1 & \sum_{i=n_0}^n b_i d_{i-1} \dots d_{n_0} \\ 0 & d_n \dots d_{n_0} \end{pmatrix}$

**Proposition 1.1.**  $M_n \dots M_{n_0}$  converges to a nonnull limit, for any fixed  $n_0$  and when  $n \rightarrow \infty$ , if and only if

– either the  $M_n$  have from a certain rank, a common right-eigenvector  $\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$  for the eigenvalue 1, with  $v_1 v_2 > 0$ , and the left-product of the stochastic matrices  $S_n = \begin{pmatrix} \frac{1}{v_1} & 0 \\ 0 & \frac{1}{v_2} \end{pmatrix} M_n \begin{pmatrix} v_1 & 0 \\ 0 & v_2 \end{pmatrix}$  from any rank  $n_0$  converges;

– or the  $M_n$  have from a certain rank  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  for common positive right-eigenvector with respect to the eigenvalue 1, the sum  $\sum_{n=n_0}^{\infty} b_n d_{n-1} \dots d_{n_0}$  is finite and  $d_n \dots d_{n_0}$  converges, for any  $n_0$ , when  $n \rightarrow \infty$ ;

– or the  $M_n$  have from a certain rank  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  for common positive right-eigenvector with respect to the eigenvalue 1, the sum  $\sum_{n=n_0}^{\infty} c_n a_{n-1} \dots a_{n_0}$  is finite and  $a_n \dots a_{n_0}$  converges, for any  $n_0$ , when  $n \rightarrow \infty$ .

**1.2. Case of stochastic matrices.** The case of the matrices

$$S_n = \begin{pmatrix} x_n & 1 - x_n \\ y_n & 1 - y_n \end{pmatrix}, \quad x_n, y_n \in [0, 1],$$

is also trivial because one can compute the left-product

$$(1) \quad Q_n := S_n \dots S_1 = \begin{pmatrix} t_n & 1 - t_n \\ s_n & 1 - s_n \end{pmatrix} \quad \text{where} \quad \begin{cases} s_n := \sum_{i=1}^n y_i \det Q_{i-1} \\ t_n := s_n + \det Q_n \\ Q_0 := I. \end{cases}$$

To find the conditions for the sequence of matrices  $(Q_n)$  to converge one can use the relation

$$(2) \quad s_n = s_{n_0-1} + s_{n_0,n} \det Q_{n_0-1}, \quad \text{where } s_{n_0,n} := \sum_{i=n_0}^n \left( y_i \prod_{n_0 \leq j < i} \det S_j \right).$$

$s_{n_0,n}$  belongs to  $[0, 1]$  because it is one of the entries of the stochastic matrix  $S_n \dots S_{n_0}$ . Hence, in case  $\det Q_n$  has limit 0 the relation (2) implies that  $(s_n)$  is Cauchy; so  $(s_n)$ ,  $(t_n)$  and  $(Q_n)$  converge.

Suppose now that  $\det Q_n$  do not have limit 0. Since  $\det Q_n = \prod_{i=1}^n (x_i - y_i)$  with  $x_i - y_i \in [-1, 1]$ , the non-increasing sequence  $(|\det Q_n|)$  has a positive limit  $\delta$  hence  $|x_n - y_n|$  has limit 1;  $(x_n, y_n)$  cannot have other limit points than  $(0, 1)$  and  $(1, 0)$ .

In case  $(0, 1)$  is one of its limit points,  $y_n |\det Q_{n-1}|$  do not tend to 0 hence the series  $\sum_n y_n \det Q_{n-1}$  diverges and  $(Q_n)$  also do.

In case  $(1, 0)$  is the unique limit point of  $(x_n, y_n)$ ,  $x_n - y_n$  is positive from a rank  $n_0$ . Since the series  $\sum_n \log |x_n - y_n|$  converges to  $\log \delta$ , the inequalities  $\sum_{n \geq n_0} \log(x_n - y_n) \leq \sum_{n \geq n_0} \log(1 - y_n) \leq -\sum_{n \geq n_0} y_n$  prove that the series  $\sum_n y_n$  converges. Since  $\det Q_n$  has limit  $\delta$  or  $-\delta$  according to the sign of  $\det Q_{n_0-1}$ , the sequences  $(s_n)$ ,  $(t_n)$  and  $(Q_n)$  converge.

Consider now the right-product  $P_n := S_1 \dots S_n$  and suppose that the  $S_n$  belong to a finite set. As noticed in [3] it is clear that the nonnull rows of the limit matrix  $P$ , if this matrix exists, are nonnegative left-eigenvectors – for the eigenvalue 1 – of each matrix  $S$  such that  $S_n = S$  for infinitely many  $n$ , because the equality  $P_n = P_{n-1}S$  implies  $P = PS$ . So  $P_n$  can converge only if the  $S_n$ , for  $n$  greater or equal to some integer  $n_0$ , have a common nonnegative left-eigenvector for the eigenvalue 1.

We suppose there exists such a left-eigenvector, let  $L$ , and we search the condition for  $(P_n)$  to converge. Notice that the  $S_n$  for  $n \geq n_0$  commute: since any  $2 \times 2$  stochastic matrix  $S$  has left-eigenvector  $(1 \ -1)$  for the eigenvalue  $\det S$ , both vectors  $L$  and  $(1 \ -1)$  are orthogonal to the columns of  $S_n S_{n'} - S_{n'} S_n$  and consequently this matrix is null. So we fall again in the case of the left-products.

In case  $S_{n_0} \dots S_n$  diverges, nevertheless the sequence of the row-vectors  $LS_{n_0} \dots S_n$  converges (it is constant). Let  $L'$  be some row-eigenvector not colinear to  $L$ ; considering the invertible matrix  $M$  whose rows are  $L$  and  $L'$ ,  $MS_{n_0} \dots S_n$  obviously diverges hence  $L'S_{n_0} \dots S_n$  also do. Consequently  $S_1 \dots S_n$  diverges if and only if at least one of the rows of  $S_1 \dots S_{n_0-1}$  is not colinear to  $L$ .

We have proved the following

**Proposition 1.2.** *Let  $(S_n)$  be a sequence of  $2 \times 2$  stochastic matrices, namely*

$$S_n = \begin{pmatrix} x_n & 1 - x_n \\ y_n & 1 - y_n \end{pmatrix}.$$

(i) *The left-product  $Q_n = S_n \dots S_1$  converges if and only if  $\prod_{k=1}^n (x_k - y_k)$  has limit 0 or  $(x_n, y_n)$  has limit  $(1, 0)$  when  $n \rightarrow +\infty$ .*

(ii) *It diverges only in the case where  $\sum_n (1 - |x_n - y_n|)$  converges and  $S_n$  do not tend to  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ .*

(iii) *If it converges,  $\lim_{n \rightarrow \infty} Q_n = \begin{pmatrix} s + q & 1 - s - q \\ s & 1 - s \end{pmatrix}$  where  $s := \sum_n y_n \det Q_{n-1}$  and  $q := \lim_{n \rightarrow \infty} (\det Q_n)$ .*

(iv) *Suppose now that the  $S_n$  belong to a finite set. Then the right-product  $P_n = S_1 \dots S_n$  converges if and only if there exists  $n_0$  such that*

- *the matrices  $S_n$  for  $n \geq n_0$  have a common left-eigenvector for the eigenvalue 1*
- *and either  $\lim_{n \rightarrow \infty} \prod_{k=n_0}^n (x_k - y_k) = 0$ , or  $\lim_{n \rightarrow \infty} (x_n, y_n) = (1, 0)$ , or the rows of  $S_1 \dots S_{n_0-1}$  are colinear to the left-eigenvector.*

(v) *Suppose the  $S_n$ ,  $n \geq 1$ , have a common left-eigenvector with respect to the eigenvalue 1. Then the  $S_n$  commute and  $P_n = Q_n$ .*

## 2. LEFT PRODUCTS OF $d \times d$ NONNEGATIVE MATRICES

Let us first give one example in order to illustrate the proposition that follows: we consider the products  $Q_n = M_n \dots M_1$ , where the  $M_i$  belong to the set of nonnegative matrices of

the form

$$M = \begin{pmatrix} a & b & 3-3a-2b & e & 4-4a-3b-5e & 1 & 0 \\ c & d & 2-3c-2d & f & 3-4c-3d-5f & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & g & 5-5g & 0 & 1 \\ 0 & 0 & 0 & h & 1-5h & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & x & 1-x \\ 0 & 0 & 0 & 0 & 0 & y & \frac{1}{2}-y \end{pmatrix}$$

where  $a, b, c, d, e, f, g, h, x, y$  are some reals such that  $M$  has exactly twenty four nonnull

entries. Since the eigenspace associated to the eigenvalue 1 is generated by  $\begin{pmatrix} 3 \\ 2 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$  and

$\begin{pmatrix} 4 \\ 3 \\ 0 \\ 5 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ , we can associate to each matrix  $M$  two submatrices with positive eigenvectors:

the submatrix  $M^{\{1,2,3\}}$  of the entries of  $M$  with row and column indexes in  $\{1, 2, 3\}$  and the submatrix  $M^{\{1,2,4,5\}}$  of the entries of  $M$  with row and column indexes in  $\{1, 2, 4, 5\}$ .

Then we associate two stochastic matrices  $S = \Delta^{-1}M^{\{1,2,3\}}\Delta$  and  $S' = \Delta'^{-1}M^{\{1,2,3\}}\Delta'$ ,

where  $\Delta = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$  and  $\Delta' = \begin{pmatrix} 4 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ ; namely

$$S = \begin{pmatrix} a & 2b/3 & 1-a-2b/3 \\ 3c/2 & d & 1-3c/2-d \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad S' = \begin{pmatrix} a & 3b/4 & 5e/4 & 1-a-3b/4-5e/4 \\ 4c/3 & d & 5f/3 & 1-4c/3-d-5f/3 \\ 0 & 0 & g & 1-g \\ 0 & 0 & 5h & 1-5h \end{pmatrix}.$$

Now we obtain 21 of the 49 entries of  $Q_n = M_n \dots M_1$  in function of two products of stochastic matrices, and the other entries in the first five columns of  $Q_n$  are null: indeed  $Q_n$  has for submatrices  $M_n^{\{1,2,3\}} \dots M_1^{\{1,2,3\}} = \Delta S_n \dots S_1 \Delta^{-1}$  and  $M_n^{\{1,2,4,5\}} \dots M_1^{\{1,2,4,5\}} = \Delta' S'_n \dots S'_1 \Delta'^{-1}$ .

The products  $S_n \dots S_1$  and  $S'_n \dots S'_1$  converge to rank 1 matrices: use for instance [5], or use the previous section and the formula for the products of triangular-by-blocks matrices that is,

$$\prod_{i=n}^1 \begin{pmatrix} A_i & B_i \\ 0 & D_i \end{pmatrix} = \begin{pmatrix} \prod_{i=n}^1 A_i & \sum_{i=1}^n A_n \dots A_{i+1} B_i D_{i-1} \dots D_1 \\ 0 & \prod_{i=n}^1 D_i \end{pmatrix}.$$

The limit of  $Q_n$  is a rank 2 matrix of the form  $\begin{pmatrix} 0 & 0 & 3 & 4\alpha & 4\beta & 4\gamma + 6 & 4\delta + 6 \\ 0 & 0 & 2 & 3\alpha & 3\beta & 3\gamma + 4 & 3\delta + 4 \\ 0 & 0 & 1 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 5\alpha & 5\beta & 5\gamma & 5\delta \\ 0 & 0 & 0 & \alpha & \beta & \gamma & \delta \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$

The following proposition generalizes what we have seen on the example. For any  $d \times d$  matrix  $M$  and for any  $K, K' \subset \{1, \dots, d\}$  we denote by  $M^K$  the submatrix of the entries of  $M$  whose row index and column index belong to  $K$  and by  $M^{K, K'}$  the submatrix of the entries of  $M$  whose row index belongs to  $K$  and column index belongs to  $K'$ . By commodity we use the same notation for the row-matrices  $L$  (resp. the column-matrices  $V$ ):  $L^K$  (resp.  $V^K$ ) is the row-matrice (resp. column-matrice) of the entries of  $L$  (resp.  $V$ ) whose index belong to  $K$ .

**Proposition 2.1.** *Let  $(M_n)$  be a sequence of nonnegative  $d \times d$  matrices that belong to a given finite set. The left-product  $Q_{n, n_0} := M_n \dots M_{n_0}$  converges to a nonnull limit when  $n \rightarrow \infty$  – for each positive integer  $n_0$  – if and only if  $(M_n)$  satisfies both conditions:*

(i) *there exist some subsets of  $\{1, \dots, d\}$ , let  $K_1, \dots, K_t$  with complementarities  $K_1^c, \dots, K_t^c$ , and some diagonal matrices with positive diagonals, let  $\Delta_1, \dots, \Delta_t$ , such that the  $S_n^{(i)} = \Delta_i^{-1} M_n^{K_i} \Delta_i$  are stochastic, the  $M_n^{K_i^c, K_i}$  are null, and  $\lim_{n \rightarrow \infty} S_n^{(i)} \dots S_{n_0}^{(i)}$  exists for any  $i$  and  $n_0$ ;*

(ii) *setting  $K = \cup_i K_i$ ,  $\lim_{n \rightarrow \infty} M_n^{K^c} \dots M_{n_0}^{K^c}$  is the null matrix for any  $n_0$  and the series  $\sum_{i=n_0}^{\infty} M_n^K \dots M_{i+1}^K M_i^{K, K^c} M_{i-1}^{K^c} \dots M_{n_0}^{K^c}$  converges.*

*Proof.* If the conditions (i) and (ii) are satisfied, the entries of  $M_n \dots M_{n_0}$  with column index in  $K$  converges either to 0 or to the entries of the matrices  $\lim_{n \rightarrow \infty} \Delta_i S_n^{(i)} \dots S_{n_0}^{(i)} \Delta_i^{-1}$ ,  $i = 1, \dots, t$ . Consequently  $M_n \dots M_{n_0}$  converges, by using the formula of product of

triangular-by-blocs matrices:

$$\prod_{i=n}^{n_0} \begin{pmatrix} A_i & B_i \\ 0 & D_i \end{pmatrix} = \begin{pmatrix} \prod_{i=n}^{n_0} A_i & \sum_{i=n_0}^n A_n \dots A_{i+1} B_i D_{i-1} \dots D_{n_0} \\ 0 & \prod_{i=n}^{n_0} D_i \end{pmatrix}.$$

Conversely suppose that  $Q_{n,n_0} = M_n \dots M_{n_0}$  converges to a nonnull limit for any  $n_0 \in \mathbb{N}$ . Since the  $M_n$  belong to a finite set, we can choose  $n_0$  large enough such that all the matrices  $M$  that are equal to  $M_n$  for at least one  $n \geq n_0$ , are also equal to  $M_n$  for infinitely many  $n$ . Then the nonnull columns of the limit matrix  $Q$  are right-eigenvectors of all the  $M_n$ ,  $n \geq n_0$ , for the eigenvalue 1, because the equality  $Q_{n,n_0} = M_n Q_{n-1,n_0} = M Q_{n-1,n_0}$  implies  $Q = MQ$ . Let  $t$  be the rank of  $Q$ ,  $t = 0$  if  $Q$  is null, and denote by  $V_1, \dots, V_t$  the linearly independent columns of  $Q$ . We denote also by  $K_i$  the set of the indexes of the nonnull entries in  $V_i$ , and by  $\Delta_i$  the diagonal matrix whose diagonal entries are the nonnull entries of  $V_i$ . Now (i) results from the equality  $M_n^{D,K_i} V_i^{K_i} = V_i$ ,  $D := \{1, \dots, d\}$ ;  $S_n^{(i)} \dots S_{n_0}^{(i)}$  converges because  $\Delta_i S_n^{(i)} \dots S_{n_0}^{(i)} \Delta_i^{-1}$  is a submatrix of  $Q_{n,n_0}$ .

On the other side, denoting by  $K$  the union of the  $K_i$ , the product  $M_n^{K^c} \dots M_{n_0}^{K^c}$  and the sum of products  $\sum_{i=n_0}^n M_n^K \dots M_{i+1}^K M_i^{K,K^c} M_{i-1}^{K^c} \dots M_{n_0}^{K^c}$  converge when  $n \rightarrow \infty$  because they are submatrices of  $Q_{n,n_0}$ . The first converges to 0: by the definitions of  $K$  and the vectors  $V_i$ , the rows of  $Q$  whose indexes belong to  $K^c$  are null.

□

**Remark 2.1.** The condition that  $\sum_{i=1}^{\infty} M_n^K \dots M_{i+1}^K M_i^{K,K^c} M_{i-1}^{K^c} \dots M_1^{K^c}$  converges cannot be avoided. Suppose for instance that  $M_n^K \dots M_1^K$  converges and that  $M_n^{K,K^c}$  is for any  $n$  the identity  $\frac{d}{2} \times \frac{d}{2}$  matrix, let  $I_{\frac{d}{2}}$ ,  $d$  even. Suppose also  $M_n^{K^c} = d_n I_{\frac{d}{2}}$  for any  $n$ , where the positive reals  $d_n$  satisfy  $\lim_{n \rightarrow \infty} d_n \dots d_1 = 0$  and  $\sum_{n=1}^{\infty} d_n \dots d_1 = \infty$ . Then if the  $M_n$  have a common right-eigenvector for the eigenvalue 1, it has the form  $\begin{pmatrix} W \\ \theta_{\frac{d}{2}} \end{pmatrix}$  where  $W$  is an eigenvector of the  $M_n^K$  and  $\theta_{\frac{d}{2}}$  is the  $\frac{d}{2}$ -dimensional null column-vector. Since  $\left( \sum_{i=1}^n M_n^K \dots M_{i+1}^K M_i^{K,K^c} M_{i-1}^{K^c} \dots M_1^{K^c} \right) W = \left( \sum_{i=1}^n d_{i-1} \dots d_1 \right) W$  diverges,  $M_n \dots M_1$  also do although  $M_n^{K^c} \dots M_1^{K^c}$  converges to the null matrix.

**Remark 2.2.** The two conditions of (ii) are satisfied – assuming that the ones of (i) are – if all the submatrices  $M_n^{K^c}$  have spectral radius less than 1, or if their eigenvalues greater or equal to 1 disappear in the product  $M_{n+h}^{K^c} \dots M_n^{K^c}$  for some fixed  $h$  and for any  $n$ .

**Remark 2.3.** Let us compare now the problem of the convergence of  $Q_n = M_n \dots M_1$  to the one of

$$R_n := \frac{M_n \dots M_1}{\|M_n \dots M_1\|}.$$

Let  $(M_n)$  be a sequence of complex-valued matrices such that  $R_n$  converges. Since the limit matrix  $R$  has norm 1, it has some nonnull columns; let us prove that they are right-eigenvectors of each matrix  $M$  that occurs infinitely many times in the sequence  $(M_n)$ . The nonnegative real  $\lambda_n := \|MR_{n-1}\|$  is bounded by  $\|M\|$  and satisfy  $MR_{n-1} = \lambda_n R_n$  for any  $n$  such that  $M_n = M$ , so it has at least one limit point  $\lambda$  that satisfy  $MR = \lambda R$ , and the columns of  $R$  are right-eigenvectors of  $M$  for the eigenvalue  $\lambda$ .

Suppose that  $\lambda \neq 0$  – perhaps it is not possible that  $\lambda = 0$ . The convergence of  $R_n$  can hold without the convergence of  $\left(\frac{1}{\lambda}M_n\right) \dots \left(\frac{1}{\lambda}M_1\right)$ , see for instance the case where all the  $M_n$  are equal to  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . But conversely the convergence of this last product to a nonnull matrix, for some  $\lambda \neq 0$ , implies obviously the one of  $R_n$ .

#### REFERENCES

- [1] I. DAUBECHIES & J. C. LAGARIAS, Sets of matrices all infinite products of which converge, *Linear Algebra and its Applications* **161** (1992), 227-263.
- [2] I. DAUBECHIES & J. C. LAGARIAS, Corrigendum/addendum to: Sets of matrices all infinite products of which converge, *Linear Algebra and its Applications* **327** (2001), 69-83.
- [3] L. ELSNER & S. FRIEDLAND, Norm conditions for convergence of infinite products, *Linear Algebra and its Applications* **250** (1997), 133-142, *download the list of publications*
- [4] P. ERDŐS, On a family of symmetric Bernoulli convolutions. *Amer. J. Math.* **61** (1939), 974-976, *download*.
- [5] J. LORENZ, Convergence of products of stochastic matrices with positive diagonal and the opinion dynamics background (2007), *download*.
- [6] A. MUKHERJEA, A. NAKASSIS & J.S. RATTI, On the distribution of the limit of products of i. i. d.  $2 \times 2$  random stochastic matrices, *J. Theor. Probab.* **12** (1999), 571-583.
- [7] A. MUKHERJEA & A. NAKASSIS, On the continuous singularity of the limit distribution of products of i. i. d.  $d \times d$  stochastic matrices, *J. Theor. Probab.* **15** (2002), 903-918.
- [8] E. OLIVIER, N. SIDOROV, & A. THOMAS, On the Gibbs properties of Bernoulli convolutions related to  $\beta$ -numeration in multinacci bases, *Monatsh. Math.* **145** (2005), 145–174, *download*.
- [9] E. SENETA, Non-negative matrices and Markov chains, *Springer Series in Statistics. New York - Heidelberg - Berlin: Springer- Verlag XV* (1981), *partial download*.

(Alain Thomas) LATP, 39, RUE JOLIOT-CURIE,  
13453 MARSEILLE, CEDEX 13, FRANCE  
*E-mail address:* thomas@cmi.univ-mrs.fr