RANK-CRANK TYPE PDE'S FOR HIGHER LEVEL APPELL FUNCTIONS

SANDER ZWEGERS

ABSTRACT. In this paper we consider level l Appell functions, and find a partial differential equation for all odd l. For l = 3 this recovers the Rank-Crank PDE, found by Atkin and Garvan, and for l = 5we get a similar PDE found by Garvan.

1. INTRODUCTION AND STATEMENT OF RESULTS

Dyson in [5] introduced the *rank* of a partition, to explain the first two of the three Ramanujancongruences

$$p(5n+4) \equiv 0 \pmod{5},$$

$$p(7n+5) \equiv 0 \pmod{7},$$

$$p(11n+6) \equiv 0 \pmod{11}.$$
(1.1)

Here p(n) denotes the number of partitions of n. He defined the rank of a partition as the largest part minus the number of its parts and conjectured that the partitions of 5n + 4 (resp. 7n + 5) form 5 (resp. 7) groups of equal size when sorted by their ranks modulo 5 (resp. 7). This was later proven by Atkin and Swinnerton-Dyer in [3]. We are interested here in the generating function

$$R(w;q) := \sum_{\lambda} w^{\operatorname{rank}(\lambda)} q^{||\lambda||} = \frac{(1-w)}{(q)_{\infty}} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{n}{2}(3n+1)}}{1 - wq^n},$$

where $(q)_{\infty} := \prod_{n=1}^{\infty} (1-q^n)$. In the first sum the λ run over all partitions, rank (λ) denotes the rank of λ and $||\lambda||$ denotes the size of the partition (the sum of all the parts).

Another partition statistic is the so called *crank* of a partition. For the generating function we have

$$\begin{split} C(w;q) &:= \sum_{\lambda} w^{\operatorname{crank}(\lambda)} q^{||\lambda||} = \prod_{n=1}^{\infty} \frac{(1-q^n)}{(1-wq^n) (1-w^{-1}q^n)} \\ &= (1-w) \frac{(q)_{\infty}^2}{\sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{1}{2}n(n-1)} w^n}. \end{split}$$

The crank was introduced by Andrews and Garvan in [1] to explain the Ramanujan congruence (1.1) with modulus 11.

Date: October 16, 2018.

²⁰⁰⁰ Mathematics Subject Classification. 11F11, 11F27, 11F50.

SANDER ZWEGERS

In the setting of Jacobi forms it is more natural to consider the following modified rank and crank generating functions

$$\mathcal{R}(z;\tau) := \frac{w^{1/2}q^{-1/24}}{1-w}R(w,q),$$
$$\mathcal{C}(z;\tau) := \frac{w^{1/2}q^{-1/24}}{1-w}C(w,q).$$

Here we use $w = \exp(2\pi i z)$ and $q = \exp(2\pi i \tau)$, with $z \in \mathbb{C}$ and τ in the complex upper half plane \mathbb{H} .

Remark 1.1. C is a meromorphic Jacobi form of weight 1/2 and index -1/2 and in [7] it is shown that \mathcal{R} is mock Jacobi form of weight 1/2 and index -3/2.

The two (modified) generating functions are related by a partial differential equation, which we will refer to as the Rank-Crank PDE.

Theorem 1.2 (see [2]). If we define the heat operator H by

$$\mathbf{H} := \frac{3}{\pi i} \frac{\partial}{\partial \tau} + \frac{1}{(2\pi i)^2} \frac{\partial^2}{\partial z^2},$$

then

$$\mathrm{H}\,\mathcal{R}=2\eta^2\mathcal{C}^3$$

where η is the Dedekind η -function, given by $\eta(\tau) = q^{1/24}(q)_{\infty}$.

Note that the identity found in [2] is slightly different, because they use a different normalization. However, the two are easily seen to be equivalent. In [4] it is explained how the Rank-Crank PDE arises naturally in the setting of certain non-holomorphic Jacobi forms and a generalization is given to partial differential equations for an infinite family of related functions.

The method in [4] works only in certain special cases and no results are found for the level l Appell functions

$$A_l(z;\tau) := w^{l/2} \sum_{n \in \mathbb{Z}} \frac{(-1)^{ln} q^{\frac{l}{2}n(n+1)}}{1 - wq^n} \qquad l \in \mathbb{Z}_{>0},$$
(1.2)

for values of l higher than 3.

Garvan ([6]), however, found the following PDE for a level 5 Appell function

Theorem 1.3 (Garvan). Let

$$G_5(z;\tau) := \frac{A_5(z;\tau)}{\eta(\tau)^3},$$

and define the heat operator

$$\mathbf{H} := \frac{5}{\pi i} \frac{\partial}{\partial \tau} + \frac{1}{(2\pi i)^2} \frac{\partial^2}{\partial z^2},$$

then

$$\left(\mathrm{H}^2 - E_4\right)G_5 = 24\eta^2 \mathcal{C}^5,$$

where E_4 is the usual Eisenstein series

$$E_4(\tau) = 1 + 240 \sum_{n=1}^{\infty} \left(\sum_{d|n} d^3\right) q^n.$$

Note that the identity found by Garvan is slightly different, because he uses a different normalization. However, the two are easily seen to be equivalent.

This theorem is a special case of the following

Theorem 1.4. Let *l* be an odd positive integer. Define

$$\mathcal{H}_k := \frac{l}{\pi i} \frac{\partial}{\partial \tau} + \frac{1}{(2\pi i)^2} \frac{\partial^2}{\partial z^2} - \frac{l(2k-1)}{12} E_2,$$

$$\mathcal{H}^k := \mathcal{H}_{2k-1} \mathcal{H}_{2k-3} \cdots \mathcal{H}_3 \mathcal{H}_1,$$

where $E_2(\tau) = 1 - 24 \sum_{n=1}^{\infty} \left(\sum_{d|n} d \right) q^n$ is the usual Eisenstein series in weight 2. Then there exist holomorphic modular forms f_j (j = 0, 2, 4, ..., l - 1) on $SL_2(\mathbb{Z})$ of weight j, such that

$$\sum_{k=0}^{(l-1)/2} f_{l-2k-1} \mathcal{H}^k A_l = (l-1)! f_0 \eta^l \mathcal{C}^l.$$

Remark 1.5. In the proof of the theorem we will see an explicit construction for the f_j 's for given l.

In the next section we will proof Theorem 1.4 and in section 3 we will look at the first few cases and in particular we'll see that the theorem for l = 5 is equivalent to Theorem 1.3.

Acknowledgements

The author wishes to thank Frank Garvan and Karl Mahlburg for helpful discussions.

2. Proof of Theorem 1.4

Throughout we assume that l is an odd positive integer. We (trivially) have

$$A_l(z+1;\tau) = -A_l(z;\tau),$$
(2.1)

and if we replace z by $z + \tau$ and n by n - 1 in (1.2) we find

$$e^{-2\pi i l z - \pi i l \tau} A_l(z + \tau; \tau) = -w^{-l/2} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{l}{2}n(n-1)}}{1 - w q^n},$$

and so

$$A_{l}(z;\tau) + e^{-2\pi i l z - \pi i l \tau} A_{l}(z+\tau;\tau) = -w^{-l/2} \sum_{n \in \mathbb{Z}} \frac{(-1)^{n} q^{\frac{l}{2}n(n-1)}}{1 - wq^{n}} \left(1 - w^{l} q^{ln}\right)$$

$$= -w^{-l/2} \sum_{n \in \mathbb{Z}} (-1)^{n} q^{\frac{l}{2}n(n-1)} \sum_{r=0}^{l-1} w^{r} q^{nr}$$

$$= -\sum_{r=0}^{l-1} w^{r-l/2} q^{-\frac{1}{2l}(r-l/2)^{2}} \sum_{n \in \mathbb{Z}} (-1)^{n} q^{\frac{l}{2}(n-1/2+r/l)^{2}}$$

$$= -\sum_{r=0}^{l-1} e^{2\pi i (r-l/2)z - \frac{\pi i}{l}(r-l/2)^{2\tau}} \vartheta_{l,r}(\tau),$$
(2.2)

with

$$\vartheta_{l,r}(\tau) := \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{l}{2}(n-1/2+r/l)^2}.$$

It is easy to check that

$$\left(\frac{l}{\pi i}\frac{\partial}{\partial \tau} + \frac{1}{(2\pi i)^2}\frac{\partial^2}{\partial z^2}\right)e^{2\pi i(r-l/2)z - \frac{\pi i}{l}(r-l/2)^2\tau} = 0,$$

and that for functions $F: \mathbb{C} \times \mathbb{H} \to \mathbb{C}$

$$\mathcal{H}_k\left(F(z+1;\tau)\right) = \left(\mathcal{H}_kF\right)(z+1;\tau),$$
$$\mathcal{H}_k\left(e^{-2\pi i l z - \pi i l \tau}F(z+\tau;\tau)\right) = e^{-2\pi i l z - \pi i l \tau}\left(\mathcal{H}_kF\right)(z+\tau;\tau),$$

with \mathcal{H}_k as in the theorem. Hence we get from applying \mathcal{H}_1 to equations (2.1) and (2.2)

$$(\mathcal{H}_1 A_l)(z+1;\tau) = -(\mathcal{H}_1 A_l)(z;\tau),$$

and

$$(\mathcal{H}_{1}A_{l})(z;\tau) + e^{-2\pi i l z - \pi i l \tau} (\mathcal{H}_{1}A_{l})(z+\tau;\tau)$$

= $-2l \sum_{r=0}^{l-1} e^{2\pi i (r-l/2)z - \frac{\pi i}{l}(r-l/2)^{2}\tau} (\mathbf{D}_{1/2} \vartheta_{l,r})(\tau),$

with the operator \mathbf{D}_k defined by

$$\mathbf{D}_k := \frac{1}{2\pi i} \frac{\partial}{\partial \tau} - \frac{k}{12} E_2.$$

If we now apply $\mathcal{H}_3, \mathcal{H}_5, \ldots$, upto \mathcal{H}_{2k-1} we find

$$(\mathcal{H}^k A_l)(z+1;\tau) = -(\mathcal{H}^k A_l)(z;\tau), \qquad (2.3)$$

and

$$(\mathcal{H}^{k}A_{l})(z;\tau) + e^{-2\pi i l z - \pi i l \tau} (\mathcal{H}^{k}A_{l})(z+\tau;\tau) = -(2l)^{k} \sum_{r=0}^{l-1} e^{2\pi i (r-l/2)z - \frac{\pi i}{l}(r-l/2)^{2}\tau} (\mathbf{D}^{k} \vartheta_{l,r})(\tau),$$
(2.4)

with

$$\mathbf{D}^k := \mathbf{D}_{2k-3/2} \, \mathbf{D}_{2k-7/2} \cdots \mathbf{D}_{5/2} \, \mathbf{D}_{1/2}$$

We need the following

Lemma 2.1. Let l be an odd positive integer, then there exist holomorphic modular forms F_j (j = 0, 2, 4, ..., l-1) on $SL_2(\mathbb{Z})$ of weight j, such that

$$\sum_{k=0}^{(l-1)/2} F_{l-2k-1} \mathbf{D}^k \,\vartheta_{l,r} = 0$$
(2.5)

for all $r \in \mathbb{Z}$.

4

If we now define

$$P = \sum_{k=0}^{(l-1)/2} f_{l-2k-1} \mathcal{H}^k A_l,$$

with $f_{l-2k-1} = (2l)^{-k} F_{l-2k-1}$ and F_j as in the lemma, then we see from equation (2.3) and (2.4)

$$P(z+1;\tau) = e^{-2\pi i l z - \pi i l \tau} P(z+\tau;\tau) = -P(z;\tau).$$
(2.6)

Now consider the Jacobi theta function

$$\begin{split} \vartheta(z;\tau) &:= \sum_{n \in \mathbb{Z}} (-1)^n w^{n+1/2} q^{\frac{1}{2}(n+1/2)^2} \\ &= w^{1/2} q^{1/8} \prod_{n=1}^\infty (1-q^n) (1-wq^n) (1-w^{-1}q^{n-1}) \\ &= -\frac{\eta(\tau)^2}{\mathcal{C}(z;\tau)}. \end{split}$$

This function satisfies

$$\vartheta(z+1;\tau) = e^{2\pi i z + \pi i \tau} \vartheta(z+\tau;\tau) = -\vartheta(z;\tau), \qquad (2.7)$$

 $z\mapsto \vartheta(z;\tau)$ has simple zeros in $\mathbb{Z}\tau+\mathbb{Z}$ and

$$\frac{1}{2\pi i} \left. \frac{\partial}{\partial z} \right|_{z=0} \vartheta(z;\tau) = \eta(\tau)^3.$$
(2.8)

Since the poles of $z \mapsto A_l(z; \tau)$ are simple poles in $\mathbb{Z}\tau + \mathbb{Z}$, the function $z \mapsto P(z; \tau)$ has poles of order l in $\mathbb{Z}\tau + \mathbb{Z}$, and so the function

$$p(z;\tau) := \vartheta(z;\tau)^l P(z;\tau),$$

is a holomorphic function as a function of z. Using (2.6) and (2.7) we find that

$$p(z+1;\tau) = p(z+\tau;\tau) = p(z;\tau),$$

from which we get that p is constant (as a function of z). To determine the constant, we consider the behaviour for $z \to 0$. From (1.2) we easily see that for $z \to 0$

$$A_l(z;\tau) = -\frac{1}{2\pi i}\frac{1}{z} + \mathcal{O}(1),$$

and so

$$P(z;\tau) = -f_0(\tau) \frac{(l-1)!}{(2\pi i)^l} \frac{1}{z^l} + \mathcal{O}\left(\frac{1}{z^{l-1}}\right)$$

Combining this with (2.8) we see

$$p(z;\tau) = -f_0(\tau)(l-1)!\eta(\tau)^{3l},$$

and so

$$P(z;\tau) = -f_0(\tau)(l-1)! \frac{\eta(\tau)^{3l}}{\vartheta(z;\tau)^l} = (l-1)! f_0(\tau)\eta(\tau)^l \mathcal{C}(z;\tau)^l,$$

which finishes the proof.

Proof of Lemma 2.1. Throughout, let l be an odd integer. Because of the trivial relations

$$\begin{aligned} \vartheta_{l,r+l} &= -\vartheta_{l,r} \\ \vartheta_{l,-r} &= -\vartheta_{l,r} \end{aligned}$$

it suffices to consider $\vartheta_{l,r}$ for $r = 1, 2, \ldots, (l-1)/2$. Define

$$\Theta_{l} = \begin{pmatrix} \vartheta_{l,1} \\ \vartheta_{l,2} \\ \vdots \\ \vartheta_{l,(l-1)/2} \end{pmatrix},$$

then Θ_l transforms as a (vector-valued) modular form of weight 1/2 on the full modular group $SL_2(\mathbb{Z})$:

$$\Theta_l(\tau+1) = \operatorname{diag}\left(\zeta_{8l}^{(l-2j)^2}\right)_{1 \le j \le (l-1)/2} \Theta_l(\tau),$$

$$\Theta_l(-1/\tau) = (-1)^{(l+1)/2} \sqrt{\tau/li} \ (2\sin 2\pi rk/l)_{1 \le r,k \le (l-1)/2} \ \Theta_l(\tau)$$

Using

$$E_2\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^2 E_2(\tau) + \frac{6}{\pi i}c(c\tau+d) \quad \text{for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}),$$

we can easily verify that

$$\mathbf{D}_k\left((c\tau+d)^{-k}f\left(\frac{a\tau+b}{c\tau+d}\right)\right) = (c\tau+d)^{-k-2}(\mathbf{D}_k f)\left(\frac{a\tau+b}{c\tau+d}\right),$$

and so

$$\mathbf{D}^{k}\left((c\tau+d)^{-1/2}\Theta_{l}\left(\frac{a\tau+b}{c\tau+d}\right)\right) = (c\tau+d)^{-2k-1/2}\left(\mathbf{D}^{k}\Theta_{l}\right)\left(\frac{a\tau+b}{c\tau+d}\right)$$

Now define the $(l-1)/2 \times (l-1)/2$ -matrix

$$T_l = \begin{pmatrix} \Theta_l & D^1 \Theta_l & D^2 \Theta_l & \cdots & D^{(l-3)/2} \Theta_l \end{pmatrix}$$

then T_l transforms as a (matrix-valued) modular form on the full modular group $SL_2(\mathbb{Z})$:

$$T_{l}(\tau+1) = \operatorname{diag}\left(\zeta_{8l}^{(l-2j)^{2}}\right)_{1 \le j \le (l-1)/2} T_{l}(\tau),$$

$$T_{l}(-1/\tau) = (-1)^{(l+1)/2} \sqrt{\tau/li} \left(2\sin 2\pi r k/l\right)_{1 \le r,k \le (l-1)/2} T_{l}(\tau) \operatorname{diag}\left(\tau^{2j-2}\right)_{1 \le j \le (l-1)/2}.$$
(2.9)

From this we see that

$$det(T_l(\tau+1)) = \zeta_{24}^{(l-1)(l-2)/2} det(T_l(\tau)),$$

$$det(T_l(-1/\tau)) = (-i\tau)^{(l-1)(l-2)/4} det(T_l(\tau)),$$

and so det (T_l) is a multiple of $\eta^{(l-1)(l-2)/2}$. We determine what that multiple is by looking at the lowest order terms:

First observe that by doing elementary column operations we get

$$\det(T_l(\tau)) = \det \left(\Theta_l \quad \partial_\tau \Theta_l \quad \partial_\tau^2 \Theta_l \quad \cdots \quad \partial_\tau^{(l-3)/2} \Theta_l \right),$$

with $\partial_{\tau} := \frac{1}{2\pi i} \frac{\partial}{\partial \tau}$.

For $1 \le r \le (l-1)/2$ we have

$$\vartheta_{l,r}(\tau) = q^{(l-2r)^2/8l} \left(1 + \mathcal{O}(q)\right),$$

 \mathbf{SO}

$$\begin{pmatrix} \Theta_l & \partial_\tau \Theta_l & \partial_\tau^2 \Theta_l & \cdots & \partial_\tau^{(l-3)/2} \Theta_l \end{pmatrix}$$

$$= \operatorname{diag} \left(q^{(l-2i)^2/8l} \right)_{1 \le i \le (l-1)/2} \cdot \left(\left(\frac{(l-2i)^2}{8l} \right)^{j-1} + \mathcal{O}(q) \right)_{1 \le i, j \le (l-1)/2},$$

$$\operatorname{det} \left(\Theta_l & \partial_\tau \Theta_l & \partial_\tau^2 \Theta_l & \cdots & \partial_\tau^{(l-3)/2} \Theta_l \right)$$

$$= q^{(l-1)(l-2)/48} \left(\operatorname{det}(B) + \mathcal{O}(q) \right),$$

and hence

$$\det(T_l(\tau)) = \det(B) \ \eta(\tau)^{(l-1)(l-2)/2}, \tag{2.10}$$

with

$$B_{ij} = \left(\frac{(l-2i)^2}{8l}\right)^{j-1}$$
 for $1 \le i, j \le (l-1)/2$.

B is a Vandermonde matrix: an $m \times n$ matrix *V*, such that $V_{ij} = \alpha_i^{j-1}$ with $\alpha_i \in \mathbb{R}$. Since a square Vandermonde matrix is invertible if and only if the α_i are distinct, we see that *B* is invertible. From (2.10) and the fact that η has no zeros on \mathbb{H} we then get that $T_l(\tau)$ is invertible for all $\tau \in \mathbb{H}$. We can rewrite the condition that (2.5) holds for $1 \leq r \leq (l-1)/2$ as

$$T_l \begin{pmatrix} F_{l-1} \\ F_{l-3} \\ \vdots \\ F_2 \end{pmatrix} + F_0 \operatorname{D}^{(l-1)/2} \Theta_l = 0.$$

If we take $F_0 = 1$ we get the other F_j 's by inverting T_l

$$\begin{pmatrix} F_{l-1} \\ F_{l-3} \\ \vdots \\ F_2 \end{pmatrix} = -T_l^{-1} \mathbf{D}^{(l-1)/2} \Theta_l.$$
 (2.11)

What remains to be shown is that the F_j found this way are holomorphic modular forms of weight j on $SL_2(\mathbb{Z})$. The modular transformation properties follow easy from (2.9) and those of $D^{(l-1)/2} \Theta_l$

$$\begin{aligned} \mathbf{D}^{(l-1)/2} \,\Theta_l(\tau+1) &= \operatorname{diag} \left(\zeta_{8l}^{(l-2j)^2} \right)_{1 \le j \le (l-1)/2} \mathbf{D}^{(l-1)/2} \,\Theta_l(\tau), \\ \mathbf{D}^{(l-1)/2} \,\Theta_l(-1/\tau) &= (-1)^{(l+1)/2} \sqrt{\tau/li} \,\,\tau^{l-1} \,\,(2\sin 2\pi rk/l)_{1 \le r,k \le (l-1)/2} \,\,\mathbf{D}^{(l-1)/2} \,\Theta_l(\tau). \end{aligned}$$

Since det T_l has no zeros on \mathbb{H} we get that F_j is a holomorphic function on \mathbb{H} . That it also doesn't have a pole at infinity follows from

$$D^{(l-1)/2} \Theta_l = \operatorname{diag} \left(q^{(l-2i)^2/8l} \right)_{1 \le i \le (l-1)/2} \cdot \begin{pmatrix} \mathcal{O}(1) \\ \mathcal{O}(1) \\ \vdots \\ \mathcal{O}(1) \end{pmatrix},$$
$$T_l = \operatorname{diag} \left(q^{(l-2i)^2/8l} \right)_{1 \le i \le (l-1)/2} \cdot (C_{ij} + \mathcal{O}(q))_{1 \le i, j \le (l-1)/2},$$
$$(l-1)/2 \operatorname{-matrix} C, \text{ with}$$
$$\operatorname{det}(C) = \operatorname{det}(B) \ne 0$$

for some (l-1)

$$\det(C) = \det(B) \neq 0.$$

3. Some examples

From (2.11) we can calculate the first few coefficients in the Fourier expansion of the F_j 's and since they are holomorphic modular forms on $SL_2(\mathbb{Z})$, that means that we can easily identify them.

 $\Theta_l = (\vartheta_{3,1}) = (\eta) \,.$

For l = 3, we have

Using

 $\mathbf{D}_{k/2}\left(\boldsymbol{\eta}^{k}\right) = \mathbf{0},$ $E_2 = \frac{12}{\pi i} \frac{\eta'}{\eta},$

we see

and so we find

which follows from

$$F_0(\tau) = 1$$
 and $F_2(\tau) = 0$.

 $D_{1/2}\Theta_l = 0,$

and

$$f_0(\tau) = 1/6$$
 and $f_2(\tau) = 0.$

If we put this into Theorem 1.4 and multiply by 6 we get

$$\mathcal{H}_1 A_3 = 2\eta^3 \mathcal{C}^3.$$

Using

$$\mathcal{R}(z;\tau) = \frac{A_3(z;\tau)}{\eta(\tau)} + e^{\pi i z - \pi i \tau/12},$$

we see

$$\mathcal{H}_{1/2}\mathcal{R} = \mathcal{H}_{1/2}\left(\frac{A_3}{\eta}\right) = \frac{\mathcal{H}_1A_3}{\eta} + 6A_3 \,\mathrm{D}_{-1/2}\left(\frac{1}{\eta}\right) = 2\eta^2 \mathcal{C}^3,$$

which is the Rank-Crank PDE.

For l = 5, we find from (2.11) ($F_0 = 1$)

$$F_4(\tau) = -\frac{11}{3600} - \frac{11}{15}q + \mathcal{O}(q^2),$$

$$F_2(\tau) = \mathcal{O}(q^2),$$

and hence we can identify them as

$$F_4 = -\frac{11}{3600}E_4$$
 and $F_2 = 0.$

 So

$$f_0 = \frac{1}{100}, \quad f_2 = 0 \quad \text{and} \quad f_4 = -\frac{11}{3600}E_4.$$

If we put this into Theorem 1.4 and multiply by 100 we get

$$\left(\mathcal{H}_3\mathcal{H}_1 - \frac{11}{36}E_4\right)A_5 = 24\eta^5\mathcal{C}^5.$$

We now rewrite this in terms of G_5 :

$$\begin{aligned} \mathcal{H}_{1}A_{5} &= \mathcal{H}_{1}\left(\eta^{3}G_{5}\right) \\ &= 10\left(\mathrm{D}_{3/2}\,\eta^{3}\right)G_{5} + \eta^{3}\mathcal{H}_{-1/2}G_{5} = \eta^{3}\mathcal{H}_{-1/2}G_{5}, \\ \mathcal{H}_{3}\mathcal{H}_{1}A_{5} &= \mathcal{H}_{3}\left(\eta^{3}\mathcal{H}_{-1/2}G_{5}\right) \\ &= 10\left(\mathrm{D}_{3/2}\,\eta^{3}\right)\mathcal{H}_{-1/2}G_{5} + \eta^{3}\mathcal{H}_{3/2}\mathcal{H}_{-1/2}G_{5} = \eta^{3}\mathcal{H}_{3/2}\mathcal{H}_{-1/2}G_{5}, \end{aligned}$$

and so we get

$$\left(\mathcal{H}_{3/2}\mathcal{H}_{-1/2} - \frac{11}{36}E_4\right)G_5 = 24\eta^2 \mathcal{C}^5.$$

Using

$$\mathcal{H}_{3/2}\mathcal{H}_{-1/2} = \mathrm{H}^2 + \frac{25}{3} \left(\frac{1}{2\pi i} E_2' - \frac{1}{12} E_2^2 \right)$$

and

$$\frac{1}{2\pi i}E_2' - \frac{1}{12}E_2^2 = -\frac{1}{12}E_4$$

we see that is equivalent to the statement of Theorem 1.3.

For l = 7 we find

$$F_6 = \frac{85}{74088}E_6$$
, $F_4 = -\frac{5}{252}E_4$, $F_2 = 0$ and $F_0 = 1$.

For l = 9

$$F_8 = -\frac{253}{559872}E_8, \qquad F_6 = \frac{53}{5832}E_6, \qquad F_4 = -\frac{13}{216}E_4, \qquad F_2 = 0, \qquad F_0 = 1.$$

For l = 11

$$F_{10} = \frac{7888}{39135393} E_{10}, \qquad F_8 = -\frac{6151}{1724976} E_8, \qquad F_6 = \frac{295}{8712} E_6,$$

$$F_4 = -\frac{53}{396} E_4, \qquad F_2 = 0, \qquad F_0 = 1.$$

And for
$$l = 13$$

$$F_{12} = -\frac{1462986875}{14412774445056}E_{12} + \frac{170060275}{5683867488}\Delta, \qquad F_{10} = \frac{377735}{296120448}E_{10},$$

$$F_{8} = -\frac{621665}{45556992}E_{8}, \qquad F_{6} = \frac{3281}{36504}E_{6}, \qquad F_{4} = -\frac{459}{1872}E_{4}, \qquad F_{2} = 0, \qquad F_{0} = 1.$$

SANDER ZWEGERS

References

- [1] G.E. Andrews and F.G. Garvan, *Dyson's crank of a partition*, Bull. Amer. Math. Soc. (N. S.) **18** No. 2 (1988), pages 167–171.
- [2] A.O.L. Atkin and F.G. Garvan, Relations between the ranks and the cranks of partitions, Ramanujan Journal 7, pages 137–152.
- [3] A.O.L. Atkin and H.P.F. Swinnerton-Dyer, *Some properties of partitions*, Proc. London Math. Soc. 4 (1954), pages 84–106.
- [4] K. Bringmann and S.P. Zwegers, Rank-crank type PDE's and non-holomorphic Jacobi forms, Math. Res. Lett., accepted for publication.
- [5] F. Dyson, Some guesses in the theory of partitions, Eureka (Cambridge) 8 (1944), pages 10–15.
- [6] F.G. Garvan, personal communication.
- [7] S.P. Zwegers, Mock theta functions I: Appell functions and the Mordell integral, in preparation.

School of Mathematical Sciences, University College Dublin, Belfield, Dublin 4, Ireland *E-mail address:* sander.zwegers@ucd.ie