

# Global Existence for Coupled Systems of Nonlinear Wave and Klein-Gordon Equations in Three Space Dimensions

Soichiro Katayama

## Abstract

We consider the Cauchy problem for coupled systems of wave and Klein-Gordon equations with quadratic nonlinearity in three space dimensions. We show global existence of small amplitude solutions under certain condition including the null condition on self-interactions between wave equations. Our condition is much weaker than the strong null condition introduced by Georgiev for this kind of coupled system. Consequently our result is applicable to certain physical systems, such as the Dirac-Klein-Gordon equations, the Dirac-Proca equations, and the Klein-Gordon-Zakharov equations.

## 1 Introduction

We consider the Cauchy problem for the following system:

$$(\square + m_i^2) u_i = F_i(u, \partial u, \partial_x \partial u), \quad i = 1, 2, \dots, N \quad (1.1)$$

in  $(0, \infty) \times \mathbb{R}^3$  with initial data

$$u(0, x) = \varepsilon f(x), \quad (\partial_t u)(0, x) = \varepsilon g(x) \text{ for } x = (x_1, x_2, x_3) \in \mathbb{R}^3, \quad (1.2)$$

where  $\square = \partial_t^2 - \Delta_x$ ,  $u = (u_j)_{1 \leq j \leq N}$ ,  $m_i \geq 0$  for  $1 \leq i \leq N$ , and  $\varepsilon$  is a small and positive parameter. Here each component  $u_j$  of  $u$  is supposed to be a real-valued unknown function of  $(t, x) \in (0, \infty) \times \mathbb{R}^3$ , and  $\Delta_x$  denotes the Laplacian in  $x$ -variables. In the above,  $\partial u$  and  $\partial_x \partial u$  are given by

$$\partial u = (\partial_a u_j)_{1 \leq j \leq N, 0 \leq a \leq 3}, \quad \partial_x \partial u = (\partial_k \partial_a u_j)_{1 \leq j \leq N, 1 \leq k \leq 3, 0 \leq a \leq 3},$$

respectively, with the notation

$$\partial_0 = \partial_t = \frac{\partial}{\partial t} \text{ and } \partial_k = \frac{\partial}{\partial x_k} \text{ for } 1 \leq k \leq 3.$$

Here, by writing  $(\partial_a u_j)_{j,a}$ , we mean that  $\partial_a u_j$ 's are arranged in dictionary order with respect to  $(j, a)$ . Similarly,  $(\partial_k \partial_a u_j)_{j,k,a}$  means that  $\partial_k \partial_a u_j$ 's are arranged

in dictionary order with respect to  $(j, k, a)$ . Similar convention will be used throughout this paper. We assume that, for  $1 \leq i \leq N$ , each  $F_i = F_i(\xi, \xi', \xi'')$  is a real-valued smooth function of  $(\xi, \xi', \xi'') \in \mathbb{R}^N \times \mathbb{R}^{4N} \times \mathbb{R}^{12N}$ , where  $\xi$ ,  $\xi'$ , and  $\xi''$  are independent variables for which  $u$ ,  $\partial u$ , and  $\partial_x \partial u$  are substituted in (1.1); more precisely, if we write

$$\begin{aligned}\xi &= (\xi_j)_{1 \leq j \leq N} \in \mathbb{R}^N, & \xi' &= (\xi'_{j,a})_{1 \leq j \leq N, 0 \leq a \leq 3} \in \mathbb{R}^{4N}, \\ \xi'' &= (\xi''_{j,k,a})_{1 \leq j \leq N, 1 \leq k \leq 3, 0 \leq a \leq 3} \in \mathbb{R}^{12N},\end{aligned}$$

then  $\xi_j$ ,  $\xi'_{j,a}$ , and  $\xi''_{j,k,a}$  are the independent variables for which  $u_j$ ,  $\partial_a u_j$ , and  $\partial_k \partial_a u_j$  are substituted in (1.1), respectively. We assume that  $F = (F_i)_{1 \leq i \leq N}$  vanishes of second order at the origin  $(\xi, \xi', \xi'') = (0, 0, 0)$ , namely

$$F(\xi, \xi', \xi'') = O(|\xi|^2 + |\xi'|^2 + |\xi''|^2) \text{ around } (\xi, \xi', \xi'') = (0, 0, 0).$$

For simplicity, we also assume that the system is quasi-linear. In other words, we assume that

$$F_i(\xi, \xi', \xi'') = \sum_{j=1}^N \sum_{k=1}^3 \sum_{a=0}^3 \gamma_{ka}^{ij}(\xi, \xi') \xi''_{j,k,a} + G_i(\xi, \xi'), \quad 1 \leq i \leq N, \quad (1.3)$$

where  $\gamma_{ka}^{ij} = \gamma_{ka}^{ij}(\xi, \xi')$  and  $G_i = G_i(\xi, \xi')$  are some functions vanishing of first and second order at the origin, respectively. Moreover, to assure the hyperbolicity of the system, we always assume the symmetricity condition

$$\gamma_{ka}^{ij}(\xi, \xi') = \gamma_{ka}^{ji}(\xi, \xi') \text{ and } \gamma_{kl}^{ij}(\xi, \xi') = \gamma_{lk}^{ij}(\xi, \xi') \quad (1.4)$$

for any  $(\xi, \xi') \in \mathbb{R}^N \times \mathbb{R}^{4N}$ ,  $1 \leq i, j \leq N$ ,  $1 \leq k, l \leq 3$ , and  $0 \leq a \leq 3$  (note that the last half of (1.4) is no restriction as far as we consider smooth solutions). For each  $i \in \{1, \dots, N\}$ , the equation (1.1) for  $u_i$  is called a (nonlinear) Klein-Gordon (resp. wave) equation if  $m_i > 0$  (resp.  $m_i = 0$ ).

For a while, we suppose that  $f = (f_i)_{1 \leq i \leq N}$ ,  $g = (g_i)_{1 \leq i \leq N} \in C_0^\infty(\mathbb{R}^3; \mathbb{R}^N)$  in (1.2). Under the conditions (1.3) and (1.4), the classical theory of nonlinear hyperbolic systems implies local existence of smooth solutions to the Cauchy problem (1.1)–(1.2) for sufficiently small  $\varepsilon$ . Hence we are interested in the sufficient condition for global existence of small amplitude solutions. Here we recall the known results briefly. If the nonlinearity  $F$  vanishes of third order at the origin, (1.1)–(1.2) admits a global solution for small  $\varepsilon$ . For arbitrary quadratic nonlinearity  $F$ , we also have global existence of small solutions if (1.1) is a system of nonlinear Klein-Gordon equations, namely if  $m_i > 0$  for all  $i = 1, \dots, N$  (see Klainerman [16] and Shatah [25]; see also Bachelot [2] and Hayashi-Naumkin-Ratno Bagus Edy Wibowo [6]). By contrast, this is not true if (1.1) is a system of wave equations (namely if  $m_1 = m_2 = \dots = m_N = 0$ ), and the solution to (1.1)–(1.2) with certain quadratic nonlinearity  $F$  may blow up in finite time no matter how small  $\varepsilon$  is (see John [10], [11]). Thus we need to put some condition

on quadratic nonlinearity, in order to obtain global solutions for wave equations. The *null condition* introduced by Klainerman [17] is one of such conditions. Before describing the null condition, we introduce the following notation: For a smooth function  $\Phi = \Phi(z)$  ( $z \in \mathbb{R}^d$ ), we write  $\Phi^{(q)}$  for the quadratic part of  $\Phi$  (“(q)” stands for “quadratic”); more precisely, for a smooth function  $\Phi = \Phi(z)$ , we define

$$\Phi^{(q)}(z) = \sum_{|\alpha|=2} \frac{(\partial_z^\alpha \Phi)(0)}{\alpha!} z^\alpha, \quad z = (z_1, \dots, z_d) \in \mathbb{R}^d, \quad (1.5)$$

where  $\partial_z = (\partial_{z_1}, \dots, \partial_{z_d})$ ,  $\alpha$  is a multi-index, and we have used the standard notation of multi-indices. The null condition can be stated as follows:

**Definition 1.1 (The null condition)** *We say that a function  $F = (F_i)_{1 \leq i \leq N}$  of  $(\xi, \xi', \xi'') \in \mathbb{R}^N \times \mathbb{R}^{4N} \times \mathbb{R}^{12N}$  satisfies the null condition if each  $F_i$  ( $1 \leq i \leq N$ ) satisfies*

$$F_i^{(q)}(\lambda, (X_a \mu_j)_{1 \leq j \leq N, 0 \leq a \leq 3}, (X_k X_a \nu_j)_{1 \leq j \leq N, 1 \leq k \leq 3, 0 \leq a \leq 3}) = 0 \quad (1.6)$$

for any  $\lambda = (\lambda_j)_{1 \leq j \leq N}$ ,  $\mu = (\mu_j)_{1 \leq j \leq N}$ ,  $\nu = (\nu_j)_{1 \leq j \leq N} \in \mathbb{R}^N$ , and any  $X = (X_a)_{0 \leq a \leq 3} \in \mathbb{R}^4$  satisfying  $X_0^2 - X_1^2 - X_2^2 - X_3^2 = 0$ , where  $F_i^{(q)} = F_i^{(q)}(\xi, \xi', \xi'')$  is the quadratic part of  $F_i$  given by (1.5) with  $\Phi = F_i$  and  $z = (\xi, \xi', \xi'')$ .

If  $F$  satisfies the null condition, then we have global existence of small solutions for systems of wave equations (1.1) with  $m_1 = \dots = m_N = 0$  (see Klainerman [17] and Christodoulou [3]).

Klainerman used the so-called vector field method in [16] and [17]. But his method is not directly applicable to systems consisting of both wave and Klein-Gordon equations, because the scaling operator  $S = t\partial_t + \sum_{k=1}^3 x_k \partial_k$ , which was used in [17], is compatible with the wave equations, but not with the Klein-Gordon equations. This causes some difficulty in the treatment of the null condition, and hence Georgiev [4] introduced the *strong null condition* to obtain global existence of small solutions for coupled systems of nonlinear wave and Klein-Gordon equations (see Section 4 below for the detail), where  $F$  is said to satisfy the strong null condition if (1.6) with  $1 \leq i \leq N$  holds for any  $\lambda, \mu, \nu \in \mathbb{R}^N$  and any  $X \in \mathbb{R}^4$  not necessarily satisfying  $X_0^2 - X_1^2 - X_2^2 - X_3^2 = 0$ .

Our aim in this paper is to establish a global existence theorem for systems of the nonlinear wave and Klein-Gordon equations under more natural and weaker condition than the strong null condition, so that it can cover the previous results for wave equations and the Klein-Gordon equations, as well as some important examples from physics.

## 2 The Main Result and Examples

First we introduce some notation. Suppose that we can take some natural number  $N_1$  such that we have

$$m_i > 0 \text{ for } 1 \leq i \leq N_1, \text{ and } m_i = 0 \text{ for } N_1 + 1 \leq i \leq N \quad (2.1)$$

in (1.1). We set

$$v = (v_j)_{1 \leq j \leq N_1} := (u_j)_{1 \leq j \leq N_1} \text{ and } w = (w_j)_{1 \leq j \leq N_2} := (u_{N_1+j})_{1 \leq j \leq N_2}, \quad (2.2)$$

where  $N_2 = N - N_1$ , so that  $u = (u_1, \dots, u_{N_1}, u_{N_1+1}, \dots, u_N) = (v, w)$ . Note that each  $v_j (= u_j)$  satisfies a nonlinear Klein-Gordon equation, while each  $w_j (= u_{N_1+j})$  satisfies a nonlinear wave equation. In accordance with (2.2), we introduce independent variables  $(\eta, \zeta) \in \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$ ,  $(\eta', \zeta') \in \mathbb{R}^{4N_1} \times \mathbb{R}^{4N_2}$ , and  $(\eta'', \zeta'') \in \mathbb{R}^{12N_1} \times \mathbb{R}^{12N_2}$  to write

$$\begin{aligned} \xi &= (\xi_j)_{1 \leq j \leq N} =: ((\eta_j)_{1 \leq j \leq N_1}, (\zeta_j)_{1 \leq j \leq N_2}) = (\eta, \zeta), \\ \xi' &= (\xi'_{j,a})_{1 \leq j \leq N, 0 \leq a \leq 3} =: ((\eta'_{j,a})_{1 \leq j \leq N_1, 0 \leq a \leq 3}, (\zeta'_{j,a})_{1 \leq j \leq N_2, 0 \leq a \leq 3}) = (\eta', \zeta'), \\ \xi'' &= (\xi''_{j,k,a})_{1 \leq j \leq N, 1 \leq k \leq 3, 0 \leq a \leq 3} \\ &=: ((\eta''_{j,k,a})_{1 \leq j \leq N_1, 1 \leq k \leq 3, 0 \leq a \leq 3}, (\zeta''_{j,k,a})_{1 \leq j \leq N_2, 1 \leq k \leq 3, 0 \leq a \leq 3}) = (\eta'', \zeta''). \end{aligned}$$

Correspondingly, we write  $\partial u = (\partial v, \partial w)$  and  $\partial_x \partial u = (\partial_x \partial v, \partial_x \partial w)$ . For a smooth function  $\Phi = \Phi(\xi, \xi', \xi'')$ , we define

$$\Phi^{(W)}(\zeta, \zeta', \zeta'') = \Phi^{(q)}((\eta, \zeta), (\eta', \zeta'), (\eta'', \zeta'')) \Big|_{(\eta, \eta', \eta'') = (0, 0, 0)} \quad (2.3)$$

for  $(\zeta, \zeta', \zeta'') \in \mathbb{R}^{N_2} \times \mathbb{R}^{4N_2} \times \mathbb{R}^{12N_2}$ , where  $\Phi^{(q)}$  is the quadratic part of  $\Phi$  given by (1.5) with  $z = (\xi, \xi', \xi'')$ . Thus  $F_i^{(W)}(w, \partial w, \partial_x \partial w)$  appearing below represents the self-interaction between the solutions to wave equations (“(W)” in the notation (2.3) stands for “wave”).

Now we are in a position to state our main result.

**Theorem 2.1** *Suppose that (1.3), (1.4), and (2.1) are fulfilled. Assume that the following two conditions (a) and (b) hold:*

- (a)  $(F_i^{(W)})_{N_1+1 \leq i \leq N}$  satisfies the null condition.
- (b) There exist two (empty or non-empty) sets  $\mathcal{I}_1$  and  $\mathcal{I}_2$  satisfying

$$\mathcal{I}_1 \cup \mathcal{I}_2 = \{1, \dots, N_2\}, \quad \mathcal{I}_1 \cap \mathcal{I}_2 = \emptyset, \quad (2.4)$$

and the following properties:

- (b-i) For any  $k \in \mathcal{I}_1$ , we have

$$\frac{\partial F_i^{(q)}}{\partial \zeta_k}((\eta, \zeta), (\eta', \zeta'), (\eta'', \zeta'')) \left( = \frac{\partial F_i^{(q)}}{\partial \xi_{N_1+k}}(\xi, \xi', \xi'') \right) = 0 \quad (2.5)$$

for all  $i = 1, \dots, N$ , and all  $(\xi, \xi', \xi'') \in \mathbb{R}^N \times \mathbb{R}^{4N} \times \mathbb{R}^{12N}$ .

(b–ii) For any  $k \in \mathcal{I}_2$ , there exist some functions  $\mathcal{G}_{k,a} = \mathcal{G}_{k,a}(\xi, \xi')$  with  $0 \leq a \leq 3$  such that

$$F_{N_1+k}(\phi, \partial\phi, \partial_x\partial\phi) = \sum_{a=0}^3 \partial_a \{ \mathcal{G}_{k,a}(\phi, \partial\phi) \} \quad (2.6)$$

holds for any  $\phi = \phi(t, x) \in C^2((0, \infty) \times \mathbb{R}^3; \mathbb{R}^N)$ , and

$$\frac{\partial \mathcal{G}_{k,a}^{(q)}}{\partial \zeta_l}((\eta, \zeta), (\eta', \zeta')) \left( = \frac{\partial \mathcal{G}_{k,a}^{(q)}}{\partial \xi_{N_1+l}}(\xi, \xi') \right) = 0, \quad 0 \leq a \leq 3 \quad (2.7)$$

holds for all  $l \in \mathcal{I}_1$ , and all  $(\xi, \xi') \in \mathbb{R}^N \times \mathbb{R}^{4N}$ , where  $\mathcal{G}_{k,a}^{(q)}$  is the quadratic part of  $\mathcal{G}_{k,a}$ .

Then, for any  $f, g \in \mathcal{S}(\mathbb{R}^3; \mathbb{R}^N)$ , there exists a positive constant  $\varepsilon_0$  such that for any  $\varepsilon \in (0, \varepsilon_0]$  the Cauchy problem (1.1)–(1.2) admits a unique global solution  $u \in C^\infty([0, \infty) \times \mathbb{R}^3; \mathbb{R}^N)$ . Here  $\mathcal{S}$  denotes the Schwartz class, the class of rapidly decreasing functions.

Here and hereafter, we say that  $(F_i^{(W)})_{N_1+1 \leq i \leq N}$  satisfies the null condition, if each  $F_i^{(W)} = F_i^{(W)}(\zeta, \zeta', \zeta'')$  with  $N_1 + 1 \leq i \leq N$  satisfies

$$F_i^{(W)}(\lambda, (X_a \mu_j)_{1 \leq j \leq N_2, 0 \leq a \leq 3}, (X_k X_a \nu_j)_{1 \leq j \leq N_2, 1 \leq k \leq 3, 0 \leq a \leq 3}) = 0$$

for any  $\lambda = (\lambda_j)_{1 \leq j \leq N_2}$ ,  $\mu = (\mu_j)_{1 \leq j \leq N_2}$ ,  $\nu = (\nu_j)_{1 \leq j \leq N_2} \in \mathbb{R}^{N_2}$ , and for any  $X = (X_a)_{0 \leq a \leq 3} \in \mathbb{R}^4$  satisfying  $X_0^2 - X_1^2 - X_2^2 - X_3^2 = 0$ . Notice that the null structure is required only for  $F_i^{(W)}$  with  $N_1 + 1 \leq i \leq N$  in Theorem 2.1. We define the null forms  $Q_0$  and  $Q_{ab}$  by

$$Q_0(\varphi, \psi) = (\partial_t \varphi)(\partial_t \psi) - (\nabla_x \varphi) \cdot (\nabla_x \psi), \quad (2.8)$$

$$Q_{ab}(\varphi, \psi) = (\partial_a \varphi)(\partial_b \psi) - (\partial_b \varphi)(\partial_a \psi), \quad 0 \leq a < b \leq 3, \quad (2.9)$$

where  $\nabla_x = (\partial_1, \partial_2, \partial_3)$ , and  $\cdot$  denotes the inner product in  $\mathbb{R}^3$ . Then we can easily check that the assumption (a) in Theorem 2.1 is equivalent to the following condition (refer to [17] for instance):

(a') There exist some constants  $A_{i,jk}^{\alpha\beta}$  and  $B_{i,jk}^{ab,\alpha\beta}$  such that

$$\begin{aligned} F_i^{(W)}(w, \partial w, \partial_x \partial w) &= \sum_{\substack{1 \leq j, k \leq N_2 \\ 0 \leq |\alpha|, |\beta| \leq 1}} A_{i,jk}^{\alpha\beta} Q_0(\partial^\alpha w_j, \partial^\beta w_k) \\ &+ \sum_{\substack{1 \leq j, k \leq N_2 \\ 0 \leq |\alpha|, |\beta| \leq 1}} \sum_{0 \leq a < b \leq 3} B_{i,jk}^{ab,\alpha\beta} Q_{ab}(\partial^\alpha w_j, \partial^\beta w_k) \end{aligned} \quad (2.10)$$

holds for any  $i = N_1 + 1, \dots, N$ , and any  $C^2$ -function  $w = (w_1, \dots, w_{N_2})$ , where  $\partial^\alpha = \partial_0^{\alpha_0} \partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}$  for a multi-index  $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3)$ .

The condition (b) in Theorem 2.1 is assumed in order to compensate for bad behavior of the solutions to the wave equations, as compared with their derivatives: Let  $u = (v, w)$  be the solution to (1.1). The condition (b-i) says that if  $k \in \mathcal{I}_1$ , then all of  $F_i^{(q)}(u, \partial u, \partial_x \partial u)$  can depend on  $(\partial w_k, \partial_x \partial w_k)$ , but *not* on  $w_k$  itself (remember that  $\zeta_k (= \xi_{N_1+k})$  is the variable corresponding to  $w_k (= u_{N_1+k})$ ), while the divergence structure in the condition (b-ii) assures that for each  $k \in \mathcal{I}_2$ ,  $w_k$  behaves better than we can expect in general (see Lemmas 3.6 and 3.7 below; observe that the equation for  $w_k (= u_{N_1+k})$  with  $k \in \mathcal{I}_2$  in (1.1) is  $\square w_k = F_{N_1+k}(u, \partial u, \partial_x \partial u) = \sum_{a=0}^3 \partial_a \{ \mathcal{G}_{k,a}(u, \partial u) \}$ ). Here we remark that the condition (b-i) does not imply (2.7) for  $(l, k) \in \mathcal{I}_1 \times \mathcal{I}_2$  in general, because we have

$$2Q_{ab}(\varphi, \psi) = \partial_a \{ \varphi(\partial_b \psi) - (\partial_b \varphi)\psi \} + \partial_b \{ (\partial_a \varphi)\psi - \varphi(\partial_a \psi) \} \quad (2.11)$$

for example (observe that  $Q_{ab}(\varphi, \psi)$  depends only on  $\partial \varphi$  and  $\partial \psi$ , while  $\varphi(\partial_b \psi) - (\partial_b \varphi)\psi$  on the right-hand side depends not only on  $\partial \varphi$  and  $\partial \psi$ , but also on  $\phi$  and  $\psi$ ).

To help the understanding of our condition, we give a typical example here. In what follows, for a finite family of functions  $\{\phi_\lambda\}_{\lambda \in \Lambda}$  and a function  $\psi$ , we write  $\psi = \sum'_{\lambda \in \Lambda} \phi_\lambda$  if there exists a family of constants  $\{c_\lambda\}_{\lambda \in \Lambda}$  such that  $\psi = \sum_{\lambda \in \Lambda} c_\lambda \phi_\lambda$ . Let  $u = (v, w) = (v, w_1, w_2)$  be an  $\mathbb{R}^3$ -valued function, and let  $m$  be a positive constant. Then the assumption in Theorem 2.1 is fulfilled with  $\mathcal{I}_1 = \{1\}$  and  $\mathcal{I}_2 = \{2\}$  for the following semilinear system:

$$\begin{aligned} (\square + m^2)v &= \sum'_{|\alpha|, |\beta| \leq 1} (\partial^\alpha v)(\partial^\beta v) + \sum'_{\substack{|\alpha| \leq 1 \\ 0 \leq b \leq 3}} (\partial^\alpha v)(\partial_b w_1) + \sum'_{|\alpha|, |\beta| \leq 1} (\partial^\alpha v)(\partial^\beta w_2) \\ &+ \sum'_{0 \leq a, b \leq 3} (\partial_a w_1)(\partial_b w_1) + \sum'_{\substack{0 \leq a \leq 3 \\ |\beta| \leq 1}} (\partial_a w_1)(\partial^\beta w_2) \\ &+ \sum'_{|\alpha|, |\beta| \leq 1} (\partial^\alpha w_2)(\partial^\beta w_2) + H_1(u, \partial u), \end{aligned} \quad (2.12)$$

$$\begin{aligned} \square w_1 &= \sum'_{|\alpha|, |\beta| \leq 1} (\partial^\alpha v)(\partial^\beta v) + \sum'_{\substack{|\alpha| \leq 1 \\ 0 \leq b \leq 3}} (\partial^\alpha v)(\partial_b w_1) + \sum'_{|\alpha|, |\beta| \leq 1} (\partial^\alpha v)(\partial^\beta w_2) \\ &+ \sum'_{j, k=1, 2} Q_0(w_j, w_k) + \sum'_{\substack{j, k=1, 2 \\ 0 \leq a < b \leq 3}} Q_{ab}(w_j, w_k) + H_2(u, \partial u), \end{aligned} \quad (2.13)$$

$$\square w_2 = \sum_{a=0}^3 \partial_a (C_{1,a} v^2 + C_{2,a} v w_2 + H_{3,a}(u)) + \sum'_{0 \leq a < b \leq 3} Q_{ab}(v, w_2), \quad (2.14)$$

where  $C_{j,a}$ 's are real constants, while  $H_1$ ,  $H_2$ , and  $H_{3,a}$  are smooth functions in their arguments satisfying  $H_1(u, \partial u)$ ,  $H_2(u, \partial u) = O(|u|^3 + |\partial u|^3)$  near  $(u, \partial u) = 0$ , and  $H_{3,a}(u) = O(|u|^3)$  near  $u = 0$ . We use (2.11) to treat  $Q_{ab}$  in (2.14).

Now we would like to see the relation between our theorem and the previous results. When  $m_i > 0$  for all  $i = 1, \dots, N$ , by regarding  $v = u$ , and by neglecting the meaningless conditions (a) and (b), Theorem 2.1 covers the previous results in [16] and [25] for nonlinear Klein-Gordon equations. Similarly, when  $m_i = 0$  for all  $i = 1, \dots, N$ , by regarding  $w = u$  and  $N_1 = 0$  (thus  $F_i^{(W)}$  is regarded as  $F_i^{(q)}$  for  $1 \leq i \leq N$ ), it also covers the previous results in [3] and [17] for nonlinear wave equations; note that the condition (b) for this case is automatically satisfied under the condition (a), because (2.10) implies (b) with the choice of  $\mathcal{I}_1 = \{1, \dots, N\}$  and  $\mathcal{I}_2 = \emptyset$ . It is easy to show that the strong null condition is satisfied if and only if each  $F_i^{(q)}$  ( $1 \leq i \leq N$ ) is a linear combination of  $Q_{ab}(\partial^\alpha u_j, \partial^\beta u_k)$  with  $1 \leq j, k \leq N$ ,  $|\alpha|, |\beta| \leq 1$  and  $0 \leq a < b \leq 3$ . Hence our conditions (a) and (b) are much weaker than the strong null condition in [4]. Note that some case of variable coefficients is also treated in [4], but we can easily modify our conditions (a) and (b) to treat such case.

The main difficulty in the proof of Theorem 2.1 lies in the fact that we can only use the vector fields which are compatible with both wave and Klein-Gordon equations. To prove Theorem 2.1, instead of the weighted  $L^2$ - $L^\infty$  estimate derived in [4], we use weighted  $L^\infty$ - $L^\infty$  estimates for wave equations (see Lemma 3.4 below), which require a smaller set of vector fields than the admissible set of vector fields for the Klein-Gordon equations. We also need some estimates for null forms without using the scaling operator  $S$ ; they will be given in Lemma 4.1 below. To treat  $F_i^{(W)}$  with  $1 \leq i \leq N_1$ , for which the null condition is *not* assumed, we adopt a technique used in Y. Tsutsumi [29], where the Dirac-Proca equations are considered (see (2.18)–(2.19) below). This technique is motivated by Bachelot [2] and Kosecki [19], and it is closely related to the normal form technique in Shatah [25]. We will prove Theorem 2.1 in Section 5.

We conclude this section with some examples from physics which can be treated by Theorem 2.1. Note that all the following examples are semilinear (or can be regarded as semilinear), and the conditions (1.3) and (1.4) in Theorem 2.1 are trivially satisfied. Thus we only have to check the conditions (a) and (b).

**Example 1 (The Dirac-Klein-Gordon equations)** Let us consider the Dirac equation coupled with the Klein-Gordon or wave equation:

$$-\sqrt{-1} \sum_{a=0}^3 \gamma_a \partial_a \psi + M\psi = \sqrt{-1} c \varphi \gamma_5 \psi, \quad (2.15)$$

$$(\square + m^2) \varphi = \psi^* H \psi \quad (2.16)$$

in  $(0, \infty) \times \mathbb{R}^3$ , where  $\sqrt{-1}$  denotes the imaginary unit,  $M, m \geq 0$ ,  $c$  is a real constant,  $H$  is a  $4 \times 4$  Hermitian matrix,  $\psi$  is a  $\mathbb{C}^4$ -valued function,  $\varphi$  is a real valued function, and  $\psi^*$  denotes the complex conjugate transpose of  $\psi$ .  $\gamma_a$  ( $0 \leq a \leq 3$ ) in the above are  $4 \times 4$  matrices satisfying  $\gamma_a \gamma_b + \gamma_b \gamma_a = 2g_{ab}I$  for  $0 \leq a, b \leq 3$ , where  $I$  is the  $4 \times 4$  identity matrix, and  $(g_{ab})_{0 \leq a, b \leq 3} = \text{diag}(1, -1, -1, -1)$ ;  $\gamma_5$

is defined by  $\gamma_5 = -\sqrt{-1}\gamma_0\gamma_1\gamma_2\gamma_3$ . The initial data are supposed to be sufficiently small.

We set  $D_M^\pm = \pm\sqrt{-1}\sum_{a=0}^3\gamma_a\partial_a + MI$ . Since we have  $\gamma_a\gamma_5 = -\gamma_5\gamma_a$  for  $0 \leq a \leq 3$ , we get  $D_M^+\gamma_5 = \gamma_5D_M^-$ . Therefore, operating  $D_M^+$  to (2.15), we get

$$\begin{aligned} (\square + M^2)\psi &= \sqrt{-1}cD_M^+(\varphi\gamma_5\psi) = -c\sum_{a=0}^3(\partial_a\varphi)\gamma_a\gamma_5\psi + \sqrt{-1}c\varphi\gamma_5(D_M^-\psi) \\ &= -c\sum_{a=0}^3(\partial_a\varphi)\gamma_a\gamma_5\psi - c^2\varphi^2\psi, \end{aligned} \quad (2.17)$$

where we have used (2.15) and  $\gamma_5\gamma_5 = I$  to obtain the last identity. If  $M > 0$  and  $m > 0$ , the system (2.16)–(2.17) is a system of the nonlinear Klein-Gordon equations, and we have the global solution. When  $M > 0$  and  $m = 0$ , putting  $u = (v, w)$  with  $v = (\operatorname{Re}\psi, \operatorname{Im}\psi)$  and  $w (= w_1) = \varphi$ , we see that the conditions (a) and (b) are satisfied for the system (2.16)–(2.17), and thus Theorem 2.1 is applicable; more precisely the condition (b) is satisfied with  $\mathcal{I}_1 = \{1\}$  and  $\mathcal{I}_2 = \emptyset$ . The global existence result for this case where  $M > 0$  and  $m = 0$ , with compactly supported initial data, has been already obtained by Bachelot [2]. Differently from [2], we can also treat the case where  $M = 0$  and  $m > 0$ . Indeed, the first identity in (2.17) with  $M = 0$  can be read as  $\square\psi = -c\sum_{a=0}^3\gamma_a\partial_a(\varphi\gamma_5\psi)$ , and putting  $u = (u_i)_{1 \leq i \leq 9} = (v, w)$  with  $v (= v_1) = \varphi$  and  $w = (w_k)_{1 \leq k \leq 8} = (\operatorname{Re}\psi, \operatorname{Im}\psi)$ , we can verify the conditions (a) and (b) with  $\mathcal{I}_1 = \emptyset$  and  $\mathcal{I}_2 = \{1, \dots, 8\}$ . This last case is closely connected to the next example, the Dirac-Proca equations.

**Example 2 (The Dirac-Proca equations)** Y. Tsutsumi [29] proved the global existence of small solutions to the Dirac-Proca equations, which can be reduced to the following coupled system of the massless Dirac and the Klein-Gordon equations:

$$-\sqrt{-1}\sum_{a=0}^3\gamma_a\partial_a\psi = -\frac{1}{2}\sum_{a=0}^3g_{aa}A_a\gamma_a(I + \gamma_5)\psi, \quad (2.18)$$

$$(\square + m^2)A_a = \frac{1}{2}\psi^*\gamma_0\gamma_a(I + \gamma_5)\psi, \quad a = 0, 1, 2, 3 \quad (2.19)$$

with the constraint  $\sum_{a=0}^3\partial_aA_a = 0$  at  $t = 0$ , where  $m > 0$ ,  $\psi$  is a  $\mathbb{C}^4$ -valued function, and  $A_a$  for  $0 \leq a \leq 3$  are real-valued functions.  $I$ ,  $\gamma_a$  ( $a = 0, 1, 2, 3, 5$ ) and  $(g_{ab})$  are as in the Dirac-Klein-Gordon equations. In a similar manner to (2.17) with  $M = 0$ , from (2.18) we obtain

$$\square\psi = -\frac{1}{2}\sqrt{-1}\sum_{b=0}^3\partial_b\left(\sum_{a=0}^3g_{aa}A_a\gamma_b\gamma_a(I + \gamma_5)\psi\right). \quad (2.20)$$

Putting  $u = (v, w) \in \mathbb{R}^4 \times \mathbb{R}^8$  with  $v = (A_a)_{0 \leq a \leq 3}$ , and  $w = (\operatorname{Re}\psi, \operatorname{Im}\psi)$ , we find that the conditions (a) and (b) hold for the system (2.19)–(2.20), and thus Theorem 2.1 is applicable; more precisely (2.20) implies that (b) is satisfied with  $\mathcal{I}_1 = \emptyset$  and  $\mathcal{I}_2 = \{1, \dots, 8\}$ .



**Example 3 (The Klein-Gordon-Zakharov equations)** Ozawa-Tsutaya-Tsutsumi [23] and Tsutaya [28] proved the global existence of small solutions to the Klein-Gordon-Zakharov equations:

$$\begin{cases} (\square + 1)\tilde{u} = -\tilde{n}\tilde{u}, \\ \square\tilde{n} = \Delta_x|\tilde{u}|^2 \end{cases} \quad (2.21)$$

in  $(0, \infty) \times \mathbb{R}^3$ , where  $\tilde{u} = (\tilde{u}_1, \tilde{u}_2, \tilde{u}_3)$  is a  $\mathbb{C}^3$ -valued function, and  $\tilde{n}$  is a real valued function (see also Ozawa-Tsutaya-Tsutsumi [24] for the multiple speed case).

By setting  $v_i = \operatorname{Re} \tilde{u}_i$ ,  $v_{i+3} = \operatorname{Im} \tilde{u}_i$  ( $1 \leq i \leq 3$ ),  $v_{3i+3+k} = \partial_k v_i$  ( $1 \leq i \leq 6$ ,  $1 \leq k \leq 3$ ), and  $w (= w_1) = \tilde{n}$ , we see that solving (2.21) is equivalent to solving

$$\begin{cases} (\square + 1)v_i = -wv_i, & 1 \leq i \leq 6, \\ (\square + 1)v_{3i+3+k} = -w(\partial_k v_i) - (\partial_k w)v_i, & 1 \leq i \leq 6, 1 \leq k \leq 3, \\ \square w = \sum_{j=1}^3 \partial_j \sum_{i=1}^3 2(v_i v_{3i+3+j} + v_{i+3} v_{3i+12+j}). \end{cases} \quad (2.22)$$

Note that the system (2.22) is a semilinear system of

$$u = (u_1, \dots, u_{25}) = (v_1, \dots, v_{24}, w_1) = (v, w).$$

The conditions (a) and (b) (with  $\mathcal{I}_1 = \emptyset$  and  $\mathcal{I}_2 = \{1\}$ ) are satisfied for (2.22). Hence we can apply Theorem 2.1 to show the global existence of small amplitude solutions to (2.21).

**Example 4** The last example is not from physics, as far as the author knows. This example shows that some change of unknowns may help us to apply our theorem. Consider

$$\begin{cases} (\square + 1)v = w^2, \\ \square w = v^2 \end{cases} \quad (2.23)$$

in  $(0, \infty) \times \mathbb{R}^3$ . We can treat this example in the following way, though it does not explicitly satisfy the assumption of Theorem 2.1: Set  $\tilde{v} = v - w^2$  (cf. (5.46) below). Then we get

$$\begin{cases} (\square + 1)\tilde{v} = -2Q_0(w, w) - 2w(\tilde{v} + w^2)^2, \\ \square w = (\tilde{v} + w^2)^2 \end{cases} \quad (2.24)$$

(cf. (5.52) below). This system (2.24) satisfies the assumption in Theorem 2.1 with  $u = (\tilde{v}, w) = (\tilde{v}_1, w_1)$ ,  $I_1 = \{1\}$ , and  $I_2 = \emptyset$ . Thus we get a global solution  $(\tilde{v}, w)$  to (2.24) for small data, and accordingly we obtain a global solution  $(v, w)$  to the original system (2.23).

### 3 Preliminary Results

In this section, we state the known estimates for the wave and Klein-Gordon equations. Throughout this paper, we write  $\langle z \rangle = \sqrt{1 + |z|^2}$  for  $z \in \mathbb{R}^d$ , where  $d$  is a positive integer.

We start this section with the energy inequality for hyperbolic systems, which can be shown easily by the standard method.

**Lemma 3.1** *Let  $m_i \geq 0$  for  $1 \leq i \leq N$ , and  $T > 0$ . Suppose that  $\tilde{\gamma} = (\tilde{\gamma}_{ka}^{ij})$  be a smooth function satisfying*

$$\tilde{\gamma}_{ka}^{ij}(t, x) = \tilde{\gamma}_{ka}^{ji}(t, x), \quad \tilde{\gamma}_{kl}^{ij}(t, x) = \tilde{\gamma}_{lk}^{ij}(t, x), \quad (t, x) \in (0, T) \times \mathbb{R}^3$$

for  $1 \leq i, j \leq N$ ,  $1 \leq k, l \leq 3$ , and  $0 \leq a \leq 3$ . We also assume that

$$\left| \sum_{1 \leq i, j \leq N} \sum_{1 \leq k, l \leq 3} \tilde{\gamma}_{kl}^{ij}(t, x) \xi'_{i,k} \xi'_{j,l} \right| \leq \frac{1}{2} \sum_{1 \leq i \leq N} \sum_{1 \leq k \leq 3} |\xi'_{i,k}|^2, \quad (t, x) \in (0, T) \times \mathbb{R}^3$$

for any  $(\xi'_{i,k})_{1 \leq i \leq N, 1 \leq k \leq 3} \in \mathbb{R}^{3N}$ .

Let  $\varphi = (\varphi_1, \dots, \varphi_N)$  be the solution to

$$(\square + m_i^2) \varphi_i - \sum_{1 \leq j \leq N} \sum_{\substack{1 \leq k \leq 3 \\ 0 \leq a \leq 3}} \tilde{\gamma}_{ka}^{ij} (\partial_k \partial_a \varphi_j) = \Phi_i \quad \text{in } (0, T) \times \mathbb{R}^3,$$

$$\varphi_i(0, x) = \varphi_i^{(0)}(x), \quad (\partial_t \varphi_i)(0, x) = \varphi_i^{(1)}(x), \quad x \in \mathbb{R}^3$$

for  $1 \leq i \leq N$ , where  $\varphi^{(0)} = (\varphi_1^{(0)}, \dots, \varphi_N^{(0)}) \in H^1(\mathbb{R}^3; \mathbb{R}^N)$ ,  $\varphi^{(1)} = (\varphi_1^{(1)}, \dots, \varphi_N^{(1)}) \in L^2(\mathbb{R}^3; \mathbb{R}^N)$ , and  $\Phi = (\Phi_1, \dots, \Phi_N) \in L^1((0, T); L^2(\mathbb{R}^3; \mathbb{R}^N))$ . Then, there exists a positive constant  $C$ , which is independent of  $T$ , such that

$$\begin{aligned} & \sum_{i=1}^N (\|\partial \varphi_i(t)\|_{L^2} + m_i \|\varphi_i(t)\|_{L^2}) \\ & \leq C \left( \|\varphi^{(0)}\|_{H^1} + \|\varphi^{(1)}\|_{L^2} + \int_0^t \|\partial \tilde{\gamma}(\tau)\|_{L^\infty} \|\partial \varphi(\tau)\|_{L^2} d\tau + \int_0^t \|\Phi(\tau)\|_{L^2} d\tau \right) \end{aligned}$$

for  $0 \leq t < T$ .

Before we proceed to the decay estimates of the solutions to the Klein-Gordon and wave equations, we introduce the vector fields  $\Omega_j$  and  $L_j$  for  $1 \leq j \leq 3$  by

$$\Omega = (\Omega_1, \Omega_2, \Omega_3) = x \times \nabla_x = (x_2 \partial_3 - x_3 \partial_2, x_3 \partial_1 - x_1 \partial_3, x_1 \partial_2 - x_2 \partial_1), \quad (3.1)$$

$$L = (L_1, L_2, L_3) = x \partial_t + t \nabla_x = (x_1 \partial_t + t \partial_1, x_2 \partial_t + t \partial_2, x_3 \partial_t + t \partial_3), \quad (3.2)$$

where  $\nabla_x = (\partial_1, \partial_2, \partial_3)$ , and  $\times$  is the external product in  $\mathbb{R}^3$ . Writing  $\partial = (\partial_a)_{0 \leq a \leq 3}$ , we set

$$Z = (Z_1, \dots, Z_{10}) = ((\Omega_j)_{1 \leq j \leq 3}, (L_j)_{1 \leq j \leq 3}, (\partial_a)_{0 \leq a \leq 3}) = (\Omega, L, \partial). \quad (3.3)$$

Note that we have

$$[L_j, \square + m^2] = [\Omega_j, \square + m^2] = [\partial_a, \square + m^2] = 0, \quad 1 \leq j \leq 3, \quad 0 \leq a \leq 3 \quad (3.4)$$

for  $m \geq 0$ , where  $[A, B] = AB - BA$  for operators  $A$  and  $B$ . Hence the vector fields in  $Z = (\Omega, L, \partial)$  are compatible with the Klein-Gordon equations, as well as the wave equations. Here we note that we have

$$[Z_j, \partial_a] = \sum_{b=0}^3 C_b^{ja} \partial_b, \quad [Z_j, Z_k] = \sum_{l=1}^{10} D_l^{jk} Z_l, \quad 1 \leq j, k \leq 10, \quad 0 \leq a \leq 3 \quad (3.5)$$

with appropriate constants  $C_b^{ja}$  and  $D_l^{jk}$ . For a multi-index  $\alpha = (\alpha_1, \dots, \alpha_{10})$ , we write  $Z^\alpha = Z_1^{\alpha_1} \cdots Z_{10}^{\alpha_{10}}$ . For a function  $\varphi = \varphi(t, x)$  and a nonnegative integer  $s$ , we define

$$|\varphi(t, x)|_s = \sum_{|\alpha| \leq s} |Z^\alpha \varphi(t, x)|, \quad \|\varphi(t)\|_s = \sum_{|\alpha| \leq s} \|Z^\alpha \varphi(t, \cdot)\|_{L^2(\mathbb{R}^3)}. \quad (3.6)$$

Using the vector fields in  $Z = (\Omega, L, \partial)$ , Klainerman [16] obtained the decay estimate for the solutions to the Klein-Gordon equations. This estimate has been modified and generalized by many authors (for instance, see Bachelot [2], Hörmander [8], Sideris [26], and Georgiev [5]). Here we state the estimate obtained in [5]: Let  $\chi_j$  ( $j \geq 0$ ) be nonnegative  $C_0^\infty(\mathbb{R})$ -functions satisfying

$$\sum_{j=0}^{\infty} \chi_j(\tau) = 1 \text{ for } \tau \geq 0, \quad (3.7)$$

$$\text{supp } \chi_j = [2^{j-1}, 2^{j+1}] \text{ for } j \geq 1, \text{ and } \text{supp } \chi_0 \cap [0, \infty) = [0, 2]. \quad (3.8)$$

**Lemma 3.2** *Let  $m > 0$ , and  $v$  be a smooth solution to*

$$(\square + m^2)v(t, x) = \Phi(t, x), \quad (t, x) \in (0, \infty) \times \mathbb{R}^3.$$

*Then there exists a positive constant  $C = C(m)$  such that we have*

$$\begin{aligned} \langle t + |x| \rangle^{3/2} |v(t, x)| \leq & C \sum_{j=0}^{\infty} \sum_{|\alpha| \leq 4} \sup_{\tau \in [0, t]} \chi_j(\tau) \|\langle \tau + |\cdot| \rangle Z^\alpha \Phi(\tau, \cdot)\|_{L^2(\mathbb{R}^3)} \\ & + C \sum_{j=0}^{\infty} \sum_{|\alpha| \leq 5} \left\| \langle \cdot \rangle^{3/2} \chi_j(|\cdot|) Z^\alpha v(0, \cdot) \right\|_{L^2(\mathbb{R}^3)} \end{aligned} \quad (3.9)$$

*for  $(t, x) \in (0, \infty) \times \mathbb{R}^3$ , provided that the right-hand side of (3.9) is finite.*

For the proof, see Georgiev [5, Theorem 1].

Now we turn our attention to the wave equations. In [17], a weighted  $L^1$ - $L^\infty$  estimate for the wave equation is derived (see also Hörmander [7]), where the scaling operator  $S = t\partial_t + x \cdot \nabla_x$  as well as  $Z = (\Omega, L, \partial)$  is used. Since we have  $[S, \square] = -2\square$ , the scaling operator  $S$  is applicable to the wave equations, but it is incompatible with the Klein-Gordon equations. Therefore Georgiev ([4]) developed a weighted  $L^2$ - $L^\infty$  estimate involving only  $Z$ . There is also a large literature on the study of systems of nonlinear wave equations with multiple speeds of the form

$$\square_{c_i} u_i = F_i(u, \partial u, \partial_x \partial u), \quad 1 \leq i \leq N,$$

where  $c_i > 0$  ( $1 \leq i \leq N$ ) and  $\square_c = \partial_t^2 - c^2 \Delta_x$  (see, for example, [13] and the references cited therein). In the study of this kind of system, the vector field method using only  $(S, \Omega, \partial)$  has been developed, because  $L = (L_j)_{1 \leq j \leq 3}$  is incompatible with such system (observe that  $[L_j, \square_c] = 2(c^2 - 1)\partial_t \partial_j$  has no good property when  $c \neq 1$ ). Especially, in Yokoyama [30] and Kubota-Yokoyama [22] (see also the author [12]), weighted  $L^\infty$ - $L^\infty$  estimates requiring only  $(\Omega, \partial)$  are adopted to prove some global existence results under the null condition (the origin of these estimates can be found in John [10] and Kovalyov [20]; see also Kovalyov-Tsutaya [21]). We will employ these  $L^\infty$ - $L^\infty$  estimates in the proof of Theorem 2.1 because they require only  $(\Omega, \partial)$  and are easily applicable to the coupled system of the wave and Klein-Gordon equations. Here we note that  $S$  is still used in the arguments in [30], [22] and [12] to treat the null forms (see (4.11) below).

To state the weighted  $L^\infty$ - $L^\infty$  estimates, we define

$$\mathcal{W}_\rho(t, r) := \begin{cases} \langle t+r \rangle^\rho & \text{if } \rho < 0, \\ \{\log(2 + \langle t+r \rangle \langle t-r \rangle^{-1})\}^{-1} & \text{if } \rho = 0, \\ \langle t-r \rangle^\rho & \text{if } \rho > 0. \end{cases} \quad (3.10)$$

We also introduce

$$W_-(t, r) := \min \{ \langle r \rangle, \langle t-r \rangle \}. \quad (3.11)$$

For the homogeneous wave equations we have the following estimate which was essentially proved in Asakura [1, Proposition 1.1] (see also [15, Lemma 3.1] for the expression below):

**Lemma 3.3** *Let  $w$  be a smooth solution to*

$$\square w(t, x) = 0, \quad (t, x) \in (0, \infty) \times \mathbb{R}^3$$

*with initial data  $w = w^{(0)}$ ,  $\partial_t w = w^{(1)}$  at  $t = 0$ .*

*Let  $\kappa > 0$ . Then, there exists a positive constant  $C = C(\kappa)$  such that*

$$\begin{aligned} & \langle t + |x| \rangle \mathcal{W}_{\kappa-1}(t, |x|) |w(t, x)| \\ & \leq C \sup_{|y-x| \leq t} \langle y \rangle^\kappa \left( \langle y \rangle \sum_{|\alpha| \leq 1} |(\partial_x^\alpha w^{(0)})(y)| + |y| |w^{(1)}(y)| \right) \end{aligned} \quad (3.12)$$

for  $(t, x) \in (0, \infty) \times \mathbb{R}^3$ . Here  $\partial_x = (\partial_1, \partial_2, \partial_3)$ , and we have used the standard notation of multi-indices.

The following weighted  $L^\infty$ - $L^\infty$  estimates are the special cases of the estimates obtained in Kubota-Yokoyama [22, Lemma 3.2] (see also Katayama-Kubo [14, Lemma 3.4] for the expression below):

**Lemma 3.4** *Let  $w$  be a smooth solution to*

$$\square w(t, x) = \Psi(t, x), \quad (t, x) \in (0, \infty) \times \mathbb{R}^3$$

with initial data  $w = \partial_t w = 0$  at  $t = 0$ .

Suppose that  $\rho \geq 0$ ,  $\kappa \geq 1$ , and  $\mu > 0$ . Then there exists a positive constant  $C = C(\rho, \kappa, \mu)$  such that

$$\begin{aligned} & \langle t + |x| \rangle^{1-\rho} \mathcal{W}_{\kappa-1}(t, |x|) |w(t, x)| \\ & \leq C \sup_{\tau \in [0, t]} \sup_{|y-x| \leq t-\tau} |y| \langle \tau + |y| \rangle^{\kappa-\rho+\mu} W_-(\tau, |y|)^{1-\mu} |\Psi(\tau, y)|, \end{aligned} \quad (3.13)$$

$$\begin{aligned} & \langle t + |x| \rangle^{-\rho} \langle x \rangle \langle t - |x| \rangle^\kappa |\partial w(t, x)| \\ & \leq C \sup_{\tau \in [0, t]} \sup_{|y-x| \leq t-\tau} |y| \langle \tau + |y| \rangle^{\kappa-\rho+\mu} W_-(\tau, |y|)^{1-\mu} \sum_{|\alpha|+|\beta| \leq 1} |\partial^\alpha \Omega^\beta \Psi(\tau, y)| \end{aligned} \quad (3.14)$$

for  $(t, x) \in (0, \infty) \times \mathbb{R}^3$ . Here  $\partial = (\partial_0, \partial_1, \partial_2, \partial_3)$ , and  $\Omega$  is given by (3.1).

The following Sobolev type inequality will be used to combine decay estimates with the energy estimates (see Klainerman [18] for the proof):

**Lemma 3.5** *For a smooth function  $\varphi$  on  $\mathbb{R}^3$ , we have*

$$\sup_{x \in \mathbb{R}^3} \langle x \rangle |\varphi(x)| \leq C \sum_{|\alpha|+|\beta| \leq 2} \|\partial_x^\alpha \Omega^\beta \varphi\|_{L^2(\mathbb{R}^3)}, \quad (3.15)$$

provided that the right-hand side of (3.15) is finite. Here  $C$  is a universal positive constant.

We conclude this section with some observation on the wave equations of the following type:

$$\begin{cases} \square \psi(t, x) = \sum_{a=0}^3 \partial_a \Psi_a(t, x), & (t, x) \in (0, \infty) \times \mathbb{R}^3, \\ \psi(0, x) = \psi^{(0)}(x), \quad (\partial_t \psi)(0, x) = \psi^{(1)}(x), & x \in \mathbb{R}^3. \end{cases} \quad (3.16)$$

For  $0 \leq a \leq 3$ , let  $\psi_a = \psi_a(t, x)$  be the solution to  $\square \psi_a = \Psi_a$  with initial data  $\psi_a = \partial_t \psi_a = 0$  at  $t = 0$ , and let  $\psi_f(t, x)$  be the solution to  $\square \psi_f = 0$  with initial data  $\psi_f = \psi^{(0)}$  and  $(\partial_t \psi_f) = \psi^{(1)} - \Psi_0(0, \cdot)$  at  $t = 0$ . It is easy to verify that the solution  $\psi$  to (3.16) can be written as  $\psi = \sum_{a=0}^3 \partial_a \psi_a + \psi_f$ . Therefore, we can essentially regard  $\psi$  as derivatives of solutions to some wave equations, and  $\psi$  enjoys better estimates than we can expect in general.

**Lemma 3.6** *Let  $\psi$  be the solution to (3.16). Then we have*

$$\begin{aligned} \|\psi(t, \cdot)\|_{L^2(\mathbb{R}^3)} &\leq C \left( \|\psi^{(0)}\|_{L^2(\mathbb{R}^3)} + \|\psi^{(1)}\|_{L^{6/5}(\mathbb{R}^3)} + \|\Psi_0(0, \cdot)\|_{L^{6/5}(\mathbb{R}^3)} \right) \\ &\quad + C \sum_{a=0}^3 \int_0^t \|\Psi_a(\tau, \cdot)\|_{L^2(\mathbb{R}^3)} d\tau, \end{aligned} \quad (3.17)$$

*provided that the right-hand side of (3.17) is finite.*

*Proof.* We have  $\|\psi\|_{L^2} \leq \sum_{a=0}^3 \|\partial \psi_a\|_{L^2} + \|\psi_f\|_{L^2}$ . Hence (3.17) follows from the energy inequality (cf. Lemma 3.1) for  $\psi_a$  ( $0 \leq a \leq 3$ ), and the  $L^2$ -estimate for  $\psi_f$  (see Strauss [27] for example).

**Lemma 3.7** *Let  $\psi$  be the smooth solution to (3.16). Suppose that  $\rho \geq 0$ ,  $\kappa \geq 1$  and  $\mu > 0$ . Then we have*

$$\begin{aligned} &\langle t + |x| \rangle^{-\rho} \langle x \rangle \langle t - |x| \rangle^\kappa |\psi(t, x)| \\ &\leq C \sup_{\tau \in [0, t]} \sup_{|y-x| \leq t-\tau} |y| \langle \tau + |y| \rangle^{\kappa-\rho+\mu} W_-(\tau, |y|)^{1-\mu} \sum_{\substack{|\alpha|+|\beta| \leq 1 \\ 0 \leq a \leq 3}} |\partial^\alpha \Omega^\beta \Psi_a(\tau, y)| \\ &\quad + C \sup_{|y-x| \leq t} \langle y \rangle^{\kappa+1-\rho} \left( \langle y \rangle \sum_{|\alpha| \leq 1} |(\partial_x^\alpha \psi^{(0)})(y)| + |y| |\psi^{(1)}(y)| \right). \end{aligned} \quad (3.18)$$

*Proof.* The estimates for  $\partial_a \psi_a$  follow from (3.14); using Lemma 3.3 to estimate  $\psi_f$ , and noting that we have

$$\langle y \rangle^{\kappa+1-\rho} |y| |\Psi_0(0, y)| = \left\{ |y| \langle \tau + |y| \rangle^{\kappa-\rho+\mu} W_-(\tau, |y|)^{1-\mu} |\Psi_0(\tau, y)| \right\} \Big|_{\tau=0},$$

we obtain the desired result immediately (more precisely, (3.18) for zero initial data is directly proved in [22], and (3.14) is obtained as its corollary in fact).

## 4 Estimates for the Null Forms

In this section, we will derive some estimates for the null forms. We set  $r = |x|$ ,  $\omega = (\omega_1, \omega_2, \omega_3)$  with  $\omega_j = x_j/r$  for  $1 \leq j \leq 3$ , and  $\partial_r = \omega \cdot \nabla_x = \sum_{j=1}^3 \omega_j \partial_j$ . Then we have

$$\nabla_x = \omega \partial_r - r^{-1}(\omega \times \Omega), \quad (4.1)$$

where  $\Omega$  is defined by (3.1). Since (3.1) and (3.2) yield  $tr^{-1}\Omega = \omega \times L$ , the expression (4.1) implies

$$(t+r)(\nabla_x - \omega \partial_r) = -\omega \times (\Omega + \omega \times L). \quad (4.2)$$

From (4.2), we obtain

$$\begin{aligned} &|Q_0(\varphi, \psi) - Q_0^{\text{rad}}(\varphi, \psi)| + \sum_{k=1}^3 |Q_{0k}(\varphi, \psi) - \omega_k Q_{0r}^{\text{rad}}(\varphi, \psi)| \\ &\quad + \sum_{1 \leq j < k \leq 3} |Q_{jk}(\varphi, \psi)| \leq C \langle t+r \rangle^{-1} (|Z\varphi| |\partial\psi| + |\partial\varphi| |Z\psi|) \end{aligned} \quad (4.3)$$

at  $(t, x) \in [0, \infty) \times \mathbb{R}^3$ , where

$$Q_0^{\text{rad}}(\varphi, \psi) := (\partial_t \varphi)(\partial_t \psi) - (\partial_r \varphi)(\partial_r \psi), \quad Q_{0r}^{\text{rad}}(\varphi, \psi) := (\partial_t \varphi)(\partial_r \psi) - (\partial_r \varphi)(\partial_t \psi),$$

$Z$  is given by (3.3), and  $C$  is a positive constant. Putting  $L_r := \omega \cdot L = r\partial_t + t\partial_r$ , we get

$$(t+r)Q_{0r}^{\text{rad}}(\varphi, \psi) = (\partial_t \varphi - \partial_r \varphi)(L_r \psi) - (L_r \varphi)(\partial_t \psi - \partial_r \psi). \quad (4.4)$$

From (4.3) and (4.4), we obtain

$$|Q_{ab}(\varphi, \psi)| \leq C \langle t+r \rangle^{-1} (|Z\varphi| |\partial\psi| + |\partial\varphi| |Z\psi|) \quad (4.5)$$

for  $0 \leq a < b \leq 3$  at  $(t, x) \in [0, \infty) \times \mathbb{R}^3$ , where  $C$  is some positive constant. Thus we only need the vector fields in  $Z = (\Omega, L, \partial)$  to obtain the extra decay factor  $\langle t+r \rangle^{-1}$  for terms satisfying the strong null condition.

To treat the null form  $Q_0$ , we introduce  $\partial_{\pm} := \partial_t \pm \partial_r$ . Then we get

$$Q_0^{\text{rad}}(\varphi, \psi) = \frac{1}{2} ((\partial_+ \varphi)(\partial_- \psi) + (\partial_- \varphi)(\partial_+ \psi)). \quad (4.6)$$

As we will see below, estimates of  $\partial_+ \varphi$  and  $\partial_+ \psi$  are important in deriving enhanced decay for  $Q_0$ . Note that we also have

$$Q_{0r}^{\text{rad}}(\varphi, \psi) = (\partial_+ \varphi)(\partial_r \psi) - (\partial_r \varphi)(\partial_+ \psi). \quad (4.7)$$

Rewriting  $\partial_+$  as

$$\partial_+ = (t+r)^{-1}(S + L_r) \quad (4.8)$$

with  $S = t\partial_t + x \cdot \nabla_x = t\partial_t + r\partial_r$ , from (4.3) and (4.6) we obtain

$$|Q_0(\varphi, \psi)| \leq C \langle t+r \rangle^{-1} (|\Gamma\varphi| |\partial\psi| + |\partial\varphi| |\Gamma\psi|), \quad (4.9)$$

where  $\Gamma := (S, Z) = (S, L, \Omega, \partial)$ . The estimate (4.9) was used in Klainerman [17], and the usage of  $S$  in (4.9) makes it difficult to treat the null form  $Q_0$  included in coupled systems of the wave and Klein-Gordon equations, and this is the reason why the notion of the strong null condition was introduced in [4].

Before we proceed to our new estimate for  $Q_0$ , we introduce another kind of known estimate for the null forms here. If we only use (4.1), then we find that the left-hand side of (4.3) is bounded from above by  $C \langle r \rangle^{-1} (|Z'\varphi| |\partial\psi| + |\partial\varphi| |Z'\psi|)$ , where  $Z' := (\Omega, \partial)$ . Hence, rewriting  $\partial_+$  as

$$\partial_+ = r^{-1}(S - (t-r)\partial_t), \quad (4.10)$$

we get

$$|Q_0(\varphi, \psi)| \leq C \langle r \rangle^{-1} (|\Gamma'\varphi| |\partial\psi| + |\partial\varphi| |\Gamma'\psi| + \langle t-r \rangle |\partial\varphi| |\partial\psi|), \quad (4.11)$$

where  $\Gamma' := (S, \Omega, \partial)$ . Similar estimate can be obtained for  $Q_{ab}$  in view of (4.7). These estimates are used in the study of systems of wave equations with multiple

speeds because  $L$  is incompatible with such systems (see Hoshiga-Kubo [9] and Yokoyama [30] for example). As we have mentioned in the previous section, the estimate (4.11) is the point where  $S$  comes in the arguments of [30], [22] and [12], though the weighted  $L^\infty$ - $L^\infty$  estimates (cf. Lemma 3.4) are free of  $S$ .

In Katayama-Kubo [13], for the  $\partial_+$ -derivative of the solution to the wave equation, a weighted  $L^\infty$ - $L^\infty$  estimate with a better decay factor than (3.14) is directly obtained through an explicit expression of the solution (without rewriting  $\partial_+$  by the other vector fields), and the null forms are treated using only  $Z' = (\Omega, \partial)$  (see also [14]). We can adopt this approach in [13] to systems of the wave and Klein-Gordon equations because the required vector fields  $\Omega$  and  $\partial$  are admissible. However we take another approach here since we can use the vector field  $L$ ; motivated by (4.10), we rewrite  $\partial_+$  as

$$\partial_+ = (t+r)^{-1} (2L_r + (t-r)\partial_t - (t-r)\partial_r). \quad (4.12)$$

Then, by (4.3), (4.6) and (4.12), we obtain

$$|Q_0(\varphi, \psi)| \leq C \langle t+r \rangle^{-1} (|Z\varphi| |\partial\psi| + |\partial\varphi| |Z\psi| + \langle t-r \rangle |\partial\varphi| |\partial\psi|) \quad (4.13)$$

at  $(t, x) \in [0, \infty) \times \mathbb{R}^3$ .

For any multi-index  $\alpha$ , we can easily check that  $Z^\alpha Q_0(\varphi, \psi)$  can be written as a linear combination of the null forms  $Q_0(Z^\beta \varphi, Z^\gamma \psi)$  and  $Q_{cd}(Z^\beta \varphi, Z^\gamma \psi)$  with  $|\beta| + |\gamma| \leq |\alpha|$  and  $0 \leq c < d \leq 3$ . The same is true for  $Z^\alpha Q_{ab}(\varphi, \psi)$  ( $0 \leq a < b \leq 3$ ). Therefore, (4.5) and (4.13) yield the following:

**Lemma 4.1** *Let  $k$  be a nonnegative integer, and let  $Q$  be one of the null forms  $Q_0$  and  $Q_{ab}$  with  $0 \leq a < b \leq 3$ . Then we have*

$$\begin{aligned} \langle t+|x| \rangle |Q(\varphi, \psi)|_k &\leq C (|\varphi|_{[k/2]+1} |\partial\psi|_k + |\varphi|_{k+1} |\partial\psi|_{[k/2]}) \\ &\quad + C (|\partial\varphi|_{[k/2]} |\psi|_{k+1} + |\partial\varphi|_k |\psi|_{[k/2]+1}) \\ &\quad + C \langle t-|x| \rangle (|\partial\varphi|_{[k/2]} |\partial\psi|_k + |\partial\varphi|_k |\partial\psi|_{[k/2]}) \end{aligned}$$

at  $(t, x) \in [0, \infty) \times \mathbb{R}^3$  for any smooth functions  $\varphi$  and  $\psi$ . Here  $C$  is a positive constant depending only on  $k$ ,  $|\cdot|_s$  is given by (3.6) for a nonnegative integer  $s$ , and  $[m]$  denotes the largest integer not exceeding the number  $m$ .

## 5 Proof of Theorem 2.1

In this section, we will prove Theorem 2.1.

Suppose that all the assumptions in Theorem 2.1 are fulfilled. The classical theory for nonlinear hyperbolic equations implies the local existence of the classical solutions to (1.1)–(1.2) for small  $\varepsilon$ . Moreover, we see that the solution  $u$  exists as long as  $\sum_{|\alpha| \leq 2} \|\partial^\alpha u(t, \cdot)\|_{L^\infty(\mathbb{R}^3)}$  stays finite (see Hörmander [8] for instance). Hence what we need for the proof of Theorem 2.1 is such an *a priori* estimate to guarantee the boundedness of  $\sum_{|\alpha| \leq 2} \|\partial^\alpha u(t, \cdot)\|_{L^\infty(\mathbb{R}^3)}$ . Let



$u = (u_i)_{1 \leq i \leq N} = (v, w)$  be the local solution to (1.1)–(1.2) for  $0 \leq t < T_0$  with some  $T_0 > 0$ , where  $v$  and  $w$  are given by (2.2). If both  $\mathcal{I}_1$  and  $\mathcal{I}_2$  in the condition (b) are non-empty, without loss of generality we may assume that  $\mathcal{I}_1 = \{1, \dots, N_3\}$ , and  $\mathcal{I}_2 = \{N_3 + 1, \dots, N_2\}$  with some positive integer  $N_3$ . Correspondingly, we write

$$w = (w_k)_{1 \leq k \leq N_2} = ((w_k^{(i)})_{1 \leq k \leq N_3}, (w_k^{(ii)})_{1 \leq k \leq N_4}) = (w^{(i)}, w^{(ii)}),$$

where  $N_4 = N_2 - N_3$ . If  $\mathcal{I}_2$  (resp.  $\mathcal{I}_1$ ) is empty, then we put  $w^{(i)} = w$  (resp.  $w^{(ii)} = w$ ), and  $w^{(ii)}$  (resp.  $w^{(i)}$ ) should be neglected in what follows.

For a nonnegative integer  $\sigma$ , and a positive constant  $p$ , we define

$$\begin{aligned} d_{\sigma,p}(t, x) &= \langle t + |x| \rangle^{3/2} |v(t, x)|_{\sigma+2} + \langle x \rangle \langle t - |x| \rangle |\partial w(t, x)|_{\sigma+1} \\ &\quad + \langle t + |x| \rangle (\mathcal{W}_0(t, |x|) |w^{(i)}(t, x)|_{\sigma+2} + W_-(t, |x|)^{1-p} |w^{(ii)}(t, x)|_{\sigma+2}) \end{aligned}$$

for  $(t, x) \in [0, T_0] \times \mathbb{R}^3$ , where  $|\cdot|_s$ ,  $\mathcal{W}_0$ , and  $W_-$  are given by (3.6), (3.10), and (3.11), respectively.

For a smooth function  $\varphi = \varphi(x)$  and a nonnegative integer  $s$ , we set

$$\|\varphi\|_{X^s} = \sum_{|\alpha| \leq s} \left( \left( \int_{\mathbb{R}^3} |\langle x \rangle^{s+2} \partial_x^\alpha \varphi(x)|^2 dx \right)^{1/2} + \left( \int_{\mathbb{R}^3} |\langle x \rangle^s \partial_x^\alpha \varphi(x)|^{6/5} dx \right)^{5/6} \right).$$

Note that the Sobolev embedding theorem implies that

$$\sup_{x \in \mathbb{R}^3} \sum_{|\alpha| \leq s-2} \langle x \rangle^{|\alpha|+4} |\partial_x^\alpha \varphi(x)| \leq \sup_{x \in \mathbb{R}^3} \sum_{|\alpha| \leq s-2} \langle x \rangle^{s+2} |\partial_x^\alpha \varphi(x)| \leq C_s \|\varphi\|_{X^s}$$

for  $s \geq 2$ , where  $C_s$  is a positive constant depending only on  $s$ .

Our aim here is to show the following:

**Proposition 5.1** *Fix some  $\sigma \geq 19$ , and  $0 < p < 1/100$ , say. Suppose that all the assumptions in Theorem 2.1 are fulfilled. Assume that  $\|f\|_{X^{2\sigma+1}} + \|g\|_{X^{2\sigma}} \leq M_0$  with some positive constant  $M_0$ . Let  $u = (v, w)$  be the local solution to (1.1)–(1.2) for  $0 \leq t < T_0$ . Then there exists a positive constant  $A_0 = A_0(M_0)$  having the following property: For any  $A \geq A_0$ , there exists a positive constant  $\varepsilon_0 = \varepsilon_0(A)$  such that*

$$\sup_{0 \leq t < T} \|d_{\sigma,p}(t, \cdot)\|_{L^\infty(\mathbb{R}^3)} \leq A\varepsilon \tag{5.1}$$

implies

$$\sup_{0 \leq t < T} \|d_{\sigma,p}(t, \cdot)\|_{L^\infty(\mathbb{R}^3)} \leq \frac{A}{2}\varepsilon, \tag{5.2}$$

provided that  $0 < \varepsilon \leq \varepsilon_0$  and  $0 < T \leq T_0$ . Here  $A_0$  and  $\varepsilon_0$  are independent of  $T_0$ .

Once Proposition 5.1 is established, by the continuity argument (or the bootstrap argument), we find that  $\|d_{\sigma,p}(t, \cdot)\|_{L^\infty(\mathbb{R}^3)}$  stays bounded as long as the solution exists, provided that  $\varepsilon$  is small enough. Indeed, suppose that  $f$  and  $g$  belong

to  $C_0^\infty(\mathbb{R}^3; \mathbb{R}^N)$  at first. Then, taking the support of  $u$  into account, we see that  $\|d_{\sigma,p}(t, \cdot)\|_{L^\infty(\mathbb{R}^3)}$  is continuous in  $t$ . We choose a large constant  $A(\geq A_0)$  to satisfy  $\|d_{\sigma,p}(0, \cdot)\|_{L^\infty(\mathbb{R}^3)} < A\varepsilon$ . Then we see that (5.1) is true for some small  $T$ . Let  $T_*( > 0)$  be the supremum of  $T (\in (0, T_0))$  for which (5.1) holds. If  $\varepsilon \in (0, \varepsilon_0]$ , then by (5.2) and the continuity of  $\|d_{\sigma,p}(t, \cdot)\|_{L^\infty}$ , we conclude that  $T_* = T_0$  (otherwise we meet a contradiction). In other words, if  $u$  is the local solution for  $0 \leq t < T_0$ , then we have  $\sup_{0 \leq t < T_0} \|d_{\sigma,p}(t, \cdot)\|_{L^\infty} \leq A\varepsilon$ , provided that  $\varepsilon \in (0, \varepsilon_0]$ . We see that the same is true for general  $f, g \in \mathcal{S}(\mathbb{R}^3; \mathbb{R}^N)$  through the approximation by  $C_0^\infty$ -functions. This *a priori* estimate implies Theorem 2.1 immediately.

Now we are going to prove Proposition 5.1. We assume that (5.1) holds. In the following, various positive constants, being independent of  $A(> 0)$ ,  $\varepsilon(\leq 1)$ ,  $T(> 0)$ , and  $M_0$ , are indicated just by the same letter  $C$ . Thus the practical value of  $C$  may change line by line. Similarly  $C_*$  stands for various positive constants depending only on  $M_0$  and the bounds for finite numbers of derivatives of  $F$  in a small neighborhood of the origin  $(\xi, \xi', \xi'') = (0, 0, 0)$ . We always assume that  $\varepsilon$  is small enough to satisfy  $A\varepsilon \leq 1$ , say.

First we remark that for any nonnegative integer  $s$ , there exists a positive constant  $C_s$  such that

$$C_s^{-1} |\partial\varphi(t, x)|_s \leq \sum_{|\alpha| \leq s} |\partial Z^\alpha \varphi(t, x)| \leq C_s |\partial\varphi(t, x)|_s \quad (5.3)$$

holds for any smooth function  $\varphi$ , because of (3.5). We also note that we have

$$\langle x \rangle^{-1} \langle t - |x| \rangle^{-1} \leq C \langle t + |x| \rangle^{-1} W_-(t, |x|)^{-1}, \quad (t, x) \in [0, \infty) \times \mathbb{R}^3. \quad (5.4)$$

We fix some small and positive constant  $\delta$ . Then we have

$$\mathcal{W}_0(t, |x|)^{-1} \leq C \langle t + |x| \rangle^\delta \langle t - |x| \rangle^{-\delta}, \quad (t, x) \in [0, \infty) \times \mathbb{R}^3. \quad (5.5)$$

We will use (5.4) and (5.5) repeatedly in the following. Note that (5.1), (5.4), and (5.5) yield

$$|u(t, x)|_{\sigma+2} \leq CA\varepsilon \langle t + |x| \rangle^{-1+\delta} \langle t - |x| \rangle^{-\delta}, \quad (t, x) \in [0, T) \times \mathbb{R}^3. \quad (5.6)$$

The proof of Proposition 5.1 is divided into several steps.

*Step 1: Energy Estimate.* Let  $0 < \lambda < p/4$ . In this step, we are going to prove that

$$\sup_{0 \leq t < T} (1+t)^{-\lambda} (\|v(t)\|_{2\sigma} + \|w^{(ii)}(t)\|_{2\sigma} + \|\partial u(t)\|_{2\sigma}) \leq C_*\varepsilon \quad (5.7)$$

holds for small  $\varepsilon$ , where  $\|\cdot\|_s$  is given by (3.6). The difficulty here is the lack of a natural estimate for  $\|w^{(i)}(t)\|_{2\sigma}$  (cf. Lemma 3.1). To overcome this difficulty, we will use the following lemma that is easily obtained from the definition of  $Z$  and (5.3):

**Lemma 5.1** *For any  $s \geq 1$ , there exists a positive constant  $C = C(s)$  such that we have*

$$|\varphi(t, x)|_s \leq C (|\varphi(t, x)| + \langle t + |x| \rangle |\partial\varphi(t, x)|_{s-1}) \quad (5.8)$$

for any smooth function  $\varphi = \varphi(t, x)$ .

In fact, for  $s \geq 1$ , we have

$$\begin{aligned} |\varphi(t, x)|_s &\leq C \left( |\varphi(t, x)| + \sum_{|\alpha|=1} \sum_{|\beta| \leq s-1} |Z^\alpha Z^\beta \varphi(t, x)| \right) \\ &\leq C \left( |\varphi(t, x)| + \langle t + |x| \rangle \sum_{|\beta| \leq s-1} |\partial(Z^\beta \varphi)(t, x)| \right), \end{aligned}$$

which leads to (5.8), thanks to (5.3).

Now we start the proof of (5.7). Let  $|\alpha| = s \leq 2\sigma$ . We set

$$F_{i,\alpha} = Z^\alpha \{F_i(u, \partial u, \partial_x \partial u)\} - \sum_{j,k,a} \gamma_{ka}^{ij}(u, \partial u) \partial_k \partial_a (Z^\alpha u_j), \quad (5.9)$$

where  $\gamma = (\gamma_{ka}^{ij})$  is from (1.3). Then we have

$$(\square + m_i^2)(Z^\alpha u_i) - \sum_{j,k,a} \gamma_{ka}^{ij}(u, \partial u) \partial_k \partial_a (Z^\alpha u_j) = F_{i,\alpha}, \quad 1 \leq i \leq N. \quad (5.10)$$

Note that we have  $||Z^\alpha, \partial_k \partial_a] u_j| \leq C|\partial u|_s$  by (3.5). Hence, in view of (1.3), (3.5), and (5.9), we obtain from the condition (b-i), (5.1), (5.4), and (5.6) that

$$\begin{aligned} |F_{i,\alpha}| &\leq C (|v|_{[s/2]} + |w^{(ii)}|_{[s/2]} + |\partial u|_{[s/2]+1}) (|v|_s + |w^{(ii)}|_s + |\partial u|_s) \\ &\quad + C|u|_{[s/2]+2}^2 (|u|_s + |\partial u|_s) \\ &\leq CA\varepsilon \langle t + |x| \rangle^{-1} (|v|_s + |w^{(ii)}|_s + |\partial u|_s) \\ &\quad + CA^2\varepsilon^2 \langle t + |x| \rangle^{-2+2\delta} \langle t - |x| \rangle^{-2\delta} \\ &\quad \times \left( A\varepsilon \langle t + |x| \rangle^{-1+\delta} \langle t - |x| \rangle^{-\delta} + \langle t + x \rangle |\partial u|_{s-1} + |\partial u|_s \right) \quad (5.11) \end{aligned}$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$ . Here we have also used (5.8) to estimate  $|u|_s$  for  $s \geq 1$ . Thus the term  $\langle t + |x| \rangle |\partial u|_{s-1}$  on the right-hand side of (5.11) should be neglected when  $s = 0$ . Since we have

$$\left\| \langle t + |\cdot| \rangle^{-3+3\delta} \langle t - |\cdot| \rangle^{-3\delta} \right\|_{L^2(\mathbb{R}^3)} \leq C(1+t)^{-3/2}$$

for  $\delta < 1/6$ , (5.11) yields

$$\begin{aligned} \|F_{i,\alpha}\|_{L^2(\mathbb{R}^3)} &\leq CA\varepsilon(1+t)^{-1} (\|v\|_s + \|w^{(ii)}\|_s + \|\partial u\|_s) \\ &\quad + CA^2\varepsilon^2(1+t)^{-1+2\delta} \|\partial u\|_{s-1} + CA^3\varepsilon^3(1+t)^{-3/2}, \quad (5.12) \end{aligned}$$

where the term  $CA^2\varepsilon^2(1+t)^{-1+2\delta} \|\partial u\|_{s-1}$  should be neglected when  $s = 0$ . In view of the condition (b-i), we also obtain from (5.1), (5.4), and (5.6) that

$$|\gamma|_1 \leq C(|v|_1 + |w^{(ii)}|_1 + |\partial u|_1 + |u|_1^2 + |\partial u|_1^2) \leq CA\varepsilon \langle t + |x| \rangle^{-1} \quad (5.13)$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$ . Because of (1.4) and (5.13), we can apply Lemma 3.1 to (5.10) for small  $\varepsilon$ .

For  $N_3 + 1 \leq k \leq N_2$  and  $0 \leq a \leq 3$ , let  $\mathcal{G}_{k,a}$  be from the condition (b-ii). Because of (3.5), we get

$$\square \left( Z^\alpha w_j^{(ii)} \right) = \sum_{a,b=0}^3 \sum_{|\beta| \leq |\alpha|} C_{ab}^{\alpha\beta} \partial_b \left( Z^\beta \mathcal{G}_{N_3+j,a}(u, \partial u) \right), \quad 1 \leq j \leq N_4 \quad (5.14)$$

with appropriate constants  $C_{ab}^{\alpha\beta}$ . Remember that each  $\mathcal{G}_{k,a}^{(q)}$  is independent of  $w^{(i)}$  itself. Thus, going similar lines to (5.11) and (5.12), we get

$$\begin{aligned} \|\mathcal{G}_{k,a}(u, \partial u)\|_s &\leq CA\varepsilon(1+t)^{-1}(\|v\|_s + \|w^{(ii)}\|_s + \|\partial u\|_s) \\ &\quad + CA^2\varepsilon^2(1+t)^{-1+2\delta}\|\partial u\|_{s-1} + CA^3\varepsilon^3(1+t)^{-3/2} \end{aligned} \quad (5.15)$$

for  $s \leq 2\sigma$ . As before, the term including  $\|\partial u\|_{s-1}$  on the right-hand side should be neglected when  $s = 0$ .

We put

$$E_s(t) = \|v(t)\|_s + \|w^{(ii)}(t)\|_s + \|\partial u(t)\|_s$$

for  $s \geq 0$ . Applying Lemma 3.1 to (5.10) with  $|\alpha| = s = 0$ , and applying Lemma 3.6 to (5.14) with  $|\alpha| = s = 0$ , we obtain from (5.12), (5.13), and (5.15) that

$$E_0(t) \leq C_*\varepsilon + CA^3\varepsilon^3 + CA\varepsilon \int_0^t (1+\tau)^{-1} E_0(\tau) d\tau.$$

The Gronwall lemma yields

$$E_0(t) \leq (C_*\varepsilon + CA^3\varepsilon^3) (1+t)^{CA\varepsilon} \leq C_*\varepsilon(1+t)^{CA\varepsilon}, \quad (5.16)$$

provided that  $\varepsilon$  is small enough to satisfy  $A^3\varepsilon^2 \leq 1$ . Starting with (5.16), we can inductively obtain

$$E_s(t) \leq C_{*,s}\varepsilon(1+t)^{2s\delta+CA\varepsilon} \quad (5.17)$$

for  $0 \leq s \leq 2\sigma$ , where  $C_{*,s}$ 's are positive constants depending on  $s$ ,  $M_0$  and the nonlinearity  $F$ . In fact, if (5.17) with  $s$  replaced by  $s-1$  is true for some  $s \geq 1$ , then applying Lemmas 3.1 and 3.6 to (5.10) and (5.14) with  $|\alpha| = s$ , respectively, and using (5.12), (5.13) and (5.15), we obtain

$$\begin{aligned} E_s(t) &\leq C_*\varepsilon + C \left( A^3\varepsilon^3 + (2s\delta + CA\varepsilon)^{-1} C_{*,s-1} A^2\varepsilon^3 (1+t)^{2s\delta+CA\varepsilon} \right) \\ &\quad + CA\varepsilon \int_0^t (1+\tau)^{-1} E_s(\tau) d\tau, \end{aligned}$$

and the Gronwall lemma leads to (5.17).

Finally, we obtain (5.7) from (5.17) with  $s = 2\sigma$ , provided that  $\delta$  in (5.5) is chosen to satisfy  $4\sigma\delta \leq \lambda/2$ , and  $\varepsilon$  is small enough to satisfy  $CA\varepsilon \leq \lambda/2$ .

*Step 2: Decay Estimates, Part 1.* By Lemma 3.5 and (5.7), we get

$$\langle x \rangle \left( |v(t, x)|_{2\sigma-2} + |w^{(ii)}(t, x)|_{2\sigma-2} + |\partial u(t, x)|_{2\sigma-2} \right) \leq C_*\varepsilon(1+t)^\lambda \quad (5.18)$$

for  $(t, x) \in [0, T) \times \mathbb{R}^3$ .

Similarly to (5.11), we get

$$\begin{aligned}
|F_i|_s &\leq C(|v|_\sigma + |w^{(ii)}|_\sigma + |\partial u|_{\sigma+1}) (|v|_s + |w^{(ii)}|_s + |\partial u|_{s+1}) \\
&\quad + C|u|_{\sigma+2}^2 (|u|_s + |\partial u|_{s+1}) \\
&\leq CA\varepsilon \left( \langle t + |x| \rangle^{-3/2} + \langle t + |x| \rangle^{-1} W_-(t, |x|)^{-1+p} \right) (|v|_s + |w^{(ii)}|_s + |\partial u|_{s+1}) \\
&\quad + CA^2\varepsilon^2 \langle t + |x| \rangle^{-2+2\delta} \langle t - |x| \rangle^{-2\delta} (|v|_s + |w^{(ii)}|_s + |\partial u|_{s+1} + |w^{(i)}|_s)
\end{aligned} \tag{5.19}$$

for  $s \leq 2\sigma$ . For  $\rho \geq 0$  and a nonnegative integer  $s$ , we set

$$M_{\rho,s}^{(i)} = \sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{1-\rho} \mathcal{W}_0(t, |x|) |w^{(i)}(t, x)|_s. \tag{5.20}$$

Then, using (5.18) and (5.19), we get

$$\begin{aligned}
\langle x \rangle |F_i|_{2\sigma-3} &\leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1+\lambda} W_-(t, |x|)^{-1/2} \\
&\quad + C_* A^2 \varepsilon^3 \langle t + |x| \rangle^{-2+2\delta+\lambda} \langle t - |x| \rangle^{-2\delta} \\
&\quad + CA^2 \varepsilon^2 M_{\lambda+(1/2), 2\sigma-3}^{(i)} \langle t + |x| \rangle^{-(3/2)+3\delta+\lambda} \langle t - |x| \rangle^{-3\delta} \\
&\leq \left( C_* A \varepsilon^2 + CA^2 \varepsilon^2 M_{\lambda+(1/2), 2\sigma-3}^{(i)} \right) \langle t + |x| \rangle^{-(1/2)+\lambda-\mu} W_-(t, |x|)^{-1+\mu},
\end{aligned}$$

where  $\mu$  is a small and positive constant. Hence, by Lemma 3.3, and also by (3.13) of Lemma 3.4 with  $(\rho, \kappa) = (\lambda + (1/2), 1)$ , we get

$$M_{\lambda+(1/2), 2\sigma-3}^{(i)} \leq C_* (\varepsilon + A\varepsilon^2) + CA^2 \varepsilon^2 M_{\lambda+(1/2), 2\sigma-3}^{(i)}.$$

Therefore, if  $\varepsilon$  is small enough to satisfy  $CA^2 \varepsilon^2 \leq 1/2$ , then we obtain

$$\begin{aligned}
\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{-\lambda+(1/2)} \mathcal{W}_0(t, |x|) |w^{(i)}(t, x)|_{2\sigma-3} &= M_{\lambda+(1/2), 2\sigma-3}^{(i)} \\
&\leq C_* \varepsilon.
\end{aligned} \tag{5.21}$$

Using (5.21), we have

$$\langle x \rangle |F_i|_{2\sigma-3} \leq C_* (A\varepsilon^2 + A^2\varepsilon^3) \langle t + |x| \rangle^{-(1/2)+\lambda-\mu} W_-(t, |x|)^{-1+\mu}. \tag{5.22}$$

Hence, similarly to (5.21), Lemma 3.3 and (3.14) of Lemma 3.4 yield

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{-\lambda-(1/2)} \langle x \rangle \langle t - |x| \rangle |\partial w(t, x)|_{2\sigma-4} \leq C_* \varepsilon, \tag{5.23}$$

provided that  $\varepsilon$  is small enough. Going similar lines to (5.19)–(5.22), we get

$$\langle x \rangle |\mathcal{G}_{k,a}(u, \partial u)|_{2\sigma-3} \leq C_* (A\varepsilon^2 + A^2\varepsilon^3) \langle t + |x| \rangle^{-(1/2)+\lambda-\mu} W_-(t, |x|)^{-1+\mu},$$

and applying Lemma 3.7 to (5.14) with  $|\alpha| \leq 2\sigma - 4$ , we get

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{-\lambda - (1/2)} \langle x \rangle \langle t - |x| \rangle |w^{(ii)}(t, x)|_{2\sigma-4} \leq C_* \varepsilon, \quad (5.24)$$

provided that  $\varepsilon$  is small enough.

Using (5.7), we obtain from (5.9), (5.11), and (5.13) that

$$\begin{aligned} \|\langle t + |\cdot| \rangle |F_i|_{2\sigma-1}\|_{L^2} &\leq C_* A \varepsilon^2 (1+t)^\lambda + C_* A^2 \varepsilon^3 (1+t)^{\lambda+2\delta} \\ &\quad + C A^3 \varepsilon^3 \left\| \langle t + |\cdot| \rangle^{-2+3\delta} \langle t - |\cdot| \rangle^{-3\delta} \right\|_{L^2} \leq C_* A \varepsilon^2 (1+t)^{2\lambda} \end{aligned}$$

for sufficiently small  $\delta$ . Hence Lemma 3.2 leads to

$$\langle t + |x| \rangle^{3/2} |v(t, x)|_{2\sigma-5} \leq C_* \left( \varepsilon + A \varepsilon^2 \sum_{j=0}^{\infty} \sup_{\tau \in (0,t)} \chi_j(\tau) (1+\tau)^{2\lambda} \right). \quad (5.25)$$

Let  $2^{J-1} \leq t < 2^J$  with some nonnegative integer  $J$ . Then we have

$$\begin{aligned} \sum_{j=0}^{\infty} \sup_{\tau \in (0,t)} \chi_j(\tau) (1+\tau)^{2\lambda} &= \sum_{j=0}^J \sup_{\tau \in (0,t)} \chi_j(\tau) (1+\tau)^{2\lambda} \\ &\leq \sum_{j=0}^J 2^{2(j+2)\lambda} = \frac{2^{4\lambda} (2^{2\lambda(J+1)} - 1)}{2^{2\lambda} - 1} \leq C (1+t)^{2\lambda}. \end{aligned}$$

A similar estimate for  $0 \leq t < 1$  is trivially obtained. Now (5.25) leads to

$$\langle t + |x| \rangle^{(3/2)-2\lambda} |v(t, x)|_{2\sigma-5} \leq C_* \varepsilon, \quad (5.26)$$

provided that  $\varepsilon$  is small enough.

*Step 3: Decay Estimates, Part 2.* We make use of the detailed structure of the nonlinearity from now on. Recall that  $\Phi^{(W)}$  is given by (2.3) for a smooth function  $\Phi = \Phi(\xi, \xi', \xi'')$ . We also define

$$\Phi^{(K)}(\eta, \eta', \eta'') := \Phi^{(q)}((\eta, \zeta), (\eta', \zeta'), (\eta'', \zeta'')) \Big|_{(\zeta, \zeta', \zeta'') = (0,0,0)}, \quad (5.27)$$

$$\begin{aligned} \Phi^{(KW)}(\xi, \xi', \xi'') &:= \Phi^{(q)}((\eta, \zeta), (\eta', \zeta'), (\eta'', \zeta'')) \\ &\quad - \Phi^{(K)}(\eta, \eta', \eta'') - \Phi^{(W)}(\zeta, \zeta', \zeta''), \end{aligned} \quad (5.28)$$

$$\Phi^{(H)}(\xi, \xi', \xi'') := \Phi(\xi, \xi', \xi'') - \Phi^{(q)}(\xi, \xi', \xi''), \quad (5.29)$$

where  $(\xi, \xi', \xi'') = ((\eta, \zeta), (\eta', \zeta'), (\eta'', \zeta''))$  as before (the letters “K”, “W”, “H” in this notation stand for “Klein-Gordon”, “wave”, “higher nonlinearity”, respectively).

Since  $(F_i^{(W)})_{N_1+1 \leq i \leq N}$  satisfies the null condition, by (2.10), Lemma 4.1, and (5.1), we get

$$\begin{aligned}
|F_i^{(W)}|_s &\leq C \langle t + |x| \rangle^{-1} (|w|_{[s/2]+2} |\partial w|_{s+1} + |\partial w|_{[s/2]+1} |w|_{s+1}) \\
&\quad + C \langle t + |x| \rangle^{-1} \langle t - |x| \rangle |\partial w|_{[s/2]+1} |\partial w|_{s+1} \\
&\leq CA\varepsilon \langle t + |x| \rangle^{-2+\delta} \langle t - |x| \rangle^{-\delta} |\partial w|_{s+1} \\
&\quad + CA\varepsilon \langle x \rangle^{-1} \langle t + |x| \rangle^{-1} (\langle t - |x| \rangle^{-1} |w|_{s+1} + |\partial w|_{s+1}) \quad (5.30)
\end{aligned}$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$  for  $N_1 + 1 \leq i \leq N$ , provided that  $s \leq 2\sigma$ .

For  $1 \leq i \leq N$ , it is easy to see that each  $F_i^{(K)}(v, \partial v, \partial_x \partial v)$  is a linear combination of  $(\partial^\alpha v_j)(\partial^\beta v_k)$  with  $|\alpha|, |\beta| \leq 2$ , and  $1 \leq j, k \leq N_1$ . Similarly we can see that each  $F_i^{(KW)}(u, \partial u, \partial_x \partial u)$  is a linear combination of  $(\partial^\alpha v_j)(\partial^\beta w_k)$  and  $(\partial^\alpha v_j)w_l^{(ii)}$  with  $|\alpha| \leq 2$ ,  $1 \leq |\beta| \leq 2$ ,  $1 \leq j \leq N_1$ ,  $1 \leq k \leq N_2$ , and  $1 \leq l \leq N_4$ , because of the condition (b-i). Therefore (5.1) yields

$$|F_i^{(K)}|_s \leq C |v|_{[s/2]+2} |v|_{s+2} \leq CA\varepsilon \langle t + |x| \rangle^{-3/2} |v|_{s+2}, \quad (5.31)$$

$$\begin{aligned}
|F_i^{(KW)}|_s &\leq C (|w^{(ii)}|_{[s/2]} + |\partial w|_{[s/2]+1}) |v|_{s+2} + |v|_{[s/2]+2} (|w^{(ii)}|_s + |\partial w|_{s+1}) \\
&\leq CA\varepsilon \langle t + |x| \rangle^{-1} W_-(t, |x|)^{-1+p} |v|_{s+2} \\
&\quad + CA\varepsilon \langle t + |x| \rangle^{-3/2} (|w^{(ii)}|_s + |\partial w|_{s+1}) \quad (5.32)
\end{aligned}$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$  for  $1 \leq i \leq N$ , provided that  $s \leq 2\sigma$ .

Since we have  $F_i^{(H)}(u, \partial u, \partial_x \partial u) = O(|u|^3 + |\partial u|^3 + |\partial_x \partial u|^3)$ , we get

$$\begin{aligned}
|F_i^{(H)}|_s &\leq C |u|_{[s/2]+2}^2 (|v|_{s+2} + |w|_s + |\partial w|_{s+1}) \\
&\leq CA^2 \varepsilon^2 \langle t + |x| \rangle^{-2+2\delta} \langle t - |x| \rangle^{-2\delta} \\
&\quad \times (|v|_{s+2} + |w^{(ii)}|_s + |\partial w|_{s+1} + |w^{(i)}|_s) \quad (5.33)
\end{aligned}$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$  for  $1 \leq i \leq N$ , provided that  $s \leq 2\sigma$ .

Let  $N_1 + 1 \leq i \leq N$  in the following. Using (5.21), (5.23), (5.24), and (5.26), we obtain from (5.30), (5.31), and (5.32) that

$$\langle x \rangle |F_i^{(W)}|_{2\sigma-7} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-(3/2)+\delta+\lambda} W_-(t, |x|)^{-1-\delta}, \quad (5.34)$$

$$\langle x \rangle |F_i^{(K)}|_{2\sigma-7} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-2+2\lambda}, \quad (5.35)$$

$$\langle x \rangle |F_i^{(KW)}|_{2\sigma-7} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1+\lambda} W_-(t, |x|)^{-1} \quad (5.36)$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$ . Similarly, by (5.33), we get

$$\begin{aligned}
\langle x \rangle |F_i^{(H)}|_{2\sigma-7} &\leq C_* A^2 \varepsilon^3 \langle t + |x| \rangle^{-(3/2)+2\delta+2\lambda} \langle t - |x| \rangle^{-1-2\delta} \\
&\quad + CA^2 \varepsilon^2 M_{3\lambda, 2\sigma-7}^{(i)} \langle t + |x| \rangle^{-2+3\delta+3\lambda} \langle t - |x| \rangle^{-3\delta} \quad (5.37)
\end{aligned}$$

at  $(t, x) \in [0, T) \times \mathbb{R}^3$ , where  $M_{3\lambda, 2\sigma-7}^{(i)}$  is given by (5.20). Since we can choose  $\delta$  as small as we wish, (5.34) – (5.37) lead to

$$\langle x \rangle |F_i|_{2\sigma-7} \leq \left( C_* A \varepsilon^2 + C A^2 \varepsilon^2 M_{3\lambda, 2\sigma-7}^{(i)} \right) \langle t + |x| \rangle^{-1+3\lambda-\mu} W_-(t, |x|)^{-1+\mu} \quad (5.38)$$

for small  $\mu > 0$ . Therefore, by (3.13) with  $(\rho, \kappa) = (3\lambda, 1)$ , and by Lemma 3.3, we obtain

$$M_{3\lambda, 2\sigma-7}^{(i)} \leq C_* (\varepsilon + A \varepsilon^2) + C A^2 \varepsilon^2 M_{3\lambda, 2\sigma-7}^{(i)},$$

which leads to

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{1-3\lambda} \mathcal{W}_0(t, |x|) |w^{(i)}(t, x)|_{2\sigma-7} = M_{3\lambda, 2\sigma-7}^{(i)} \leq C_* \varepsilon, \quad (5.39)$$

provided that  $\varepsilon$  is sufficiently small. Now (5.38) and (5.39) yield

$$\langle x \rangle |F_i|_{2\sigma-7} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1+3\lambda-\mu} W_-(t, |x|)^{-1+\mu}. \quad (5.40)$$

Hence Lemma 3.3 and (3.14) imply

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{-3\lambda} \langle x \rangle \langle t - |x| \rangle |\partial w(t, x)|_{2\sigma-8} \leq C_* \varepsilon, \quad (5.41)$$

provided that  $\varepsilon$  is sufficiently small.

Now we are going to estimate  $w^{(ii)}$ . Suppose that  $N_3 + 1 \leq k \leq N_2$ , and  $0 \leq a \leq 3$ . Let  $\mathcal{G}_{k,a}^{(W)}(\zeta, \zeta')$ ,  $\mathcal{G}_{k,a}^{(K)}(\eta, \eta')$ ,  $\mathcal{G}_{k,a}^{(KW)}(\xi, \xi')$ , and  $\mathcal{G}_{k,a}^{(H)}(\xi, \xi')$  be given by (2.3), (5.27), (5.28), and (5.29), respectively, with  $\Phi = \mathcal{G}_{k,a}(\xi, \xi')$ . We can easily verify that  $\mathcal{G}_{k,a}^{(K)}$ ,  $\mathcal{G}_{k,a}^{(KW)}$ , and  $\mathcal{G}_{k,a}^{(H)}$  have similar structures to  $F_i^{(K)}$ ,  $F_i^{(KW)}$ , and  $F_i^{(H)}$ , respectively. By contrast, in view of (2.11), we find that  $\mathcal{G}_{k,a}^{(W)}$  may not be written in terms of the null forms, differently from  $F_i^{(W)}$  with  $N_1 + 1 \leq i \leq N$ . Hence we divide  $w^{(ii)}$  into two parts  $w^{(iii)}$  and  $w^{(iv)}$ : For  $1 \leq l \leq N_4$ , let  $w_l^{(iii)}$  and  $w_l^{(iv)}$  be the solutions to

$$\begin{cases} \square w_l^{(iii)} = \sum_{a=0}^3 \partial_a \left\{ \mathcal{G}_{N_3+l,a}(u, \partial u) - \mathcal{G}_{N_3+l,a}^{(W)}(w, \partial w) \right\}, \\ w_l^{(iii)}(0, x) = w_l^{(ii)}(0, x), \quad \partial_t w_l^{(iii)}(0, x) = \partial_t w_l^{(ii)}(0, x), \end{cases}$$

and

$$\begin{cases} \square w_l^{(iv)} = F_{N_1+N_3+l}^{(W)}(w, \partial w, \partial_x \partial w), \\ w_l^{(iv)}(0, x) = \partial_t w_l^{(iv)}(0, x) = 0, \end{cases}$$

respectively. Since we have

$$F_{N_1+N_3+l}^{(W)}(w, \partial w, \partial_x \partial w) = \sum_{a=0}^3 \partial_a \left( \mathcal{G}_{N_3+l,a}^{(W)}(w, \partial w) \right), \quad 1 \leq l \leq N_4,$$



we get  $w_l^{(ii)} (= w_{N_3+l}) = w_l^{(iii)} + w_l^{(iv)}$ . We put  $w^{(iii)} = (w_l^{(iii)})$  and  $w^{(iv)} = (w_l^{(iv)})$  with  $1 \leq l \leq N_4$ .

It is easy to see that  $\mathcal{G}_{k,a}^{(K)}$ ,  $\mathcal{G}_{k,a}^{(KW)}$ , and  $\mathcal{G}_{k,a}^{(H)}$  enjoy the estimates corresponding to (5.31), (5.32) and (5.33), respectively. Since we have

$$\mathcal{G}_{k,a} - \mathcal{G}_{k,a}^{(W)} = \mathcal{G}_{k,a}^{(K)} + \mathcal{G}_{k,a}^{(KW)} + \mathcal{G}_{k,a}^{(H)},$$

similarly to (5.40) we get

$$\langle x \rangle \left| \mathcal{G}_{k,a} - \mathcal{G}_{k,a}^{(W)} \right|_{2\sigma-7} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1+3\lambda-\mu} W_-(t, |x|)^{-1+\mu}, \quad (5.42)$$

and Lemma 3.7 yields

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{-3\lambda} \langle x \rangle \langle t - |x| \rangle |w^{(iii)}(t, x)|_{2\sigma-8} \leq C_* \varepsilon. \quad (5.43)$$

In view of (5.34), using (3.13) with  $(\rho, \kappa) = (0, (3/2) - 2\lambda)$ , we get

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle \langle t - |x| \rangle^{(1/2)-2\lambda} |w^{(iv)}(t, x)|_{2\sigma-8} \leq C_* \varepsilon, \quad (5.44)$$

since we may assume  $\mu + \delta - \lambda < 0$  for small  $\mu > 0$ . Summing up the estimates, we obtain

$$\sup_{(t,x) \in [0,T) \times \mathbb{R}^3} \langle t + |x| \rangle^{1-3\lambda} W_-(t, |x|)^{(1/2)+\lambda} |w^{(ii)}(t, x)|_{2\sigma-8} \leq C_* \varepsilon, \quad (5.45)$$

provided that  $\varepsilon$  is sufficiently small.

*Step 4: Decay Estimates, Part 3.* Let  $1 \leq i \leq N_1$  in this step. Motivated by the technique in [2], [19] and [29], we introduce

$$\tilde{v}_i = v_i - m_i^{-2} F_i^{(W)}(w, \partial w, \partial_x \partial w) \quad (5.46)$$

in order to treat  $F_i^{(W)}$  for which the null condition is not assumed. Then we get

$$\begin{aligned} (\square + m_i^2) \tilde{v}_i &= \left\{ (\square + m_i^2) v_i - F_i^{(W)}(w, \partial w, \partial_x \partial w) \right\} \\ &\quad - m_i^{-2} \square \left\{ F_i^{(W)}(w, \partial w, \partial_x \partial w) \right\}. \end{aligned} \quad (5.47)$$

From the condition (b-i), we can write

$$F_i^{(W)}(w, \partial w, \partial_x \partial w) = \sum_{|\alpha|, |\beta| \leq 2} \sum_{1 \leq j, k \leq N_2} P_i^{jk\alpha\beta} (\partial^\alpha w_j) (\partial^\beta w_k) \quad (5.48)$$

with appropriate constants  $P_i^{jk\alpha\beta}$ , where  $P_i^{jk\alpha\beta}$  vanishes either when  $1 \leq j \leq N_3$  and  $|\alpha| = 0$ , or when  $1 \leq k \leq N_3$  and  $|\beta| = 0$ . Since we have

$$\square(\varphi\psi) = 2Q_0(\varphi, \psi) + (\square\varphi)\psi + \varphi(\square\psi)$$

for any smooth functions  $\varphi$  and  $\psi$ , from (5.48) we get

$$\square F_i^{(W)} = \tilde{F}_i^{(W)} + \tilde{F}_i^{(H)}, \quad (5.49)$$

where

$$\tilde{F}_i^{(W)} = 2 \sum_{|\alpha|, |\beta| \leq 2} \sum_{1 \leq j, k \leq N_2} P_i^{jk\alpha\beta} Q_0(\partial^\alpha w_j, \partial^\beta w_k), \quad (5.50)$$

$$\tilde{F}_i^{(H)} = \sum_{|\alpha|, |\beta| \leq 2} \sum_{1 \leq j, k \leq N_2} P_i^{jk\alpha\beta} \{(\partial^\alpha F_{N_1+j})(\partial^\beta w_k) + (\partial^\alpha w_j)(\partial^\beta F_{N_1+k})\}. \quad (5.51)$$

Note that each  $\tilde{F}_i^{(W)}$  is written in terms of the null forms, and we can expect extra decay for  $\tilde{F}_i^{(W)}$ . Note also that each  $\tilde{F}_i^{(H)}$  is a function of cubic order with respect to  $\partial^\alpha u_j$  with  $|\alpha| \leq 4$  and  $1 \leq j \leq N$ . From (5.47) and (5.49), we obtain

$$(\square + m_i^2) \tilde{v}_i = F_i^{(K)} + F_i^{(KW)} - m_i^{-2} \tilde{F}_i^{(W)} + (F_i^{(H)} - m_i^{-2} \tilde{F}_i^{(H)}). \quad (5.52)$$

By (5.31) and (5.7), we have

$$\begin{aligned} \left\| \langle t + |\cdot| \rangle |F_i^{(K)}|_{2\sigma-10} \right\|_{L^2(\mathbb{R}^3)} &\leq C A \varepsilon (1+t)^{-1/2} \|v\|_{2\sigma-8} \\ &\leq C_* A \varepsilon^2 (1+t)^{\lambda-(1/2)}. \end{aligned} \quad (5.53)$$

Similarly to (5.34) and (5.36), but using (5.39), (5.41) and (5.45) instead of (5.21), (5.23) and (5.24), we obtain

$$\langle t + |x| \rangle | \tilde{F}_i^{(W)} |_{2\sigma-10} \leq C_* A \varepsilon^2 \langle x \rangle^{-1} \langle t + |x| \rangle^{-1+\delta+3\lambda} W_-(t, |x|)^{-1-\delta}, \quad (5.54)$$

$$\langle t + |x| \rangle |F_i^{(KW)}|_{2\sigma-10} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-(3/2)+3\lambda} W_-(t, |x|)^{-(1/2)-\lambda} \quad (5.55)$$

at  $(t, x) \in [0, \infty) \times \mathbb{R}^3$ . Since we may assume  $\delta < 1/2$ , it follows from (5.54) and (5.55) that

$$\begin{aligned} &\left\| \langle t + |\cdot| \rangle \left( | \tilde{F}_i^{(W)} |_{2\sigma-10} + |F_i^{(KW)}|_{2\sigma-10} \right) \right\|_{L^2(\mathbb{R}^3)} \\ &\leq C_* A \varepsilon^2 \left\| \langle \cdot \rangle^{-1} \langle t + |\cdot| \rangle^{-(1/2)+3\lambda} W_-(t, |\cdot|)^{-(1/2)-\lambda} \right\|_{L^2(\mathbb{R}^3)} \\ &\leq C_* A \varepsilon^2 (1+t)^{-(1/2)+3\lambda}. \end{aligned} \quad (5.56)$$

Going similar lines to (5.33), and then using (5.7) and (5.39), we obtain

$$\begin{aligned} &\left\| \langle t + |\cdot| \rangle |F_i^{(H)} + m_i^{-2} \tilde{F}_i^{(H)}|_{2\sigma-10} \right\|_{L^2(\mathbb{R}^3)} \\ &\leq C A^2 \varepsilon^3 (1+t)^{-1+2\delta} (\|v\|_{2\sigma-10} + \|w^{(ii)}\|_{2\sigma-10} + \|\partial u\|_{2\sigma-7}) \\ &\quad + C_* A^2 \varepsilon^3 \left\| \langle t + |\cdot| \rangle^{-2+3\delta+3\lambda} \langle t - |\cdot| \rangle^{-3\delta} \right\|_{L^2(\mathbb{R}^3)} \\ &\leq C_* A^2 \varepsilon^3 \left( (1+t)^{-1+\lambda+2\delta} + (1+t)^{-(1/2)+3\lambda} \right). \end{aligned} \quad (5.57)$$

To sum up, we have proved

$$\left\| \langle t + |\cdot| \rangle |(\square + m_i^2) \tilde{v}_i|_{2\sigma-10} \right\|_{L^2} \leq C_* A \varepsilon^2 (1+t)^{-1/4}, \quad (5.58)$$

because  $\lambda \leq 1/12$  and  $\delta \ll 1$ . Now Lemma 3.2 implies

$$\begin{aligned} & \sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle^{3/2} |\tilde{v}(t,x)|_{2\sigma-14} \\ & \leq C_* \left( \varepsilon + A \varepsilon^2 \sum_{j=0}^{\infty} \sup_{\tau \in (0,t)} \chi_j(\tau) (1+\tau)^{-1/4} \right) \leq C_* \varepsilon, \end{aligned} \quad (5.59)$$

because we have  $\sup_{\tau \in (0,t)} \chi_j(\tau) (1+\tau)^{-1/4} \leq 2^{-(j-1)/4}$  for  $j \geq 1$ . From (5.41), (5.45) and (5.48), we get

$$\begin{aligned} |F_i^{(W)}|_{2\sigma-14} & \leq C (|\partial w|_{\sigma+1} + |w^{(ii)}|_{\sigma+2}) (|\partial w|_{2\sigma-13} + |w^{(ii)}|_{2\sigma-14}) \\ & \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{3\lambda-2}, \end{aligned}$$

which, together with (5.46) and (5.59), yields

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle^{3/2} |v(t,x)|_{2\sigma-14} \leq C_* \varepsilon \quad (5.60)$$

for small  $\varepsilon$ .

*Step 5: Decay Estimates, the Final Part.* Let  $N_1 + 1 \leq i \leq N$ . Similarly to (5.54), we get

$$\langle x \rangle |F_i^{(W)}|_{2\sigma-9} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-2+\delta+3\lambda} W_-(t, |x|)^{-1-\delta}. \quad (5.61)$$

Therefore, using (3.13) with  $(\rho, \kappa) = (0, 2 - 4\lambda)$ , we get

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle \langle t - |x| \rangle^{1-4\lambda} |w^{(iv)}(t,x)|_{2\sigma-9} \leq C_* \varepsilon, \quad (5.62)$$

which, together with (5.43), yields

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle^{1-3\lambda} W_-(t, |x|)^{1-\lambda} |w^{(ii)}(t,x)|_{2\sigma-9} \leq C_* \varepsilon. \quad (5.63)$$

By (5.31) and (5.60), we obtain

$$\langle x \rangle |F_i^{(K)}|_{2\sigma-16} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-2} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1-\mu} W_-(t, |x|)^{\mu-1} \quad (5.64)$$

at  $(t,x) \in [0,T] \times \mathbb{R}^3$  for small  $\mu > 0$ . From (5.32), (5.41), (5.60), and (5.63) we get

$$\begin{aligned} \langle x \rangle |F_i^{(KW)}|_{2\sigma-16} & \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{3\lambda-(3/2)} W_-(t, |x|)^{-1+p} \\ & \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1-\mu} W_-(t, |x|)^{\mu-1} \end{aligned} \quad (5.65)$$

for small  $\mu > 0$ . (5.33), (5.41), (5.60), and (5.63) yield

$$\begin{aligned} \langle x \rangle |F_i^{(H)}|_{2\sigma-16} &\leq C_* A^2 \varepsilon^3 \langle t + |x| \rangle^{-(5/2)+2\delta} \langle t - |x| \rangle^{-2\delta} \\ &\quad + C_* A^2 \varepsilon^3 \langle t + |x| \rangle^{-2+2\delta+3\lambda} W_-(t, |x|)^{\lambda-2\delta-1} \\ &\quad + C A^2 \varepsilon^2 M_{0,2\sigma-16}^{(i)} \langle t + |x| \rangle^{-2+3\delta} \langle t - |x| \rangle^{-3\delta} \\ &\leq (C_* A^2 \varepsilon^3 + C A^2 \varepsilon^2 M_{0,2\sigma-16}^{(i)}) \langle t + |x| \rangle^{-1-\mu} W_-(t, |x|)^{\mu-1} \end{aligned} \quad (5.66)$$

at  $(t, x) \in [0, T] \times \mathbb{R}^3$ . Since we may assume  $\delta + 3\lambda \leq 1/4$ , say, it follows from (5.61), (5.64), (5.65), and (5.66) that

$$\langle x \rangle |F_i|_{2\sigma-16} \leq (C_* A \varepsilon^2 + C A^2 \varepsilon^2 M_{0,2\sigma-16}^{(i)}) \langle t + |x| \rangle^{-1-\mu} W_-(t, |x|)^{\mu-1}. \quad (5.67)$$

Now Lemma 3.3 and (3.13) with  $(\rho, \kappa) = (0, 1)$  imply

$$M_{0,2\sigma-16}^{(i)} \leq C_*(\varepsilon + A\varepsilon^2) + C A^2 \varepsilon^2 M_{0,2\sigma-16}^{(i)},$$

which leads to

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle \mathcal{W}_0(t, |x|) |w^{(i)}(t, x)|_{2\sigma-16} = M_{0,2\sigma-16}^{(i)} \leq C_* \varepsilon, \quad (5.68)$$

provided that  $\varepsilon$  is small enough. By (5.67) and (5.68), using Lemma 3.3 and (3.14) with  $(\rho, \kappa) = (0, 1)$ , we obtain

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle x \rangle \langle t - |x| \rangle |\partial w(t, x)|_{2\sigma-17} \leq C_* \varepsilon, \quad (5.69)$$

provided that  $\varepsilon$  is small enough.

Going similar lines to (5.64)–(5.67), and using (5.68), we obtain

$$\langle x \rangle |\mathcal{G}_{k,a} - \mathcal{G}_{k,a}^{(W)}|_{2\sigma-16} \leq C_* A \varepsilon^2 \langle t + |x| \rangle^{-1-\mu} W_-(t, |x|)^{\mu-1}, \quad (5.70)$$

and Lemma 3.7 yields

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle x \rangle \langle t - |x| \rangle |w^{(iii)}(t, x)|_{2\sigma-17} \leq C_* \varepsilon. \quad (5.71)$$

Now, (5.62) and (5.71) imply

$$\sup_{(t,x) \in [0,T] \times \mathbb{R}^3} \langle t + |x| \rangle W_-(t, |x|)^{1-p} |w^{(ii)}(t, x)|_{2\sigma-17} \leq C_* \varepsilon, \quad (5.72)$$

provided that  $\varepsilon$  is small enough, since we have assumed  $4\lambda < p$  at the beginning.

*Step 6: Conclusion.* Finally, (5.60), (5.68), (5.69), and (5.72) yield

$$\sup_{0 \leq t < T} \|d_{\sigma,p}(t, \cdot)\|_{L^\infty(\mathbb{R}^3)} \leq C_0 \varepsilon \quad (5.73)$$

for  $\varepsilon \leq \varepsilon_0(A)$ , where  $\varepsilon_0(A)$  is a positive constant depending on  $A$ , and  $C_0$  is some positive constant which depends on  $M_0$  and  $F$ , but is independent of  $A$ ,  $\varepsilon$  and  $T$ . We put  $A_0 = 2C_0$ . Now (5.73) implies (5.2) for  $A \geq A_0$  and  $\varepsilon \leq \varepsilon_0(A)$ . This completes the proof of Proposition 5.1.

## Acknowledgments

The author is grateful to Professors Masahito Ohta and Hideaki Sunagawa for giving the author valuable informations on the previous works in the related field. The author would also like to thank Professor Hideo Kubo for fruitful conversations. This research is partially supported by Grant-in-Aid for Scientific Research (C) (No. 20540211), Japan Society for the Promotion of Science.

## References

- [1] F. Asakura, *Existence of a global solution to a semi-linear wave equation with slowly decreasing initial data in three space dimensions*, Comm. Partial Differential Equations **11**, 1459–1487 (1986).
- [2] A. Bachelot, *Problème de Cauchy global pour des systèmes de Dirac-Klein-Gordon*, Ann. Inst. Henri Poincaré **48**, 387–422 (1988).
- [3] D. Christodoulou, *Global solutions of nonlinear hyperbolic equations for small initial data*, Comm. Pure Appl. Math. **39**, 267–282 (1986).
- [4] V. Georgiev, *Global solution of the system of wave and Klein-Gordon equations*, Math. Z. **203**, 683–698 (1990).
- [5] V. Georgiev, *Decay estimates for the Klein-Gordon equation*, Comm. Partial Differential Equations **17**, 1111–1139 (1992).
- [6] N. Hayashi, P. I. Naumkin, and Ratno Bagus Edy Wibowo, *Nonlinear scattering for a system of nonlinear Klein-Gordon equations*, J. Math. Phys. **49**, 103501 (2008).
- [7] L. Hörmander,  $L^1, L^\infty$  estimates for the wave operator, in: Analyse Mathématique et Applications, Contributions en l’Honneur de J. L. Lions, pp. 211–234, Gauthier-Villars, Paris (1988).
- [8] L. Hörmander, Lectures on Nonlinear Hyperbolic Differential Equations, Springer-Verlag, Berlin (1997).
- [9] A. Hoshiga, and H. Kubo, *Global small amplitude solutions of nonlinear hyperbolic systems with a critical exponent under the null condition*, SIAM J. Math. Anal. **31**, 486–513 (2000).
- [10] F. John, *Blow-up of solutions of nonlinear wave equations in three space dimensions*, Manuscripta Math. **28**, 235–268 (1979).
- [11] F. John, *Blow-up of solutions for quasi-linear wave equations in three space dimensions*, Comm. Pure Appl. Math. **34**, 29–51 (1981).
- [12] S. Katayama, *Global and almost-global existence for systems of nonlinear wave equations with different propagation speeds*, Diff. Integral Eqs. **17**, 1043–1078 (2004).

- [13] S. Katayama, and H. Kubo, *Decay estimates of a tangential derivative to the light cone for the wave equation and their application*, SIAM J. Math. Anal. **39**, 1851–1862 (2008).
- [14] S. Katayama, and H. Kubo, *An alternative proof of global existence for nonlinear wave equations in an exterior domain*, J. Math. Soc. Japan **60**, 1135–1170 (2008).
- [15] S. Katayama, and K. Yokoyama, *Global small amplitude solutions to systems of nonlinear wave equations with multiple speeds*, Osaka J. Math. **43**, 283–326 (2006).
- [16] S. Klainerman, *Global existence of small amplitude solutions to nonlinear Klein-Gordon equations in four space-time dimensions*, Comm. Pure Appl. Math. **38**, 631–641 (1985).
- [17] S. Klainerman, *The null condition and global existence to nonlinear wave equations*, in: Nonlinear Systems of Partial Differential Equations in Applied Mathematics, Part 1, Lectures in Applied Math. **23**, pp. 293–326, AMS, Providence, RI (1986).
- [18] S. Klainerman, *Remarks on the global Sobolev inequalities in the Minkowski space  $\mathbf{R}^{n+1}$* , Comm. Pure Appl. Math. **40**, 111–117 (1987).
- [19] R. Kosecki, *The unit condition and global existence for a class of nonlinear Klein-Gordon equations*, J. Differential Equations **100**, 257–268 (1992).
- [20] M. Kovalyov, *Long-time behavior of solutions of a system of nonlinear wave equations*, Comm. Partial Differential Equations **12**, 471–501 (1987).
- [21] M. Kovalyov, and K. Tsutaya, *Erratum: “Long-time behavior of solutions of a system of nonlinear wave equations”*, Comm. Partial Differential Equations **18**, 1971–1976 (1993).
- [22] K. Kubota, and K. Yokoyama, *Global existence of classical solutions to systems of nonlinear wave equations with different speeds of propagation*, Japanese J. Math. **27**, 113–202 (2001).
- [23] T. Ozawa, K. Tsutaya, and Y. Tsutsumi, *Normal form and global solutions for the Klein-Gordon-Zakharov equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **12**, 459–503 (1995).
- [24] T. Ozawa, K. Tsutaya, and Y. Tsutsumi, *Well-posedness in energy space for the Cauchy problem of the Klein-Gordon-Zakharov equations with different propagation speeds in three space dimensions*, Math. Ann. **313**, 127–140 (1999).
- [25] J. Shatah, *Normal forms and quadratic nonlinear Klein-Gordon Equations*, Comm. Pure Appl. Math. **38**, 685–696 (1985).
- [26] T. C. Sideris, *Decay estimates for the three space dimensional inhomogeneous Klein-Gordon equation and applications*, Comm. Partial Differential Equations **14**, 1421–1455 (1989).
- [27] W. A. Strauss, *Decay and asymptotics for  $\square u = F(u)$* , J. Funct. Anal. **2**, 409–457 (1968).

- [28] K. Tsutaya, *Global existence of small amplitude solutions for the Klein-Gordon-Zakharov equations*, *Nonlinear Anal.* **27**, 1373–1380 (1996).
- [29] Y. Tsutsumi, *Global solutions for the Dirac-Proca equations with small initial data in  $3 + 1$  space time dimensions*, *J. Math. Anal. Appl.* **278**, 485–499 (2003).
- [30] K. Yokoyama, *Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions*, *J. Math. Soc. Japan* **52**, 609–632 (2000).

DEPARTMENT OF MATHEMATICS, WAKAYAMA UNIVERSITY  
930 SAKAEDANI, WAKAYAMA 640-8510, JAPAN  
e-mail: katayama@center.wakayama-u.ac.jp