

Fermi-Dirac Integrals in terms of Zeta Functions

Michael Morales*

Department of Physics, Universidad del Valle de Guatemala,
Guatemala City 01015, Guatemala.

Abstract

A constructive method to write Fermi-Dirac integrals in terms of Riemann and Hurwitz zeta functions is presented. By using of an auxiliary function we can decompose Fermi-Dirac integrals in terms of well-known special functions, which evades the necessity of iterative methods. The procedure is presented for a particular example; nevertheless, its extension to general Fermi-Dirac integrals is straightforward. The final expression obtained along with Sommerfeld's lemma gives nearly a complete representation of Fermi-Dirac integrals.

1 Introduction

Fermi-Dirac integrals $F_{\frac{k}{2}}(\eta)$ are defined as

$$F_{\frac{k}{2}}(\eta) = \int_0^{\infty} \frac{\xi^{\frac{k}{2}}}{1 + e^{\xi - \eta}} d\xi \quad (1)$$

appear in a variety of physical disciplines including partial degenerate stars (Clayton, 1983), thermal conductions by electrons (Khalifaoui & Bennaceur, 1997), and condensed matter physics in general, where $\eta = \mu/kT$, k is the Boltzmann constant, T is the temperature and μ the chemical potential. In this work we determine expressions that allow us to calculate Fermi-Dirac integrals specifically for $\eta \gg 5$, values that so far have been calculated using iterative methods as the trapezoidal method (Mohankumar & Natarajan, 1996). In the following sections we present the case $k=1$; nonetheless, the presented procedure can be extended to any integer k .

This procedure involves a basic auxiliary function that relates $F_{\frac{1}{2}}(\eta)$ with $F_{\frac{1}{2}}(0)$, which in practical terms is a model based on their graphical behavior. This makes the integral $F_{\frac{1}{2}}(\eta)$ easier to calculate obtaining Riemann and Hurwitz Zeta functions, with almost no need for iterative methods.

*Currently member of Department of Mathematics, Ohio University, OH 45701, USA

2 Sommerfeld's lemma applied to $F_{\frac{1}{2}}(\eta)$

Let $\psi(u)$ be a sufficiently regular function which vanishes for $u = 0$, then we have the asymptotic formula (Chandrasekhar, 1958)

$$\int_0^\infty \frac{du}{\frac{1}{\Lambda}e^u + 1} \frac{d\psi(u)}{du} = \psi(u_0) + 2 \left[c_2 \psi''(u_0) + c_4 \psi^{(iv)}(u_0) + \dots \right] \quad (2)$$

where $u_0 = \log[\Lambda]$ and c_2, c_4, \dots are numerical coefficients defined by

$$c_\nu = 1 - \frac{1}{2^\nu} + \frac{1}{3^\nu} - \frac{1}{4^\nu} + \dots \quad (3)$$

It can be show that the Lemma applied to $F_{\frac{1}{2}}(\eta)$ in terms of Riemann Zeta function $\zeta(z)$, with $\psi(u) = u^{1/2}$, and $u_0 = \eta$, results in

$$\int_0^\infty \frac{u^{1/2}}{1 + e^{u-\eta}} du = \frac{2}{3} \eta^{3/2} + 2 \sum_{n=1}^\infty (1 - 2^{1-2n}) \zeta(2n) \frac{(4n-5)!!}{2^{2n-1}} \eta^{-\frac{4n-3}{2}} \quad (4)$$

which is valid when we neglect values of order $e^{-\eta}$. In fact, it can be shown that (4) is valid for $\eta > 3$.

3 $F_{\frac{1}{2}}(\eta)$ and the auxiliary function

As mentioned, we analyze the Fermi-Dirac integral $F_{\frac{1}{2}}(\eta)$

$$F_{\frac{1}{2}}(\eta) = \int_0^\infty \frac{\xi^{\frac{1}{2}}}{1 + e^{\xi-\eta}} d\xi \quad (5)$$

by focusing on the integrand

$$f_{\frac{1}{2}}(\xi, \eta) = \frac{\xi^{\frac{1}{2}}}{1 + e^{\xi-\eta}} \quad (6)$$

and the related function $f_{\frac{1}{2}}(\xi)$

$$f_{\frac{1}{2}}(\xi) = \frac{\xi^{\frac{1}{2}}}{1 + e^\xi} \quad (7)$$

We also define a function, denoted as $f(\xi, \eta)$, as the ratio of both functions, $f_{\frac{1}{2}}(\xi, \eta)$ and $f_{\frac{1}{2}}(0)$

$$f(\xi, \eta) = \frac{f_{\frac{1}{2}}(\eta)}{f_{\frac{1}{2}}(0)} = \frac{1 + e^\xi}{1 + e^{\xi - \eta}} \quad (8)$$

this function, graphically, behaves as $a - be^{-c\xi}$. This suggests that we can write (??eqn:ratio) as

$$f(\xi, \eta) = a - be^{-c\xi} \quad (9)$$

where a, b , and c are constants. The first two can be found considering some analytical aspects of equation (8):

$$f(0, \eta) = \frac{2}{1 + e^{-\eta}} \quad (10)$$

and

$$\lim_{x \rightarrow \infty} f(0, \eta) = e^\eta \quad (11)$$

this implies that $a = e^\eta$ and $b = \frac{e^\eta - 1}{e^{-\eta} + 1}$. With this we obtain

$$f(\xi, \eta) = e^\eta - \frac{e^\eta - 1}{e^{-\eta} + 1} e^{-c\xi} \quad (12)$$

This leaves the issue of finding c . We take the following approach, perhaps the only one in this procedure that needs iterative methods: we find the extreme point of $f_{\frac{1}{2}}(\xi, \eta)$, equation (6). After some algebra, we obtain the following equation which must be solved for ξ

$$e^\eta + e^\xi(1 - 2\xi) = 0 \quad (13)$$

We call the solution of (13) ξ_m , which depends on η . Using equations (8) and (12), and substituting ξ_m , we obtain:

$$c = -\frac{1}{\xi_m} \ln \left[\frac{(e^{-\eta} + 1) \left(e^\eta - \frac{1 + e^{\xi_m}}{1 + e^{\xi_m - \eta}} \right)}{e^\eta - 1} \right] \quad (14)$$

It can be shown that calculating the maxima of equation (6) and the product $f(\xi, \eta)f_{\frac{1}{2}}(\xi)$ leads to the same result. After we determine the constants in equation (9), we can write equation (5) as

$$\int_0^\infty \frac{\xi^{\frac{1}{2}}}{1+e^{\xi-\eta}} d\xi = \int_0^\infty \left(e^\eta - \frac{e^\eta - 1}{e^{-\eta} + 1} e^{-c\xi} \right) \frac{\xi^{\frac{1}{2}}}{1+e^\xi} d\xi = e^\eta \int_0^\infty \frac{\xi^{\frac{1}{2}}}{1+e^\xi} d\xi - \frac{e^\eta - 1}{e^{-\eta} + 1} \int_0^\infty \frac{e^{-c\xi} \xi^{\frac{1}{2}}}{1+e^\xi} d\xi \quad (15)$$

where the first integral in the right hand side can be expressed as Arfken (2005)

$$\int_0^\infty \frac{\xi^{\frac{1}{2}}}{1+e^\xi} d\xi = \Gamma\left(\frac{3}{2}\right) \left(1 - 2^{-\frac{1}{2}}\right) \zeta\left(1 + \frac{1}{2}\right) \quad (16)$$

The second term in (refeqn:preintegral) after expanding some terms takes the form

$$\int_0^\infty \frac{e^{-c\xi} \xi^{\frac{1}{2}}}{1+e^\xi} d\xi = \Gamma\left(\frac{3}{2}\right) \sum_{n=0}^\infty \frac{(-1)^n}{(c+1+n)^{1+\frac{1}{2}}} = \Gamma\left(\frac{3}{2}\right) \left[2^{-\frac{1}{2}} \zeta\left(1 + \frac{1}{2}, \frac{c+1}{2}\right) - \zeta\left(1 + \frac{1}{2}, c+1\right) \right] \quad (17)$$

In this expression we recognize $\zeta(p, q)$ as the Hurwitz zeta function, where we have used the result in Williams & Nan-yue (1993, p3). Using these results we can finally write $F_{\frac{1}{2}}(\eta)$ as

$$F_{\frac{1}{2}}(\eta) = \Gamma\left(\frac{3}{2}\right) \left[e^\eta \left(1 - 2^{-\frac{1}{2}}\right) \zeta\left(1 + \frac{1}{2}\right) - \frac{e^\eta - 1}{e^{-\eta} + 1} \left[2^{-\frac{1}{2}} \zeta\left(1 + \frac{1}{2}, \frac{c+1}{2}\right) - \zeta\left(1 + \frac{1}{2}, c+1\right) \right] \right] \quad (18)$$

4 Some results

Using equation (18) we construct Table 1 for some specific values of η which allows a comparison with values taken from McDougall & Stoner (1938), including some values for positive $\eta < 1$. These values have been used in electron conduction opacity, (Khalfaoui & Bennaceur, 1997), showing how reliable and potentially applicable this method is.

Table 1: $F_{\frac{1}{2}}(\eta)$ values comparison

η	$F_{\frac{1}{2}}$ eq. (18)	$F_{\frac{1}{2}}$ McDougall & Stoner	difference (%)
-4.0	0.01614	0.01613	0.06
-3.0	0.04345	0.04337	0.18
-2.0	0.11506	0.11459	0.41
-1.0	0.29241	0.29050	0.66
0.0	0.67809	0.67810	0.00
0.1	0.73203	0.73340	0.19
0.5	0.97795	0.99021	1.24
1.0	1.35129	1.39638	3.23
2.0	2.30003	2.50246	8.09
3.0	3.58315	3.97699	9.90
4.0	5.54950	5.77073	3.83
5.0	8.99919	7.83797	14.82

5 Observations

The model according to (9) is reliable for $\eta < 5$, specially with for values close to zero and negative η . Yet, as the ratio of the functions in equation (8) shows a higher inflexion point for $\eta > 5$, the model needs an improvement. The reason for the limitation around this value appears when analyzing the equation (18), where a simple examination shows a dependency of order e^η ; recall that, in Sommerfeld lemma, this dependence is of order $e^{-\eta}$.

The use of the auxiliary function, a simple exponential model, practically evades the use of iterative methods or pole expansion methods. Numerical methods are needed only for the simple equation (13) and to determine Riemann and Hurwitz zeta functions.

6 Conclusion

Equation (18) gives an expression for calculating Fermi-Dirac integrals $F_{\frac{1}{2}}(\eta)$ in terms of the Riemann and Hurwitz Zeta functions. This approach complements Sommerfeld's lemma, resulting in a nearly complete expression for Fermi-Dirac integrals.

Using the same procedure we can find expressions the corresponding expression $F_{\frac{3}{2}}(\eta)$

$$F_{\frac{3}{2}}(\eta) = \Gamma\left(\frac{5}{2}\right) \left[e^\eta \left(1 - 2^{-\frac{3}{2}}\right) \zeta\left(1 + \frac{3}{2}\right) - \frac{e^\eta - 1}{e^{-\eta} + 1} \left[2^{-\frac{3}{2}} \zeta\left(1 + \frac{3}{2}, \frac{c+1}{2}\right) - \zeta\left(1 + \frac{3}{2}, c+1\right) \right] \right] \quad (19)$$

in general, the procedure presented here can be extended for any integer k to determine $F_{\frac{k}{2}}(\eta)$,

$$F_{\frac{k}{2}}(\eta) = \Gamma\left(1 + \frac{k}{2}\right) \left[e^\eta \left(1 - 2^{-\frac{k}{2}}\right) \zeta\left(1 + \frac{k}{2}\right) - \frac{e^\eta - 1}{e^{-\eta} + 1} \left[2^{-\frac{k}{2}} \zeta\left(1 + \frac{k}{2}, \frac{c+1}{2}\right) - \zeta\left(1 + \frac{k}{2}, c+1\right) \right] \right] \quad (20)$$

with the remark that ξ_m , in order to find c (notice that c and $f(\xi, \eta)$ are k -independent) equation (13) must also be generalized:

$$ke^\eta + e^\xi(k - 2\xi) = 0 \quad (21)$$

Although is not a complete close form expression, by using an elementary methods, we obtained a single expression with which when combined with Sommerfeld's lemma, we have a complete accurate description for Fermi-Dirac integrals $F_{\frac{k}{2}}(\eta)$.

References

- Arfken, G. & Weber, H. *Mathematical Methods for physicists*, sixth ed., Elsevier Academic Press, 2005.
- Chandrasekhar, S. *An introduction to the study of stellar structure*. Dover publications, 1958.
- Clayton, D. *Principles of stellar evolution and nucleosynthesis*, 2nd ed., McGraw-Hill, 1983.
- Gong, Z., Zejda, L., Däppen, W. & Aparicio, J. *Generalized Fermi-Dirac Functions and derivatives: properties and evaluation* Comp. Phys. Commun. **136**(2001), 294-309
- Khalifaoui, A. & Bennaceur, D. *Thermal conduction by electrons in hot dense plasmas*. Astrophys. J. March 20; 478(1997), 326-331.
- McDougall, J. and Stoner, E. Mohankumar, N. *The computation of Fermi-Dirac Functions*, Phil. Trans. Roy. Soc. **237**:67(1938), 67-104.

Mohankumar, N. & Natarajan, A., *A note on the evaluation of the generalized Fermi-Dirac integral*, *Astrophys. J.* Feb. 10; 458(1996), 233-235.

Williams, K & Nan-yue, Z., *Special values of the Lerch Zeta Function and the evaluation of certain integrals*, *Proc. Amer. Math. Soc.* **119** (1993), 35-49.