

SYMMETRY PROPERTIES OF THE GENERALIZED HIGHER-ORDER EULER POLYNOMIALS

TAEKYUN KIM

ABSTRACT. The purpose of this paper is to generalize this relation of symmetry between the power sum polynomials and the generalized Euler polynomials to the relation between the power sum polynomials and the generalized higher-order Euler polynomials.

§1. Introduction

Let d be a fixed positive odd integer and let χ be the Dirichlet's character with conductor d . Then the generalized Euler numbers and polynomials attached to χ are defined as

$$(1) \quad \begin{aligned} \frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{at}}{e^{dt} + 1} &= \sum_{n=0}^{\infty} E_{n,\chi} \frac{t^n}{n!}, \\ \frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{at}}{e^{dt} + 1} e^{xt} &= \sum_{n=0}^{\infty} E_{n,\chi}(x) \frac{t^n}{n!}, \text{ for } |t| < \frac{\pi}{d}, \text{ (see [1-14])}. \end{aligned}$$

For a real or complex parameter α , we define the generalized higher-order Euler numbers and polynomials attached to χ as follows:

$$(2) \quad \begin{aligned} \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{at}}{e^{dt} + 1} \right)^\alpha &= \sum_{n=0}^{\infty} E_{n,\chi}^{(\alpha)} \frac{t^n}{n!}, \text{ and} \\ \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{at}}{e^{dt} + 1} \right)^\alpha e^{xt} &= \sum_{n=0}^{\infty} E_{n,\chi}^{(\alpha)}(x) \frac{t^n}{n!}, \text{ where } |t| < \frac{\pi}{d}. \end{aligned}$$

From (2), we note that

$$(3) \quad E_{n,\chi}^{(\alpha)}(x) = \sum_{l=0}^n \binom{n}{l} E_{l,\chi}^{(\alpha)} x^{n-l}.$$

Key words and phrases. Euler numbers and polynomials, Symmetry.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2009

By (1), we can also easily see that

$$(4) \quad E_{k,\chi}(nd) + E_{k,\chi} = 2T_{k,\chi}(nd - 1), \text{ where } T_{k,\chi}(n) = \sum_{l=0}^n (-1)^l \chi(l) l^k.$$

Recently, Kurt (see [10]) and several authors (see [5-8], [11-13]) have studied the symmetry property for the Bernoulli and Euler polynomials. The main purpose of this paper is to prove an identity of symmetry for the generalized higher-order Euler polynomials by using the method of Kurt. It turns out that the recurrence relation and multiplication theorem for the generalized Euler polynomials attached to χ .

§2. Identities related to the generalized higher-order Euler polynomials

For $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1, w_2 \equiv 1 \pmod{2}$, let us consider the following functional equation:

$$\begin{aligned} I &= \frac{1}{2} \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{w_1 at}}{e^{dw_1 t} + 1} \right)^m e^{w_1 w_2 xt} (e^{dw_1 w_2 t} + 1) \\ &\quad \times \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{w_2 at}}{e^{dw_2 t} + 1} \right)^m e^{w_1 w_2 yt} \\ (5) \quad &= \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{w_1 at}}{e^{dw_1 t} + 1} \right)^m e^{w_1 w_2 xt} \left(\frac{e^{dw_1 w_2 t} + 1}{e^{w_2 dt} + 1} \right) \left(\sum_{b=0}^{d-1} \chi(b) (-1)^b e^{w_2 bt} \right) \\ &\quad \times \left(\frac{2 \sum_{a=0}^{d-1} (-1)^a \chi(a) e^{w_2 at}}{e^{dw_2 t} + 1} \right)^{m-1} e^{w_1 w_2 yt}. \end{aligned}$$

It is easy to see that

$$\begin{aligned} (6) \quad &\left(\frac{e^{w_1 dt} + 1}{e^{dt} + 1} \right) \left(\sum_{i=0}^{d-1} \chi(i) e^{it} (-1)^i \right) = \left(\sum_{l=0}^{w_1 d-1} e^{ldt} (-1)^l \right) \left(\sum_{i=0}^{d-1} \chi(i) (-1)^i e^{it} \right) \\ &= \sum_{i=0}^{w_1 d-1} \chi(i) (-1)^i e^{it} = \sum_{k=0}^{\infty} \left(\sum_{i=0}^{w_1 d-1} \chi(i) (-1)^i i^k \right) \frac{t^k}{k!} = \sum_{k=0}^{\infty} T_{k,\chi}(w_1 d - 1) \frac{t^k}{k!}. \end{aligned}$$

By (5) and (6), we have

$$\begin{aligned} (7) \quad I &= \left(\sum_{i=0}^{\infty} E_{i,\chi}^{(m)}(w_2 x) \frac{w_1^i t^i}{i!} \right) \left(\sum_{l=0}^{\infty} T_{l,\chi}(w_1 d - 1) \frac{w_2^l t^l}{l!} \right) \left(\sum_{k=0}^{\infty} E_{k,\chi}^{(m-1)}(w_1 y) \frac{w_2^k t^k}{k!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{j=0}^n \binom{n}{j} w_2^j w_1^{n-j} E_{n-j,\chi}^{(m)}(w_2 x) \sum_{k=0}^j T_{k,\chi}(w_1 d - 1) \binom{j}{k} E_{j-k,\chi}^{(m-1)}(w_1 y) \right) \frac{t^n}{n!}. \end{aligned}$$

By the symmetry of I in w_1 and w_2 , we also see that

$$(8) \quad I = \sum_{n=0}^{\infty} \left(\sum_{j=0}^n \binom{n}{j} w_1^j w_2^{n-j} E_{n-j,\chi}^{(m)}(w_1 x) \sum_{k=0}^j \binom{j}{k} T_{k,\chi}(w_2 d - 1) E_{j-k,\chi}^{(m-1)}(w_2 y) \right) \frac{t^n}{n!}.$$

By comparing the coefficients on the both sides of (7) and (8), we obtain the following theorem.

Theorem 1. *Let χ be the Dirichlet's character with an odd conductor $d \in \mathbb{N}$. For $w_1, w_2, d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$, $w_1 \equiv 1 \pmod{2}$, $w_2 \equiv 1 \pmod{2}$, we have*

$$(9) \quad \begin{aligned} & \sum_{j=0}^n \binom{n}{j} w_2^j w_1^{n-j} E_{n-j,\chi}^{(m)}(w_2 x) \sum_{k=0}^j \binom{j}{k} T_{k,\chi}(w_1 d - 1) E_{j-k,\chi}^{(m-1)}(w_1 y) \\ &= \sum_{j=0}^n \binom{n}{j} w_1^j w_2^{n-j} E_{n-j,\chi}^{(m)}(w_1 x) \sum_{k=0}^j \binom{j}{k} T_{k,\chi}(w_2 d - 1) E_{j-k,\chi}^{(m-1)}(w_2 y). \end{aligned}$$

Let $y = 0$ and $m = 1$ in (9). Then we have the following corollary.

Corollary 2. *For $n \in \mathbb{Z}_+$, we have*

$$(10) \quad \begin{aligned} & \sum_{j=0}^n \binom{n}{j} w_2^j w_1^{n-j} E_{n-j,\chi}(w_2 x) T_{j,\chi}(w_1 d - 1) \\ &= \sum_{j=0}^n \binom{n}{j} w_1^j w_2^{n-j} E_{n-j,\chi}(w_1 x) T_{j,\chi}(w_2 d - 1). \end{aligned}$$

From (5) and (6), we can also derive the following Eq.(11).

$$(11) \quad \begin{aligned} I &= \left(\frac{2 \sum_{a=0}^{d-1} \chi(a) (-1)^a e^{w_1 a t}}{e^{dw_1 t} + 1} \right)^m e^{w_1 w_2 x t} \left(\sum_{i=0}^{w_1 d - 1} \chi(i) (-1)^i e^{i w_2 t} \right) \\ &\times \left(\frac{2 \sum_{b=0}^{d-1} \chi(b) (-1)^b e^{w_2 b t}}{e^{dw_2 t} + 1} \right)^{m-1} e^{w_1 w_2 y t} \\ &= \left(\sum_{i=0}^{w_1 d - 1} \chi(i) (-1)^i \sum_{k=0}^{\infty} E_{k,\chi}^{(m)}(w_2 x + \frac{w_2}{w_1} i) \frac{w_1^k t^k}{k!} \right) \left(\sum_{l=0}^{\infty} E_{l,\chi}^{(m-1)}(w_1 y) \frac{w_2^l t^l}{l!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} w_1^k w_2^{n-k} E_{n-k,\chi}^{(m-1)}(w_1 y) \sum_{i=0}^{w_1 d - 1} \chi(i) (-1)^i E_{k,\chi}^{(m)}(w_2 x + \frac{w_2}{w_1} i) \right) \frac{t^n}{n!}. \end{aligned}$$

By the symmetry of I in w_1, w_2 , we also see that

$$(12) \quad I = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} w_2^k w_1^{n-k} E_{n-k, \chi}^{(m-1)}(w_2 y) \sum_{i=0}^{w_2 d-1} \chi(i) (-1)^i E_{k, \chi}^{(m)}(w_1 x + \frac{w_1}{w_2} i) \right) \frac{t^n}{n!}.$$

By comparing the coefficients on the both sides of (11) and (12), we obtain the following theorem.

Theorem 3. *Let w_1, w_2 be the odd natural numbers. For $n \in \mathbb{Z}_+, m \in \mathbb{N}$, we have*

$$(13) \quad \begin{aligned} & \sum_{k=0}^n \binom{n}{k} w_1^k w_2^{n-k} E_{n-k, \chi}^{(m-1)}(w_1 y) \sum_{i=0}^{w_1 d-1} \chi(i) (-1)^i E_{k, \chi}^{(m)}(w_2 x + \frac{w_2}{w_1} i) \\ &= \sum_{k=0}^n \binom{n}{k} w_2^k w_1^{n-k} E_{n-k, \chi}^{(m-1)}(w_2 y) \sum_{i=0}^{w_2 d-1} \chi(i) (-1)^i E_{k, \chi}^{(m)}(w_1 x + \frac{w_1}{w_2} i) \end{aligned}$$

Let $m = 1$ in (13). Then we have

$$w_1^n \sum_{i=0}^{w_1 d-1} \chi(i) (-1)^i E_{n, \chi}(w_2 x + \frac{w_2}{w_1} i) = w_2^n \sum_{i=0}^{w_2 d-1} \chi(i) (-1)^i E_{n, \chi}(w_1 x + \frac{w_1}{w_2} i)$$

REFERENCES

- [1] M. Cenkci, M. Can and V. Kurt, *p-adic interpolation functions and Kummer-type congruences for q-twisted Euler numbers*, Adv. Stud. Contemp. Math. **9** (2004), 203–216.
- [2] M. Cenkci, M. Can and V. Kurt, Y. Simsek, *Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Euler l-functions*, Adv. Stud. Contemp. Math. **18** (2009), 135–160.
- [3] L. C. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht, 1974.
- [4] E. Deeba, D. Rodriguez, *Stirling's series and Bernoulli numbers*, Amer. Math. Monthly **98** (1991), 423-426.
- [5] T. Kim, *A symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on \mathbb{Z}_p* , Russian J. Math. Phys. **16** (2009), 51-54.
- [6] T. Kim, *Symmetry p-adic invariant integral on \mathbb{Z}_p for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. **14** (2008), 1267-1277.
- [7] T. Kim, *Symmetry identities for the twisted generalized Euler polynomials*, Adv. Stud. Contemp. Math. **19** (2009), 111-118.
- [8] Y. H. Kim, K.-W. Hwang, *A symmetry of power sum and twisted Bernoulli polynomials*, Adv. Stud. Contemp. Math. **18** (2009), 127-133.
- [9] Y. H. Jang, M.S. Kim, J. W. Son, *An analogue of Bernoulli numbers and their congruences*, Proceedings of the Jangjeon Mathematical Society **1** (2000), 133-144.
- [10] V. Kurt, *A Further symmetric relation on the analogue of the Apostol-Bernoulli and the analogue of the Apostol-Genocchi polynomials*, Appl. Math. Sciences **3, no.56** (2009), 2357-2764.
- [11] L.C. Jang, *A study on the distribution of twisted q-Genocchi polynomials*, Adv. Stud. Contemp. Math. **18** (2009), 181–189.

- [12] H. Ozden, Y. Simsek, S.-H. Rim, I.N. Cangul, *A note on p -adic q -Euler measure*, Adv. Stud. Contemp. Math. **14** (2007), 233–239.
- [13] H. Ozden, Y. Simsek, *Remarks on q -Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), 41–48.
- [14] K. Shiratani, S. Yamamoto, *On a p -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci., Kyushu University Ser. A **39** (1985), 113-125.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University, Seoul
139-701, S. Korea e-mail: tkkim@kw.ac.kr