Compact Lie algebras, transversely Lie foliations and fibrations

M. Tavares

Contents

1	Introduction	1
2	Lie foliations	3
3	Foliations invariant under transverse actions	6
4	Lie foliations with compact Lie algebra	6
5	Lie foliations on manifolds with amenable fundamental group	12
6	Subexponential growth and Lie foliations	13
7	Extension to transversely homogeneous foliations	14
8	A remark about Riemannian foliations	15

1 Introduction

An important theorem due to Tischler states that a smooth closed manifold fibers over the qdimensional torus if and only if admits a foliation given by q linearly independent closed forms of degree one. Indeed, given q linearly independent closed forms of degree one, it is possible to find new closed forms arbitrarily close to the originals and having *rational periods*. The foliation defined by these new forms has all leaves compact and these leaves are the fibers of a locally trivial fibration over the torus. All this can be seen as embedded in the theory of Lie-foliations [10], [12]. Let \mathfrak{g} be a Lie algebra of dimension q and let M be a smooth closed manifold. We say that a foliation \mathcal{F} of codimension q on M is a \mathfrak{g} -Lie foliation if, given structure constants $\{c_{ij}^k\}$ of \mathfrak{g} , there exists a system $\{\omega_1,\ldots,\omega_q\}$ of linearly independent smooth forms of degree one on M which satisfies $d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$ and whose kernel is tangent to \mathcal{F} . Let G be a connected Lie group having \mathfrak{g} as Lie algebra. A foliation \mathcal{F} on M is a Lie \mathfrak{g} -foliation if and only if \mathcal{F} admits a Lie transverse structure modeled by G (see [12]), that is, \mathcal{F} is defined by local submersions taking values on G with transition maps given by left translations on G. The existence of Lie foliation does not guarantee the existence of a global submersion of M in G. The difficulty in obtaining such a submersion from a Lie foliation \mathcal{F} can be measured by the *global* holonomy group of \mathcal{F} with respect to G, which is a representation of the fundamental group of M in G. In his thesis, Fedida proves that, in fact, the structure of \mathcal{F} is strongly related to its global holonomy ([10], [12]). In this work, we obtain a generalization of the Tischler fibration Theorem for \mathfrak{g} -Lie foliations, when \mathfrak{g} is a *compact* Lie algebra, in the sense that \mathfrak{g} is the Lie algebra of some compact Lie group:

Theorem 1. Let \mathfrak{g} be a compact Lie algebra, M a smooth closed manifold and \mathcal{F} a Lie \mathfrak{g} foliation on M. Let G be a connected compact Lie group having \mathfrak{g} as Lie algebra. If the closure
of the global holonomy of \mathcal{F} with respect to G has Abelian Lie algebra, then M possesses a finite
covering which fibers over G. Indeed, there is a finite covering of M where the lifting of \mathcal{F} can
be C^0 -approximated by Lie \mathfrak{g} -foliations having all leaves compact.

The proof of Theorem 1 gives a description of the structure of the foliation, and this allows us to well understand the finite covering mentioned. Combining our result with a result due to Caron and Carrière ([6]), we have

Corollary 1. Let \mathfrak{g} be a semisimple compact Lie algebra and let \mathcal{F} be a (not necessarily minimal) one-dimensional Lie \mathfrak{g} -foliation on a smooth closed manifold M. We have

- 1. there is a finite covering of M where the lifting of \mathcal{F} can be C^0 -approximated by Lie \mathfrak{g} -foliations having all leaves compact;
- 2. If G is the simply connected Lie group having \mathfrak{g} as Lie algebra, then M possesses a finite covering which fibers over G.

An important remark in the theory is that the closure of the global holonomy of a Lie foliation must be an Abelian group provided that the fundamental group of M is Abelian. In this work, we extend this observation by proving that if the fundamental group of M is amenable and G is compact, then the closure of the global holonomy of \mathcal{F} has Abelian Lie algebra. Thus, we can apply Theorem A to obtain the following:

Theorem 2. Let \mathfrak{g} be a compact Lie algebra and M a smooth closed manifold with amenable fundamental group. If M admits a Lie \mathfrak{g} -foliation, then there is a finite covering of M where the lifted foliation can be C^0 -approximated by Lie \mathfrak{g} -foliations having all leaves compact. If Gis a compact Lie group having \mathfrak{g} as Lie algebra, then M possesses a finite covering which fibers over G.

We also studied the particular case where the fundamental group of the ambient manifold has subexponential growth and \mathfrak{g} is not necessarily compact:

Theorem 3. Let M be a smooth closed manifold with fundamental group of subexponential growth and suppose that M admits a Lie g-foliation. We have:

- 1. If G is a connected Lie group having \mathfrak{g} as Lie algebra, then G has polynomial volume growth;
- 2. If $\mathfrak{s} < \mathfrak{g}$ is a Levi factor of \mathfrak{g} and S is a connected Lie group having \mathfrak{s} as Lie algebra, then S is compact and M possesses a finite covering which fibers over S.

In the semisimple case, the result above has the following statement:

Corollary 2. Let M be a smooth closed manifold which admits a Lie \mathfrak{g} -foliation where \mathfrak{g} is a semisimple Lie algebra. If \mathfrak{g} is non-compact then the fundamental group of M has exponential growth. If the fundamental group of M has subexponential growth then any connected Lie group G having \mathfrak{g} as Lie algebra is compact and M possesses a finite covering which fibers over G.

The concept of Lie foliations admits a natural generalization: Let G be a connected Lie group of dimension n and let K be a connected closed Lie subgroup of G of dimension n-q. Let \mathfrak{g} and \mathfrak{k} be the Lie algebras of G and K, respectively. Let $\beta = \{X_1, \ldots, X_q, X_{q+1}, \ldots, X_n\}$ be a basis of \mathfrak{g} such that X_{q+1}, \ldots, X_n is a basis of \mathfrak{k} and denote by $\{c_{ij}^k\}$ the structure constants of β . Suppose that a smooth manifold M admits n smooth forms of degree one $\omega_1, \ldots, \omega_q, \omega_{q+1}, \ldots, \omega_n$ satisfying $d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$. If the forms $\omega_1, \ldots, \omega_q$ are linearly independent at each point of M, then they define a foliation \mathcal{F} of codimension q on M and, for this case, we can adapt Theorem A. Such a foliation admits a structure of *transversely homogeneous foliation modeled by* G/K (see [2], [12]), that is, \mathcal{F} is defined by local submersions taking values on G/K with transition maps given by left translations on G. Similar structures are constructed, as a global holonomy group ([12], [2]). We should note, however, that not every transversely homogeneous foliation is obtained from global forms of degree one. Our preceding results together with results of Blumenthal ([2], [3]), give the following fibration result for transversely homogeneous foliations:

Theorem 4. Let G be a connected compact Lie group, K a closed and connected subgroup of G and let M be a smooth closed manifold with amenable fundamental group. Given a G/K-transversely homogeneous foliation \mathcal{F} on M, suppose one of the following conditions is satisfied

- 1. $H^1(M, K)$ is trivial;
- 2. the normal bundle of \mathcal{F} is trivial and G/K is a simply-connected Riemannian homogeneous space of constant curvature.

Then there is a finite covering of M where the lifting of \mathcal{F} can be C^0 -approximated by G/Ktransversely homogeneous foliations. Moreover, M possesses a finite covering which fibers over G/K.

We stress that conditions (1) and (2) in Theorem 4 above are used in order to assure that the homogeneous transverse structure is given by globally defined one-forms and can be replaced by this hypothesis, which may also verified in some other situations.

Corollary 3. Let \mathcal{F} be an one-dimensional SO(2q+1)/SO(2q)-transversely homogeneous foliation with trivial normal bundle on a smooth closed manifold M. Then there is a finite covering of M where the lifting of \mathcal{F} can be C^0 -approximated by SO(2q+1)/SO(2q)-foliations having all leaves compact. If M is orientable and q = 1, then the finite covering mentioned is either the identity or a double covering of M.

2 Lie foliations

Let us recall the formal definition and main properties we need concerning Lie foliations (see [12], [10] for more):

Definition 1. Let M be a smooth manifold and \mathfrak{g} a Lie algebra of dimension q. We say that a codimension q foliation \mathcal{F} on M is a *Lie* \mathfrak{g} -*foliation* if \mathcal{F} is tangent to the kernel of a smooth form of degree one Ω on M with values in \mathfrak{g} such that

- 1. $\Omega(x): T_x M \to \mathfrak{g}$ is surjective, for each $x \in M$;
- 2. $d\Omega + \frac{1}{2}[\Omega, \Omega] = 0.$

Let X_1, \ldots, X_q be a basis of a Lie algebra \mathfrak{g} with structure constants $\{c_{ij}^k\}$. If there are linearly independent forms of degree one $\omega_1, \ldots, \omega_q$ on a smooth manifold M which satisfy $d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$, then $\Omega = \sum_{l=1}^q \omega_l X_l$ is a form of degree one on M with values in \mathfrak{g} satisfying the second item of Definition 1. Conversely, let Ω be a smooth form of degree one on M with values in \mathfrak{g} satisfying (2). If X_1, \ldots, X_q is a basis of \mathfrak{g} and $\Omega = \sum_{l=1}^q \omega_l X_l$, then $d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$. The form Ω is surjective in each point of M if and only if the forms $\omega_1, \ldots, \omega_q$ are linearly independent in each point of M. Moreover, the kernel of Ω coincides with the kernel of the system $\{\omega_1, \ldots, \omega_q\}$. Therefore, a codimension q foliation \mathcal{F} on M is a Lie \mathfrak{g} -foliation if and only if \mathcal{F} is tangent to the kernel of a system $\{\omega_1, \ldots, \omega_q\}$ of smooth linearly independent forms of degree one on M which satisfies

$$d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j.$$

Example 1. A Lie \mathbb{R}^r -foliation on a smooth manifold M is a foliation defined by the kernel of r smooth linearly independent closed forms of degree one on M.

Example 2. Let $\pi : E \to M$ be a *G*-principal bundle. If a connection form Ω one *E* is flat, that is, has zero curvature, then Ω satisfies $d\Omega = -\frac{1}{2}[\Omega, \Omega]$ (see [17], pg 78, 92-94). Therefore, Ω defines a Lie g-foliation \mathcal{F} on *E* which is transverse to the fibers.

We also recall the concept of Lie transverse structure for foliations:

Definition 2. Let \mathcal{F} be a codimension q foliation on a differentiable manifold M and let G be a q-dimensional Lie group. A Lie transverse structure of model G for \mathcal{F} is given by a family of submersions $\{f_i : U_i \to G\}_{i \in \Lambda}$ and a family of locally constant maps $g_{ij} : U_i \cap U_j \to \{\text{left translations on } G\}$ such that

- 1. $\{U_i\}_{i \in \Lambda}$ is an open cover of M;
- 2. the plaques of \mathcal{F} in U_i are given by $f_i = \text{constant}$;
- 3. $f_i(x) = g_{ij}(x) \circ f_j(x), \forall x \in U_i \cap U_j.$

The result below allow us to understand the relation between Lie \mathfrak{g} -foliations and a foliation with a Lie transverse structure:

Darboux-Fedida's Theorem. Let M be a smooth manifold and let $\mathcal{F} = \mathcal{F}_{\Omega}$ be a Lie \mathfrak{g} foliation on M. Let G be a connected Lie group having \mathfrak{g} as Lie algebra and denote by α its
Maurer-Cartan form. Denote by ϕ the natural action of G on $M \times G$ by left translations and
let $p_1 : M \times G \to M$ and $p_2 : M \times G \to G$ be the canonical projections. We have that:

1. The form Θ on $M \times G$ with values in \mathfrak{g} defined by

$$\Theta_{(x,q)}(u,v) = \Omega(x)u - \alpha(g)v$$

induces on $M \times G$ a codimension q foliation \mathcal{G} ;

- 2. \mathcal{G} is invariant by ϕ , that is, for each $g \in G$, the diffeomorphism ϕ^g take leaves of \mathcal{G} over leaves of \mathcal{G} .
- 3. The leaves of \mathcal{G} are transversal to the orbits of ϕ .
- 4. If M' is a leaf of \mathcal{G} , the map $p_1|_{M'}: M' \to M$ is a Galoisian covering. The automorphism group of this covering is associated with a homomorphism $h: \pi_1(M) \to G$ which satisfies

 $\alpha(p) = \phi^{h(\alpha)}(p), \ \forall \alpha \in \pi_1(M) \text{ and } p \in M'$

and whose image is $h(\pi_1(M)) = \{g \in G; \phi^g(M') = M'\};$

- 5. The map $p_2|_{M'}: M' \to G$ is a submersion;
- 6. $p_1^*(\Omega)|_{M'} \equiv p_2^*(\alpha)|_{M'}$.

Consider the notations used in the Darboux-Fedida's Theorem. The group $\Gamma = h(\pi_1(M)) < G$ is called global holonomy group of \mathcal{F} with respect to G, or, for simplicity, holonomy global of \mathcal{F} . We observe that, fixed the form Ω and the Lie group G, the global holonomy group of \mathcal{F} with respect to G is well defined up to a conjugation by an element of G, because it also depends on the choice of the leaf M' of foliation \mathcal{G} . Throughout the work, if $\mathcal{F} = \mathcal{F}_{\Omega}$ is a Lie \mathfrak{g} -foliation on M and G is a connected Lie group which has \mathfrak{g} as Lie algebra, we denote by $\Gamma(\Omega, G)$ the global holonomy group of \mathcal{F} constructed according to Darboux-Fedida' Theorem, after having chosen a leaf M' of the foliation \mathcal{G} . Still using the notations of the Darboux-Fedida Theorem, consider the Galoisian covering $P_1 = p_1|_{M'} : M' \to M$, the submersion $P_2 = p_2|_{M'} : M' \to G$ and the homomorphism o $h : \pi_1(M) \to G$ associated to P_1 . We have that

- i. P_2 is equivariant by h, that is, $P_2(\alpha(x)) = h(\alpha) * P_2(x)$;
- ii. $P_1^*(\mathcal{F})$ is the foliation defined by P_2 ;

The data (M', P_2, G, h) is a *G*-development for \mathcal{F} (see [12]). If we choose another leaf M'' of \mathcal{G} different of M', we obtain another *G*-development $(M'', \tilde{P}_2, \tilde{h}, G)$ for \mathcal{F} . The two developments are related as follows: $\tilde{h} = g * h * g^{-1}$ e $\tilde{P}_2 = g * P_2$, for an element $g \in G$ such that $\phi^g(M') = \tilde{M}$. Using a development, it is easy to obtain a Lie transverse structure of model G for the foliation. Conversely, let \mathcal{F} be a foliation that admits a Lie transverse structure of model G. If $\alpha_1, \dots, \alpha_q$ are linearly independent forms of degree one in G invariant by left translations with structure constants $\{c_{ij}^k\}$, then we can use the local submersions that define \mathcal{F} to obtain smooth linearly independent global forms of degree one $\omega_1, \dots, \omega_q$ on M which satisfy $d\omega_k = \sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$. Therefore, a foliation \mathcal{F} is a Lie \mathfrak{g} -foliation if and only if \mathcal{F} admits a G-development. Moreover, \mathcal{F} is a Lie \mathfrak{g} -foliation, if and only if \mathcal{F} admits a Lie transverse structure of model G.

We say that a Lie algebra \mathfrak{g} is *compact* if it is the Lie algebra of some compact connected Lie group. The example bellow describes a known method of obtaining examples of Lie \mathfrak{g} -foliations on compact manifolds when \mathfrak{g} is compact:

Example 3. Let G be a connected compact Lie group, N a compact manifold and $h: \pi_1(N) \to G$ a homomorphism of the fundamental group of N in G. Denote by \tilde{N} the universal covering of N and consider the diagonal action of $\pi_1(N)$ on $\tilde{N} \times G$ defined by $\alpha(x,g) \mapsto ((x)\alpha, h(\alpha)^{-1}g)$. Such action is properly discontinuous and this allow us to give a smooth structure which turns $E = (\tilde{N} \times G) / \pi_1(N)$ a (compact) manifold. Moreover, we can define a foliation \mathcal{F} on E which is transverse to the fibers of E. Such foliation admits a Lie transverse structure of model G and thus is a Lie g-foliation.

The result below is contained in Fedida's work:

Fedida's Theorem ([10]). Let M be a compact manifold and G a connected Lie group with Lie algebra \mathfrak{g} . If $\mathcal{F} = \mathcal{F}_{\Omega}$ is a Lie \mathfrak{g} -foliation on M with global holonomy $\Gamma = \Gamma(\Omega, G)$, then there is a locally trivial fibration $P: M \to \overline{\Gamma} \setminus G$ such that each fiber \mathcal{F} -saturated subset of M. In particular, $\overline{\Gamma} \setminus G$ is compact.

Remark 1. When G is a simply-connected Lie group, there exists a more complete version of the Fedida's Theorem. In this case, the fibration $P: M \to \overline{\Gamma} \setminus G$ has connected fibers and \mathcal{F} induces on each fiber of P a minimal Lie \mathfrak{h} -foliation, where \mathfrak{h} is the Lie algebra of $\overline{\Gamma}$.

3 Foliations invariant under transverse actions

The Lie \mathfrak{g} -foliation in Example 2 has a special property: there is a smooth action ϕ on the ambient manifold which preserve the leaves of \mathcal{F} and whose orbits are transversal to \mathcal{F} . The same is true for the foliation \mathcal{G} defined on $M \times G$ in Darboux-Fedida's Theorem. They are part of a special class of foliations called G-*i.u.t.a.* foliations ([1]), which we describe more formally: Let G be a q-dimensional connected Lie group and \mathcal{G} a codimension q foliation on a smooth manifold E. We say that \mathcal{G} is a G-*i.u.t.a.* foliation (or *invariant under transverse action of* G) if there is a smooth locally free action ϕ of G on E whose orbits are transverse to \mathcal{G} and such that, for each $g \in G$, ϕ^g take leaves of \mathcal{G} diffeomorphically over leaves of \mathcal{G} . In [1], the authors construct a development for this class of foliations. In particular, if \mathfrak{g} is the Lie algebra of G, then a G-i.u.t.a. foliation is a Lie \mathfrak{g} -foliation. The converse is not necessarily true (see [1]).

Proposition 1 (Fedida's Theorem for G-i.u.t.a. foliations). Let E be a smooth closed manifold let and \mathcal{G} be a G-i.u.t.a. foliation on E. Let L be a leaf of \mathcal{G} and consider the subgroup $\Gamma = \{g \in G; \phi^g(L) = L\}$. There is a locally trivial fibration with connected fibers $P : E \to G/\overline{\Gamma}$ such that each fiber is a \mathcal{G} -saturated subset of E. Moreover, \mathcal{G} induces on the fiber containing L a minimal Lie \mathfrak{h} -foliation, where \mathfrak{h} is the Lie algebra of de $\overline{\Gamma}$.

Proof. Let ϕ be the *G*-action on *E* which preserves \mathcal{G} . Let $p_1 : G \times L \to G$ be the canonical projection. It follows from Theorem 1 of [1] that the map

$$\phi|_{G \times L} : G \times L \to E$$

is a Galoisian covering map and that there is a unique $P: E \to G/\overline{\Gamma}$ such that the diagram

$$\begin{array}{c} G \times L \xrightarrow{\phi|_{G \times L}} E \\ p_1 \\ \downarrow \\ G \xrightarrow{\pi} G / \overline{\Gamma} \end{array}$$
(1)

is commutative. We conclude that P is a submersion and, since E is compact, P is a locally trivial fibration. We affirm that $\overline{\Gamma} = H = \{g \in G; \phi^g(\overline{L}) = \overline{L}\}$. In fact, the closed set $P^{-1}([e])$ contains \overline{L} . Therefore, by Diagram 1, we have that $H \subset \overline{\Gamma}$. Since E is compact, if $g \in \overline{\Gamma}$, then $\phi^g(\overline{L}) \subset \overline{L}$. On the other hand, if $g \in \overline{\Gamma}$, then $g^{-1} \in \overline{\Gamma}$. Thus $\phi^{g^{-1}}(\overline{L}) \subset \overline{L}$ and $\overline{\Gamma} \subset H$. Let L_1 be a leaf of \mathcal{G} contained in $P^{-1}([e])$. There is $g \in G$ such that $\phi^g(L) = L_1$. Since $L_1 \subset P^{-1}([e])$, we have that $g \in \overline{\Gamma}$. Therefore, $L_1 \subset \phi^g(\overline{L}) = \overline{L}$, and we conclude that $P^{-1}([e]) = \overline{L}$. Since ϕ preserves \mathcal{G} , we have $\overline{L} = \phi^g(\overline{L}) = \overline{\phi^g(L)} = \overline{L_1}$. Therefore, \mathcal{G} induces on \overline{L} a minimal foliation. If H_0 is the connected component of H containing the identity, \mathcal{G} induces on \overline{L} a H_0 -i.u.t.a. foliation and thus a Lie \mathfrak{h} -foliation.

4 Lie foliations with compact Lie algebra

In this section we study the structure of Lie \mathfrak{g} -foliations when \mathfrak{g} is compact proving Theorem A.

Proof of the Theorem A. We can suppose that M is orientable, taking the orientable double covering if necessary. Let \mathcal{G} be the foliation on $E = M \times G$ constructed in Darboux-Fedida Theorem. If $p_1 : M \times G \to M$ and $p_2 : M \times G \to G$ are the canonical projections and M' is a

leaf of \mathcal{G} , we have that $p_1|_{M'}: M' \to M$ is a Galoisian covering. The automorphism group of this covering is associated with a homomorphism $h: \pi_1(M) \to G$ whose image is

$$\Gamma = \{ g \in G; \ \phi^g(M') = M' \},\$$

the global holonomy of \mathcal{F} . Moreover, the natural action ϕ on $E = M \times G$ by left tranlations turns \mathcal{G} a G-i.u.t.a. foliation. From Fedida's Theorem (for G-i.u.t.a. foliations), there is a fibration $P: E \to G/\overline{\Gamma}$ such that the fiber $N = P^{-1}([e])$ is the closure of the leaf M'. Moreover, we have that

- 1. $\overline{\Gamma} = H = \{g \in G; \phi^g(N) = N\};$
- 2. the foliation on N induced for \mathcal{G} is a minimal \mathfrak{h} -Lie foliation, where \mathfrak{h} is a Lie algebra of $\overline{\Gamma}$.

By hypothesis, $\overline{\Gamma}_0$ is Abelian and, thus, \mathcal{G} induces on N a minimal Lie \mathbb{R}^r -foliation \mathcal{T} , for some $r \in \mathbb{N}$. now we divide the proof in two cases:

First case. *H* is connected.

For our proposes, we could consider only the restriction of \mathcal{G} on N. However, we will construct a perturbation of the foliation \mathcal{G} in E. For this, we adapt a beautiful argument presented by E. Ghys in the proof of Theorem A of [11]. The homogeneous space G/H is orientable (see Prop. 5.15 of [4]). Since E is orientable, each fiber of P is orientable. Fix an orientation in N. Dado $[g] \in G/H$, we can consider an orientation in $N_g = \phi^g(N)$ induced by ϕ^g . Since H is connected, such a orientation is consistent, that is, it does not depend on the choice of $g \in [g]$. Consider E endowed with a Riemannian metric invariant by ϕ and on each fiber of P, consider the metric induced by E. Let $\omega_1, \ldots, \omega_r$ be smooth linearly independent closed forms on N which define \mathcal{T} . Given $[g] \in G/H$, consider the fiber $N_g = \phi^g(N)$ of P. We can use the diffeomorphism ϕ^g and the forms $\omega_1, \ldots, \omega_r$ to obtain smooth closed forms $\omega_{(1,q)}, \ldots, \omega_{(r,q)}$ on N_g . These forms are well defined, that is, they does not depend on the choice of $g \in [g]$. In fact, if $l \in H$ and $f = \phi^l|_N : N \to N$, then f is isotopic to identity, because H is connected conexo. Therefore, for each $i \in \{1, \ldots, r\}$, there is a function $h_i : N \to \mathbb{R}$ such that $f_*\omega_i - \omega_i = dh_i$. On the other hand, since f preserves \mathcal{T} , we have that h_i is a *basic function* for \mathcal{T} (see [20], pag 34). This implies that h_i is constant along the leaves of \mathcal{T} . Since \mathcal{T} has dense leaves, h_i is constant and thus $f_*(\omega_i) = \omega_i$ for each $i \in \{1, \ldots, r\}$, and the assertion is proved. Since ϕ preserves the leaves of \mathcal{G} , we have that $\omega_{(1,g)}, \ldots, \omega_{(r,g)}$ define the foliation induced by \mathcal{G} on N_g . From Tischler's Theorem, there are arbitrarily small harmonic closed forms of degree one u_1, \ldots, u_r on N such that

$$\alpha_1 = (\omega_1 + u_1), \dots, \alpha_r = (\omega_r + u_r)$$

are linearly independent and define a \mathbb{R}^r -Lie foliation on N having all leaves compact. Using the diffeomorphism ϕ^g and the forms u_1, \ldots, u_r we obtain De Rham cohomology classes

$$[u_{(1,g)}], \ldots, [u_{(r,g)}]$$

on N_g . Since H is connected, this can be done consistently. For each $k > k_0$ and each $i \in \{1, \ldots, r\}$, consider the only harmonic form $u_{(i,g)}$ in the cohomology class $[u_{(i,g)}]$. For each $g \in G$, we have that

$$\phi_*^g(\omega_i + u_i) = \phi_*^g(\omega_i + u_{(i,e)}) = (\omega_{(i,g)} + u_{(i,g)}).$$
(2)

In fact, $\phi_*^g(\omega_i) = \omega_{(i,g)}$ and ϕ^g takes the cohomology class of $u_{(i,e)}$ in the cohomology class of $u_{(i,g)}$. But there is a only harmonic representative in the class $[u_{(i,g)}]$ and ϕ^g is a positive isometry. Therefore, $\phi_*^g(u_{(i,e)}) = u_{(i,g)}$ and assertion is proved. The forms

$$\omega_{(1,g)} := \left(\omega_{(1,g)} + u_{(1,g)}\right), \dots, \omega_{(r,g)} := \left(\omega_{(r,g)} + u_{(r,g)}\right)$$

define on N_g a Lie \mathbb{R}^r -foliation having all leaves compact. Since ϕ acts transitively on G/H, we have a new foliation $\tilde{\mathcal{G}}$ on E satisfying:

- In the fiber N_g , the foliation $\tilde{\mathcal{G}}$ é defined by the closed forms $(\omega_{(i,g)} + u_{(i,g)})$, para $i = 1, \ldots, r$;
- $\tilde{\mathcal{G}}$ is invariant by ϕ and has all leaves compact.

Since E is compact, if u_1, \ldots, u_r are sufficiently small, the foliation $\tilde{\mathcal{G}}$ remains transversal to ϕ . If M'' is a leaf of $\tilde{\mathcal{G}}$, we have that

$$p_1|_{M''}: M'' \to M$$

is a finite Galoisian covering and the group

$$\tilde{\Gamma} = \{ g \in G; \ \phi^g(M'') = M'' \}$$

is finite. Moreover, since N is compact, if u_1, \ldots, u_r are sufficiently small, we have that

$$p_2|_{M''}: M'' \to G$$

is a submersion and, since M'' is compact, $p_2|_{M''}$ is a locally trivial fibration . The diagram

$$\begin{array}{c}
M'' \stackrel{p_1|_{M''}}{\longrightarrow} M \\
 p_2|_{M''} \downarrow \\
G
\end{array} \tag{3}$$

provides a development to a Lie \mathfrak{g} -foliation $\tilde{\mathcal{F}}$ on M having all leaves compact. Taking the forms u_1, \ldots, u_r sufficiently small, the foliation $\tilde{\mathcal{F}}$ is C^0 -close to \mathcal{F} .

Second case. H is not connected.

In this case, we construct the following structures:

1. Two finite coverings maps $\xi : \tilde{E} \to E, \eta : \tilde{M} \to M$ and a locally trivial fibration $\tilde{p}_1 : \tilde{E} \to \tilde{M}$ such that the diagram

$$\begin{array}{c} \tilde{E} \xrightarrow{\tilde{p}_1} \tilde{M} \\ \xi & & & & \\ \xi & & & & \\ \xi & & & & \\ E \xrightarrow{p_1} M \end{array}$$

is commutative;

- 2. A free action $\tilde{\phi}$ of G on \tilde{E} such that
 - $\xi \circ \tilde{\phi}^g \equiv \phi^g \circ \xi$ for all $g \in G$;

- $\tilde{p}_1(x) = \tilde{p}_1(y)$ if and only if x and y are in the same orbit of $\tilde{\phi}$;
- $\tilde{\phi}$ turns \tilde{p}_1 a *G*-principal bundle;

3. A locally trivial fibration $\tilde{P}: \tilde{E} \to G/H_0$ with connected fibers such that

- the restriction of ξ to a fiber of \tilde{P} defines a diffeomorphism with a fiber of P;
- \tilde{P} possesses a fiber \tilde{N} which projects by ξ over N, for which

$$\{g \in G; \ \tilde{\phi}^g(\tilde{N}) = \tilde{N}\} = H_0 = \overline{\Gamma}_0.$$

After obtaining such structures, we can consider the foliations $\xi^*(\mathcal{G})$ and $\eta^*(\mathcal{F})$ and proceed as in the previous case. We go on to describe such structures. Fix the leaf $N = P^{-1}([e])$ of \mathcal{G} . the action

$$\begin{array}{rcl} H_0 \times (G \times N) & \to & (G \times N) \\ (h_0, (g, p))) & \mapsto & \left(gh_0^{-1}, \phi^{h_0}(p)\right) \end{array}$$

is proper and free, thus, there is a unique smooth structure of $H_0 \setminus (G \times N)$ which turs the canonical projection

$$\beta: (G \times N) \to H_0 \setminus (G \times N)$$

a principal bundle. Consider the manifold $\tilde{E} := H_0 \setminus (G \times N)$ endowed with this structure. The action

$$\varphi : (H_0 \setminus H) \times \tilde{E} \to \tilde{E}$$
$$((h), [(g, p)]) \mapsto \left[\left(gh^{-1}, \phi^h(p) \right) \right]$$

is well-defined. In fact, if $\tilde{h} = h_0 h$, $\tilde{g} = g h_0^{-1}$ and $\tilde{p} = \phi^{h_0}(p)$, then

$$\tilde{g}\tilde{h}^{-1} = gh^{-1}\underbrace{(hh_0^{-1}h^{-1})}_{l_0^{-1}\in H_0}h_0^{-1} = gh^{-1}(h_0l_0)^{-1}$$

and

$$\phi^{\tilde{h}}(\tilde{p}) = \phi^{h_0 h h_0}(p) = \phi^{h_0 h h_0 h^{-1} h}(p) = \phi^{h_0 l_0} \circ \phi^{h}(p).$$

Therefore, φ is well-defined. Moreover, φ is free. Since $H_0 \setminus H$ is a finite group, there is a unique smooth structure for $\varphi \setminus \tilde{E}$ which turns

$$q: \tilde{E} \to \varphi \setminus \tilde{E}$$

a finite Galoisian map. The map

$$\begin{array}{rccc} \xi : E & \to & E \\ [(g,p)] & \mapsto & \phi^g(p) \end{array}$$

is well-defined and a submersion, because $\phi = \xi \circ \beta : G \times N \to E$. There is a unique map $f : \varphi \setminus \tilde{E} \to E$ which which make the diagram below to commute.



and, thus, f is smooth. Moreover, f is bijective and this implies that $\xi: \tilde{E} \to E$ is a finite covering map. Consider the free action

$$\begin{array}{rccc} H_0 \times N & \to & N \\ (h_0, p) & \mapsto & \phi^{h_0}(p) \end{array}$$

and the unique smooth structure of $H_0 \setminus N$ which turns the canonical projection

$$i: N \to H_0 \setminus N$$

a principal bundle. Write $\tilde{M} := H_0 \setminus N$. The action

$$\begin{split} \psi : (H_0 \setminus H) \times \tilde{M} &\to \tilde{M} \\ ((h), [p]) &\mapsto & \left[\phi^h(p) \right] \end{split}$$

is well-defined and free. Consider the unique structure of $\psi \setminus \tilde{M}$ which turns the projection

$$j: \tilde{M} \to \psi \setminus \tilde{M}$$

a fibration. Such a projection is a finite covering map, because $H_0 \setminus H$ is finite. It follows from definition of ϕ that the map

$$\eta: H_0 \setminus N = M \to M$$
$$[p = (x, g)] \mapsto x$$

is well defined. Moreover, η is smooth, because

$$p_1|_N = \eta \circ i : N \to M.$$

There is an unique smooth map $\theta:\psi\setminus \tilde{M}\to M$ which make the diagram



to commute. This implies that θ is smooth. Moreover, θ is bijective and then η is a finite covering map. The map

$$\tilde{p}_1 : \tilde{E} \to \tilde{M} [(p,g)] \mapsto [p]$$

is a submersion, because the diagram



is commutative. Since \tilde{E} is compact, we have that \tilde{p}_1 is a locally trivial fibration which make the diagram of item 1 to commute and we conclude item 1. By construction, the action

$$\begin{array}{lll} \tilde{\phi}:G\times\tilde{E} & \to & \tilde{E} \\ (l,[(g,p)]) & \mapsto & [(lg,p)] \end{array}$$

satisfies the conditions in item 2. The map

$$\tilde{P}: \tilde{E} \to G/H_0$$

 $[(g,p)] \mapsto [g]$

is well-defined and a submersion, because the diagram

$$\begin{array}{c} G \times N \xrightarrow{\beta} \tilde{E} \\ \downarrow & \downarrow^{\tilde{P}} \\ G \longrightarrow G/H_0 \end{array}$$

is commutative. Since \tilde{E} is compact, \tilde{P} is a locally trivial fibration. We have that

$$\tilde{N} = \{ [(h_0, p)]; h_0 \in H_0 \text{ and } p \in N \} = \tilde{P}^{-1}([e]) = \beta(H_0 \times N)$$

is a connected set, because N is connected. Moreover, $\xi|_{\tilde{N}} : \tilde{N} \to N$ is a diffeomorphism and

$$\{g \in G; \ \tilde{\phi}^g(\tilde{N}) = \tilde{N}\} = H_0,$$

and this completes the proof.

Proof of Corollary 1. In [6], Caron and Carrière prove that a minimal one-dimensional Lie foliation on a closed manifold M^n has Abelian global holonomy (see [5] for a detailed proof). Since the foliation is minimal, the global holonomy is dense, and thus, such foliation is a Lie \mathbb{R}^{n-1} -foliation. If \mathcal{F}_{Ω} is a (not necessarily minimal) one-dimensional Lie \mathfrak{g} -foliation on M and G is the simply-connected Lie group having \mathfrak{g} as Lie algebra, then, by Fedida Theorem, $\overline{\Gamma(\Omega, G)}$ has Abelian Lie algebra. Since \mathfrak{g} é semisimple and compact, G is compact and we obtain Corollary 1 from Theorem 1.

Theorem 1 was obtained from a perturbation of the foliation G-i.u.t.a. on the compact manifold $E = M \times G$ constructed from original Lie foliation on M. We can use the constructions developed during the proof of Theorem 1 to obtain a fact about G-i.u.t.a. foliations:

Proposition 2. Let E be a smooth orientable closed manifold and \mathcal{G} a G-i.u.t.a. foliation on E, where G is a compact Lie group. Using the notations of Proposition 1, suppose that $\overline{\Gamma}_0$ is Abelian. Then, up to a finite equivariant covering, \mathcal{G} can be C^0 -approximated by a G-i.u.t.a. foliation having all leaves compact.

Proof. We can construct \tilde{E} and $\tilde{\phi}$ as in the proof of Theorem 1. Taking a Riemannian metric on \tilde{E} invariant by the action of G, we can proceed as in the first case in the proof of Theorem 1 and to obtain a G-i.u.t.a. foliation on \tilde{E} having all leaves compact.

5 Lie foliations on manifolds with amenable fundamental group

Let A be a (discrete) group and let $\mathcal{B}(A)$ be the vector space of bounded real functions on A endowed with the sup norm. A mean on A is a left-invariant linear functional $\eta : \mathcal{B}(A) \to \mathbb{R}$ satisfying $\eta(f) \geq 0$ and $\eta(1) = 1$. The class of amenable groups is closed under operations of taking subgroups, extensions, quotient groups and direct limits. Solvable groups are amenable. In contrast, a group having a non-cyclic free subgroup is not amenable. For more details, see [13] and [8]. By Theorem 4.1 of [16], a finitely generated group having subexponential growth is amenable.

Lemma 1. Let G be a Lie group and $\Gamma < G$ a subgroup of G. If Q is a subgroup of Γ such that $[\Gamma:Q] < \infty$, then $[\overline{\Gamma}:\overline{Q}] \leq [\Gamma:Q] < \infty$.

Proof. Write $\Gamma/Q = \{(y_1), \ldots, (y_l)\}$ and suppose, by contradiction, that $[\overline{\Gamma} : \overline{Q}] > [\Gamma : Q] = l$. In this case, there are distinct elements $[g_1], \ldots, [g_l], [g_{l+1}]$ in $\overline{\Gamma}/\overline{Q}$. For each $k \in \{1, \ldots, l+1\}$, there is a sequence x_n^k in Γ which converges to g_k . Taking subsequences if necessary, we can suppose that $(x_n^k) = (y_{j_k}) \in \Gamma/Q$. Therefore, for each $k \in \{1, \ldots, l+1\}$, there is a sequence q_n^k in Q such that $x_n^k = y_{j_k} * q_n^k$ and, thus, $(y_{j_k}^{-1} * g_k) \in \overline{Q}$. There are $k_1, k_2 \in \{1, \ldots, l+1\}$ such that $k_1 \neq k_2$ and $y_{j_{k_1}} = y_{j_{k_2}} = y$. Then

$$(y^{-1} * g_{k_1})^{-1} * (y^{-1} * g_{k_2}) = g_{k_1}^{-1} * g_{k_2} \in \overline{Q}.$$

This is a contradiction, because $[g_{k_1}] \neq [g_{k_2}]$.

Lemma 2. Let G be a connected Lie group. If a finitely generated subgroup $\Gamma < G$ is amenable, then there is a solvable subgroup $Q < \Gamma$ such that $[\Gamma : Q] < \infty$.

Proof. Let $Ad : G \to GL(\mathfrak{g}, \mathbb{R})$ be the adjoint homomorphism. Since Γ is finitely generated, either $Ad(\Gamma)$ has a non-cyclic free subgroup or $Ad(\Gamma)$ is virtually solvable (Tits Alternative [24]). Since $Ad(\Gamma)$ is an amenable group, the first option does not occur. Therefore, there is a solvable subgroup E of $Ad(\Gamma)$ such that $[Ad(\Gamma) : E] < \infty$. Writing $T = Ad^{-1}(E)$, we have that

$$\frac{T}{\ker(Ad|_T)} \simeq E$$

Since $\ker(Ad|_T) < \ker Ad$ is Abelian (G is connected), we conclude that T is solvable. Therefore, $Q = \Gamma \cap T$ is a solvable group. Since $[Ad(\Gamma) : E] < \infty$, we have that $[\Gamma : Q] < \infty$ and the proof is concluded.

Proposition 3. Let M be a smooth closed manifold with amenable fundamental group and let $\mathcal{F} = \mathcal{F}_{\Omega}$ be a Lie \mathfrak{g} -foliation on M. Let G be a connected Lie group having \mathfrak{g} as Lie algebra and let $\Gamma = \Gamma(\Omega, G)$ be the global holonomy of \mathcal{F} . The connected component of $\overline{\Gamma}$ containing the identity is a solvable Lie subgroup of G. In particular, if G is compact, then the connected component of $\overline{\Gamma}$ containing the identity is Abelian.

Proof. Since $\pi_1(M)$ is amenable, Γ is amenable. Denote by $\overline{\Gamma}_0$ the connected component of $\overline{\Gamma}$ containing the identity. Let Q be the solvable subgroup of Γ given by Lemma 2. Taking the closure of the derived series of Q, we conclude that \overline{Q} is solvable. Therefore $\overline{\Gamma}_0 \cap \overline{Q}$ is a closed subgroup of $\overline{\Gamma}_0$ and a solvable group. By Lemma 1, $[\overline{\Gamma}, \overline{Q}] < \infty$ and this implies that $[\overline{\Gamma}_0, \overline{\Gamma}_0 \cap \overline{Q}] < \infty$. Since $\overline{\Gamma}_0$ is connected, we conclude that $\overline{\Gamma}_0 = \overline{\Gamma}_0 \cap \overline{Q}$ is solvable. Then, $\overline{\Gamma}_0$ is a connected solvable Lie group. If G is compact, then $\overline{\Gamma}_0$ is compact and, thus, Abelian.

Proof of Theorem 2. It follows from Theorem 1 and Proposition 3.

Using the notations of Proposition 1, we obtain

Proposition 4. Let E be a smooth orientable closed manifold with amenable fundamental group and let G a compact Lie group. If G is a G-i.u.t.a. foliation on E then, up to a finite equivariant covering, \mathcal{G} can be C^0 -approximated by a G-i.u.t.a. foliation having all leaves compact.

Proof. It follows from Proposition 2 and Proposition 3.

6 Subexponential growth and Lie foliations

In this section, we study Lie foliations on manifolds with fundamental growth of subexponential growth.

Lemma 3. Let G be a connected Lie group and $\Gamma < G$ a finitely generated subgroup of subexponential growth. There is a nilpotent subgroup $T < \Gamma$ such that $[\overline{\Gamma} : \overline{T}] < \infty$.

Proof. Since Γ has subexponential growth, Γ is amenable. Let $Q < \Gamma$ be the solvable subgroup given by Lemma 2. Since $[\Gamma : Q] < \infty$, Q is a finitely generated solvable group of subexponential growth. By Milnor-Wolf Theorem ([19], [26]) Q is virtually nilpotent, that is, there is a nilpotent subgroup T of Q $[Q : T] < \infty$. We have $[\Gamma : T] < \infty$ and, by Lemma 1, $[\overline{\Gamma} : \overline{T}] < \infty$.

A Lie group (not necessarily connected) is *compactly generated* if G is generated by some symmetric and compact neighborhood W of the identity of G. If W is such a neighborhood, we write

$$W^n = \underbrace{W \cdot W \cdots W \cdot W}_{n-\text{times}}.$$

Using the Haar measure of G, we can consider the volume growth of G and this does not depend on the choice of W. In particular, we say that G has polynomial volume growth if there are constants c > 0 and $d \in \mathbb{N}^*$ such that $\mu(W^n) \leq cn^d$, $\forall n \in \mathbb{N}$. For more details, see [14].

Proposition 5. Let M be a smooth closed manifold with fundamental group of subexponential growth. If M admits a Lie \mathfrak{g} -foliation and G is a connected Lie group having \mathfrak{g} as Lie algebra, then G has polynomial volume growth.

Proof. Let $\mathcal{F} = \mathcal{F}_{\Omega}$ be such foliation and let $\Gamma = \Gamma(\Omega, \mathcal{F})$ be the global holonomy of \mathcal{F} . Using Fedida Fibration Theorem, we conclude that $G/\overline{\Gamma}$ is compact. By [18], $\overline{\Gamma}$ is compactly generated. It follows from Theorem I.4 of [14] that G and $\overline{\Gamma}$ has the same volume growth. Let T be the nilpotent subgroup of Γ given by Lemma 3. Since $[\overline{\Gamma}:\overline{T}] < \infty$, \overline{T} is compactly generated and has the same volume growth of of $\overline{\Gamma}$ and, thus, of G. Taking the closure of the lower central series of T, we obtain a central series for \overline{T} and, thus \overline{T} is a nilpotent group. By Theorem II.4 of [14], \overline{T} (and thus G) has polynomial volume growth.

Given a Lie algebra \mathfrak{g} , there is a solvable ideal of \mathfrak{g} that contains all the solvable ideals of \mathfrak{g} . This ideal is called *radical of* \mathfrak{g} and we will denote it by $\mathfrak{r}(\mathfrak{g})$. We have that \mathfrak{g} is solvable if and only if $\mathfrak{r}(\mathfrak{g}) = \mathfrak{g}$; \mathfrak{g} is *semisimple* if and only if $\mathfrak{r}(\mathfrak{g}) = 0$, that is, if and only if \mathfrak{g} has no solvable ideal. If G is a connected Lie group having \mathfrak{g} as Lie algebra, then the *radical of* G is the Lie subgroup of G having $\mathfrak{r}(\mathfrak{g})$ as Lie algebra.

Proposition 6. Let M be a smooth closed manifold with fundamental group of subexponential growth. If M admits a Lie \mathfrak{g} -foliation, and G is a connected Lie group having \mathfrak{g} as Lie algebra, then G/R is compact, where R is the radical of G.

Proof. By Proposition 5, G has polynomial volume growth. By Theorem 6.39 of [21], the adjoint representation of \mathfrak{g} has only pure imaginary eigenvalues (see (6.35) of [21]). It follows from 6.32 of [21] that G/R is compact.

In order to obtain Theorem 3 in its most general form, we use the Levi-Malcev decomposition for Lie algebras [25]:

Levi-Malcev's decomposition. Let \mathfrak{g} be a Lie algebra and $\mathfrak{r}(\mathfrak{g}) \subset \mathfrak{g}$ its radical. There is a semisimple subalgebra $\mathfrak{s} \subset \mathfrak{g}$ such that \mathfrak{g} is a direct sum (as vector space) of \mathfrak{s} and $\mathfrak{r}(\mathfrak{g})$.

A Lie subalgebra $\mathfrak{s} < \mathfrak{g}$ given by Levi-Malvec decomposition is called it Levi factor of \mathfrak{g} . Let G be a connected Lie group having \mathfrak{g} as Lie algebra. If S < G is the Lie subgroup of G that has \mathfrak{s} as Lie algebra, then S is called of *Levi factor of G*.

Lemma 4. Let \mathcal{F} be a Lie \mathfrak{g} -foliation on a smooth manifold M and $\mathfrak{s} < \mathfrak{g}$ a Levi factor of \mathfrak{g} . There is a Lie \mathfrak{s} -foliation \mathcal{S} on M. Each leaf of \mathcal{S} is a \mathcal{F} -saturated subset of M.

Proof. Consider a Levi-Malcev decomposition $\mathfrak{g} = \mathfrak{s} + \mathfrak{r}(\mathfrak{g})$. Take a basis $\{X_1, \ldots, X_s\}$ of \mathfrak{s} and a basis $\{X_{s+1}, \ldots, X_q\}$ of $\mathfrak{r}(\mathfrak{g})$. Let c_{ij}^k be the structure constants of $\beta = \{X_1, \ldots, X_s, X_{s+1}, \ldots, X_q\}$. Since $\mathfrak{r}(\mathfrak{g})$ is an ideal of \mathfrak{g} , we have that

if $1 \le k \le s$ and either *i* or *j* are are larger that *s*, then $c_{ij}^k = 0.$ (4)

Since \mathcal{F} is a Lie g-foliation, there are linearly independent forms of degree one

$$\omega_1,\ldots,\omega_s,\omega_{s+1},\ldots,\omega_q$$

on M which are tangent to \mathcal{F} and satisfy the relations involving $\{c_{ij}^k\}$. By (4), $\omega_1, \ldots, \omega_s$ define a Lie \mathfrak{s} -foliation \mathcal{S} on M.

Proof of Theorem 3. The first item is the Proposition 5. Let \mathfrak{s} be a Levi factor of \mathfrak{g} . By Lemma 4, there is a Lie \mathfrak{s} -foliation on M. If S is a connected Lie group having \mathfrak{s} as Lie algebra, then S is semisimple. By Proposition 6, S is compact. By Theorem 1, there is a finite covering of M which fibers over S and, thus, we have item 2.

7 Extension to transversely homogeneous foliations

Proof of Theorem 4. Let \mathfrak{g} and \mathfrak{k} be the Lie algebras of G and K, respectively. By Theorem 2 and Corollary 3.2 of [2], there are n smooth forms of degree one $\omega_1, \ldots, \omega_q, \omega_{q+1}, \ldots, \omega_n$ on M such that

- i. $\omega_1, \ldots, \omega_q$ are linearly independent at each point of M;
- ii. $d\omega_k = -\sum_{i < j} c_{ij}^k \omega_i \wedge \omega_j$, where $\{c_{ij}^k\}$ are the structure constants of a basis $\beta = \{X_1, \ldots, X_n\}$ of \mathfrak{g} such that X_{q+1}, \ldots, X_n is a basis of \mathfrak{k} .

Consider the form $\Omega = \sum_{l=1}^{n} \omega_l X_l$ of degree one on M with values in \mathfrak{g} and let α be the Maurer-Cartan form of G. Consider the form Θ on $M \times G$ with values in \mathfrak{g} given by

$$\Theta_{(x,q)}(u,v) = \Omega(x)u - \alpha(g)v,$$

as in Darboux-Fedida Theorem. The form Ω is singular in M, however, the form Θ has rank n in each point of $M \times G$. Therefore, as in Darboux-Fedida's Theorem, Θ induces a foliation G-i.u.t.a. \mathcal{G} on $M \times G$ whose dimension is the dimension of M. Let $p_2 : M \times G \to G$ and $\pi : G \to G/K$ be the canonical projections. By (i), if M' is a leaf of \mathcal{G} , the map $\pi \circ p_2|_{M'} : M' \to G/K$ is a submersion. Then, we can follow the proof of Theorem 1 and use the Theorem 2 to obtain Theorem 4.

Proof of Corollary 2. By Corollary 2.1 of [3], the fundamental group of M has polynomial volume growth. Then, the first statement follows from Theorem 4. In the case q = 1, suppose that the global holonomy Γ of \mathcal{F} is not finite. Since $\overline{\Gamma}_0$ is Abelian, then $\overline{\Gamma}_0$ contains (a subgroup of SO(3) isomorphic to) SO(2). We have that

$$SO(3) \supseteq \overline{\Gamma} \supset \overline{\Gamma}_0 \supset SO(2).$$

Therefore, either $\overline{\Gamma} = SO(2)$ or $\overline{\Gamma} = O(2)$. It follows from construction in Theorem 1 that, in the first case, the finite covering mentioned is the identity and, in the second case, it has two folds.

8 A remark about Riemannian foliations

This section is concerned with the study of Riemannian foliations and contains results already found in [22]. Let M be a smooth manifold and \mathcal{F} a smooth foliation on M of codimension q. Denote by $\mathcal{X}(\mathcal{F})$ the set of smooth vector fields on M which are tangent to \mathcal{F} . We say that a vector field X on M is *foliated with respect to* \mathcal{F} if

$$[X, Y] \in \mathcal{X}(\mathcal{F}), \text{ for all } Y \in \mathcal{X}(\mathcal{F}).$$

If $N\mathcal{F}$ is the normal bundle of \mathcal{F} and X is a foliated vector field, we can project X over $N\mathcal{F}$, obtaining a vector field \overline{X} tangent to $N\mathcal{F}$. We say that \overline{X} is the *transversal vector field* associated with X. We say that a foliation \mathcal{F} is *transversely parallelizable* if its normal bundle admits a trivialization by q linearly independent transversal foliated vector fields. Every Lie foliation is a transversely parallelizable foliation, however, the reciprocal is not necessarily true.

We say that a Riemannian metric g on M is bundle-like with respect to a foliation \mathcal{F} if, given an open $U \subset M$ and foliated vector fields X, Y on U orthogonal to the leaves of \mathcal{F} , the function g(X, Y) is constant along the leaves of $\mathcal{F}|_U$. For all details, see [20]. We say that \mathcal{F} is a *Riemannian foliation* if there exists a bundle-like metric with respect to \mathcal{F} . A deep result due to Molino describes the structure of transversely parallelizable and Riemannian foliations and proves that the study of such foliations is closely related to the study of Lie foliations:

Molino's structure theorem ([12, 20]). Let M be a connected closed manifold and let \mathcal{F} be a foliation on M.

1. If \mathcal{F} is transversally parallelizable, then the closures of the leaves of \mathcal{F} are the fibers of a locally trivial basic fibration $\pi : M \to B$. The restriction of \mathcal{F} to each fiber defines a Lie g-foliation with dense leaves. The Lie algebra g is an algebraic invariant of \mathcal{F} called structural Lie algebra of \mathcal{F} .

- 2. If \mathcal{F} is a transversally orientable Riemannian foliation and $\tau : M' \to M$ is the principal SO(q)-bundle of orthonormal frames transverse to \mathcal{F} , then \mathcal{F} lifts to a transversally parallelizable foliation \mathcal{F}' on M' such that
 - dim $\mathcal{F}' = \dim \mathcal{F};$
 - \mathcal{F}' is invariant by the action of SO(q).

The structural Lie algebra of \mathcal{F}' is an invariant of \mathcal{F} called structural Lie algebra of \mathcal{F} .

Based on Molino Theorem, we obtain the following:

Proposition 7. Let M be a smooth closed manifold and let \mathcal{F} be a Riemannian foliation on M. If the fundamental group of M is amenable, then the structural Lie algebra of \mathcal{F} is solvable.

Proof. Let

$$\cdots \longrightarrow \pi_2(M) \xrightarrow{\phi} \pi_1(SO(q)) \xrightarrow{\varphi} \pi_1(M') \xrightarrow{\psi} \pi_1(M) \longrightarrow \cdots$$

be the homotopy exact sequence associated with the principal SO(q)-bundle of orthonormal frames transverse to \mathcal{F} . We have that ker $\psi = \text{Im}\varphi$ is Abelian and thus amenable. Moreover, Im ψ is a subgroup of $\pi_1(M)$ and thus it is amenable. Since

$$\pi_1(M')/\ker\psi\simeq\operatorname{Im}\psi,$$

we conclude that $\pi_1(M')$ is amenable, because amenable groups are closed to extensions. Let $\pi : M' \to B$ the locally trivial basic fibration of \mathcal{F}' and let $N = \pi^{-1}(y)$ be a fibre of π . From homotopy exact sequence

$$\cdots \longrightarrow \pi_2(B) \xrightarrow{\phi} \pi_1(N) \xrightarrow{\varphi} \pi_1(M') \xrightarrow{\psi} \pi_1(B) \longrightarrow \cdots$$

of π , we have that $\operatorname{Im} \varphi = \ker \psi$ is a subgroup of $\pi_1(M')$, so it is amenable. Moreover, $\operatorname{Im} \phi = \ker \varphi$ is an Abelian group, so it is amenable. Since

$$\pi_1(N)/\ker\varphi\simeq\operatorname{Im}\varphi,$$

we conclude that $\pi_1(N)$ is amenable. From Molino Theorem, \mathcal{F} induces a minimal Lie \mathfrak{g} -foliation on N. Let G be a connected lie group having \mathfrak{g} as Lie algebra. By Fedida theorem, the global holonomy of \mathcal{F} is dense in G. It follows from Proposition 3 that the structural Lie algebra of \mathcal{F} is solvable.

References

- A. Behague and B. Scárdua Foliations invariant under Lie group transverse actions. Monatsh. Math. 153 (2008), 4, 295–308.
- [2] R. A. Blumenthal Transversely homogeneous foliations. Annales de l'Institut Fourier, 29 (1979), 143–158.
- [3] R. A. Blumenthal Riemannian homogeneous foliations without holonomy. Nagoya Math. J. 83 (1981), 197–201.

- [4] T. Brocker and T. t. Dieck: Representations of compact Lie groups, Springer-Verlag, New York, 1985.
- [5] P. Caron Flots transversalement de Lie. Thèse de 3ème cycle, Université de Lille (1980).
- [6] P. Caron e Y. Carrière Flots transversalement de Lie ℝⁿ, flots de Lie minimaux. C. R. Ac. Sci. Paris, 280 (9) (1980), 477-478.
- [7] L. Corwin e F.P Greenleaf Character formulas and spectra of compact nilmanifolds. J. Funct. Anal. 21 (1976), 123–154.
- [8] M. Day: Amenable semigroups. Ill. J. Math 1 (1957), 509–544.
- [9] J. J. Duistemaat e J. A. C. Kolk: *Lie Groups*, Springer-Verlag, Berlin Heidelberg New York, 2000.
- [10] E. Fedida: Feuilletages du plan Feuilleteges de Lie. thèse Strasbourg, 1973, Lecture Notes in Mathematics, 352, 183–195. 205-228 (1971).
- [11] E. Ghys: Feuilletages Riemanniens sur les variétés simplement connexes. Ann. Inst. Fourier, Grenoble 34, 4 (1984), 203–223.
- [12] C. Godbillon: Feuilletages, Études geométriques, Birkhäuser. MR1120547, Zbl 0724.58002.
- [13] F. P. Greenleaf: Invariant means on topological groups and their applications, American Book Company, 1969.
- [14] Y. Guivarc'h: Croissance polynomiale, Bulletin de la société mathématique de France, 101, 4, 1973.
- [15] P. de la Harpe: Topics in geometric group theory, University of Chicago Press, 2000.
- [16] M. W. Hirsch and W. P. Thurston: Foliated bundles, invariant measures and flat manifolds. Ann. of Math, 101 (1975), 369–390.
- [17] S. Kobayashi and K. Nomizu: Foundations of Differential Geometry, Interscience Publishers, 1963.
- [18] A. M. Macbeath and S. Swierczkowski: On the set of generators of a subgroup. Indag. Math, 21, (1959) 280-281.
- [19] J. Milnor: Growth of finitely generated solvable groups. J. Differential Geometry, 2 (1968) 447-449.
- [20] P. Molino: *Riemannian Foliations*. Progress in mathematics, vol 73, Birkhäuser, Boston, 1988.
- [21] A. T. Paterson: *Amenability*. Math. Surveys and Monographs, vol 29, Amer. Math. Society, Providence, RI.
- [22] Tavares and B. Scardua: Lie and Riemannian foliations on manifolds with amenable fundamental group. Pre-print Sept/2009, submitted.
- [23] D. Tischler: On fibering certain foliated manifolds over S^1 . Topology, 9 (1970) 153–154.

- [24] J. Tits: Free subgroups in linear groups. J. Algebra, 20 (1972) 250–270.
- [25] V.S. Varadarajan: Lie Groups, Lie Algebras, and Their Representations, Springer-Verlag, New York, 1974, 1984.
- [26] J. A. Wolf: Growth of finitely generated solvable groups and curvature of Riemannian manifolds. J. Differential Geometry, 2 (1968) 421-446.

Marcelo Tavares: marcelotrls@gmail.com Instituto de Matemática - Universidade Federal do Rio de Janeiro Rio de Janeiro - RJ, Caixa Postal 68530 21.945-970 Rio de Janeiro-RJ BRAZIL