

Groups with regular free length function in Λ

Olga Kharlampovich* Alexei Myasnikov† Denis Serbin

October 30, 2009

Abstract

A group G has a regular free Lyndon length function in an ordered abelian group Λ if and only if it has a free isometric action on a Λ -tree such that all branch points are in the same orbit as well as the base point. In this paper we prove that every finitely presented group G with a regular free Lyndon length function in Λ has an index two subgroup with free Lyndon length function in \mathbb{Z}^k ordered lexicographically for an appropriate $k \in \mathbb{N}$.

1 Introduction

This is the third paper in a series where we investigate non-Archimedean group actions, length functions and infinite words. There are two classical approaches to the fundamental groups of graphs of groups: one uses group actions on trees, and another one - length functions.

Lyndon introduced groups equipped with length functions with values in an ordered abelian group Λ as a tool to carry over Nielsen cancellation theory from free groups to a much more general setting [23] (see also [24]). Some partial results (for $\Lambda = \mathbb{Z}$ or $\Lambda = \mathbb{R}$) were obtained in [15, 16, 14, 29, 2]. In [5] Chiswell described a crucial construction that shows that a group with a length function with values in \mathbb{R} comes naturally from some action of the group on an \mathbb{R} -tree, and vice versa. It was realized later that a similar construction holds for arbitrary group with a free Lyndon length function with values in Λ [28]. Thus free group actions and free Lyndon length functions are two equivalent languages describing the same objects. We refer to the book [6] for a detailed discussion on the subject.

One of the major events in combinatorial group theory in 1970's was the development of Bass-Serre theory. We refer to the book [31] where Serre laid down fundamentals of the theory of groups acting freely on simplicial trees. In 1980's Morgan and Shalen introduced group actions on Λ -trees for an arbitrary ordered abelian group Λ [28]. In particular, they studied actions on \mathbb{R} -trees in relation with Thurston's Geometrization Program.

Alperin and Bass [1] developed the initial framework of the theory of group actions on Λ -trees and stated the fundamental research goals: find the group theoretic information carried by an action (by isometries) on a Λ -tree; generalize Bass-Serre theory to actions on arbitrary Λ -trees.

A joint effort of several researchers culminated in a description of finitely generated groups acting freely on \mathbb{R} -trees [4, 9], which is now known as Rips' theorem: a finitely generated group acts freely on an \mathbb{R} -tree if and only if it is a free product of free abelian groups and surface groups (with exception of non-orientable groups of genus 1, 2, and 3). The key ingredient of this theory is the so-called "Rips machine", the idea of which comes from Makanin's algorithm for solving equations in free groups [25]. The Rips machines appear in applications as a general tool that takes a sequence of isometric actions of a group G on some "negatively curved spaces" and produces an isometric action of G on an \mathbb{R} -tree in the Gromov-Hausdorff limit. Free actions on \mathbb{R} -trees cover all Archimedean

*Supported by NSERC grant

†Supported by NSERC grant

actions, since every group acting freely on a Λ -tree for an Archimedean ordered abelian group Λ acts freely also on an \mathbb{R} -tree.

The case of non-Archimedean free actions is open. In [3] Bass studied finitely generated groups acting freely on $(\Lambda \oplus \mathbb{Z})$ -trees with respect to the lexicographic order on $\Lambda \oplus \mathbb{Z}$. Recently, Guirardel (see [12]) studied finitely generated groups acting freely on an \mathbb{R}^n -tree (with the lexicographic order). In [20] we described the algebraic structure of finitely generated groups acting freely and regularly on \mathbb{Z}^n trees in terms of HNN-extensions of a very particular type. In [21] we proved that a finitely generated group acting freely on a \mathbb{Z}^n -tree is a subgroup of a group acting freely and regularly on a \mathbb{Z}^m -tree. However, the following main problem of the Alperin-Bass program remains largely intact.

Problem *Describe finitely presented (finitely generated) groups acting freely on an arbitrary Λ -tree.*

Nowadays, the geometric method of group actions seems much more in use than the length functions, even though the Nielsen theory still gives very powerful results (see the discussion in [20]). There are several reasons for it. Firstly, the Lyndon's abstract axiomatic approach to length functions is less intuitive than group actions. Secondly, the current development of the abstract Nielsen theory is incomplete and insufficient - one of the principal notions of complete actions and regular length functions were not in use until recently. Thirdly, and this is crucial, an abstract analog of the Rips machine for length functions with values in \mathbb{R} (or an arbitrary Λ) was not introduced or developed enough to be used easily in applications. Notice, that in case of $\Lambda = \mathbb{Z}$ the completeness and regularity come for free, so there was no need for axiomatic formalization, - it went mostly unnoticed. The regularity axiom appeared first in [26, 27] as a tool to deal with length functions in \mathbb{Z}^n (with respect to the lexicographic order). On the other hand, an analog of the Rips machine for $\Lambda = \mathbb{Z}$ did exist for a long time - the original Makanin-Razborov process [30] for solving equations in free groups. But it was not recognized as such until [18, 19], where it was used systematically to get splittings of groups.

Introduction of infinite Λ -words in [27] gives another approach to group actions. Every group admitting a faithful representation by Λ words acts freely on a Λ -tree. The converse is proved in [5] Formulation of the regularity axiom is another important ingredient. This axiom ensures that the group is complete with respect to the Gromov's inner product, i.e., the Gromov's product of two elements of the group can be realized as the length of a particular element of the group (not only as an abstract length). In the language of actions this means that all branch points of the group action are in one same orbit, as well as the base point. This allows one to use the Nielsen cancelation argument very similar to the case of a free group. The regularity (completeness) condition is crucial for existence of Makanin-Razborov's type processes over groups with Λ -length functions.

In this paper we will prove the following result.

Theorem 1. *Any f.p. group G with a regular free length function in an ordered abelian group Λ has an index two subgroup with free length function in \mathbb{Z}^k ordered lexicographically for an appropriate $k \in \mathbb{N}$.*

2 Length functions, actions, infinite words

Here we introduce basic definitions and notations which are to be used throughout the whole paper.

2.1 Lyndon length functions and free actions

Let G be a group and A an ordered abelian group. Then a function $l : G \rightarrow A$ is called a (*Lyndon*) *length function* on G if the following conditions hold:

$$(L1) \quad \forall g \in G : l(g) \geq 0 \text{ and } l(1) = 0;$$

$$(L2) \quad \forall g \in G : l(g) = l(g^{-1});$$

$$(L3) \quad \forall g, f, h \in G : c(g, f) > c(g, h) \rightarrow c(g, h) = c(f, h),$$

where $c(g, f) = \frac{1}{2}(l(g) + l(f) - l(g^{-1}f))$.

Sometimes we refer to length functions with values in A as to A -length functions. Observe, that the standard length function on a finitely generated group (with respect to a fixed finite generating set) is not necessary a Lyndon length function. In this paper all length functions are Lyndon's ones.

Notice that $c(g, f)$ may not be defined in A (if $l(g) + l(f) - l(g^{-1}f)$ is not divisible by 2), so in the axiom (L3) we assume that A is canonically embedded into a divisible ordered abelian group $A_{\mathbb{Q}} = A \otimes_{\mathbb{Z}} \mathbb{Q}$ (see [27] for details).

It is not difficult to derive the following two properties of length functions from the axioms (L1)-(L3):

- $\forall g, f \in G : l(gf) \leq l(g) + l(f)$;
- $\forall g, f \in G : 0 \leq c(g, f) \leq \min\{l(g), l(f)\}$.

A length function $l : G \rightarrow A$ is called *free* if it satisfies the following two axioms.

$$(L4) \quad \forall g, f \in G : c(g, f) \in A.$$

$$(L5) \quad \forall g \in G : g \neq 1 \rightarrow l(g^2) > l(g).$$

As we have mentioned in the introduction group actions on A -trees provide a geometric counterpart to A -length functions. To explain we need the following definitions, which we also use later in the text.

Let X be a non-empty set, A an ordered abelian group. An A -metric on X is a mapping $p : X \times X \rightarrow A$ such that for all $x, y, z \in X$:

$$(M1) \quad p(x, y) \geq 0 \text{ and } p(x, y) = 0 \text{ if and only if } x = y,$$

$$(M2) \quad p(x, y) = p(y, x),$$

$$(M3) \quad p(x, y) \leq p(x, z) + p(y, z).$$

In this case the pair (X, p) is an A -metric space. If (X, p) and (X', p') are A -metric spaces, an *isometry* from (X, p) to (X', p') is a mapping $f : X \rightarrow X'$ such that $p(x, y) = p'(f(x), f(y))$ for all $x, y \in X$.

For elements $a, b \in A$ the *closed segment* $[a, b]$ is defined by

$$[a, b] = \{c \in A \mid a \leq c \leq b\}.$$

More generally, a *segment* in an A -metric space is the image of an isometry $\alpha : [a, b] \rightarrow X$ for some $a, b \in A$. The endpoints of the segment are $\alpha(a), \alpha(b)$. A segment with endpoints $x, y \in X$ is denoted by $[x, y]$. An A -metric space (X, p) is *geodesic* if for all $x, y \in X$, there is a segment in X with endpoints x, y .

An A -tree is an A -metric space (X, p) such that:

$$(T1) \quad (X, p) \text{ is geodesic,}$$

$$(T2) \quad \text{if two segments of } (X, p) \text{ intersect in a single point, which is an endpoint of both, then their union is a segment,}$$

$$(T3) \quad \text{the intersection of two segments with a common endpoint is also a segment.}$$

We say that a group G acts on an A -tree X if it acts on X by isometries, i.e., an embedding of G into the group of isometries of X is fixed. An action of G on X is termed *free* if for every $1 \neq g \in G$ not g nor g^2 has a fixed point in X .

The following fundamental result relates free actions with free Lyndon length functions: a group G has a free Lyndon length function with values in A if and only if G acts freely on an A -tree [6, 28].

If a group G acts freely on an A -tree X , then the length function on G based at x , defined as $L_x(g) = p(x, gx)$, $g \in G$, is a free Lyndon length function on G .

For elements $g_1, \dots, g_n \in G$ we write

$$g = g_1 \circ \dots \circ g_n$$

if $g = g_1 \dots g_n$ and $l(g) = l(g_1) + \dots + l(g_n)$. Also, for $\alpha \in A$ we write $g = g_1 \circ_\alpha g_2$ if $g = g_1 g_2$ and $c(g_1^{-1}, g_2) < \alpha$.

A length function $l : G \rightarrow A$ is called *regular* if it satisfies the *regularity* axiom:

$$(L6) \quad \forall g, f \in G, \exists u, g_1, f_1 \in G :$$

$$g = u \circ g_1 \ \& \ f = u \circ f_1 \ \& \ l(u) = c(g, f).$$

Many examples of groups with regular free length functions are given in [20].

2.2 Regular actions

In this section we give a geometric characterization of group actions that come from the regular length functions.

Definition 1. *Let G act on a Λ -tree Γ . The action is regular with respect to $x \in \Gamma$ if for any $g, h \in G$ there exists $f \in G$ such that $[x, fx] = [x, gx] \cap [x, hx]$.*

Lemma 1. [20] *Let G act on a Λ -tree Γ . Then the action of G is regular with respect to $x \in \Gamma$ if and only if the length function $L_x : G \rightarrow \Lambda$ based at x is regular.*

Proof. Let d be the Λ -metric on Γ . By definition, the length function L_x is regular if for every $g, h \in G$ there exists $f \in G$ such that $g = fg_1$, $h = fh_1$, where $L_x(f) = c(g, h)$ and $L_x(g) = L_x(f) + L_x(g_1)$, $L_x(h) = L_x(f) + L_x(h_1)$.

Suppose the action of G is regular with respect to x . Then for $g, h \in G$ there exists $f \in G$ such that $[x, fx] = [x, gx] \cap [x, hx]$. We have $[x, gx] = [x, fx] \cup [fx, gx]$, $[x, hx] = [x, fx] \cup [fx, hx]$ and $[fx, gx] = [x, (f^{-1}g)x] = L_x(f^{-1}g)$, $[fx, hx] = [x, (f^{-1}h)x] = L_x(f^{-1}h)$. Taking $g_1 = f^{-1}g$, $h_1 = f^{-1}h$ we have $L_x(g) = L_x(f) + L_x(g_1)$, $L_x(h) = L_x(f) + L_x(h_1)$. Finally, since $c(g, h) = \frac{1}{2}(L_x(g) + L_x(h) - L_x(g^{-1}h))$ and $L_x(g^{-1}h) = d(x, (g^{-1}h)x) = d(gx, hx) = d(fx, gx) + d(fx, hx)$ we get $L_x(f) = c(g, h)$.

Suppose that L_x is regular. Then from $g = fg_1$, $h = fh_1$, where $L_x(g) = L_x(f) + L_x(g_1)$, $L_x(h) = L_x(f) + L_x(h_1)$ it follows that $[x, gx] = [x, fx] \cup [fx, gx]$, $[x, hx] = [x, fx] \cup [fx, hx]$. $L_x(f) = c(g, h) = \frac{1}{2}(L_x(g) + L_x(h) - L_x(g^{-1}h))$, so $2d(x, fx) = d(x, gx) + d(x, hx) - d(x, (g^{-1}h)x) = d(x, gx) + d(x, hx) - d(gx, hx)$. In other words,

$$d(gx, hx) = d(x, gx) + d(x, hx) - 2d(x, fx) = (d(x, gx) - d(x, fx)) +$$

$$(d(x, hx) - d(x, fx)) = d(fx, gx) + d(fx, hx)$$

which is equivalent to $[x, fx] = [x, gx] \cap [x, hx]$. □

Lemma 2. *Let G act minimally on a Λ -tree Γ . If the action of G is regular with respect to $x \in \Gamma$ then all branch points of Γ are G -equivalent.*

Proof. From minimality of the action it follows that Γ is spanned by the set of points $Gx = \{gx \mid g \in G\}$.

Now let y be a branch point in Γ . It follows that there exist (not unique in general) $g, h \in G$ such that $[x, y] = [x, gx] \cap [x, hx]$. From regularity of the action it follows that there exists $f \in G$ such that $y = fx$. Hence, every branch point is G -equivalent to x and the statement of the lemma follows. \square

Lemma 3. *Let G act on a Λ -tree Γ . If the action of G is regular with respect to $x \in \Gamma$ then it is regular with respect to any $y \in Gx$.*

Proof. We have to show that for every $g, h \in G$ there exists $f \in G$ such that $[y, fy] = [y, gy] \cap [y, hy]$. Since $y = tx$ for some $t \in G$ then we have to prove that $[tx, (ft)x] = [tx, (gt)x] \cap [tx, (ht)x]$. The latter equality is equivalent to $[x, (t^{-1}ft)x] = [x, (t^{-1}gt)x] \cap [x, (t^{-1}ht)x]$ which follows from regularity of the action with respect to x . \square

Lemma 4. *Let G act freely on a Λ -tree Γ so that all branch points of Γ are G -equivalent. Then the action of G is regular with respect to any branch point in Γ .*

Proof. Let x be a branch point in Γ and $g, h \in G$. If $g = h$ then $[x, gx] \cap [x, hx] = [x, gx]$ and g is the required element. Suppose $g \neq h$. Since the action is free then $gx \neq hx$ and we consider the tripod formed by x, gx, hx . Hence, $y = Y(x, gx, hx)$ is a branch point in Γ and by the assumption there exists $f \in G$ such that $y = fx$. \square

2.3 Infinite words and length functions

In this subsection at first we recall some notions from the theory of ordered abelian groups (for all the details refer to the books [10] and [22]) and then following [27] describe the construction of infinite words.

Let A be an ordered abelian group. A subset $C \subset A$ is called *convex* if for every $a, b \in C$ the set C contains $[a, b]$. In particular, a subgroup B of A is convex if $[0, b] \subset B$ for every positive $b \in B$.

An ordered abelian group A is called *discretely ordered* if A has a minimal positive element (we denote it by 1_A). In this event, for any $a \in A$ the following hold:

- 1) $a + 1_A = \min\{b \mid b > a\}$,
- 2) $a - 1_A = \max\{b \mid b < a\}$.

Observe that if A is any ordered abelian group then $\mathbb{Z} \oplus A$ is discretely ordered with respect to the right lexicographic order.

Let A be a discretely ordered abelian group and let $X = \{x_i \mid i \in I\}$ be a set. Put $X^{-1} = \{x_i^{-1} \mid i \in I\}$ and $X^\pm = X \cup X^{-1}$. An *A-word* is a function of the type

$$w : [1_A, \alpha_w] \rightarrow X^\pm,$$

where $\alpha_w \in A$, $\alpha_w \geq 0$. The element α_w is called the *length* $|w|$ of w .

By $W(A, X)$ we denote the set of all A -words. Observe, that $W(A, X)$ contains an empty A -word which we denote by ε .

Concatenation uv of two A -words $u, v \in W(A, X)$ is an A -word of length $|u| + |v|$ and such that:

$$(uv)(a) = \begin{cases} u(a) & \text{if } 1_A \leq a \leq |u| \\ v(a - |u|) & \text{if } |u| < a \leq |u| + |v| \end{cases}$$

An A -word w is *reduced* if $w(\beta + 1_A) \neq w(\beta)^{-1}$ for each $1_A \leq \beta < |w|$. We denote by $R(A, X)$ the set of all reduced A -words. Clearly, $\varepsilon \in R(A, X)$.

For $u \in W(A, X)$ and $\beta \in [1_A, \alpha_u]$ by u_β we denote the restriction of u on $[1_A, \beta]$. If $u \in R(A, X)$ and $\beta \in [1_A, \alpha_u]$ then

$$u = u_\beta \circ \tilde{u}_\beta,$$

for some uniquely defined \tilde{u}_β .

An element $\text{com}(u, v) \in R(A, X)$ is called the (*longest*) *common initial segment* of A -words u and v if

$$u = \text{com}(u, v) \circ \tilde{u}, \quad v = \text{com}(u, v) \circ \tilde{v}$$

for some (uniquely defined) A -words \tilde{u}, \tilde{v} such that $\tilde{u}(1_A) \neq \tilde{v}(1_A)$.

Now, we can define the product of two A -words. Let $u, v \in R(A, X)$. If $\text{com}(u^{-1}, v)$ is defined then

$$u^{-1} = \text{com}(u^{-1}, v) \circ \tilde{u}, \quad v = \text{com}(u^{-1}, v) \circ \tilde{v},$$

for some uniquely defined \tilde{u} and \tilde{v} . In this event put

$$u * v = \tilde{u}^{-1} \circ \tilde{v}.$$

The product $*$ is a partial binary operation on $R(A, X)$.

An element $v \in R(A, X)$ is termed *cyclically reduced* if $v(1_A)^{-1} \neq v(|v|)$. We say that an element $v \in R(A, X)$ admits a *cyclic decomposition* if $v = c^{-1} \circ u \circ c$, where $c, u \in R(A, X)$ and u is cyclically reduced. Observe that a cyclic decomposition is unique (whenever it exists). We denote by $CR(A, X)$ the set of all cyclically reduced words in $R(A, X)$ and by $CDR(A, X)$ the set of all words from $R(A, X)$ which admit a cyclic decomposition.

Below we refer to A -words as *infinite words* usually omitting A whenever it does not produce any ambiguity.

The following result establishes the connection between infinite words and length functions.

Theorem 2. [27] *Let A be a discretely ordered abelian group and X be a set. Then any subgroup G of $CDR(A, X)$ has a free Lyndon length function with values in A – the restriction $L|_G$ on G of the standard length function L on $CDR(A, X)$.*

Theorem 3. [5] *Let G have a free Lyndon length function $L : G \rightarrow A$, where A is a discretely ordered abelian group. Then there exists a length preserving embedding $\phi : G \rightarrow CDR(A, X)$, that is, $|\phi(g)| = L(g)$ for any $g \in G$.*

Corollary 1. [5] *Let G have a free Lyndon length function $L : G \rightarrow A$, where A is an arbitrary ordered abelian group. Then there exists an embedding $\phi : G \rightarrow CDR(A', X)$, where $A' = \mathbb{Z} \oplus A$ is discretely ordered with respect to the right lexicographic order and X is some set, such that, $|\phi(g)| = (0, L(g))$ for any $g \in G$.*

Corollary 2. [17] *Let G have a free regular Lyndon length function $L : G \rightarrow A$, where A is an arbitrary ordered abelian group. Then there exists an embedding $\phi : G \rightarrow CDR(A', X)$, where A' is a discretely ordered abelian group and X is some set, such that, the Lyndon length function on $\phi(G)$ induced from $R(A', X)$ is regular.*


Theorems 2 and 3, and Corollary 1 show that a group has a free Lyndon length function if and only if it embeds into a set of infinite words and this embedding preserves the length. Moreover, it is not hard to show that this embedding also preserves regularity of the length function.

3 Generalized equations

3.1 The notion of a generalied equation

We are going to represent the group G by infinite words and, therefore, as a group of a generalized equation, and use the elimination process. We will define a generalized equation here.

Definition 2. A combinatorial generalized equation Ω (which is convenient to visualize as on the picture)



consists of the following objects:

1. A finite set of **bases** $\mathcal{M} = BS(\Omega)$. The set of bases \mathcal{M} consists of $2n$ elements $\mathcal{M} = \{\mu_1, \dots, \mu_{2n}\}$. The set \mathcal{M} comes equipped with two functions: a function $\varepsilon : \mathcal{M} \rightarrow \{1, -1\}$ and an involution $\Delta : \mathcal{M} \rightarrow \mathcal{M}$ (that is, Δ is a bijection such that Δ^2 is an identity on \mathcal{M}). Bases μ and $\Delta(\mu)$ (or $\bar{\mu}$) are called dual bases. We denote bases by letters μ, λ , etc.
2. A set of **boundaries** $BD = BD(\Omega)$. BD is a finite initial segment of the set of positive integers $BD = \{1, 2, \dots, \rho + 1\}$. We use letters i, j , etc. for boundaries.
3. Two functions $\alpha : BS \rightarrow BD$ and $\beta : BS \rightarrow BD$. We call $\alpha(\mu)$ and $\beta(\mu)$ the initial and terminal boundaries of the base μ (or endpoints of μ). These functions satisfy the following conditions: for every base $\mu \in BS$: $\alpha(\mu) < \beta(\mu)$ if $\varepsilon(\mu) = 1$ and $\alpha(\mu) > \beta(\mu)$ if $\varepsilon(\mu) = -1$.
4. A set of boundary connections (p, λ, q) , where p is a boundary on λ (that is a number between $\alpha(\lambda)$ and $\beta(\lambda)$) and q on $\bar{\lambda}$. In this case we say that p and q are λ -tied. If (p, λ, q) is a boundary connection then $(q, \bar{\lambda}, p)$ is also a boundary connection. (The meaning of the boundary connections will be explained in ET5).

For a combinatorial generalized equation Ω , one can canonically associate a system of equations in **variables** h_1, \dots, h_ρ over $F(Z)$ (variables h_i are sometimes called *items*). This system is called a **generalized equation**, and (slightly abusing the language) we denote it by the same symbol Ω . The generalized equation Ω consists of the following two types of equations.

1. Each pair of dual bases $(\lambda, \Delta(\lambda))$ provides an equation

$$[h_{\alpha(\lambda)} h_{\alpha(\lambda)+1} \cdots h_{\beta(\lambda)-1}]^{\varepsilon(\lambda)} = [h_{\alpha(\Delta(\lambda))} h_{\alpha(\Delta(\lambda))+1} \cdots h_{\beta(\Delta(\lambda))-1}]^{\varepsilon(\Delta(\lambda))}.$$

These equations are called **basic equations**.

2. Every boundary connection (p, λ, q) gives rise to a **boundary equation**

$$[h_{\alpha(\lambda)} h_{\alpha(\lambda)+1} \cdots h_{p-1}] = [h_{\alpha(\Delta(\lambda))} h_{\alpha(\Delta(\lambda))+1} \cdots h_{q-1}],$$

if $\varepsilon(\lambda) = \varepsilon(\Delta(\lambda))$ and

$$[h_{\alpha(\lambda)} h_{\alpha(\lambda)+1} \cdots h_{p-1}] = [h_q h_{q+1} \cdots h_{\beta(\Delta(\lambda))-1}]^{-1},$$

if $\varepsilon(\lambda) = -\varepsilon(\Delta(\lambda))$.

Remark 1. We assume that every generalized equation comes associated with a combinatorial one.

Definition 3. Let $\Omega(h) = \{L_1(h) = R_1(h), \dots, L_s(h) = R_s(h)\}$ be a generalized equation in variables $h = (h_1, \dots, h_\rho)$. A sequence of reduced nonempty Λ -words $H = (H_1(Z), \dots, H_\rho(Z))$ in the alphabet $Z^{\pm 1}$ is a solution of Ω if:

1. all words $L_i(H), R_i(H)$ are reduced,
2. $L_i(H) = R_i(H)$, $i \in [1, s]$.

If we specify a particular solution H of a generalized equation Ω then we use a pair (Ω, H) . The interval on which the bases are situated is denoted by I .

Definition 4. A cancellation table $C(U)$ of a solution U is defined as follows

$$C(U) = \{h_i^\epsilon h_j^\sigma, \epsilon, \sigma = \pm 1, \text{ such that there is a cancellation between } H_i^\epsilon \text{ and } H_j^\sigma\}.$$

Definition 5. A solution H^+ in Λ words of a generalized equation Ω is called consistent with a system of linear equations $\Sigma \alpha_i \ell(h_i) = 0$ and solution H if it satisfies this system of linear equations and $C(H^+) \leq C(H)$

3.2 From a f.p. group to a generalized equation

Let $G = \langle X \mid R \rangle$ be a finitely presented group that has a regular free length function in an ordered abelian group Λ . By Corollary 2, G can be viewed as a subgroup of $CDR(\Lambda, Z)$, where Λ is a discretely ordered abelian group and Z is a set. Let us fix the embedding $\pi : G \hookrightarrow CDR(\Lambda, Z)$ for the rest of this section.

Let $X = \{x_1, \dots, x_n\}$ and $R = \{r_1(X), \dots, r_m(X)\}$. Adding, if necessary, additional generators, we can assume that every relation involves at most three generators. Since π is a homomorphism it follows that after substitution $x_i \rightarrow \pi(x_i)$, $i \in [1, n]$ all products $r_i(\pi(X))$, $i \in [1, m]$ cancel out. Hence, we have a cancellation diagram over $CDR(\Lambda, Z)$ which produce a generalized equation Ω corresponding to the embedding $\pi : G \hookrightarrow CDR(\Lambda, Z)$.

We can assume that each generator $x \in X$ of G occurs in the relations at least twice, otherwise we can eliminate it. Each generator x of G is represented in Ω as a product of bases without cancellation. For each occurrence of x in the relations R there is different such a representation. Since the action is regular, $G = G_\Omega = \langle \mathcal{M} \mid \Omega(\mathcal{M}) \rangle$ is the group of the generalized equation Ω . Generators here are bases, and each relation either has form $\lambda = \bar{\lambda}$, where $\lambda \in \mathcal{M}$ or $\lambda_{i_1} \dots \lambda_{i_k} = \lambda_{j_1} \dots \lambda_{j_m}$, where left and right hand sides represent the same generator $x \in X$ as product of bases without cancellation corresponding to different occurrences of x in the relations from R . From now on we will always consider \mathcal{M} as a generating set for G .

Let B be a CW 2-complex with fundamental group G with generating set \mathcal{M} .

Let $C(\Omega)$ be a CW 2-complex obtained from G_Ω by creating for each relation a polygon with boundary labeled by this relation and gluing these polygons together along edges labeled by the same letter. Then joining some pairs of vertices in $C(\Omega)$ by arcs we can obtain a CW complex $D(\Omega)$ that has the same fundamental group as B (namely, G). This set of pairs may happen to be empty.

Lemma 5. G is a free product of $\pi_1(C(\Omega))$ and, possibly, a free group.

The proof follows from the construction.

We say that bases λ and μ are oppositely oriented if $\varepsilon(\lambda) = -\varepsilon(\mu)$.

Proposition 1. Either for each pair of bases $\lambda, \bar{\lambda}$ in the generalized equation Ω , $\varepsilon(\lambda) = \varepsilon(\bar{\lambda})$ or one can construct an index two subgroup H in G such that for any pair of bases $\lambda, \bar{\lambda}$ in the generalized equation Ω_H for H , $\varepsilon(\lambda) = \varepsilon(\bar{\lambda})$.




Figure 1: From the cancellation diagram for the relation $xyz = 1$ to the generalized equation ($x = \lambda_1 \circ \lambda_2$, $y = \lambda_2^{-1} \circ \lambda_3$, $z = \lambda_3^{-1} \circ \lambda_1^{-1}$).

Proof. Let pairs of bases $\lambda_1, \bar{\lambda}_1, \dots, \lambda_k, \bar{\lambda}_k$ be oppositely oriented in Ω . Let $\{\lambda_1, \bar{\lambda}_1, \dots, \lambda_n, \bar{\lambda}_n\}$ be the set of all bases of Ω . Take a mirror reflection of Ω in the axis orthogonal to the main interval I of the generalized equation Ω , denote it by Ω_δ . Denote a mirror image of each base λ_i by δ_i (λ_i and δ_i are oppositely oriented). Now we take the disjoint union of Ω and Ω_δ , switch the names of $\bar{\lambda}_i$ and δ_i for $i = 1, \dots, k$, and denote by Ω_H the resulting generalized equation. It does not have pairs $\lambda, \bar{\lambda}$ that are oppositely oriented. Let $C(\Omega_H)$ be the CW 2-complex constructed from Ω_H the same way as $C(\Omega)$ was constructed from Ω . Similarly, let $D(\Omega_H)$ be the CW 2-complex constructed from Ω_H the same way as $D(\Omega)$ was constructed from Ω . Then $C(\Omega_H)$ is a double cover of $C(\Omega)$ and $D(\Omega_H)$ is a double cover of $D(\Omega)$. The fundamental group of $D(\Omega_H)$ is an index two subgroup of G . \square

Lemma 6. *If $Y = \mathcal{M} \cup \Delta$ is the set of bases of the generalized equation Ω_H , then the group of Ω_H , $G_{\Omega_H} = \langle Y | \Omega_H(Y) \rangle$ is a free product of H and an infinite cyclic group.*

Proof. Since complex $D(\Omega_H)$ is a double cover of $D(\Omega)$ which can be contracted to B , $D(\Omega_H)$ can be contracted to a complex with two vertices and 2-cells with boundaries labeled by the relations of G_{Ω_H} . Adding an edge joining these two vertices we obtain a CW complex with fundamental group G_{Ω_H} . \square

For the rest of the paper we replace Ω by Ω_H . Therefore we will assume that Ω does not have flipped (oppositely oriented) pairs of bases and that G is the group of Ω , $G = \langle \mathcal{M} | \Omega(\mathcal{M}) \rangle$. Then G is also generated by variables $h_i, i = 1, \dots, \rho$ and the relations of G in this set of generators are the basic equations and the boundary equations.

We will frequently use the following result.

Proposition 2. [13] *Any two nontrivial ordered abelian groups satisfy the same existential sentences in the language $L = \{0, +, -, <\}$.*

4 Elimination process

4.1 Elementary transformations

In this section we describe *elementary transformations* of generalized equations. Let (Ω, U) be a generalized equation with a solution U . An elementary transformation (*ET*) associates to a generalized equation (Ω, U) a generalized equation $ET(\Omega, U) = (\Omega_1, U_1)$ and an epimorphism $\pi : G_{R(\Omega)} \rightarrow G_{R(\Omega_1)}$ such that for the solution π_U the following diagram commutes.

$$\begin{array}{ccc} G_{R(\Omega)} & \xrightarrow{\pi} & G_{R(\Omega_1)} \\ \pi_U \downarrow & & \swarrow \pi_{U_1} \\ G & & \end{array}$$

We have a well-defined map $ET : (\Omega, U) \rightarrow (\Omega_1, U_1)$.

- (ET1) (*Cutting a base (see Fig. 1)*). Let λ be a base in Ω and p an internal boundary of λ (that is $p \neq \alpha(\lambda), \beta(\lambda)$) with a boundary connection (p, λ, q) . Then we cut the base λ in p into two new bases λ_1 and λ_2 and cut $\bar{\lambda}$ in q into the bases $\bar{\lambda}_1, \bar{\lambda}_2$.
- (ET2) (*Transferring a base (see Fig. 2)*). If a base λ of Ω contains a base μ (that is, $\alpha(\lambda) \leq \alpha(\mu) < \beta(\mu) \leq \beta(\lambda)$) and all boundaries on μ are λ -tied by boundary connections then we transfer μ from its location on the base λ to the corresponding location on the base $\bar{\lambda}$.
- (ET3) (*Removal of a pair of matched bases (see Fig. 3)*). If the bases λ and $\bar{\lambda}$ are *matched* (that is, $\alpha(\lambda) = \alpha(\bar{\lambda}), \beta(\lambda) = \beta(\bar{\lambda})$) then we remove $\lambda, \bar{\lambda}$ from Ω .

Remark 2. Observe, that for $i = 1, 2, 3$, $ETi(\Omega)$ and Ω have the same set of variables H , and the identity map $F[H] \rightarrow F[H]$ induces an isomorphism $G_{R(\Omega)} \rightarrow G_{R(\Omega')}$. Moreover, U is a solution of Ω if and only if U is a solution of Ω' .

- (ET4) (*Removal of a lone base (see Fig. 4)*). Suppose, a base λ in Ω does not *intersect* any other base, that is, the items $h_{\alpha(\lambda)}, \dots, h_{\beta(\lambda)-1}$ are contained only on the base λ . Suppose also that all boundaries in λ are λ -tied, i.e., for every i ($\alpha(\lambda) < i \leq \beta(\lambda) - 1$) there exists a boundary $b(i)$ such that $(i, \lambda, b(i))$ is a boundary connection in Ω . Then we remove the pair of bases λ and $\bar{\lambda}$ together with all the boundaries $\alpha(\lambda) + 1, \dots, \beta(\lambda) - 1$ (and rename the rest $\beta(\lambda) - \alpha(\lambda) - 1$ boundaries correspondingly).

We define the isomorphism $\theta : G_{R(\Omega)} \rightarrow G_{R(\Omega')}$ as follows:

$$\begin{aligned} \theta(h_j) &= h_j \text{ if } j < \alpha(\lambda) \text{ or } j \geq \beta(\lambda) \\ \theta(h_i) &= \begin{cases} h_{b(i)} \dots h_{b(i)-1}, & \text{if } \varepsilon(\lambda) = \varepsilon(\bar{\lambda}), \\ h_{b(i)} \dots h_{b(i-1)-1}, & \text{if } \varepsilon(\lambda) = -\varepsilon(\bar{\lambda}) \end{cases} \end{aligned}$$

for $\alpha + 1 \leq i \leq \beta(\lambda) - 1$.

- (ET5) (*Introduction of a boundary (see Fig. 5)*). Suppose a point p in a base λ is not λ -tied. The transformation (ET5) λ -ties it. To this end, denote by U_λ the word corresponding to λ , and let U'_λ be the beginning of this word ending at p . Then we perform one of the following two transformations according to where the end of U'_λ on $\bar{\lambda}$ is situated:




Figure 2: Elementary transformation (ET1).

- (a) If the end of U'_λ on $\bar{\lambda}$ is situated on the boundary q , we introduce the boundary connection $\langle p, \lambda, q \rangle$. In this case the corresponding isomorphism $\theta_q : G_{R(\Omega)} \rightarrow G_{R(\Omega_q)}$ is induced by the identity isomorphism on $F[H]$.
- (b) If the end of U'_λ on $\bar{\lambda}$ is situated between q and $q + 1$, we introduce a new boundary q' between q and $q + 1$ (and rename all the boundaries); introduce a new boundary connection $\langle p, \lambda, q' \rangle$. Denote the resulting equation by Ω'_q . In this case the corresponding isomorphism $\theta_{q'} : G_{R(\Omega)} \rightarrow G_{R(\Omega_{q'})}$ is induced by the map $\theta_{q'}(h) = h$, if $h \neq h_q$, and $\theta_{q'}(h_q) = h_{q'}h_{q'+1}$.

4.2 Derived transformations and auxiliary transformations

In this section we define complexity of a generalized equation and describe several useful “derived” transformations of generalized equations. Some of them can be realized as finite sequences of elementary transformations, others result in equivalent generalized equations but cannot be realized by finite sequences of elementary moves.




Figure 3: Elementary transformation (ET2).

A boundary is open if it is an internal boundary of some base, otherwise it is closed. A section $\sigma = [i, \dots, i + k]$ is said to be closed if boundaries i and $i + k$ are closed and all the boundaries between them are open.

Sometimes it will be convenient to subdivide all sections of Ω into **active** (denoted $A\Sigma_\sigma$) and **non-active sections**. Constant section will always be non-active. A variable h_q is called free if it meets no base. Free variables are transported to the very end of the interval behind all items in Ω and become non-active.

- (D1) (*Closing a section (see Fig. 6)*). Let σ be a section of Ω . The transformation (D1) makes the section σ closed. Namely, (D1) cuts all bases in Ω through the end-points of σ .
- (D2) (*Transporting a closed section (see Fig. 9)*). Let σ be a closed section of a generalized equation Ω . We cut σ out of the interval $[1, \rho_\Omega]$ together with all the bases on σ and put σ at the end of the interval or between any two consecutive closed sections of Ω . After that we correspondingly re-enumerate all the items and boundaries of the latter equation to bring it to the proper form. Clearly, the original equation Ω and the new one Ω' have the same solution sets and their coordinate groups are isomorphic.
- (D3) (*Moving free variables to the right*). Suppose that Ω contains a free variable h_q in an active section. Here we close the section $[q, q + 1]$ using D1, transport it to the very end of the interval behind all items in Ω using D2. In the resulting generalized equation Ω' the transported section becomes a non-active section.
- (D4) (*Deleting a complete base*). A base μ of Ω is called *complete* if there exists a closed section σ in Ω such that $\sigma = [\alpha(\mu), \beta(\mu)]$.

Suppose μ is an active complete base of Ω and σ is a closed section such that $\sigma = [\alpha(\mu), \beta(\mu)]$. In this case using ET5, we transfer all bases from μ to $\bar{\mu}$; using ET4, we remove the lone base μ together with the section $\sigma(\mu)$.




Figure 4: Elementary transformation (ET3).

(D5) (*Finding the kernel*).

Suppose that the equation Ω does not have any boundary connections. An active base $\mu \in A\Sigma_\Omega$ is called *eliminable* if at least one of the following holds:

- a) μ contains an item h_i with $\gamma(h_i) = 1$, where $\gamma(h_i)$ is the number of bases containing h_i .
- b) at least one of the boundaries $\alpha(\mu), \beta(\mu)$ is different from $1, \rho + 1$ and does not touch any other base (except μ).

The process of finding the kernel works as follows. We cut the bases of Ω along all the boundary connections then consequently remove eliminable bases until no eliminable bases left in the equation. The resulting generalized equation is called a *kernel* of Ω and we denote it by $Ker(\Omega)$. It is easy to see that $Ker(\Omega)$ does not depend on a particular removal process. Indeed, if Ω has two different eliminable bases μ_1, μ_2 , and deletion of μ_i results in an equation Ω_i then by induction (on the number of eliminations) $Ker(\Omega_i)$ is uniquely defined for $i = 1, 2$. Obviously, μ_1 is still eliminable in Ω_2 , as well as μ_2 is eliminable in Ω_1 . Now eliminating μ_1 and μ_2 from Ω_2 and Ω_1 we get one and the same equation Ω_0 . By induction $Ker(\Omega_1) = Ker(\Omega_0) = Ker(\Omega_2)$ hence the result. We say that a variable h_i belongs to the kernel ($h_i \in Ker(\Omega)$), if either h_i belongs to at least one base in the kernel, or it is constant.

Also, for an equation Ω by $\bar{\Omega}$ we denote the equation which is obtained from Ω by deleting all free variables. Obviously,

$$G_{R(\Omega)} = G_{R(\bar{\Omega})} * F(\bar{Y})$$

where \bar{Y} is the set of free variables in Ω .

Let us consider what happens on the group level in this process.

We start with the case when just one base is eliminated. Let μ be an eliminable base in $\Omega = \Omega(h_1, \dots, h_\rho)$. Denote by Ω_1 the equation resulting from Ω by eliminating μ .

(a) Suppose $h_i \in \mu$ and $\gamma(h_i) = 1$. Then the variable h_i occurs only once in Ω - precisely in the equation $s_\mu = 1$ corresponding to the base μ . Therefore, in the coordinate group $G_{R(\Omega)}$




Figure 5: Elementary transformation (ET4).

the relation $s_\mu = 1$ can be written as $h_i = w$, where w does not contain h_i . Using Tietze transformations we can rewrite the presentation of $F_{R(\Omega)}$ as $F_{R(\Omega')}$, where Ω' is obtained from Ω by deleting s_μ and the item h_i . It follows immediately that

$$G_{R(\Omega_1)} \simeq G_{R(\Omega')} * \langle h_i \rangle$$

and

$$G_{R(\Omega)} \simeq G_{R(\Omega')} \simeq G_{R(\overline{\Omega_1})} * F(B) \quad (1)$$

for some free group $F(B)$.

(b) Suppose now that μ satisfies case b) above with respect to a boundary i . Then in the equation $s_\mu = 1$ the variable h_{i-1} either occurs only once or it occurs precisely twice and in this event the second occurrence of h_{i-1} (in $\Delta(\mu)$) is a part of the subword $(h_{i-1}h_i)^{\pm 1}$. In both cases it is easy to see that the tuple

$$(h_1, \dots, h_{i-2}, s_\mu, h_{i-1}h_i, h_{i+1}, \dots, h_\rho)$$

forms a basis of the ambient free group generated by (h_1, \dots, h_ρ) and constants from A . Therefore, eliminating the relation $s_\mu = 1$, we can rewrite the presentation of $G_{R(\Omega)}$ in generators $\bar{Y} = (h_1, \dots, h_{i-2}, h_{i-1}h_i, h_{i+1}, \dots, h_\rho)$. Observe also that any other equation $s_\lambda = 1$ ($\lambda \neq \mu$) of Ω either does not contain variables h_{i-1}, h_i or it contains them as parts of the subword $(h_{i-1}h_i)^{\pm 1}$, i.e., any such a word s_λ can be expressed as a word $w_\lambda(\bar{Y})$ in terms of generators \bar{Y} . This shows that

$$G_{R(\Omega)} \simeq G(\bar{Y})_{R(w_\lambda(\bar{Y})|\lambda \neq \mu)} \simeq G_{R(\Omega')},$$

where Ω' is a generalized equation obtained from Ω_1 by deleting the boundary i . Denote by Ω'' an equation obtained from Ω' by adding a free variable z to the right end of Ω' . It follows now that

$$G_{\Omega_1} \simeq G_{\Omega''} \simeq G_\Omega * \langle z \rangle$$




Figure 6: Elementary transformation (ET5).

and

$$G_\Omega \simeq G_{\overline{\Omega'}} * F(Z) \tag{2}$$

for some free group $F(Z)$. Notice that all the groups and equations which occur above can be found effectively.

By induction on the number of steps in a process we obtain the following lemma.

Lemma 7. *If Ω is a generalized equation, then*

$$F_{R(\Omega)} \simeq F_{R(\overline{Ker\Omega})} * F(Z)$$

where $F(Z)$ is a free group on Z .

Proof. Let

$$\Omega = \Omega_0 \rightarrow \Omega_1 \rightarrow \dots \rightarrow \Omega_l = Ker\Omega$$

be a cleaning process for Ω . It is easy to see (by induction on l) that for every $j = 0, \dots, l - 1$

$$\overline{Ker\Omega_j} = \overline{Ker\overline{\Omega_j}}$$

Moreover, if Ω_{j+1} is obtained from Ω_j as in the case 2) above, then (in the notation above)

$$\overline{Ker(\Omega_j)_1} = \overline{Ker\Omega'_j}$$

Now the statement of the lemma follows from the remarks above and equalities (1) and (2).

D6 (*Linear elimination.*)

Suppose that in Ω_v there is h_i in an active section with $\gamma_i = 1$ and such that $\ell(h_i)$ is comparable with the length of the active section. In this case we say that Ω is linear in h_i .

If Ω is linear in h_i in an active section such that both boundaries i and $i + 1$ are closed, then we remove the closed section $[i, i + 1]$ together with the lone base using ET4.

If there is no such h_i but Ω_v is linear in some h_i in an active section such that one of the boundaries $i, i + 1$ is open, say $i + 1$, and the other is closed, then we perform ET5 and μ -tie $i + 1$ through the only base μ it intersects; using ET1 we cut μ in $i + 1$; and then we delete the closed section $[i, i + 1]$ by ET4.

Suppose there is no h_i as above but Ω_v is linear in some h_i in an active section such that both boundaries i and $i + 1$ are open. In addition, assume that there is a closed section σ containing exactly two (not matched) bases μ_1 and μ_2 , such that $\sigma = \sigma(\mu_1) = \sigma(\mu_2)$ and in the generalized equation $\tilde{\Omega}_v$ (see the derived transformation D3) all the bases obtained from μ_1, μ_2 by ET1 in constructing $\tilde{\Omega}_v$ from Ω_v , do not belong to the kernel of $\tilde{\Omega}_v$. Here, using ET5, we μ_1 -tie all the boundaries inside μ_1 ; using ET2, we transfer μ_2 onto $\Delta(\mu_1)$; and remove μ_1 together with the closed section σ using ET4.

Suppose now that Ω_v satisfies the first assumption of the previous paragraph and does not satisfy the second one. In this event we close the section $[i, i + 1]$ using D1 and remove it using ET4.

Lemma 8. *Suppose that the process of linear elimination continues infinitely and there is a corresponding sequence of generalized equations*

$$\Omega \rightarrow \Omega_1 \rightarrow \Omega_k \rightarrow \dots$$

1. ([18], Lemma 15) *The number of different generalized equations that appear in the process is finite. Therefore some generalized equation appears in this process infinitely many times.*
2. ([18], Lemma 15) *If $\Omega_j = \Omega_k, i, j < k$, then $\pi(j, k)$ is an isomorphism invariant with respect to the kernel, namely $\pi(j, k)(h_i) = h_i$ for any variable h_i that belongs to some base in $\ker(\Omega)$.*
3. *The interval for the equation Ω_j can be divided into two disjoint parts, each being the union of closed sections, such that one part is a generalized equation $\ker(\Omega)$ and the other part is non-empty and corresponds to a generalized equation Ω' such that $G_{\Omega'} = F(Z)$ is a free group on variables Z , and $G_{\Omega} = G_{\ker(\Omega)} * F(Z)$.*

Proof. We will only prove the third statement. In the process of linear elimination we do not remove or cut variables h_i that belong to the kernel. Since we do not cut variables that belong to the kernel, we can remove parts of bases that cover some of these variables only finitely many times. Therefore, bases that are cut in the process belong to the part of the interval that is disjoint from the $\ker(\Omega)$. Together with Lemma 7 this proves the third statement. \square

- (D7) (*Tietze cleaning*). This transformation consists of four transformations performed consecutively:
- (a) Linear elimination; if the process of linear elimination goes infinitely we replace the equation by its kernel;
 - (b) deleting all pairs of matched bases;
 - (c) deleting all complete bases;
 - (d) moving all free variables to the right.
- (D8) (*Entire transformation*). We need a few further definitions. A base μ of the equation Ω is called a *leading base* if $\alpha(\mu) = 1$. A leading base is said to be *maximal* (or a *carrier*) if

$\beta(\lambda) \leq \beta(\mu)$, for any other leading base λ . Let μ be a carrier base of Ω . Any active base $\lambda \neq \mu$ with $\beta(\lambda) \leq \beta(\mu)$ is called a *transfer* base (with respect to μ).

Suppose now that Ω is a generalized equation with $\gamma(h_i) \geq 2$ for each h_i in the active part of Ω and such that $\ell(h_i)$ is comparable with the length of the active part. An *entire transformation* is a sequence of elementary transformations which are performed as follows. We fix a carrier base μ of Ω . We transfer all transfer bases from μ onto $\bar{\mu}$. Now, there exists some $i < \beta(\mu)$ such that h_1, \dots, h_i belong to only one base μ , while h_{i+1} belongs to at least two bases. Applying (ET1) we cut μ along the boundary $i + 1$. Finally, applying (ET4) we delete the section $[1, i + 1]$.

4.3 Complexity of a generalized equation

Denote by ρ_A the number of variables h_i in all active sections of Ω , by $n_A = n_A(\Omega)$ the number of bases in active sections of Ω , by ν' the number of open boundaries in the active sections, by σ' the number of closed boundaries in the active sections.

For a closed section $\sigma \in \Sigma_\Omega$ denote by $n(\sigma), \rho(\sigma)$ the number of bases and, respectively, variables in σ .

$$\rho_A = \rho_A(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} \rho(\sigma),$$

$$n_A = n_A(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} n(\sigma).$$

The *complexity* of the generalized equation Ω is the number

$$\tau = \tau(\Omega) = \sum_{\sigma \in A\Sigma_\Omega} \max\{0, n(\sigma) - 2\}.$$

Notice that the entire transformation (D5) as well as the cleaning process (D4) do not increase complexity of equations.

4.4 Rewriting process for Ω and reassigning lengths

In this section we describe a rewriting process for a generalized equation Ω . We assume that Ω is in standard form, namely, that transformations ET3, D3 and D4 have been applied to Ω and that on each step we apply them to the generalized equation before applying any other transformation.

Let Ω be a generalized equation. We construct a path $T(\Omega)$ (with associated structures), as a directed path oriented from the root v_0 , starting at v_0 and proceeding by induction on the distance n from the root.

We start with a general description of the path $T(\Omega)$. For each vertex v in $T(\Omega)$ there exists a unique generalized equation Ω_v associated with v . The initial equation Ω is associated with the root v_0 , $\Omega_{v_0} = \Omega$. In addition there is a homogeneous system of linear equations Σ_v with integer coefficients on the lengths of some variables of the equation Ω_v such the length of these variables in \mathcal{M} satisfy this system. We take Σ_{v_0} to be empty. For each edge $v \rightarrow v'$ (here v and v' are the origin and the terminus of the edge) there exists an epimorphism that is an isomorphism on H , $\pi(v, v') : G_{\Omega_v} \rightarrow G_{\Omega_{v'}}$ associated with $v \rightarrow v'$.

If

$$v \rightarrow v_1 \rightarrow \dots \rightarrow v_s \rightarrow u$$

is a path in $T(\Omega)$, then by $\pi(v, u)$ we denote composition of corresponding epimorphisms

$$\pi(v, u) = \pi(v, v_1) \cdots \pi(v_s, u).$$

If $v \rightarrow v'$ is an edge then there exists a finite sequence of elementary or derived transformations from Ω_v to $\Omega_{v'}$ and the epimorphism $\pi(v, v')$ is composition of the epimorphisms corresponding to these transformations. We also assume that active [non-active] sections in $\Omega_{v'}$ are naturally inherited from Ω_v , if not said otherwise.

Suppose the path $T(\Omega)$ is constructed by induction up to a level n , and suppose v is a vertex at distance n from the root v_0 . We describe now how to extend the path from v . The construction of the outgoing edge at v depends on which case described below takes place at the vertex v . There are three possible cases:

Linear case: there exists h_i in the active part such that $\ell(h_i)$ is comparable with the length of the active part and $\gamma(h_i) = 1$

Quadratic and almost quadratic case: $\gamma_i = 2$ for all h_i in the active part such that $\ell(h_i)$ is comparable with the length of the active part.

General JSJ case: $\gamma_i \geq 2$ for all h_i in the active part such that $\ell(h_i)$ is comparable with the length of the active part, and there exists such h_i that $\gamma_i > 2$.

4.4.1 Linear case

We apply the Tietze cleaning at the vertex v_n if it is possible. The system Σ_{v_n} implies a new system $\Sigma_{v_{n+1}}$, and assign the result to v_{n+1} .

If $\Omega_{v_{n+1}}$ splits into two parts, $\Omega_{v_{n+1}}^{(1)} = Ker(\Omega_{v_n})$ and $\Omega_{v_{n+1}}^{(2)}$ that corresponds to a free group $F(Z)$, there are two possible cases:

Case 1. $\Sigma_{v_{n+1}}$ also splits into two parts, $\Sigma_{v_{n+1}}^{(1)}$ involves only variables from $\Omega_{v_{n+1}}^{(1)}$ and $\Sigma_{v_{n+1}}^{(2)}$ involves only variables from $\Omega_{v_{n+1}}^{(2)}$.

Case 2. $\Sigma_{v_{n+1}}$ does not split into two such parts.

Suppose h_1, \dots, h_m are variables of $\Omega_{v_{n+1}}^{(2)}$. In the first case we solve the system $\Sigma_{v_{n+1}}^{(2)}$ over \mathbb{Z} (there must be a non-zero solution because the system has integer coefficients and a non-zero solution over Λ .) If n_1, \dots, n_m is a solution, we will be looking for lengths of h_1, \dots, h_m in the form $(\dots, n_1), \dots, (\dots, n_m)$. First, we assign non-negative not all zero integer values n_1, \dots, n_m to these variables and continue running the process for $\Omega_{v_{n+1}}^{(2)}$ with these lengths. The process will eventually stop with one or several matching pairs, we will not have a reflection because we do not have pairs of oppositely oriented dual bases. Then we take another integer solution k_1, \dots, k_m of $\Sigma_{v_{n+1}}^{(2)}$ such that when we run the process with lengths of variables $(k_1, n_1), \dots, (k_m, n_m)$ we do not get a proper quotient of the group of $\Omega_{v_{n+1}}^{(2)}$. Such solution always exists because it exists in Λ . For these lengths in \mathbb{Z}^2 the process ends with matching pairs becoming overlapping pairs. So we have a series of HNN-extensions over elements of equal length.

In the second case we do not change the length of variables from $\Omega_{v_{n+1}}^{(2)}$ at this step because later we will have some additional constrains for a solution of $\Sigma_{v_{n+1}}$, but later we will similarly assign to h_1, \dots, h_m lengths from \mathbb{Z}^2 .

In both cases we put the free group section $\Omega_{v_{n+1}}^{(2)}$ into a non-active part and thus decrease the complexity.

If it is impossible to apply Tietze cleaning, that is $(\gamma(h_i) \geq 2$ for any h_i in the active part of Ω_v comparable to the length of the active part), we apply the entire transformation.

Termination condition: 1. Ω_v does not contain active sections. In this case the vertex v is called a *leaf* or an *end vertex*.

4.4.2 Quadratic case

Suppose Ω_v satisfies the condition $\gamma_i \leq 2$ for each h_i in the active part and $\gamma_i = 2$ for each h_i in the active part comparable with the length of the active part. First of all, we fill in all the h'_i 's in the

active part such that $\gamma_i = 1$ by new (infinitely short) bases μ such that $\bar{\mu}$ covers a new free variable in the non-active part.

We apply the entire transformation (D8) and then apply Tietze cleaning.

Let us analyze the structure of $G_{R(\Omega_v)}$ in this case. Let σ be a closed section in the active part $\sigma = [h_1, \dots, h_i]$. Let F_1 be a free group with basis $\{h_1, \dots, h_i\}$. If σ contains an open boundary j , then we can consider a new generalized equation Ω'_v obtained from Ω_v by replacing the product $h_{j-1}h_j$ by a new variable h'_{j-1} and represent $G_{R(\Omega_v)} = G_{R(\Omega'_v)} * \langle h_j \rangle$. Therefore we can suppose that σ does not contain open boundaries. We say that μ is a *quadratic* base if σ contains μ and $\bar{\mu}$, otherwise μ is a *quadratic-coefficient base*. Denote the set of quadratic-coefficient bases by C . Suppose that σ contains quadratic bases. Let F_1/σ be the quotient of F_1 over the normal closure of elements $h[\alpha(\mu), \beta(\mu)]h[\alpha(\bar{\mu}), \beta(\bar{\mu})]^{-1}$. Let \mathcal{M} be a set that contains exactly one representative of each pair of double bases on σ and contains also each base μ such that $\mu \in \sigma$ and $\bar{\mu} \notin \sigma$. If we identify each base on σ with its double, then the product $h_1 \cdots h_i$ can be written as a product of bases from \mathcal{M} in exactly two different ways $\mu_{i_1} \cdots \mu_{i_k}$, and $\mu_{j_1} \cdots \mu_{j_t}$. Then F_1/σ is isomorphic to the quotient of the free group $F(\mathcal{M})$ over the relation $\mu_{i_1} \cdots \mu_{i_k} = \mu_{j_1} \cdots \mu_{j_t}$. Every element of \mathcal{M} occurs in this relation at most twice. Applying an automorphism of $F(\mathcal{M})$ identical on C , we can obtain another basis $X \cup T \cup C$ of this group such that in this basis the relation has form of a standard quadratic equation in variables from X with coefficients in $F(C)$ [24], Section 1.7, (variables from T do not participate in any relations). The quadratic equation corresponding to σ can be written in the standard form with coefficients expressed in terms of non-active variables. If the equation is regular, then $G_{R(\Omega_v)}$ has a QH subgroup corresponding to this equation. If the equation is not regular and there are quadratic bases, then $G_{R(\Omega_v)}$ splits as an HNN extension over an abelian subgroup. If there are infinitely small h_i 's in the quadratic part such that $\gamma(h_i) = 1$ then $G_{R(\Omega_v)}$ splits as a free product of the group of the generalized equation on the non-quadratic part and a free group.

We say that a base *participates* in the infinite process of entire transformation if it participates infinitely many times as a leading or transfer base.

Lemma 9. *If σ has quadratic-coefficient bases, and the entire transformation goes infinitely, then after a finite number of steps there will be quadratic bases that have length infinitely larger than all participating quadratic coefficient bases.*

Proof. After finite number of steps quadratic-coefficient bases can only be used as transfer bases (otherwise quadratic bases would be leaving the quadratic part). Every time such a base of length ℓ used as a transfer base, the length of the quadratic part decreases by ℓ . For the process to go infinitely, ℓ must be infinitely less than the length of the quadratic part. Therefore there is a quadratic base that have length infinitely larger than all participating quadratic coefficient bases. \square

If σ does not have quadratic-coefficient bases, then $G_{R(\Omega_v)}$ splits as a free product with one factor being a closed surface group or a free group.

We move σ into a non-active part and thus decrease the complexity.

For the system of linear equations $\Sigma_{v_{n+1}}$ the same two cases as in the previous subsection may occur.

4.4.3 Almost Quadratic case

In this case $\gamma_i = 2$ for all h_i in the active part such that $\ell(h_i)$ is comparable with the length of the active part. This case is, actually, a part of the case considered in the next subsection, but we consider it separately to make the subdivision into cases depend only on γ_i for the oldest variables.

In this case we apply the entire transformation. If it goes infinitely, then after a finite number of steps participating quadratic-coefficient bases that are comparable with the length of the active part can only be used as transfer bases. And, therefore, there will be participating quadratic bases that have length infinitely larger than all participating quadratic coefficient bases.

We fill in all the h'_i 's in the active part such that $\gamma_i = 1$ by new (infinitely short) bases appearing only once. we perform the entire transformation of the quadratic part until we obtain an overlapping pair section where the shift is infinitely shorter than the overlapping base λ . Therefore $G_{R(\Omega)}$ is an HNN-extension with stable letter corresponding to λ . We introduce the new base ρ such that $\alpha(\rho) = \alpha(\lambda)$, $\beta(\rho) = \beta(\bar{\lambda})$, and $\bar{\rho}$ is a non-active base. Then we apply the entire transformation to the new equation and move the overlapping pair section to $\bar{\rho}$. We repeat the described transformation until there is no quadratic base on the active section that has length comparable with the length of λ .

Then we remove the overlapping pairs of bases and consider the remaining generalized equation. We have to add an equation to $\Sigma_{v_{i+1}}$ that guarantees that the associated cyclic subgroups are generated by elements of the same length. We still have to assign lengths in \mathbb{Z}^n to bases in the quadratic part after we finish the process. In this case the subgroup of $G_{R(\Omega)}$ that is isomorphic to the coordinate group of the new generalized equation is a vertex group in an abelian splitting of $G_{R(\Omega)}$ and is smaller in Delzant-Potyagailo hierarchy [7].

4.4.4 General JSJ-case

Generalized equation Ω_v satisfies the condition $\gamma_i \geq 2$ for each h_i in the active part such that $\ell(h_i)$ is comparable with the length of the active part, and $\gamma_i > 2$ for at least one h_i .

(a) QH-subgroup case. First of all, we fill in all the h'_i 's in the active part such that $\gamma_i = 1$ by new (infinitely short) bases appearing only once. We apply transformation (D1) to close the quadratic part and put it in front of the interval. If the quadratic part corresponds to QH-vertex or to the representation of $G_{R(\Omega)}$ as an HNN extension, and there is a quadratic base that is infinitely longer than all the quadratic coefficient bases. In this case we perform the entire transformation of the quadratic part until we obtain an overlapping pair section where the shift is infinitely shorter than the overlapping base λ . Therefore $G_{R(\Omega)}$ is an HNN-extension with stable letter corresponding to λ . We introduce the new base ρ such that $\alpha(\rho) = \alpha(\lambda)$, $\beta(\rho) = \beta(\bar{\lambda})$, and $\bar{\rho}$ is a non-active base. Then we apply the entire transformation to the new equation and move the overlapping pair section to $\bar{\rho}$. We repeat the described transformation until there is no quadratic base on the quadratic section that has length comparable with the length of λ . Then we remove the quadratic section together with doubles of all the bases on it and consider the remaining generalized equation. We have to add an equation to $\Sigma_{v_{i+1}}$ that guarantees that the associated cyclic subgroups are generated by elements of the same length. We still have to assign lengths in \mathbb{Z}^n to bases in the quadratic part after we finish the process. In this case the subgroup of $G_{R(\Omega)}$ that is isomorphic to the coordinate group of the new generalized equation is a vertex group in an abelian splitting of $G_{R(\Omega)}$ and is smaller in Delzant-Potyagailo hierarchy.

(b) QH-shortening

Lemma 10. *If the quadratic part doesn't correspond to the HNN-splitting of G (there are only quadratic coefficient bases) or there is no quadratic base that is infinitely longer than all the quadratic coefficient bases, then one can replace (if necessary) the words corresponding to the quadratic bases in the quadratic part by shorter words so that the length of the quadratic part is bounded by some function $f_1(\Omega)$ times the length of the non-quadratic part.*

Proof. Indeed, some of the bases will eventually be transferred to the non-quadratic part (therefore, their length is bounded by the length of the non-quadratic part), reincarnations of the others will form matching pairs, therefore can be replaced by letters. \square

(c) **Abelian splitting: short shift.** Suppose Ω_v satisfies the following condition: the carrier base μ of the equation Ω_v intersects with its dual $\bar{\mu}$ (form an overlapping pair) and is at least twice longer than $|\alpha(\bar{\mu}) - \alpha(\mu)|$.

Here we construct an auxiliary equation $\hat{\Omega}_v$ (which does not occur in $T(\Omega)$) as follows. Firstly, we add a new constant section $[\rho_v + 1, \rho_v + 2]$ to the right of all sections in Ω_v (in particular, h_{ρ_v+1} is a new free variable). Secondly, we introduce a new pair of bases $(\lambda, \bar{\lambda})$ such that

$$\alpha(\lambda) = 1, \beta(\lambda) = \beta(\bar{\mu}), \alpha(\bar{\lambda}) = \rho_v + 1, \beta(\bar{\lambda}) = \rho_v + 2.$$

Notice that Ω_v can be obtained from $\hat{\Omega}_v$ by (ET4): deleting $\bar{\lambda}$ together with the closed section $[\rho_v + 1, \rho_v + 2]$. Let

$$\hat{\pi}_v : G_{R(\Omega_v)} \rightarrow G_{R(\hat{\Omega}_v)}$$

be the isomorphism induced by (ET4). General JSJ case still holds for $\hat{\Omega}_v$, but now λ is the carrier base. Applying to $\hat{\Omega}_v$ the entire transformation, we obtain a new vertex $\Omega_{v'}$ together with the isomorphism

$$\eta_{v'} : G_{R(\hat{\Omega}_v)} \rightarrow G_{R(\Omega_{v'})}.$$

As a culmination of the above process of adding auxiliary edges we get a generalized equation Ω' which consists of finitely many closed sections $\sigma_1, \dots, \sigma_p$ such that there exists an overlapping pair $(\mu_i, \bar{\mu}_i)$ for each σ_i , $i \in [1, p]$.

Let $\sigma = \sigma_{i_0}$, where $i_0 \in [1, p]$ have the overlapping pair $(\mu, \bar{\mu})$ such that

$$|[\alpha(\mu), \alpha(\bar{\mu})]| = \max_{i \in [1, p]} |[\alpha(\mu_i), \alpha(\bar{\mu}_i)]|.$$

Without loss of generality we can assume $i_0 = 1$, that is, $\sigma = [1, \beta(\bar{\mu})]$.

A *periodic structure* on Ω' is a pair $\langle \mathcal{P}, R \rangle$, where

1. \mathcal{P} is a set consisting of some items h_i , some bases λ , and some closed sections σ from Ω' such that the following conditions are satisfied:
 - (a) if $h_i \in \mathcal{P}$ and $h_i \in \lambda$ then $\lambda \in \mathcal{P}$,
 - (b) if $\lambda \in \mathcal{P}$, then $\bar{\lambda} \in \mathcal{P}$,
 - (c) if $\lambda \in \mathcal{P}$ and $\lambda \in \sigma$, then $\sigma \in \mathcal{P}$,
 - (d) there exists a function \mathcal{X} mapping the set of closed sections from \mathcal{P} into $\{-1, +1\}$ such that for every $\lambda, \sigma_1, \sigma_2 \in \mathcal{P}$, the condition that $\lambda \in \sigma_1$ and $\bar{\lambda} \in \sigma_2$ implies $\varepsilon(\lambda) \cdot \varepsilon(\bar{\lambda}) = \mathcal{X}(\sigma_1) \cdot \mathcal{X}(\sigma_2)$.
2. R is an equivalence relation on the set of boundaries \mathcal{B} belonging to closed sections from \mathcal{P} (if a boundary l belongs to two closed section $\sigma_{left}(l) = [i, l]$, $\sigma_{right}(l) = [l, j]$ from \mathcal{P} then instead of l we add to \mathcal{B} formal copies of l_{left}, l_{right} which are boundaries of $\sigma_{left}(l)$ and $\sigma_{right}(l)$ respectively), defined as follows: if $\lambda \in \mathcal{P}$ then

$$\alpha(\lambda) \sim_R \alpha(\bar{\lambda}), \beta(\lambda) \sim_R \beta(\bar{\lambda}) \text{ if } \varepsilon(\lambda) = \varepsilon(\bar{\lambda})$$

$$\alpha(\lambda) \sim_R \beta(\bar{\lambda}), \beta(\lambda) \sim_R \alpha(\bar{\lambda}) \text{ if } \varepsilon(\lambda) = -\varepsilon(\bar{\lambda}).$$

We call items, bases and sections *long* if they belong to \mathcal{P} (the rest items, bases and sections of Ω' are called *short* respectively).

A reduced infinite word u is called a *period* if it is cyclically reduced. A reduced infinite word w is called *u-periodic* for a period u if $w^{-1}uw = v$, where $|u| = |v|$. Since u is cyclically reduced it follows that either $w^{-1}u = w^{-1} \circ u$ or $uw = u \circ w$. We are going to distinguish two cases of periodicity:

1. w is unbounded u -periodic if $ht(w) \geq ht(u)$,
2. w is bounded u -periodic if $ht(w) = ht(u)$ and $w = u^k \circ u_1$, where $k \geq 2$ and $u = u_1 \circ u_2$.

Observe that in both cases w begins with a power of u but in the former case it is unbounded while bounded in the latter one (hence, the terms used).

Lemma 11. *Let $w_1, w_2 \in CDR(A, Z)$, where $|w_1| \geq |w_2|$, be u -periodic for some period $u \in CDR(A, Z)$. Assume that $w_1^{-1}uw_1 = v_1$, $w_2^{-1}uw_2 = v_2$. If $w_1^{-1}w_2$ is defined in $CDR(A, Z)$ then w_2 cancels completely in this product. Moreover, $w_1^{-1}w_2$ is v_1^{-1} -periodic.*

Proof. Follows from the definition of u -periodic words. □

Lemma 12. *If $w \in CDR(A, Z)$ is both u - and v -periodic for some periods $u, v \in CDR(A, Z)$ then $[u, v] = \varepsilon$. In particular, u is unbounded v -periodic if $ht(u) > ht(v)$.*

Proof. If $ht(u) = ht(v)$ then commutativity follows from Lemma 3.3 [17] since powers of u and v have a common initial segment of length greater than $|u| + |v|$. If $ht(u) > ht(v)$ then from v -periodicity of w it follows that u is v -periodic. Now, since w contains any natural power of u as an initial subword it follows that the terminal subword of u of length $|v|$ must coincide with v and commutativity follows. □

Let U be a solution of the equation Ω' , where $h_i \rightarrow U_i$. According to our assumption, X_μ is P -periodic, where $P = U[1, \alpha(\bar{\mu})]$. Since μ and $\bar{\mu}$ are products of items then every U_i , $i \in [\alpha(\bar{\mu}), \beta(\mu)]$, such that $|U_i| \geq 2|P|$, is Q_i -periodic, where Q_i is an infinite word such that $|Q_i| = |P|$. Moreover, we can associate with U a periodic structure $\mathcal{P}(U, P) = \langle \mathcal{P}, R \rangle$ as follows. We construct \mathcal{P} as a union of the chain of sets

$$\mathcal{P}_1 \subseteq \mathcal{P}_2 \subseteq \dots \subseteq \mathcal{P}_k,$$

where $k \in \mathbb{N}$. To construct \mathcal{P}_0 we add to it σ_1 , and all Q -periodic items $h \in \sigma_1$, where $|Q| = |P|$, together with any base λ (and its dual) containing h . Now, assume that \mathcal{P}_i is constructed. If there exists a base $\lambda \in \mathcal{P}_i$ such that $\bar{\lambda} \in \sigma_j$ then we set

$$\mathcal{P}_{i+1} = \mathcal{P}_i \cup \{\sigma_j\} \cup \{h, \lambda \in \sigma_j \mid h \text{ is a } Q\text{-periodic item where } |Q| = |P|, \lambda \text{ is a base containing } h\}.$$

Since Ω' contains only finitely many closed sections then this process stops.

Lemma 13. *Let U be a solution of the equation Ω' and $\mathcal{P}(U, P)$ be a periodic structure corresponding to U . Then every base(item) $\lambda \in \mathcal{P}$ is Q_λ -periodic, where $|Q_\lambda| = |P|$ and Q_λ is conjugate to P in G .*

Proof. It is enough to prove the statement for bases - the argument for items is similar. We are going to use the induction on k , where k is the number of sections in \mathcal{P}

By our assumption (short shift) we have $\mu \in \mathcal{P}$. It follows that any base $\lambda \in \sigma_1$, such that $|\lambda| \geq 2|P|$, is Q_λ -periodic, where $|Q_\lambda| = |P|$. Moreover, if $w = U[\alpha(\bar{\mu}), \alpha\lambda]$ then $w \in G$ and $w^{-1}Pw = Q_\lambda$. Hence, all bases from \mathcal{P}_1 have the properties stated in the lemma.

Suppose we have the statement of the lemma for all bases from \mathcal{P}_{k-1} . By definition, \mathcal{P}_k is obtained from \mathcal{P}_{k-1} by adding a section σ_k , such that there exists $\lambda \in \mathcal{P}_{k-1}$ and $\bar{\lambda} \in \sigma_k$, as well as all appropriate items and bases of σ_k . Observe that since μ_k is Q -periodic, where $Q = U[\alpha(\mu_k), \alpha(\bar{\mu}_k)]$, it follows that $\bar{\lambda}$ is Q' -periodic, where $|Q'| = |Q|$ and Q' is conjugate to Q by means of $g \in G$. At the same time, by the induction hypothesis, λ is Q_λ -periodic, where $|Q_\lambda| = |P|$. Hence, by Lemma 12 we have $[Q', Q_\lambda] = \varepsilon$ and it follows that $[Q, g^{-1}Q_\lambda g] = \varepsilon$, where $|g^{-1}Q_\lambda g| = |P|$. Now, since $|P| \geq |Q|$ and $|\mu_k| \geq |\lambda|$ it follows that μ_k is $g^{-1}Q_\lambda g$ -periodic. The statement of the lemma for all bases in σ_k of the length at least $2|P|$ follows automatically now. □

Remark 3. δ From the construction of $\mathcal{P}(U, P)$ and Lemma 13 it follows that if $\sigma_i \in \mathcal{P}$ with an overlapping pair $(\mu_i, \bar{\mu}_i)$ and if $l \in [\alpha(\sigma_i), \beta(\sigma_i)]$ then there exists a period Q_l conjugate in G either to P or P^{-1} such that

- (i) if $ht([\alpha(\sigma_i), l]) > ht(P)$ then $U[\alpha(\sigma_i), l]$ has any natural power of Q_l as a terminal subword,
- (ii) if $ht([l, \beta(\sigma_i)]) > ht(P)$ then $U[1, \beta(\sigma_i)]$ has any natural power of Q_l as an initial subword,
- (iii) if $ht([\alpha(\sigma_i), l]) \leq ht(P)$ then $U[\alpha(\sigma_i), l] = Q'' \circ Q_l^{k_l}$, where $Q_l = Q' \circ Q''$,
- (iv) if $ht([l, \beta(\sigma_i)]) \leq ht(P)$ then $U[1, \beta(\sigma_i)] = Q_l^{k_l} \circ Q'$, where $Q_l = Q' \circ Q''$.

Lemma 14. $\mathcal{P}(U, P)$ corresponding to a solution U of Ω' is a periodic structure on Ω' .

Proof. Parts 1(a), 1(b) and 1(c) of the definition of periodic structure hold by construction.

Set $\mathcal{X}(\sigma_1) = 1$. If there is a base $\lambda \in \sigma_1$ such that $\bar{\lambda} \in \sigma_2$ then we set $\mathcal{X}(\sigma_2) = \varepsilon(\lambda)\varepsilon(\bar{\lambda})$. Observe that the sign of $\mathcal{X}(\sigma_2)$ is well-defined since if there exists another base $\lambda_1 \in \sigma_1$ such that $\bar{\lambda}_1 \in \sigma_2$ and $\varepsilon(\lambda)\varepsilon(\bar{\lambda}) \neq \varepsilon(\lambda_1)\varepsilon(\bar{\lambda}_1)$ then it follows that there exists a group element conjugating P into P^{-1} which is impossible. Hence 1(d) holds.

By Remark 3 if boundaries l_1, l_2 belong to sections from \mathcal{P} then there exist Q_{l_1}, Q_{l_2} conjugate in G either to P or P^{-1} such that some of the conditions listed in the remark hold. Thus, we define a relation R as follows: $l_1 \sim_R l_2$ if and only if $Q_{l_1} = Q_{l_2}$, and part 2 follows immediately. \square

Now let us fix a nonempty periodic structure $\langle \mathcal{P}, R \rangle$. Item (d) allows us to assume (after replacing the variables h_i, \dots, h_{j-1} by $h_{j-1}^{-1}, \dots, h_i^{-1}$ on those sections $[i, j] \in \mathcal{P}$ for which $\mathcal{X}([i, j]) = -1$) that $\varepsilon(\mu) = 1$ for all $\mu \in \mathcal{P}$. For a boundary k , we will denote by (k) the equivalence class of the relation R to which it belongs.

Let us construct an oriented graph Γ whose set of vertices is the set of R -equivalence classes. For each long h_k we introduce an oriented edge e leading from (k) to $(k+1)$ and an inverse edge e^{-1} leading from $(k+1)$ to (k) . This edge e is assigned the label $h(e) = h_k$ (respectively, $h(e^{-1}) = h_k^{-1}$). For every path $r = e_1^{\pm 1} \cdots e_s^{\pm 1}$ in the graph Γ , we denote by $h(r)$ its label $h(e_1^{\pm 1}) \cdots h(e_s^{\pm 1})$.

The periodic structure $\langle \mathcal{P}, R \rangle$ is called *connected*, if the graph Γ is connected. Suppose first that $\langle \mathcal{P}, R \rangle$ is connected. We can also suppose that each boundary of Ω' is a boundary between two bases.

Lemma 15. Let U be a solution of Ω' and $\langle \mathcal{P}, R \rangle = \mathcal{P}(U, P)$. If p is a path in the graph Γ from (i) to (j) then $U[p]^{-1}Q_i U[p] = Q_j$ in G .

Proof. If $p = h_{i_1} \cdots h_{i_k}$ then all products between U_j , $j \in \{i_1, \dots, i_k\}$ are defined and the required result follows from Lemma 11. \square

Lemma 16. Let U be a solution of Ω' and $\langle \mathcal{P}, R \rangle = \mathcal{P}(U, P)$. If c_1, c_2 are cycles in the graph Γ at the vertex $v = (k)$ then $[U[c_1], U[c_2]] = \varepsilon$ in G .

Proof. Assume at first that $(k) = (1)$. There exists a loop c at (1) such that $U[c] = P$. By Lemma 15 $U[p]^{-1}PU[p] = P$ in G for every path p from (1) to (1) , so $[U[c_1], P] = [U[c_2], P] = 1$ and from CSA-property of G the required follows.

If $(k) \neq (1)$ then, since Γ is connected it follows that there exists a path p from v to v_0 . Moreover, by Lemma 15 we have $U[p]^{-1}Q_k U[p] = P$ in G . So, by CSA-property commutativity of cycles at (k) follows from commutativity of cycles at (1) which is already proved. \square

Ω' is called *periodized* with respect to a given periodic structure $\langle \mathcal{P}, R \rangle$ if for every two cycles c_1 and c_2 with the same initial vertex in the graph Γ there is a relation $[h(c_1), h(c_2)] = 1$ in $G_{R(\Omega')}$. Observe that according to Lemma 16 Ω' is periodized with respect to $\mathcal{P}(U, P)$ for any solution U .

Let Γ_0 be the subgraph of the graph Γ having the same set of vertices and consisting of the edges e whose labels do not belong to \mathcal{P} . Choose a maximal subforest T_0 in the graph Γ_0 and extend it to a maximal subforest T of the graph Γ . Since $\langle \mathcal{P}, R \rangle$ is connected by assumption, it follows that T is a tree. Let v_0 be an arbitrary vertex of the graph Γ and $r(v_0, v)$ the (unique) path from v_0 to v all of whose vertices belong to T . For every edge $e : v \rightarrow v'$ not lying in T , we introduce a cycle $c_e = r(v_0, v)e(r(v_0, v'))^{-1}$. Then the fundamental group $\pi_1(\Gamma, v_0)$ is generated by the cycles c_e (see, for example, the proof of Proposition 3.2.1 [24]).

Furthermore, the set of elements

$$\{h(e) \mid e \in T\} \cup \{h(c_e) \mid e \notin T\} \quad (3)$$

forms a basis of the free group with the set of generators $\{h_k \mid h_k \in \sigma\}$. If $\lambda \in \mathcal{P}$, then $(\beta(\lambda)) = (\beta(\bar{\lambda}))$, $(\alpha(\lambda)) = (\alpha(\bar{\lambda}))$ by the definition of R and, consequently, the word

$$h[\alpha(\lambda), \beta(\lambda)] h[\alpha(\bar{\lambda}), \beta(\bar{\lambda})]^{-1}$$

is the label of a cycle $c'(\lambda)$ from $\pi_1(\Gamma, (\alpha(\lambda)))$. Let $c(\lambda) = r(v_0, (\alpha(\lambda))) c'(\lambda) r(v_0, (\alpha(\lambda)))^{-1}$. Then

$$h(c(\lambda)) = u h[\alpha(\lambda), \beta(\lambda)] h[\alpha(\bar{\lambda}), \beta(\bar{\lambda})]^{-1} u^{-1}, \quad (4)$$

where u is a certain word. Since $c(\lambda) \in \pi_1(\Gamma, v_0)$, it follows that $c(\lambda) = b_\lambda(\{c_e \mid e \notin T\})$, where b_λ is a certain word in the indicated generators which can be effectively constructed (see Proposition 3.2.1 [24]).

Let \tilde{b}_λ denote the image of the word b_λ in the abelianization of $\pi(\Gamma, v_0)$. Denote by \tilde{Z} the free abelian group consisting of formal linear combinations $\sum_{e \notin T} n_e \tilde{c}_e$ ($n_e \in \mathbb{Z}$), and by \tilde{B} its subgroup generated by the elements \tilde{b}_λ ($\lambda \in \mathcal{P}$) and the elements \tilde{c}_e ($e \notin T$, $h(e) \notin \mathcal{P}$). Let $\tilde{A} = \tilde{Z}/\tilde{B}$, $T(\tilde{A})$ the torsion subgroups of the group \tilde{A} , and \tilde{Z}_1 the preimage of $T(\tilde{A})$ in \tilde{Z} . The group \tilde{Z}/\tilde{Z}_1 is free; therefore, there exists a decomposition of the form

$$\tilde{Z} = \tilde{Z}_1 \oplus \tilde{Z}_2, \quad \tilde{B} \subseteq \tilde{Z}_1, \quad (\tilde{Z}_1 : \tilde{B}) < \infty. \quad (5)$$

Note that it is possible to express effectively a certain basis $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ of the group \tilde{Z} in terms of the generators \tilde{c}_e so that for the subgroups \tilde{Z}_1 and \tilde{Z}_2 generated respectively by $\tilde{c}^{(1)}$ and $\tilde{c}^{(2)}$, the relation (5) holds. It suffices, for instance, to look through the bases one by one, using the fact that under the condition $\tilde{Z} = \tilde{Z}_1 \oplus \tilde{Z}_2$ the relations $\tilde{B} \subseteq \tilde{Z}_1$, $(\tilde{Z}_1 : \tilde{B}) < \infty$ hold if and only if the generators of the groups \tilde{B} and \tilde{Z}_1 generate the same linear subspace over \mathbf{Q} , and the latter is easily verified algorithmically. Notice, that a more economical algorithm can be constructed by analyzing the proof of the classification theorem for finitely generated abelian groups. By Proposition 1.4.4 [24], one can effectively construct a basis $\bar{c}^{(1)}, \bar{c}^{(2)}$ of the free (non-abelian) group $\pi_1(\Gamma, v_0)$ so that $\tilde{c}^{(1)}, \tilde{c}^{(2)}$ are the natural images of the elements $\bar{c}^{(1)}, \bar{c}^{(2)}$ in \tilde{Z} .

Now assume that $\langle \mathcal{P}, R \rangle$ is an arbitrary periodic structure on Ω' , not necessarily connected. Let $\Gamma_1, \dots, \Gamma_r$ be the connected components of the graph Γ . The labels of edges of the component Γ_i form in Ω' a union of closed sections from \mathcal{P} ; moreover, if a base $\lambda \in \mathcal{P}$ belongs to such a section, then its dual $\bar{\lambda}$ also does by definition of periodic structure. Therefore, by taking for \mathcal{P}_i the set of labels of edges from Γ_i belonging to \mathcal{P} , sections to which these labels belong, and bases $\lambda \in \mathcal{P}$ belonging to these sections, and restricting in the corresponding way the relation R , we obtain a periodic connected structure $\langle \mathcal{P}_i, R_i \rangle$ with the graph Γ_i .

The notation $\langle \mathcal{P}', R' \rangle \subseteq \langle \mathcal{P}, R \rangle$ means that $\mathcal{P}' \subseteq \mathcal{P}$, and the relation R' is a restriction of the relation R . In particular, $\langle \mathcal{P}_i, R_i \rangle \subseteq \langle \mathcal{P}, R \rangle$ in the situation described in the previous paragraph. Since Ω' is periodized, the periodic structure must be connected.

Let e_1, \dots, e_m be all the edges of the graph Γ from $T - T_0$. Since T_0 is the spanning forest of the graph Γ_0 , it follows that $h(e_1), \dots, h(e_m) \in \mathcal{P}$. Let $F(\Omega')$ be a free group generated by the variables of Ω' . Consider in the group $F(\Omega')$ a new basis

$$Y = \{\bar{t}, \{h(e) \mid e \in T\}, h(\bar{c}^{(1)}), h(\bar{c}^{(2)})\},$$

where variables \bar{t} do not belong to the closed sections from \mathcal{P} . Let v_i be the initial vertex of the edge e_i . We introduce new variables $\bar{u}^{(i)} = \{u_{ie} \mid e \notin T, e \notin \mathcal{P}\}$, $\bar{z}^{(i)} = \{z_{ie} \mid e \notin T, e \notin \mathcal{P}\}$ for $1 \leq i \leq m$, as follows

$$u_{ie} = h(r(v_0, v_i))^{-1} h(c_e) h(r(v_0, v_i)), \quad (6)$$

$$h(e_i)^{-1} u_{ie} h(e_i) = z_{ie}. \quad (7)$$

Without loss of generality we can assume $v_0 = (1)$.

Lemma 17. *Let Ω' be periodized with respect to a periodic structure $\langle \mathcal{P}, R \rangle$. Let K be the subgroup of $G_{R(\Omega')}$ generated by*

$$Y_0 = \{\bar{t}, \{h(e) \mid e \in T_0\}, h(\bar{c}^{(1)}), \bar{u}^{(i)}, \bar{z}^{(i)}, i \in [1, m]\}.$$

If $|\bar{c}^{(2)}| = s \geq 1$ then the group $G_{R(\Omega')}$ splits as a fundamental group of a graph of groups with two vertices, where one vertex group is K and the other is a free abelian group generated by $h(\bar{c}^{(2)}), h(\bar{c}^{(1)})$. The corresponding edge group is generated by $h(\bar{c}^{(1)})$. The other edges are loops at the vertex with vertex group K , have stable letters $h(e_i)$, $i \in [1, m]$, and associated subgroups $\langle \bar{u}^i \rangle$, $\langle \bar{z}^i \rangle$. If $\bar{c}^{(2)} = \emptyset$ then there is no vertex with abelian vertex group.

Proof. We are going to study in more detail how the variables $h(e_i)$, $i \in [1, m]$ can participate in the equations from Ω' rewritten in the set of variables Y .

If h_k does not lie on a closed section from \mathcal{P} , or $h_k \notin \mathcal{P}$, but $e \in T$ (where $h(e) = h_k$), then h_k belongs to the basis Y and is distinct from each of $h(e_1), \dots, h(e_m)$. Now let $h(e) = h_k$, $h_k \notin \mathcal{P}$ and $e \notin T$. Then $e = r_1 c_e r_2$, where r_1, r_2 are paths in T . Since $e \in \Gamma_0$, $h(c_e)$ belongs to $\langle c^{(1)} \rangle$ modulo commutation of cycles. The vertices (k) and $(k+1)$ lie in the same connected component of the graph Γ_0 , and hence they are connected by a path s in the forest T_0 . Furthermore, r_1 and $s r_2^{-1}$ are paths in the tree T connecting the vertices (k) and v_0 ; consequently, $r_1 = s r_2^{-1}$. Thus, $e = s r_2^{-1} c_e r_2$ and $h_k = h(s) h(r_2)^{-1} h(c_e) h(r_2)$. The variable $h(e_i)$, $i \in [1, m]$ can occur in the right-hand side of the expression obtained (written in the basis Y) only in $h(r_2)$ and at most once. Moreover, the sign of this occurrence (if it exists) depends only on the orientation of the edge e_i with respect to the root v_0 of the tree T . If $r_2 = r_2' e_i^{\pm 1} r_2''$, then all the occurrences of the variable $h(e_i)$ in the words h_k written in the basis Y , with $h_k \notin \mathcal{P}$, are contained in the occurrences of words of the form $h(e_i)^{\mp 1} h((r_2')^{-1} c_e r_2') h(e_i)^{\pm 1}$, that is, in occurrences of the form $h(e_i)^{\mp 1} h(c) h(e_i)^{\pm 1}$, where c is a certain cycle of the graph Γ starting at the initial vertex of the edge $e_i^{\pm 1}$.

Therefore all the occurrences of $h(e_i)$, $i \in [1, m]$ in the equations corresponding to $\lambda \notin \mathcal{P}$ are of the form $h(e_i^{-1}) h(c) h(e_i)$. Also, $h(e_i)$ does not occur in the equations corresponding to $\lambda \in \mathcal{P}$ in the basis Y . The system Ω' is equivalent to the following system in the variables Y :

$$u_{ie} = h(r(v_0, v_i))^{-1} h(c_e) h(r(v_0, v_i)),$$

$$h(e_i)^{-1} u_{ie} h(e_i) = z_{ie},$$

$$[u_{ie_1}, u_{ie_2}] = 1,$$

$$[h(c_1), h(c_2)] = 1, \quad c_1, c_2 \in c^{(1)}, c^{(2)},$$

$$\bar{\psi}(Y_0) = 1,$$

where $\bar{\psi}(Y_0)$ does not contain $h(e_i)$, $\bar{c}^{(2)}$. Let $K = G_{R(\bar{\psi})}$. Then to obtain $G_{R(\Omega)}$ we first take an HNN extension of the group K with abelian associated subgroups generated by $\bar{u}^{(i)}$ and $\bar{z}^{(i)}$ and stable letters $h(e_i)$, and then extend the centralizer of the image of $\langle \bar{c}^{(1)} \rangle$ by the free abelian subgroup generated by the images of $\bar{c}^{(2)}$. \square

(d) Abelian splitting: long shift. If Ω does not satisfy (a), (c) we perform QH-shortening, then apply the entire transformation and then, if possible, (D7).

Lemma 18. *Let*

$$v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_r \rightarrow \cdots$$

be an infinite path in $T(\Omega)$. Then there exists a natural number N such that all the generalized equations in vertices v_n , $n \geq N$ satisfy the general JSJ case (d).

Proof. Indeed, the Tietze cleaning either replaces the group by its proper free factor or decreases the complexity. Every time when the case (a) holds we replace G by some vertex group in a non-trivial abelian splitting of G . This can be done only finitely many times [DP]. Every time when case c) takes place, we decrease the complexity.

Proposition 3. *The general JSJ case (d) cannot be repeated infinitely many times.*

Proof. Consider an infinite path

$$r = v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_m. \quad (8)$$

We have $\tau'_{v_i} = \tau'_v$.

Denote by μ_i the carrier base of the equation Ω_{v_i} . The path (8) will be called μ -reducing if $\mu_1 = \mu$ and either μ_2 does not overlap with its double and μ occurs in the sequence μ_1, \dots, μ_{m-1} at least twice.

The path (8) will be called *prohibited*, if it can be represented in the form

$$r = r_1 s_1 \cdots r_l s_l r', \quad (9)$$

such that for some sequence of bases η_1, \dots, η_l the following three properties hold:

- 1) every base occurring at least once in the sequence μ_1, \dots, μ_{m-1} occurs at least $4n(1 + f_1)(\Omega_{v_2})$ times in the sequence η_1, \dots, η_l , where n is the number of pairs of bases in equations Ω_{v_i} ,
- 2) the path r_i is η_i -reducing,
- 3) every transfer base of some equation of path r is a transfer base of some equation of path r' .

Every infinite path contains a prohibited subpath. Indeed, let ω be the set of all bases occurring in the sequence $\mu_1, \dots, \mu_m, \dots$ infinitely many times, and $\tilde{\omega}$ the set of all bases, that are transfer bases of infinitely many equations Ω_{v_i} . If one cuts out some finite part in the beginning of this infinite path, one can suppose that all the bases in the sequence $\mu_1, \dots, \mu_m, \dots$ belong to ω and each base that is a transfer base of at least one equation, belongs to $\tilde{\omega}$. Such an infinite path for any $\mu \in \omega$ contains infinitely many non-intersecting μ -reducing finite subpaths. Hence it is possible to construct a subpath (9) of this path satisfying the first two conditions in the definition of a prohibited subpath. Making r' longer, one obtains a prohibited subpath.

Suppose we have a prohibited path (8). As before, let μ_i denote the carrier base of Ω_{v_i} , and $\omega = \{\mu_1, \dots, \mu_{m-1}\}$, and $\tilde{\omega}$ denote the set of such bases which are transfer bases for at least one equation in (8). By ω_1 denote the set of such bases μ for which either μ or $\bar{\mu}$ belongs to $\omega \cup \tilde{\omega}$; by ω_2 denote the set of all the other bases. Let

$$\alpha(\omega) = \min(\min_{\mu \in \omega_2} \alpha(\mu), j),$$

where j is the boundary between active and non-active sections.

Let

$$X_\mu \triangleq U[\alpha(\mu), \beta(\mu)].$$

If (Ω_{v_i}, U) corresponds to a vertex in the sequence (8), then denote

$$|U_\omega| = \sum_{i=1}^{\alpha(\omega)-1} |U_i|, \quad (10)$$

$$\psi_\omega(U) = \sum_{\mu \in \omega_1} |X_\mu| - 2|U|_\omega. \quad (11)$$

We call $\psi_\omega(U)$ the *excess* of Ω . Since case (a) is not applicable, $\psi_\omega(U) \geq 0$ and is comparable with the length of the section $[1, \alpha(\omega)]$. Consider the quadratic part of Ω_{v_1} which is situated to the left of $\alpha(\omega)$. Let A be the length of the quadratic part and B be the length of the non-quadratic part.

Lemma 19. *Suppose there exists a function $f_1(\Omega_{v_1})$ such that we have an inequality*

$$A \leq f_1 B. \quad (12)$$

Then there is a function $f_2(\Omega_{v_1})$ such that

$$|U_\omega| \leq f_2 \psi_\omega(U). \quad (13)$$

Proof. $|U_\omega| = A + B \leq B(1 + f_1) \leq (1 + f_1)\psi_\omega(U)$. \square

From the definition of the process it follows that all the words $U^{(i)}[1, \rho_i + 1]$ are the ends of the word $U^{(1)}[1, \rho_1 + 1]$, that is

$$U^{(1)}[1, \rho_1 + 1] \doteq u_i U^{(i)}[1, \rho_i + 1]. \quad (14)$$

On the other hand bases $\mu \in \omega_2$ participate in these transformations neither as carrier bases nor as transfer bases; hence $U^{(1)}[\alpha(\omega), \rho_1 + 1]$ is the end of the word $U^{(i)}[1, \rho_i + 1]$, that is

$$U^{(i)}[1, \rho_i + 1] \doteq v_i U^{(1)}[\alpha(\omega), \rho_1 + 1]. \quad (15)$$

So we have

$$|U_\omega^{(i)}| - |U_\omega^{(i+1)}| = |v_i| - |v_{i+1}| = |u_{i+1}| - |u_i| = |X_{\mu_i}^{(i)}| - |X_{\mu_i}^{(i+1)}|. \quad (16)$$

In particular (11),(16) imply that $\psi_\omega(U^{(1)}) = \psi_\omega(U^{(2)}) = \dots = \psi_\omega(U^{(m)}) = \psi_\omega$. Denote the number (16) by δ_i .

Let the path (8) be μ -reducing, Estimate $d(U_m) = \sum_{i=1}^{m-1} \delta_i$ from below. First notice that if $\mu_{i_1} = \mu_{i_2} = \mu$ ($i_1 < i_2$) and $\mu_i \neq \mu$ for $i_1 < i < i_2$, then

$$\sum_{i=i_1}^{i_2-1} \delta_i \geq |U^{i_1+1}[1, \alpha(\Delta(\mu_{i_1+1}))]|. \quad (17)$$

Indeed, if $i_2 = i_1 + 1$, then $\delta_{i_1} = d(U^{(i_1)}[1, \alpha(\Delta(\mu))]) = d(U^{(i_1+1)}[1, \alpha(\Delta(\mu))])$. If $i_2 > i_1 + 1$, then $\mu_{i_1+1} \neq \mu$ and μ is a transfer base in the equation $\Omega_{v_{i_1+1}}$. Hence $\delta_{i_1+1} + |U^{(i_1+2)}[1, \alpha(\mu)]| = |U^{(i_1+1)}[1, \alpha(\mu_{i_1+1})]|$. Now (17) follows from

$$\sum_{i=i_1+2}^{i_2-1} \delta_i \geq |U^{(i_1+2)}[1, \alpha(\mu)]|.$$

So if the bases μ_2 and $\bar{\mu}_2$ do not intersect in the equation Ω_{v_2} ; then (17) implies that

$$\sum_{i=1}^{m-1} \delta_i \geq |U^{(2)}[1, \alpha(\Delta\mu_2)]| \geq |X_{\mu_2}^{(2)}| \geq |X_\mu^{(2)}| = |X_\mu^{(1)}| - \delta_1,$$

which implies that

$$\sum_{i=1}^{m-1} \delta_i \geq \frac{1}{2} |X_\mu^{(1)}|. \quad (18)$$

We obtain the same inequality if μ_2 overlaps with $\bar{\mu}_2$ but $|\mu_2| \leq |\alpha(\mu_2) - \alpha(\bar{\mu}_2)|$.

Suppose now that the path (8) is prohibited; hence it can be represented in the form (9). From definition (11) we have $\sum_{\mu \in \omega_1} d(X_\mu^{(m)}) \geq \psi_\omega$; so at least for one base $\mu \in \omega_1$ the inequality $d(X_\mu^{(m)}) \geq \frac{1}{2n} \psi_\omega$ holds. Because $X_\mu^{(m)} \doteq (X_{\Delta(\mu)}^{(m)})^{\pm 1}$, we can suppose that $\mu \in \omega \cup \tilde{\omega}$. Let m_1 be the length of the path $r_1 s_1 \cdots r_l s_l$ in (9). If $\mu \in \tilde{\omega}$ then by the third part of the definition of a prohibited path there exists $m_1 \leq i \leq m$ such that μ is a transfer base of Ω_{v_i} . Hence, $d(X_{\mu_i}^{(m_1)}) \geq d(X_{\mu_i}^{(i)}) \geq d(X_{\mu_i}^{(i)}) \geq d(X_\mu^{(m)}) \geq \frac{1}{2n} \psi_\omega$. If $\mu \in \omega$, then take μ instead of μ_i . We proved the existence of a base $\mu \in \omega$ such that

$$|X_\mu^{(m_1)}| \geq \frac{1}{2n} \psi_\omega. \quad (19)$$

By the definition of a prohibited path, inequality (18) and inequality $d(X_\mu^{(i)}) \geq d(X_\mu^{(m_1)})$ ($1 \leq i \leq m_1$), we obtain

$$\sum_{i=1}^{m_1-1} \delta_i \geq \max\left\{\frac{1}{4n} \psi_\omega, 1\right\} (4n f_2). \quad (20)$$

By (16) the sum in the left part of the inequality (20) equals $d_\omega(U^{(1)}) - d_\omega(U^{(m_1)})$; hence

$$|U_\omega^{(1)}| \geq \max\left\{\frac{1}{4n} \psi_\omega, 1\right\} (4n f_2) = \psi_\omega f_2,$$

which, in case of inequality 12, contradicts (13). This contradiction was obtained from the supposition that there are prohibited paths (8). Hence there are no prohibited paths. The proposition is proved.

Proof of Theorem 1. We perform the elimination process for the generalized equation Ω corresponding to G and obtain a decomposition of G as the fundamental group of a graph of groups with QH vertex groups corresponding to quadratic sections (4.4.3 (a)), abelian vertex groups corresponding to periodic structures (4.4.3 (c)) and HNN-extensions with stable letters infinitely longer than the generators of the abelian associated subgroups (4.4.3 (c) and 4.4.1 for discrete case). We also can get free splittings of G with non-trivial free factors (4.4.1) and surface group factors (4.4.2). On this step we will have a system Σ of homogeneous linear equations with integer coefficients on the lengths of certain items. Then we continue the elimination process with the generalized equation corresponding to rigid subgroups. They are smaller in Delzant-Potyagailo hierarchy. At the end we obtain some generalized equation Ω_{fin} . Continuing the elimination process till the end, we obtain the complete system of linear equations with integer coefficients $\Sigma_{complete}$ on the lengths of certain items.

Lemma 20. *A free length function on $G_{\Omega_{fin}}$ can be defined into \mathbb{Z}^n .*

Proof. Recall that in case (a) there is a quadratic base that is infinitely longer than all the quadratic coefficient bases. In case 1 of (c), there is also a long base that is infinitely longer than all the coefficient bases. In case 2 of (c), either the length of the overlapping base is not connected by any equation in $\Sigma_{complete}$ with the length of the shift, or this case can be treated as case (d). We first find a non-negative solution ϕ of the system $\Sigma_{complete}$ in \mathbb{Z}^n satisfying the following conditions:

1. For each quadratic section and overlapping section there exists a variable h such that $\phi(h)$ is infinitely larger than all the coefficient variables on this section;

2. For each section corresponding to a free component and infinite linear elimination there exists a variable h such that $\phi(h) \neq 0$;

3. If $\ell(h_i)$ is infinitely smaller than $\ell(h_j)$, then $\phi(h_i)$ is infinitely smaller than $\phi(h_j)$.

Then we adjust other components of new lengths so that the group were isomorphic to G . This is possible because to make $G \cong G_\Omega$ we just have to make sure that some boundaries of the generalized equations with the new length function do not coincide. This fact can be expressed by existential sentences. And we can use proposition 2.


Since $G_{\Omega_{fin}}$ is obtained from a free group by a series of extensions of HNN-extensions such that associated subgroups are abelian and length isomorphic, and stable letters have length infinitely larger than generators of the associated subgroups, the free regular length function into \mathbb{Z}^m can be constructed as in [20]. If Ω does not have pairs of flipped bases, $G = G_\Omega = G_{\Omega_{fin}}$ and G has a free regular length function into \mathbb{Z}^m .

If Ω has pairs of flipped bases, then for a subgroup H of G of index 2, $H * C = G_{\Omega_H}$, where C is an infinite cyclic group. Moreover, there is an epimorphism $\phi : G_{\Omega_H} \rightarrow G_{\Omega_{fin}}$ and the restriction of ϕ to H is a monomorphism. This epimorphism is not necessary an isomorphism because we apply the elimination process to a solution of Ω_H in infinite words, and the same words represent bases λ_i and δ_i for $i = 1, \dots, k$ from proposition 1. But the natural homomorphism from G_{Ω_H} onto G identifying λ_i and δ_i for $i = 1, \dots, k$ factors through ϕ and maps H to the isomorphic index two subgroup of G . Therefore H has a free Lyndon's length function in \mathbb{Z}^m for some m . Theorem 1 is proved. \square


References


- [1] R. Alperin and H. Bass, *Length functions of group actions on Λ -trees*. Combinatorial group theory and topology, (Ed. S. M. Gersten and J. R. Stallings), Annals of Math. Studies **111**, 265–378. Princeton University Press, 1987.
- [2] R. Alperin and K. Moss, *Complete trees for groups with a real length function*. J. London Math. Soc. (2) **31**, 1985, 55–68.
- [3] H. Bass *Groups acting on n -non-archimedean trees*. Arboreal group theory, 1991, 69–130.
- [4] M. Bestvina and M. Feighn, *Stable actions of groups on real trees*. Invent. Math., **121** no. 2 (1995), 287–321.
- [5] I. Chiswell, *Abstract length functions in groups*. Math. Proc. Cambridge Philos. Soc., **80** no. 3 (1976), 451–463.
- [6] I. Chiswell, *Introduction to Λ -trees*. World Scientific, 2001.
- [7] T. Delzant and L. Potyagailo, *Accessibilité hiérarchique des groupes de présentation finie*. (French) Topology **40** no. 3 (2001), 617–629.
- [8] D. Gildenhuys, O. Kharlampovich and A. Myasnikov, *CSA-groups and separated free constructions*. Bull. Austral. Math. Soc., **52** no. 1 (1995), 63–84.
- [9] D. Gaboriau, G. Levitt and F. Paulin, *Pseudogroups of isometries of \mathbb{R} and Rips' Theorem on free actions on \mathbb{R} -trees*. Israel. J. Math., **87** (1994), 403–428.
- [10] A. M. W. Glass, *Partially ordered groups*. Series in Algebra, **7**, World Scientific, 1999.
- [11] D. Groves, *Makanin-Razborov diagrams for relatively hyperbolic groups*. Preprint.
- [12] V. Guirardel, *Limit groups and groups acting freely on \mathbb{R}^n -trees*. Geom. Topol., **8** (2004), 1427–1470.

- [13] Ju. Gurevich, A.I. Kokorin, *Universal equivalence of ordered Abelian groups. (Russian)*, Algebra i Logika Sem., 2 1963 no. 1, p. 37–39.
- [14] N. Harrison, *Real length funtions in groups*. Trans. Amer. Math. Soc. **174** (1972), 77–106.
- [15] A. H. M. Hoare, *On length functions and Nielsen methods in free groups*. J. London Math. Soc. (2) **14** (1976), 188–192.
- [16] A. H. M. Hoare, *Nielsen method in groups with a length function*. Math. Scand. **48** (1981), 153–164.
- [17] B. Khan, A.G. Myasnikov and D. Serbin, *On positive theories of groups with regular free length functions*. to appear in Internat. J. Algebra and Comput.
- [18] O. Kharlampovich, A.G. Myasnikov, *Irreducible affine varieties over a free group. II: Systems in triangular quasi-quadratic form and description of residually free groups*. J. of Algebra **200** no. 2 (1998), 517–570.
- [19] O. Kharlampovich, A.G. Myasnikov, *Implicit function theorems over free groups*. J. of Algebra, **290** no. 1 (2005), 1–203.
- [20] O. Kharlampovich, A. Myasnikov, V. Remeslennikov, and D. Serbin *Groups with free regular length functions in \mathbb{Z}^n* . arXiv:0907.2356.
- [21] O. Kharlampovich, A. Myasnikov, D. Serbin *Free regular length functions in \mathbb{Z}^n II: Geometric approach*. Preprint, 2009.
- [22] V. Kopytov and N. Medvedev, *Right-ordered groups*. Siberian School of Algebra and Logic. Consultants Bureau, New York, 1996.
- [23] R. Lyndon, *Length functions in groups*. Math. Scand., **12** (1963), 209–234.
- [24] R.C. Lyndon, and P.E. Schupp, *Combinatorial group theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete 89, Springer-Verlag, Berlin, Heidelberg, New York, 1977.
- [25] G.S. Makanin, *Equations in a free group*. (Russian), Izv. Akad. Nauk SSSR, Ser. Mat., **46** (1982), 1199–1273 (transl. in Math. USSR Izv. **21** (1983)).
- [26] A. Myasnikov, V. Remeslennikov, *Length functions on free exponential groups*. Proc. N 26. IITPM SO RAN, Omsk, 1996, 1–34.
- [27] A. Myasnikov, V. Remeslennikov and D. Serbin, *Regular free length functions on Lyndon’s free $\mathbb{Z}[t]$ -group $F^{\mathbb{Z}[t]}$, Algorithms, Languages, Logic* (Ed. A. Borovik), Contemp. Math., Amer. Math. Soc. **378** (2005), 37–77.
- [28] J. Morgan and P. Shalen, *Valuations, Trees, and Degenerations of Hyperbolic Structures, I*. Annals of Math, 2nd Ser., **120** no. 3. (1984), 401–476.
- [29] D. Promislow, *Equivalence classes of length functions on groups*. Proc. London Math. Soc. (3) **51** (1985), 449–477.
- [30] A. Razborov, *On systems of equations in a free group*. Math. USSR Izvestiya **25** no. 1 (1985), 115–162.
- [31] J.-P. Serre, *Trees*. New York, Springer, 1980.





"TRANSFER BASE"





ET3





"DELETE LONE BASE"

