

# On several families of elliptic curves with arbitrary large Selmer groups

Fei Li and Derong Qiu <sup>\*</sup>

(School of Mathematical Sciences,  
Institute of Mathematics and Interdisciplinary Science,  
Capital Normal University, Beijing 100048, P.R.China )

## 1 Introduction and Main Results

In this paper, we consider the elliptic curve

$$E = E_\varepsilon : y^2 = x(x + \varepsilon pD)(x + \varepsilon qD), \quad (\varepsilon = \pm 1) \quad (1.1)$$

where  $p$  and  $q$  are odd prime numbers with  $q - p = 2$ , and  $D = D_1 \cdots D_n$  is a square-free integer with distinct primes  $D_1, \dots, D_n$ . Moreover,  $2 \nmid D$ ,  $p \nmid D$  and  $q \nmid D$ . For each  $D_i$ , denote  $\widehat{D}_i = D/D_i$  ( $\widehat{D}_1 = 1$  if  $D = D_1$ ). We write  $E = E_+$  if  $\varepsilon = 1$  and  $E = E_-$  if  $\varepsilon = -1$ .

Take the elliptic curve

$$E' = E'_\varepsilon : y^2 = x^3 - 2\varepsilon(p + q)Dx^2 + 4D^2x. \quad (1.2)$$

There is an isogeny  $\varphi$  of degree 2 between  $E$  and  $E'$  as follows:

$$\varphi : E \longrightarrow E', \quad (x, y) \longmapsto (y^2/x^2, y(pqD^2 - x^2)/x^2).$$

The kernel is  $E[\varphi] = \{O, (0, 0)\}$ , and the dual isogeny of  $\varphi$  is

$$\widehat{\varphi} : E' \longrightarrow E, \quad (x, y) \longmapsto (y^2/4x^2, y(4D^2 - x^2)/8x^2)$$

---

<sup>\*</sup> E-mail: derong@mail.cnu.edu.cn

with kernel  $E'[\widehat{\varphi}] = \{O, (0, 0)\}$  (see [S, p.74]).

In this paper, we calculate the  $\varphi(\widehat{\varphi})$ -Selmer groups  $S^{(\varphi)}(E/\mathbb{Q})$  and  $S^{(\widehat{\varphi})}(E'/\mathbb{Q})$  of the above elliptic curves via descent theory (see [S, Chapter X]), in particular, we obtain that the Selmer groups of several families of such elliptic curves can be arbitrary large. There are many literature studying Selmer groups of elliptic curves ( see e.g., [KS], [QZ], [Sch], [ST]). The main results in the present paper are stated as follows:

### Part 1 Results for the case $\varepsilon = +1$

**Theorem 1.1.** Let  $E = E_+$  be the elliptic curve in (1.1) with  $\varepsilon = +1$ . For each  $i \in \{1, \dots, n\}$ , denote

$$\Pi_i^+(D) = \left(1 - \left(\frac{-1}{D_i}\right)\right) + \left(1 - \left(\frac{p}{D_i}\right)\right) + \left(1 - \left(\frac{q}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_j}{D_i}\right)\right),$$

where  $(-)$  is the ( Legendre ) quadratic residue symbol. And define a function  $\rho^+(D)$  by

$$\rho^+(D) = \sum_{i=1}^n \left[ \frac{1}{1 + \Pi_i^+(D)} \right],$$

where  $[x]$  is the greatest integer  $\leq x$ . Then we have

(1) There exists a subset  $T \subset \{D_1, \dots, D_n\}$  with cardinal  $\#T = \rho^+(D)$  such that  $S^{(\varphi)}(E/\mathbb{Q}) \supset \langle T \bmod(\mathbb{Q}^{\star^2}) \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)}$ . In particular,  $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^+(D)$ .

(2) If  $p \equiv 7 \pmod{8}$  and  $D_i \equiv 1, 7 \pmod{8}$  ( $1 \leq i \leq n$ ), then  $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^+(D) + 1$  and  $S^{(\varphi)}(E/\mathbb{Q}) \supset \langle T \cup \{2\} \bmod(\mathbb{Q}^{\star^2}) \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)+1}$  for some subset  $T \subset \{D_1, \dots, D_n\}$  with  $\#T = \rho^+(D)$ .

**Corollary 1.2.** For elliptic curve  $E = E_+$  as in Theorem 1.1, we have

$S^{(\varphi)}(E/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+1}$ . Furthermore,

(A) If the following conditions hold:

- (1)  $D_i \equiv 1 \pmod{4}$ ,  $1 \leq i \leq n$ ;      (2)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (3)  $\left(\frac{D_i}{D_j}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^n$ , so  $2^n \leq \#S^{(\varphi)}(E/\mathbb{Q}) \leq 2^{n+1}$ .

(B) If the following conditions hold:

- (1)  $D_i \equiv 1 \pmod{4}$ ,  $1 \leq i \leq n$ ;
- (2)  $D_k \equiv 5 \pmod{8}$  for at least one  $k \in \{1, \dots, n\}$ ;
- (3)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_i}{D_j}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$ , and  $\#S^{(\varphi)}(E/\mathbb{Q}) = 2^n$ .

(C) If the following conditions hold:

- (1)  $p \equiv 7 \pmod{8}$ ;      (2)  $D_i \equiv 1 \pmod{8}$ ,  $1 \leq i \leq n$ ;
- (3)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_i}{D_j}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$ , and  $\#S^{(\varphi)}(E/\mathbb{Q}) = 2^{n+1}$ .

**Theorem 1.3.** Let  $E' = E'_+$  be the elliptic curve in (1.2) with  $\varepsilon = +1$ . For each  $i \in \{1, \dots, n\}$ , denote

$$\Pi_i^+(D)' = \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right),$$

where  $(-)$  is the ( Legendre ) quadratic residue symbol. Take a subset  $I$  of  $Z(n) = \{1, \dots, n\}$  as follows:  $I = \{i \in Z(n) : D_i \equiv 1 \pmod{8}\} \cup \{i \in Z(n) : (1 + p\widehat{D_i})(1 +$

$\widehat{qD_i} \equiv 0(\text{mod } 16)\} \cup \{i \in Z(n) : D_i \equiv 3(\text{mod } 8) \text{ and } p \equiv 1(\text{mod } 4)\} \cup \{i \in Z(n) : D_i \equiv 7(\text{mod } 8) \text{ and } p \equiv 3(\text{mod } 4)\}$ , and define a function  $\rho^+(D)'$  by

$$\rho^+(D)' = \sum_{i \in I} \left[ \frac{1}{1 + \Pi_i^+(D)'} \right],$$

where  $[x]$  is the greatest integer  $\leq x$ . Then there exists a subset  $T \subset \{D_1, \dots, D_n\}$  with cardinal  $\#T = \rho^+(D)'$  such that  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset < T \text{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)'}$ .

In particular,  $\dim_2 S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \geq \rho^+(D)'$ .

**Corollary 1.4.** For elliptic curve  $E' = E'_+$  as in Theorem 1.3, we have  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+3}$ . Furthermore, if for each  $i \in \{1, \dots, n\}$ ,  $\Pi_i^+(D)' = 0$  and one of the following conditions hold:

- (1)  $D_i \equiv 1(\text{mod } 8)$ ; (2)  $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0(\text{mod } 16)$ ;
- (3)  $\begin{cases} D_i \equiv 3(\text{mod } 8), \\ p \equiv 1(\text{mod } 4); \end{cases}$  (4)  $\begin{cases} D_i \equiv 7(\text{mod } 8), \\ p \equiv 3(\text{mod } 4). \end{cases}$

Then  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^{n+2}$ , so  $2^{n+2} \leq \#S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \leq 2^{n+3}$ .

Furthermore, if  $\begin{cases} p \equiv 3(\text{mod } 4), \\ \alpha(-pq)' = 0 \end{cases}$  or  $\begin{cases} D - p \equiv 0, 2(\text{mod } 8), \\ \alpha(-pq)' = 0, \end{cases}$

then  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+3}$ , so  $\#S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+3}$ ,

where  $\alpha(-pq)' = \sum_{i=1}^n \left(1 - \left(\frac{-1}{D_i}\right)\right) \left(1 - \left(\frac{-pq}{D_i}\right)\right)$ .

For example, if the following conditions hold:

- (1)  $D_i \equiv 1(\text{mod } 8)$ ,  $1 \leq i \leq n$ ; (2)  $\left(\frac{D_i}{p}\right) + \left(\frac{D_i}{q}\right) = 0$ ,  $1 \leq i \leq n$ ;
- (3)  $p \equiv 3(\text{mod } 4)$ .

Then  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+3}$ , so  $\#S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+3}$ .

**Theorem 1.5.** For elliptic curves  $E = E_+$  and  $E' = E'_+$  in (1.1) and (1.2)

with  $\varepsilon = +1$ , we have

(A) If the following conditions hold:

- (1)  $D_i \equiv 1 \pmod{4}$ ,  $1 \leq i \leq n$ ;
- (2)  $D_k \equiv 5 \pmod{8}$  for at least one  $k \in \{1, \dots, n\}$ ;
- (3)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_i}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ ;
- (5)  $p - D \equiv 0, 2 \pmod{8}$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$ ,  $(\mathbb{Z}/2\mathbb{Z})^{n+2} \subset S^{(\hat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+3}$ .

Furthermore, if  $p \equiv 3 \pmod{4}$  or  $p - D \equiv 0 \pmod{8}$ , then

$S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$ ,  $S^{(\hat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+3}$ , and

$$\text{rank}(E(\mathbb{Q})) + \dim_2 \text{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \text{TS}(E'/\mathbb{Q})[\hat{\varphi}] = 2n + 1,$$

where  $\text{TS}(E/\mathbb{Q})$  and  $\text{TS}(E'/\mathbb{Q})$  are the Tate-Shafarevich groups of  $E$  and  $E'$  over  $\mathbb{Q}$  respectively (See [S, Chapter X]).

(B) If the following conditions hold:

- (1)  $p \equiv 7 \pmod{8}$ ;      (2)  $D_i \equiv 1 \pmod{8}$ ,  $1 \leq i \leq n$ ;
- (3)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_i}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$ ,  $S^{(\hat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+3}$ , and

$$\text{rank}(E(\mathbb{Q})) + \dim_2 \text{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \text{TS}(E'/\mathbb{Q})[\hat{\varphi}] = 2n + 2.$$

## Part 2 Results for the case $\varepsilon = -1$

**Theorem 1.6.** Let  $E = E_-$  be the elliptic curve in (1.1) with  $\varepsilon = -1$ . For each  $i \in \{1, \dots, n\}$ , denote

$$\Pi_i^-(D) = \left(1 - \left(\frac{p}{D_i}\right)\right) + \left(1 - \left(\frac{q}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_j}{D_i}\right)\right),$$

and define a function  $\rho^-(D)$  by

$$\rho^-(D) = \sum_{i=1}^n \left[ \frac{1}{1 + \Pi_i^-(D)} \right],$$

where  $[x]$  is the greatest integer  $\leq x$ . Then we have

(1) There exists a subset  $T \subset \{D_1, \dots, D_n\}$  with cardinal  $\#T = \rho^-(D)$  such that  $S^{(\varphi)}(E/\mathbb{Q}) \supset \langle \{D_i^* : D_i \in T\} \bmod(\mathbb{Q}^{\star^2}) \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)}$ , where  $D_i^* \in \{D_i, -D_i\}$ . In particular,  $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^-(D)$ .

(2) Furthermore, if one of the following conditions hold:

- (a)  $p \equiv 7 \pmod{8}$  and  $D_i \equiv 1, 7 \pmod{8}$  ( $1 \leq i \leq n$ );
- (b)  $p \equiv 1 \pmod{8}$  and  $D_i \equiv 1, 3 \pmod{8}$  ( $1 \leq i \leq n$ ).

Then for  $T$  as in (1), we have  $S^{(\varphi)}(E/\mathbb{Q}) \supset \langle 2^* \rangle \times \langle \{D_i^* : D_i \in T\} \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)+1}$ , where  $2^* \in \{2, -2\}$ . In particular,  $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^-(D) + 1$ .

**Corollary 1.7.** For elliptic curve  $E = E_-$  as in Theorem 1.6, we have  $S^{(\varphi)}(E/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+1}$ . Furthermore,

(A) If the following conditions hold:

- (1)  $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (2)  $\left(\frac{D_j}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^n$ , so  $2^n \leq \#S^{(\varphi)}(E/\mathbb{Q}) \leq 2^{n+1}$ .

(B) If the following conditions hold:

- (1)  $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (2)  $\left(\frac{D_j}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ ;
- (3)  $\begin{cases} p \equiv 7 \pmod{8} \\ D_i \equiv 1, 7 \pmod{8} \end{cases} (1 \leq i \leq n)$ , or  $\begin{cases} p \equiv 1 \pmod{8} \\ D_i \equiv 1, 3 \pmod{8} \end{cases} (1 \leq i \leq n)$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$ , and  $\#S^{(\varphi)}(E/\mathbb{Q}) = 2^{n+1}$ .

**Theorem 1.8.** Let  $E' = E'_-$  be the elliptic curve in (1.2) with  $\varepsilon = -1$ . For each  $i \in \{1, \dots, n\}$ , denote

$$\Pi_i^-(D)' = \left(1 - \left(\frac{p\widehat{D}_i}{D_i}\right)\right) \left(1 - \left(\frac{q\widehat{D}_i}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right),$$

where  $(-)$  is the ( Legendre ) quadratic residue symbol. Take a subset  $I$  of  $Z(n) = \{1, \dots, n\}$  as follows:  $I = \{i \in Z(n) : D_i \equiv 1(\text{mod}8)\} \cup \{i \in Z(n) : (1 - p\widehat{D}_i)(1 - q\widehat{D}_i) \equiv 0(\text{mod}16)\} \cup \{i \in Z(n) : D_i \equiv 3(\text{mod}8) \text{ and } p \equiv 1(\text{mod}4)\} \cup \{i \in Z(n) : D_i \equiv 7(\text{mod}8) \text{ and } p \equiv 3(\text{mod}4)\}$ , and define a function  $\rho^-(D)'$  by

$$\rho^-(D)' = \sum_{i \in I} \left[ \frac{1}{1 + \Pi_i^-(D)'} \right],$$

where  $[x]$  is the greatest integer  $\leq x$ . Then there exists a subset  $T \subset \{D_1, \dots, D_n\}$  with cardinal  $\#T = \rho^-(D)'$  such that  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset < T \text{mod}(\mathbb{Q}^{\star 2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)'}$ . In particular,  $\dim_2 S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \geq \rho^-(D)'$ .

**Corollary 1.9.** For elliptic curve  $E' = E'_-$  as in Theorem 1.8, we have  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+2}$ . Furthermore, if for each  $i \in \{1, \dots, n\}$ ,  $\Pi_i^-(D)' = 0$  and one of the following conditions hold:

- (1)  $D_i \equiv 1(\text{mod}8)$ ; (2)  $(1 - p\widehat{D}_i)(1 - q\widehat{D}_i) \equiv 0(\text{mod}16)$ ;
- (3)  $\begin{cases} D_i \equiv 3(\text{mod}8), \\ p \equiv 1(\text{mod}4); \end{cases}$  (4)  $\begin{cases} D_i \equiv 7(\text{mod}8), \\ p \equiv 3(\text{mod}4). \end{cases}$

Then  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$ , so  $\#S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+2}$ .

For example, if the following conditions hold:

- (1)  $D_i \equiv 1(\text{mod}8)$ ,  $1 \leq i \leq n$ ; (2)  $\left(\frac{D_i}{p}\right) + \left(\frac{D_i}{q}\right) = 0$ ,  $1 \leq i \leq n$ .

Then  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+2}$ , so  $\#S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+2}$ .

**Theorem 1.10** For elliptic curves  $E = E_-$  and  $E' = E'_-$  in (1.1) and (1.2) with  $\varepsilon = -1$ , we have

(A) If the following conditions hold:

- (1)  $D_i \equiv 1 \pmod{4}$ ,  $1 \leq i \leq n$ ;
- (2)  $D_k \equiv 5 \pmod{8}$  for at least one  $k \in \{1, \dots, n\}$ ;
- (3)  $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_j}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ ;
- (5)  $p - D \equiv 2, 4 \pmod{8}$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$ ,  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$ , and

$$\text{rank}(E(\mathbb{Q})) + \dim_2 \text{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \text{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n.$$

(B) If the following conditions hold:

- (1)  $p \equiv 1, 7 \pmod{8}$ ; (2)  $D_i \equiv 1 \pmod{8}$ ,  $1 \leq i \leq n$ ;
- (3)  $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ ,  $1 \leq i \leq n$ ;
- (4)  $\left(\frac{D_j}{D_i}\right) = 1$ ,  $1 \leq i, j \leq n$  and  $i \neq j$ .

Then  $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$ ,  $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$ , and

$$\text{rank}(E(\mathbb{Q})) + \dim_2 \text{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \text{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n + 1.$$

**Remark 1.11** For each pair  $(p, q)$  above, by Chinese Remainder Theorem (See [L]) and Dirichlet Theorem on primes in arithmetic progressions [see[IK]], it is easy to verify that there exist infinitely many  $D$  satisfying the conditions in every Corollaries above.

## 2 Proofs of Theorems

Let  $M_{\mathbb{Q}}$  be the set of all places of  $\mathbb{Q}$ , including the infinite  $\infty$ . For each place  $p$ , denote by  $\mathbb{Q}_p$  the completion of  $\mathbb{Q}$  at  $p$ , and if  $p$  is finite, denote by  $v_p$  the corresponding normalized additive valuation, so  $v_p(p) = 1$ . Let  $S = \{\infty, 2, p, q, D_1, \dots, D_n\}$ ,

and define a subgroup of  $\mathbb{Q}^*/\mathbb{Q}^{\star 2}$  as follows:  $\mathbb{Q}(S, 2) = \langle -1, 2, p, q, D_1, \dots, D_n \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{n+4}$ . For any subset  $A \subset \mathbb{Q}^*$ , we write  $\langle A \rangle$  for the subgroup of  $\mathbb{Q}^*/\mathbb{Q}^{\star 2}$  generated by all the elements in  $A$ . For each  $d \in \mathbb{Q}(S, 2)$ , define the curves

$$C_d: dw^2 = d^2 - 2\varepsilon(p+q)Ddz^2 + 4D^2z^4, \text{ and}$$

$$C'_d: dw^2 = d^2 + \varepsilon(p+q)Ddz^2 + pqD^2z^4.$$

Then by the theory of descent via two-isogeny ( see [S, chapter X]), the  $\varphi$ -Selmer group for the elliptic curve  $E/\mathbb{Q}$  in (1.1) is

$S^{(\varphi)}(E/\mathbb{Q}) \cong \{d \in \mathbb{Q}(S, 2) : C_d(\mathbb{Q}_v) \neq \emptyset \text{ for all } v \in S\}$ , and the  $\widehat{\varphi}$ -Selmer group for the elliptic curve  $E'/\mathbb{Q}$  in (1.2) is

$$S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong \{d \in \mathbb{Q}(S, 2) : C'_d(\mathbb{Q}_v) \neq \emptyset \text{ for all } v \in S\}.$$

**Proposition 2.1** We assume  $\varepsilon = 1$  and the elliptic curve  $E = E_+$  be as in (1.1).

(A) For  $d \in \mathbb{Q}(S, 2)$ , if one of the following conditions holds:

- (1)  $d < 0$ ; (2)  $p \mid d$ ; (3)  $q \mid d$ .

Then  $d \notin S^{(\varphi)}(E/\mathbb{Q})$ . Moreover, if  $d > 0$ , then  $C_d(\mathbb{R}) \neq \emptyset$ .

(B) For  $d = 2 \in \mathbb{Q}(S, 2)$ , we have

- (1)  $C_2(\mathbb{Q}_2) \neq \emptyset \iff D(D - 2p - 2) \equiv 1 \pmod{16}$ .
- (2) For each prime number  $l \mid pqD$ ,  $C_2(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{2}{l}\right) = 1$ , i.e.,  $l \equiv 1, 7 \pmod{8}$ .

(C) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

- (1)  $C_{D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 1 \pmod{4}$ .
- (2) For each prime number  $l \mid pq\widehat{D_i}$ ,  $C_{D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{D_i}{l}\right) = 1$ .
- (3)  $C_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{p\widehat{D_i}}{D_i}\right) = \left(\frac{q\widehat{D_i}}{D_i}\right) = 1$ .

**Proof.** Conclusion (A) follows directly by the valuation property.

(B) In this case,  $C_2 : w^2 = 2 - 2(p+q)Dz^2 + 2D^2z^4$ .

(1) If  $C_2(\mathbb{Q}_2) \neq \emptyset$ , then there exists  $(z_0, w_0) \in \mathbb{Q}_2^2$  such that  $w_0^2 = 2 - 2(p+q)Dz_0^2 + 2D^2z_0^4$ , from which we have  $v_2(w_0) \geq 1$  and  $v_2(z_0) = 0$ . Let  $w_0 = 2w_1$  with  $w_1 \in \mathbb{Q}_2$  and  $v_2(w_1) \geq 0$ . Then  $2w_1^2 = 1 - (p+q)Dz_0^2 + D^2z_0^4$ . Note that  $v_2(Dz_0) = 0$ , so by lemma in [Rob], we have  $z_0^2 = 8s+1$  and  $D^2z_0^4 = 8t+1$  for some  $s, t \in \mathbb{Q}_2$  with  $v_2(s) \geq 0$  and  $v_2(t) \geq 0$ . So  $2w_1^2 = 2 - 2(p+1)D(8s+1) + 8t$ . Thus  $v_2(2w_1^2) = v_2(2) = 1$ , so  $v_2(w_1) = 0$ , and we may assume that  $(w_1)^2 = 8r+1$  for some  $r \in \mathbb{Q}_2$  and  $v_2(r) \geq 0$ . So  $2(8r+1) = 1 - 2(p+1)D(8s+1) + D^2(8s+1)^2$ , from which we get  $D^2 - 2(p+1)D \equiv 1 \pmod{16}$ , i.e.,  $D(D-2p-2) \equiv 1 \pmod{16}$ . Conversely, if  $D(D-2p-2) \equiv 1 \pmod{16}$ , let  $f(z, w) = w^2 - 2 + 2(p+q)Dz^2 - 2D^2z^4$ , then  $f'_w = 2w$ , and  $v_2(f(1, 2)) \geq 5 > 2v_2(f'_w(1, 2))$ , so by Hensel's lemma (see [S], p.322), we have  $C_2(\mathbb{Q}_2) \neq \emptyset$ .

(2) For  $l = p$ , if  $C_2(\mathbb{Q}_p) \neq \emptyset$ , then  $\exists (z_0, w_0) \in \mathbb{Q}_p^2$  s.t.  $w_0^2 = 2 - 2(p+q)Dz_0^2 + 2D^2z_0^4$ . If  $v_p(z_0) > 0$ , then  $v_p(w_0^2 - 2) \geq 1$ , so  $\left(\frac{2}{p}\right) = 1$ . If  $v_p(z_0) = 0$  then we can obtain that  $v_p(Dz_0^2 - 1) = 0$ . Let  $w_1 = w_0(Dz_0^2 - 1)^{-1} \in \mathbb{Q}_p^*$ , then  $v_p(w_1^2 - 2) \geq 1$ , so  $\left(\frac{2}{p}\right) = 1$ . If  $v_p(z_0) < 0$  then  $v_p(z_0^{-1}) > 0$ , so by the equality  $(\frac{w_0}{z_0^2})^2 = 2(z_0^{-1})^4 - 4(p+1)Dz_0^{-2} + 2D^2$  we get  $v_p(w_1) \geq 0$  and  $v_p(w_1^2 - 2D^2) \geq 1$  with  $w_1 = \frac{w_0}{z_0^2}$ . So  $\left(\frac{2D^2}{p}\right) = 1$ , and then  $\left(\frac{2}{p}\right) = 1$ . To sum up, we obtain  $\left(\frac{2}{p}\right) = 1$ .

Conversely, if  $\left(\frac{2}{p}\right) = 1$ , then  $2 \equiv a^2 \pmod{p}$  for some  $a \in \mathbb{Z}$  with  $p \nmid a$ . Then for the above  $f(z, w)$ , we have  $v_p(f(0, a)) \geq 1 > 0 = 2v_p(f'_w(0, a))$ . So by Hensel's lemma, we have  $C_2(\mathbb{Q}_p) \neq \emptyset$ . The case  $l = q$ ,  $D_i$  ( $1 \leq i \leq n$ ) are similar. This proves (B).

(C) In this case,  $C_{D_i} : w^2 = D_i - 2(p+q)Dz^2 + 4D\widehat{D_i}z^4$ .

(1) If  $C_{D_i}(\mathbb{Q}_2) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_2^2$  s. t.  $w_0^2 = D_i - 2(p+q)Dz_0^2 + 4D\widehat{D_i}z_0^4$ .

If  $v_2(z_0) \geq 0$ , then  $v_2(w_0^2) = v_2(D_i)$ ,  $v_2(w_0) = 0$  and  $v_2(w_0^2 - D_i) \geq 2$ . Since  $v_2(w_0^2 - D_i) \geq 3$  ( see lemma in [Rob]), we have  $v_2(D_i - 1) = v_2((w_0^2 - 1) + (D_i - w_0^2)) \geq 2$ . So  $D_i \equiv 1 \pmod{4}$ .

If  $v_2(z_0) < 0$ , then  $v_2(z_0^{-1}) \geq 1$ , so by  $(\frac{w_0}{2z_0^2})^2 = D_i(2z_0^2)^{-2} - (p+1)Dz_0^{-2} + D\widehat{D_i}$ , we get  $v_2((\frac{w_0}{2z_0^2})^2) = v_2(D\widehat{D_i}) = 0$ , so  $v_2((\frac{w_0}{2z_0^2})^2) = 0$  and  $v_2(D_i\widehat{D_i}^2 - (\frac{w_0}{2z_0^2})^2) \geq 2$ , hence  $v_2(D_i - (\frac{w_0}{2z_0^2\widehat{D_i}})^2) \geq 2$ . Since  $v_2((\frac{w_0}{2z_0^2\widehat{D_i}})^2 - 1) \geq 3$ , we get  $v_2(D_i - 1) = v_2((D_i - (\frac{w_0}{2z_0^2\widehat{D_i}})^2) + ((\frac{w_0}{2z_0^2\widehat{D_i}})^2 - 1)) \geq 2$ . So  $D_i \equiv 1 \pmod{4}$ .

Conversely, if  $D_i \equiv 1 \pmod{4}$ , then for the polynomial  $f(z, w) = w^2 - D_i + 2(p+q)Dz^2 - 4D\widehat{D_i}z^4$ , we have  $f'_w = 2w$ . So if  $D_i \equiv 1 \pmod{8}$ , then  $v_2(f(0, 1)) \geq 3 > 2 = 2v_2(f'_w(0, 1))$ ; If  $D_i \equiv 5 \pmod{8}$ , then  $v_2(f(1, 1)) \geq 3 > 2 = 2v_2(f'_w(1, 1))$ . So by Hensel's lemma, we have  $C_{D_i}(\mathbb{Q}_2) \neq \emptyset$ .

(2) For  $l = p$ , if  $C_{D_i}(\mathbb{Q}_p) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_p^2$  s. t.  $w_0^2 = D_i - 2(p+q)Dz_0^2 + 4D\widehat{D_i}z_0^4 = -4pDz_0^2 + D_i(2\widehat{D_i}z_0^2 - 1)^2$ .

If  $v_p(z_0) > 0$ , then  $v_p(w_0^2) = v_p(D_i) = 0$ , so  $v_p(w_0) = 0$  and  $v_p(w_0^2 - D_i) \geq 1$ , so  $\left(\frac{D_i}{p}\right) = 1$ .

If  $v_p(z_0) = 0$ , then  $v_p(w_0) \geq 0$ . If  $v_p(2\widehat{D_i}z_0^2 - 1) \geq 1$ , then  $v_p((2\widehat{D_i}z_0^2 - 1)^2) \geq 2 > 1 = v_p(-4pDz_0^2)$ , so  $v_p(w_0^2) = v_p(-4pDz_0^2)$ , i.e.,  $2v_p(w_0^2) = 1$ , which is impossible !

So  $v_p(2\widehat{D_i}z_0^2 - 1) = 0$ , and then we have  $v_p(D_i - (\frac{w_0}{2\widehat{D_i}z_0^2 - 1})^2) \geq 1$ , which shows that  $\left(\frac{D_i}{p}\right) = 1$ .

If  $v_p(z_0) < 0$ , then  $v_p(z_0^{-1}) \geq 1$ . By the above equation of  $C_{D_i}$  we have  $(\frac{w_0}{2\widehat{D_i}z_0^2})^2 = D_i(2\widehat{D_i}z_0^2)^{-2} - \frac{(p+1)D}{(\widehat{D_i}z_0^2)^2} + D_i$ . So  $v_p((\frac{w_0}{2\widehat{D_i}z_0^2})^2) = v_p(D_i) = 0$ , hence  $v_p(D_i - \frac{w_0^2}{2\widehat{D_i}z_0^2}) \geq 1$ , which shows  $\left(\frac{D_i}{p}\right) = 1$ .

Conversely, if  $\left(\frac{D_i}{p}\right) = 1$ , then  $D_i \equiv a^2 \pmod{p}$  for some  $a \in \mathbb{Z}$  with  $v_p(a) = 0$ . Then  $v_p(f(0, a)) \geq 1 > 0 = 2v_p(f'_w(0, a))$ . By Hensel's lemma, we get  $C_{D_i}(\mathbb{Q}_p) \neq \emptyset$ . The other cases follow the same line. We omit the details. The proof of Proposition 2.1 is completed.  $\square$

**Corollary 2.2** We assume  $\varepsilon = 1$  and the elliptic curve  $E = E_+$  be as in (1.1).

$$(A) \quad 2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 7 \pmod{8} \text{ and } D_i \equiv 1, 7 \pmod{8} \quad (1 \leq i \leq n).$$

(B) For each  $D_i$  ( $1 \leq i \leq n$ ), we have  $D_i \in S^{(\varphi)}(E/\mathbb{Q})$  if and only if

$$D_i \equiv 1 \pmod{4} \text{ and } \left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1 \quad (1 \leq j \leq n, \text{ and } j \neq i).$$

**Proof.** The results follow easily from Proposition 2.1.  $\square$

**Proposition 2.3** We assume  $\varepsilon = 1$  and the elliptic curve  $E' = E'_+$  be as in (1.2).

(A) (1) For any  $d \in \mathbb{Q}(S, 2)$ ,  $C'_d(\mathbb{R}) \neq \emptyset$ . If  $2 \mid d$ , then  $d \notin S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ .

(2)  $\{1, pq, -pD, -qD\} \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ .

(B) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

(1)  $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$  if and only if one of the following conditions holds:

(a)  $D_i \equiv 1 \pmod{8}$ ; (b)  $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 \pmod{16}$ ;

(c)  $D_i \equiv 3 \pmod{8}$  and  $p \equiv 1 \pmod{4}$ ; (d)  $D_i \equiv 7 \pmod{8}$  and  $p \equiv 3 \pmod{4}$ .

(2)  $C'_{D_i}(\mathbb{Q}_p) \neq \emptyset$  and  $C'_{D_i}(\mathbb{Q}_q) \neq \emptyset$ .

(3) For each  $j \neq i$ ,  $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset \iff \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$ .

(4)  $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$ .

(C) For  $d = -pq \in \mathbb{Q}(S, 2)$ , we have

(1)  $C'_{-pq}(\mathbb{Q}_2) \neq \emptyset \iff p \equiv 3 \pmod{4} \text{ or } D - p \equiv 0, 2 \pmod{8}$ .

(2)  $C'_{-pq}(\mathbb{Q}_p) \neq \emptyset$  and  $C'_{-pq}(\mathbb{Q}_q) \neq \emptyset$ .

(3) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

$$C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(1 - \left(\frac{-1}{D_i}\right)\right) \left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0.$$

**Proof.** (A) (1) Obvious.

(2) For  $C'_d : dw^2 = (d + pDz^2)(d + qDz^2)$ , if  $d = -pD$ , then  $(1, 0) \in C'_d(\mathbb{Q})$ ; and if  $d = -qD$ , then  $(1, 0) \in C'_d(\mathbb{Q})$ , so the group  $\langle -pD, -qD \rangle = \{1, pq, -pD, -qD\} \subseteq E(\mathbb{Q})/\widehat{\varphi}(E'(\mathbb{Q})) \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ .

(B)  $C'_{D_i} : w^2 = D_i + (p + q)Dz^2 + pqD\widehat{D_i}z^4 = D_i(1 + p\widehat{D_i}z^2)(1 + q\widehat{D_i}z^2)$ .

(1) If  $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_{D_i}^2$  s. t.  $w_0^2 = D_i + (p + q)Dz_0^2 + pqD\widehat{D_i}z_0^4$ .

If  $v_2(z_0) > 0$ , then  $v_2(w_0) = 0$ . By Lemma 4 in [C, p.11], we get  $v_2(D_i - 1) \geq 3$  and then  $D_i \equiv 1 \pmod{8}$ .

If  $v_2(z_0) \leq 0$ , then by the equality  $(\frac{w_0}{z_0^2})^2 = D_i(z_0^{-2} + p\widehat{D_i})(z_0^{-2} + q\widehat{D_i})$ , we have  $v_2(\frac{w_0}{z_0^2}) \geq 0$ . Denote  $W = \frac{w_0}{z_0^2}$ ,  $Z = z_0^{-1}$ , then  $v_2(Z) \geq 0$ ,  $v_2(W) \geq 0$  and  $W^2 = D_i(Z^2 + p\widehat{D_i})(Z^2 + q\widehat{D_i}) = D_i(V^2 - \widehat{D_i}^2)$  with  $V = Z^2 + (p + 1)\widehat{D_i} \in \mathbb{Q}_2$ . Obviously,  $v_2(V) \geq 0$ .

i) If  $v_2(V) = 0$ , then  $v_2(V^2 - 1) \geq 3$  and  $v_2(Z) = 0$ . So  $v_2(V^2 - \widehat{D_i}^2) = v_2((V^2 - 1) - (\widehat{D_i}^2 - 1)) \geq 3$ . Thus  $v_2(W^2) = v_2(V^2 - \widehat{D_i}^2) \geq 3$ , so  $v_2(W^2) \geq 2$ , and then  $v_2((1 + p\widehat{D_i})(1 + q\widehat{D_i})) \geq 4$ , i.e.,  $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 \pmod{16}$ .

ii) If  $v_2(V) \geq 1$ , then  $v_2(W) = 0$ .

If  $v_2(V) = 1$ , then  $v_2(V^2 - 4) \geq 5$ . So  $v_2(W^2 - 3D_i) = v_2((V^2 - 4) - (\widehat{D_i}^2 - 1)) \geq 3$ .

Thus  $v_2(1 - 3D_i) \geq 3$ , hence  $D_i \equiv 3 \pmod{8}$ . Since  $V = Z^2 + (p + 1)\widehat{D_i}$ , we get  $v_2(p + 1) = v_2(V - Z^2) = 1$ , so  $p \equiv 1 \pmod{4}$ .

If  $v_2(V) \geq 2$ , then  $v_2(V^2) \geq 4$ , so  $W^2 + D_i = D_i(V^2 + 1 - \widehat{D_i}^2)$ , hence  $v_2(W^2 + D_i) \geq 3$ ,

and then  $v_2(1 + D_i) = v_2(W^2 + D_i + 1 - W^2) \geq 3$ , so  $D_i \equiv 7 \pmod{8}$ . Since  $V = Z^2 + (p+1)\widehat{D_i}$ , we get  $v_2(p+1) = v_2(V - Z^2) \geq 2$ , which shows  $p \equiv 3 \pmod{4}$ . To sum up, we have proved that if  $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$ , then one of the following conditions holds:

- (a)  $D_i \equiv 1 \pmod{8}$ ; (b)  $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 \pmod{16}$ ;
- (c)  $D_i \equiv 3 \pmod{8}$  and  $p \equiv 1 \pmod{4}$ ; (d)  $D_i \equiv 7 \pmod{8}$  and  $p \equiv 3 \pmod{4}$ .

Conversely, we assume one of the above conditions holds. Let  $f(z, w) = D_i + (p + q)Dz^2 + pqD\widehat{D_i}z^4 - w^2$ , then  $f'_w(z, w) = -2w$ .

(a) If  $D_i \equiv 1 \pmod{8}$ , then  $v_2(f(0, 1)) \geq 3 > 2 = 2v_2(f'_w(0, 1))$ . So by Hensel's lemma, we get  $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$ .

(b) If  $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 \pmod{16}$ , then  $v_2(1 + p\widehat{D_i}) \geq 3$  or  $v_2(1 + q\widehat{D_i}) \geq 3$ . Let  $g(z) = 1 + l\widehat{D_i}z^2$ , where  $l = p$  or  $q$  according to the above first or second inequalities. Then  $g'(z) = 2l\widehat{D_i}z$ . So  $v_2(g(1)) \geq 3 > 2 = 2v_2(g'(1))$ , and by Hensel's lemma,  $g(z_0) = 0$  for some  $z_0 \in \mathbb{Q}_2$ . Therefore  $(z_0, 0) \in C'_{D_i}(\mathbb{Q}_2)$ .

(c) If  $D_i \equiv 3 \pmod{8}$  and  $p \equiv 1 \pmod{4}$ , let  $g(w) = w^2 - D_i(2^2 + p\widehat{D_i})(2^2 + q\widehat{D_i})$ , then  $g'(w) = 2w$ , and  $v_2(g(1)) > 2 = 2v_2(g'(1))$ . So by Hensel's lemma,  $g(w_0) = 0$  for some  $w_0 \in \mathbb{Q}_2$ . Thus  $(\frac{1}{2}, \frac{w_0}{4}) \in C'_{D_i}(\mathbb{Q}_2)$ .

(d) similar to (c) above. This proves (1).

(2) Let  $f(z) = D_i + (p + q)Dz^2 + pqD\widehat{D_i}z^4$ , then  $f(z) \equiv D_i(1 + 2\widehat{D_i}z^2) \pmod{p}$ , also  $f(z) \equiv D_i(1 - 2\widehat{D_i}z^2) \pmod{q}$ . Take  $\bar{g} = 1$ ,  $\bar{h} = \bar{f} \pmod{p}$  (if  $D_i$  is a square mod  $p$ , then take  $\bar{g} = D_i \pmod{p}$ ), then  $\deg \bar{g} = 0$  and it is easy to verify that  $\bar{h}$  is square free. So by lemma 14 in [MSS],  $w^2 = f(z)$  has solutions in  $\mathbb{Q}_p$ , i.e.,  $C'_{D_i}(\mathbb{Q}_p) \neq \emptyset$ . Similarly,  $C'_{D_i}(\mathbb{Q}_q) \neq \emptyset$ . This proves (2).

(3) If  $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset$  ( $j \neq i$ ), then  $\exists(z_0, w_0) \in \mathbb{Q}_{D_j}^2$  s. t.  $w_0^2 = D_i(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$ .

i) If  $v_{D_j}(z_0) \geq 0$ , then  $v_{D_j}(w_0) = 0$  and  $D_i \equiv w_0^2 \pmod{D_j}$ , so  $\left(\frac{D_i}{D_j}\right) = 1$ .

ii) If  $v_{D_j}(z_0) < 0$ , then  $v_{D_j}(z_0^{-1}) \geq 1$  and  $v_{D_j}(w_0) = 1 + 2v_{D_j}(z_0) < 0$ . Now

$(\frac{w_0}{z_0^2})^2 = D_i z_0^{-4} + (p+q)D z_0^{-2} + pqD_i \widehat{D_i}^2$ . By the above discussion, we get  $pqD_i \equiv (\frac{w_0}{z_0^2})^2 \pmod{D_j}$  with  $v_{D_j}(\frac{w_0}{z_0^2}) < 0$ , so  $\left(\frac{pqD_i}{D_j}\right) = 1$ . To sum up, we get  $\left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$ .

Conversely, if  $\left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$ , then  $D_i \equiv a^2 \pmod{D_j}$  or  $pqD_i \equiv b^2 \pmod{D_j}$  for some  $a, b \in \mathbb{Z}$  and  $D_j \nmid ab$ .

For the former, let  $f(z, w) = w^2 - D_i - (p+q)Dz^2 + pqD_i \widehat{D_i}^2 z^4$ , then  $f'_w(z, w) = 2w$  and  $v_{D_j}(f(0, a)) > 0 = 2v_{D_j}(f'_w(0, a))$ . So by Hensel's lemma,  $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset$ .

For the later, let  $g(z, w) = w^2 - D_i \widehat{D_i}^2 - (p+q)\widehat{D_i}^2 Dz^2 - pqD_i \widehat{D_i}^4 z^4$ , then  $g'_w(z, w) = 2w$  and  $v_{D_j}(g(\widehat{D_i}^{-1}, b)) > 0 = 2v_{D_j}(g'_w(\widehat{D_i}^{-1}, b))$ . So by Hensel's lemma,  $\exists(z_0, w_0) \in \mathbb{Q}_{D_j}^2$  s. t.  $g(z_0, w_0) = 0$ , and so  $(z_0, \frac{w_0}{\widehat{D_i}}) \in C'_{D_i}(\mathbb{Q}_{D_j})$ . This proves (3).

(4) If  $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_{D_i}^2$  s. t.  $w_0^2 = D_i(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$ .

It is easy to see that  $v_{D_i}(z_0) = 0$  and  $v_{D_i}(w_0) > 0$ . Set  $w_0 = D_i w_1$  with  $w_1 \in \mathbb{Q}_{D_i}$  and  $v_{D_i}(w_1) \geq 0$ . Then  $D_i w_1^2 = (1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$ , so  $(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2) \equiv 0 \pmod{D_i}$ . Thus  $\left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$ .

Conversely, if  $\left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$ , then  $\exists a \in \mathbb{Z}$  such that  $-l\widehat{D_i} \equiv a^2 \pmod{D_i}$  for  $l = p$  or  $q$ . Obviously,  $D_i \nmid a$ . Let  $g(z) = 1 + l\widehat{D_i}z^2$ , then  $g'(z) = 2l\widehat{D_i}z$ .

So  $v_{D_i}(g(a^{-1})) > 0 = 2v_{D_i}(g'(a^{-1}))$ , by Hensel's lemma,  $g(z_0) = 0$  for some  $z_0 \in \mathbb{Q}_{D_i}$ .

Thus  $(z_0, 0) \in C'_{D_i}(\mathbb{Q}_{D_i})$ , so  $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset$ . This proves (4), and (B) is proved.

(C) (1) If  $C'_{-pq}(\mathbb{Q}_2) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_2^2$  s. t.  $w_0^2 = -pq + (p+q)Dz_0^2 - D^2z_0^4$ .

i) If  $v_2(z_0) < 0$ , then  $v_2(z_0^{-1}) \geq 1$ , so by the equality  $r_0^2 = -pqz_0^{-4} + (p+q)Dz_0^{-2} - D^2$  with  $r_0 = \frac{w_0}{z_0^2}$ , we get  $v_2(r_0) = 0$ ,  $v_2(r_0^2 + D^2) \geq 4$ . Thus  $v_2(r_0^2 + D^2) = v_2((r_0^2 - D^2) + 2D^2) = 1$ , a contradiction! So we must have  $v_2(z_0) \geq 0$ .

ii) If  $v_2(z_0) > 0$ , then  $v_2(w_0) = 0$ ,  $v_2(w_0^2 + pq) \geq 4$ . So we get  $v_2(1 + pq) = v_2(1 - w_0^2 + w_0^2 + pq) \geq 3$ , which shows  $p \equiv 3 \pmod{4}$ .

iii) If  $v_2(z_0) = 0$ , then  $v_2(z_0^2 - 1) \geq 3$ ,  $v_2(z_0^4 - 1) \geq 4$ . Since  $w_0^2 = -(D-p)(D-q) + (p+q)D(z_0^2 - 1) - D^2(z_0^4 - 1)$ , we have  $v_2(w_0^2 + (D-p)(D-q)) \geq 4$ . But  $(D-p)(D-q) \equiv 0 \pmod{8}$ , so  $v_2(w_0) \geq 2$ . Thus  $v_2((D-p)(D-q)) \geq 4$ , which shows  $D-p \equiv 0 \pmod{8}$ .

Conversely, if  $p \equiv 3 \pmod{4}$ , we get  $-pq \equiv 1 \pmod{8}$ . So  $-pq = a^2$  for some  $a \in \mathbb{Q}_2$ .

Hence  $(0, a) \in C'_{-pq}(\mathbb{Q}_2)$ .

If  $D-p \equiv 0 \pmod{8}$ , then  $(D-p-1)^2 \equiv 1 \pmod{16}$ . Let  $f(z, w) = -[(p+1)z^2 - D]^2 + z^4 - w^2$ , then  $f'_z(z, w) = -4(p+1)[(p+1)z^2 - D]z + 4z^3$ .

If  $v_2((D-p-1)^2 - 1) > 4$ , then  $v_2(f(1, 0)) > 4 = 2v_2(f'_z(1, 0))$ ;

If  $v_2((D-p-1)^2 - 1) = 4$ , then  $v_2(f(3, 0)) \geq 5 > 4 = 2v_2(f'_z(3, 0))$ . So by Hensel's lemma,  $\exists(z_0, w_0) \in \mathbb{Q}_2^2$  s. t.  $f(z_0, w_0) = 0$ . It is easy to prove  $z_0 \neq 0$ . Hence  $(z_0^{-1}, w_0 z_0^{-1}) \in C'_{-pq}(\mathbb{Q}_2)$ .

(2) is similar to that of (B)(2).

(3) If  $C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset$ , then  $\exists(z_0, w_0) \in \mathbb{Q}_{D_i}^2$  s. t.  $w_0^2 = -pq + (p+q)Dz_0^2 - D^2z_0^4$ .

i) If  $v_{D_i}(z_0) \geq 0$ , then we get  $\left(\frac{-pq}{D_i}\right) = 1$ .

ii) If  $v_{D_i}(z_0) < 0$ , then  $v_{D_i}(z_0^{-1}) \geq 1$ . Since  $(\frac{w_0}{D_i z_0^2})^2 = -D^{-2}pqz_0^{-4} + (p+q)D^{-1}z_0^{-2} - 1$ , we have  $\left(\frac{-1}{D_i}\right) = 1$ . To sum up, we get  $\left(1 - \left(\frac{-1}{D_i}\right)\right) \left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0$ .

Conversely, if  $\left(1 - \left(\frac{-1}{D_i}\right)\right) \left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0$ , then  $\exists a, b \in \mathbb{Z}$  s. t.  $-pq \equiv a^2 \pmod{D_i}$

or  $-1 \equiv b^2 \pmod{D_i}$  with  $D_i \nmid ab$ . For the former, let  $f(z, w) = w^2 + pq - (p + q)Dz^2 + D^2z^4$ , then  $f'_w(z, w) = 2w$  and  $v_{D_i}(f(0, a)) > 0 = 2v_{D_i}(f'_w(0, a))$ . So by Hensel's lemma, we get  $C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset$ .

For the later, let  $f(z, w) = w^2 + D^2pq - (p + q)D^3z^2 + D^4z^4$ , then  $f'_w(z, w) = 2w$  and  $v_{D_i}(f(D^{-1}, b)) > 0 = 2v_{D_i}(f'_w(D^{-1}, b))$ . So by Hensel's lemma,  $\exists(z_0, w_0) \in \mathbb{Q}_{D_i}^2$  s. t.  $f(z_0, w_0) = 0$ . Thus  $\exists(z_0, w_0 D^{-1}) \in C'_{-pq}(\mathbb{Q}_{D_i})$ . This proves (C), and the proof of proposition 2.3 is completed.  $\square$

**Corollary 2.4** We assume  $\varepsilon = 1$  and the elliptic curve  $E' = E'_+$  be as in (1.2).

- (A) Denote  $\alpha(-pq)' = \sum_{i=1}^n \left(1 - \left(\frac{-1}{D_i}\right)\right) \left(1 - \left(\frac{-pq}{D_i}\right)\right)$ . Then  $-pq \in S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \iff \begin{cases} p \equiv 3 \pmod{4}, \\ \alpha(-pq)' = 0; \end{cases} \text{ or } \begin{cases} D - p \equiv 0, 2 \pmod{8}, \\ \alpha(-pq)' = 0. \end{cases}$
- (B) Denote  $\beta(-D)' = \sum_{i=1}^n \left(1 - \left(\frac{p}{D_i}\right)\right) \left(1 - \left(\frac{q}{D_i}\right)\right)$ . Then  $-D \in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$

if and only if one of the following conditions holds:

- (1)  $\begin{cases} D \equiv 7 \pmod{8}, \\ \beta(-D)' = 0; \end{cases}$  (2)  $\begin{cases} p \equiv 1, 7 \pmod{8}, \\ \beta(-D)' = 0; \end{cases}$
- (3)  $\begin{cases} D \equiv 1 \pmod{8}, \\ p \equiv 3 \pmod{8}, \\ \beta(-D)' = 0; \end{cases}$  (4)  $\begin{cases} D \equiv 5 \pmod{8}, \\ p \equiv 5 \pmod{8}, \\ \beta(-D)' = 0. \end{cases}$
- (C) For each  $D_i$  ( $1 \leq i \leq n$ ), denote

$$\Pi_i^+(D)' = \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right).$$

Then  $D_i \in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$  if and only if one of the following conditions holds:

- (1)  $\begin{cases} D_i \equiv 1 \pmod{8}, \\ \Pi_i^+(D)' = 0; \end{cases}$  (2)  $\begin{cases} (1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 \pmod{16}, \\ \Pi_i^+(D)' = 0; \end{cases}$
- (3)  $\begin{cases} D_i \equiv 3 \pmod{8}, \\ p \equiv 1 \pmod{4}, \\ \Pi_i^+(D)' = 0; \end{cases}$  (4)  $\begin{cases} D_i \equiv 7 \pmod{8}, \\ p \equiv 3 \pmod{4}, \\ \Pi_i^+(D)' = 0. \end{cases}$

**Proof.** The results follow easily from Proposition 2.3.  $\square$

**Proposition 2.5** We assume  $\varepsilon = -1$  and the elliptic curve  $E = E_-$  be as in

(1.1).

(A) For  $d \in \mathbb{Q}(S, 2)$ , if one of the following conditions holds:

- (1)  $p \mid d$ ; (2)  $q \mid d$ ; (3)  $d = -1$ .

Then  $d \notin S^{(\varphi)}(E/\mathbb{Q})$ .

(B<sub>1</sub>) For  $d = 2 \in \mathbb{Q}(S, 2)$ , we have

- (1)  $C_2(\mathbb{Q}_2) \neq \emptyset \iff D(D + 2p + 2) \equiv 1 \pmod{16}$ .  
(2) For each prime number  $l \mid pqD$ ,  $C_2(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{2}{l}\right) = 1$ , i.e.,  $l \equiv 1, 7 \pmod{8}$ .

(B<sub>2</sub>) For  $d = -2 \in \mathbb{Q}(S, 2)$ , we have

- (1)  $C_{-2}(\mathbb{Q}_2) \neq \emptyset \iff D(-D + 2p + 2) \equiv 3 \pmod{16}$ .  
(2) For each prime number  $l \mid pqD$ ,  $C_{-2}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{-2}{l}\right) = 1$ , i.e.,  $l \equiv 1, 3 \pmod{8}$ .

(C<sub>1</sub>) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

- (1)  $C_{D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 1 \pmod{4}$ .  
(2) For each prime number  $l \mid pq\widehat{D_i}$ ,  $C_{D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{D_i}{l}\right) = 1$ .  
(3)  $C_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{-p\widehat{D_i}}{D_i}\right) = \left(\frac{-q\widehat{D_i}}{D_i}\right) = 1$ .

(C<sub>2</sub>) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

- (1)  $C_{-D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 3 \pmod{4}$ .  
(2) For each prime number  $l \mid pq\widehat{D_i}$ ,  $C_{-D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{-D_i}{l}\right) = 1$ .  
(3)  $C_{-D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{p\widehat{D_i}}{D_i}\right) = \left(\frac{q\widehat{D_i}}{D_i}\right) = 1$ .

**Proof.** The proof is similar to Proposition 2.1, we omit the details.

**Corollary 2.6** We assume  $\varepsilon = -1$  and the elliptic curve  $E = E_-$  be as in

(1.1).

$$(A_1) \quad 2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 7(\text{mod}8) \text{ and } D_i \equiv 1, 7(\text{mod}8) \ (1 \leq i \leq n).$$

$$(A_2) \quad -2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 1(\text{mod}8) \text{ and } D_i \equiv 1, 3(\text{mod}8) \ (1 \leq i \leq n).$$

(B<sub>1</sub>) For each  $D_i$  ( $1 \leq i \leq n$ ), we have  $D_i \in S^{(\varphi)}(E/\mathbb{Q})$  if and only if

$$D_i \equiv 1(\text{mod}4) \text{ and } \left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1 \quad (1 \leq j \leq n, \text{ and } j \neq i).$$

(B<sub>2</sub>) For each  $D_i$  ( $1 \leq i \leq n$ ), we have  $-D_i \in S^{(\varphi)}(E/\mathbb{Q})$  if and only if

$$D_i \equiv 3(\text{mod}4) \text{ and } \left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1 \quad (1 \leq j \leq n, \text{ and } j \neq i).$$

**Proof.** The results follow easily from Proposition 2.5.  $\square$

**Proposition 2.7** We assume  $\varepsilon = -1$  and the elliptic curve  $E' = E'_-$  be as in

(1.2).

(A) (1) For any  $d \in \mathbb{Q}(S, 2)$  and  $d > 0$ ,  $C'_d(\mathbb{R}) \neq \emptyset$ . If  $2 \mid d$  or  $d < 0$ , then  $d \notin S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ .

(2)  $\{1, pq, pD, qD\} \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ .

(B) For each  $D_i$  ( $1 \leq i \leq n$ ), we have

(1)  $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$  if and only if one of the following conditions holds:

(a)  $D_i \equiv 1(\text{mod}8)$ ; (b)  $(1 - p\widehat{D_i})(1 - q\widehat{D_i}) \equiv 0(\text{mod}16)$ ;

(c)  $D_i \equiv 3(\text{mod}8)$  and  $p \equiv 1(\text{mod}4)$ ; (d)  $D_i \equiv 7(\text{mod}8)$  and  $p \equiv 3(\text{mod}4)$ .

(2)  $C'_{D_i}(\mathbb{Q}_p) \neq \emptyset$  and  $C'_{D_i}(\mathbb{Q}_q) \neq \emptyset$ .

(3) For each  $j \neq i$ ,  $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset \iff \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$ .

(4)  $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(1 - \left(\frac{p\widehat{D_i}}{D_i}\right)\right) \left(1 - \left(\frac{q\widehat{D_i}}{D_i}\right)\right) = 0$ .

(C) (1)  $C'_D(\mathbb{Q}_2) \neq \emptyset$  if and only if one of the following conditions holds:

(a)  $D \equiv 1(\text{mod}8)$ ; (b)  $p \equiv 1, 7(\text{mod}8)$ ;

(c)  $D \equiv 3(\text{mod}8)$  and  $p \equiv 5(\text{mod}8)$ ; (d)  $D \equiv 7(\text{mod}8)$  and  $p \equiv 3(\text{mod}8)$ .

(2)  $C'_D(\mathbb{Q}_p) \neq \emptyset$  and  $C'_D(\mathbb{Q}_q) \neq \emptyset$ .

(3) For each  $i$ ,  $C'_D(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(1 - \left(\frac{p}{D_i}\right)\right) \left(1 - \left(\frac{q}{D_i}\right)\right) = 0$ .

**Proof.** The proof is similar to Proposition 2.3, we omit the details.

**Corollary 2.8** We assume  $\varepsilon = -1$  and the elliptic curve  $E' = E'_-$  be as in (1.2).

For each  $D_i$  ( $1 \leq i \leq n$ ), denote

$$\Pi_i^-(D)' = \left(1 - \left(\frac{p\widehat{D}_i}{D_i}\right)\right) \left(1 - \left(\frac{q\widehat{D}_i}{D_i}\right)\right) + \sum_{j=1, j \neq i}^n \left(1 - \left(\frac{D_i}{D_j}\right)\right) \left(1 - \left(\frac{pqD_i}{D_j}\right)\right).$$

Then  $D_i \in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$  if and only if one of the following conditions holds:

$$\begin{aligned} (1) \quad & \begin{cases} D_i \equiv 1(\text{mod}8), \\ \Pi_i^-(D)' = 0; \end{cases} & (2) \quad & \begin{cases} (1 - p\widehat{D}_i)(1 - q\widehat{D}_i) \equiv 0(\text{mod}16), \\ \Pi_i^-(D)' = 0; \end{cases} \\ (3) \quad & \begin{cases} D_i \equiv 3(\text{mod}8), \\ p \equiv 1(\text{mod}4), \\ \Pi_i^-(D)' = 0; \end{cases} & (4) \quad & \begin{cases} D_i \equiv 7(\text{mod}8), \\ p \equiv 3(\text{mod}4), \\ \Pi_i^-(D)' = 0. \end{cases} \end{aligned}$$

**Proof.** The results follow easily from Proposition 2.7.  $\square$

Now we come to prove the Theorems.

**Proof of Theorem 1.1.** (1) For each  $i \in \{1, \dots, n\}$ , we have  $\left[\frac{1}{1+\Pi_i^+(D)}\right] = 0, 1$ . Furthermore,  $\left[\frac{1}{1+\Pi_i^+(D)}\right] = 1 \iff \Pi_i^+(D) = 0 \iff \left(\frac{-1}{D_i}\right) = 1$  and  $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = \left(\frac{D_i}{D_i}\right) = 1$  ( $1 \leq j \leq n$  and  $j \neq i$ )  $\iff D_i \in S^{(\varphi)}(E/\mathbb{Q})$  by Proposition 2.1. Then the conclusion (1) follows.

(2) Conclusion (2) follows easily from the above (1) and Corollary 2.2 (A). This proves Theorem 2.1.  $\square$

**Proof of Corollary 1.2.** By Proposition 2.1, we know that  $d \notin S^{(\varphi)}(E/\mathbb{Q})$  for any  $d \in \mathbb{Q}(S, 2)$  satisfying one of the following conditions holds: (1)  $d < 0$ ;

(2)  $p \mid d$ ; (3)  $q \mid d$ . Since  $S^{(\varphi)}(E/\mathbb{Q}) \subset \mathbb{Q}(S, 2) = \langle -1, 2, p, q, D_1, \dots, D_n \rangle$ , we get  $S^{(\varphi)}(E/\mathbb{Q}) \subset \langle 2, D_1, \dots, D_n \rangle$ . Furthermore, if condition (A) holds, then by Proposition 2.1, we have  $D_1, \dots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$ , so  $\langle D_1, \dots, D_n \rangle \subset S^{(\varphi)}(E/\mathbb{Q}) \subset \langle 2, D_1, \dots, D_n \rangle$ . If condition (B) holds, then by Proposition 2.1, we have  $D_1, \dots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$  but  $2 \notin S^{(\varphi)}(E/\mathbb{Q})$ , hence  $S^{(\varphi)}(E/\mathbb{Q}) \cong \langle D_1, \dots, D_n \rangle \cong (\mathbb{Z}/2\mathbb{Z})^n$ . If condition (C) holds, then by Proposition 2.1, we have  $2, D_1, \dots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$ , hence  $S^{(\varphi)}(E/\mathbb{Q}) \cong \langle 2, D_1, \dots, D_n \rangle \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$ . This proves Corollary 1.2.  $\square$

**Proof of Theorem 1.5.** The results of the Selmer groups are given in Corollaries 1.2 and 1.4 above.

Now  $E(\mathbb{Q})[2] = \{O, (0, 0), (-\varepsilon p D, 0), (-\varepsilon q D, 0)\}$ ,  $\varphi(E(\mathbb{Q})[2]) = \{O, (0, 0)\} = E'(\mathbb{Q})[\widehat{\varphi}]$ ,

by the exact sequences ([S] p.298, 314, 301 )

$$\begin{aligned} 0 &\longrightarrow \frac{E'(\mathbb{Q})}{\varphi(E(\mathbb{Q}))} \longrightarrow S^{(\varphi)}(E/\mathbb{Q}) \longrightarrow \text{TS}(E/\mathbb{Q})[\varphi] \longrightarrow 0, \\ 0 &\longrightarrow \frac{E(\mathbb{Q})}{\widehat{\varphi}(E'(\mathbb{Q}))} \longrightarrow S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \longrightarrow \text{TS}(E'/\mathbb{Q})[\widehat{\varphi}] \longrightarrow 0, \\ 0 &\longrightarrow \frac{E'(\mathbb{Q})[\widehat{\varphi}]}{\varphi(E(\mathbb{Q})[2])} \longrightarrow \frac{E'(\mathbb{Q})}{\varphi(E(\mathbb{Q}))} \longrightarrow \frac{E(\mathbb{Q})}{2E(\mathbb{Q})} \longrightarrow \frac{E(\mathbb{Q})}{\widehat{\varphi}(E'(\mathbb{Q}))} \longrightarrow 0, \quad \text{we get} \\ &\text{rank}(E(\mathbb{Q})) + \dim_2(\text{TS}(E/\mathbb{Q})[\varphi]) + \dim_2(\text{TS}(E'/\mathbb{Q})[\widehat{\varphi}]) \\ &= \dim_2(S^{(\varphi)}(E/\mathbb{Q})) + \dim_2(S^{(\widehat{\varphi})}(E'/\mathbb{Q})) - 2. \end{aligned}$$

From this and Corollaries 1.2 and 1.4, the results in Theorem 1.5 follow, and the proof is completed.  $\square$

Depending on the corresponding results in Propositions 2.3, 2.5, 2.7, the Theorems 1.3, 1.6, 1.8, 1.10 and their Corollaries can be similarly proved.

## References

[C ] J.W.S. Cassels, Lectures on Elliptic Curves, LMS 24, Cambridge University Press, New York, 1995.

- [**IK**] H. Iwaniec and E. Kowalski, *Analytic Number Theory*, American Mathematical Society, Providence, Rhode Island, 2004.
- [**KS**] R. Kloosterman and E. F. Schaefer, Selmer groups of elliptic curves that can be arbitrarily large, *Journal of Number Theory*, 99: 148-163 (2003).
- [**L**] S. Lang, *Algebra*, Revised Third Edition, GTM 211, Springer-Verlag, New York, 2002.
- [**MSS**] J.R. Merriman, S. Siksek and N.P.Smart, Explicit 4-descents on an elliptic curve, *Acta Arithmetica* lxxvii.(4):385-404(1996).
- [**QZ**] D. Qiu, X. Zhang, Mordell-weil groups and selmer groups of two types of elliptic curves, *Science in China (series A)*, 2002, Vol.45, No.11, 1372-1380.
- [**Sch**] S. Schmitt, Computation of the Selmer groups of certain parametrized elliptic curve, *Acta Arithmetica* lxxviii.(3):241-254(1997).
- [**S**] J.H.Silverman, *The Arithmetic of Elliptic Curves*, GTM 106, Springer-Verlag, New York, 1986.
- [**ST**] R.J.Strocker and J.Top, On the equation  $y^2 = (x + p)(x^2 + p^2)$ , *Rocky Mountain J. of Math.* 24(1994), 1135-1161.