

# Comment on “Rovibrational quantum interferometers and gravitational waves”

I. B. Khriplovich,<sup>1,\*</sup> S. K. Lamoreaux,<sup>2,†</sup> A. O. Sushkov,<sup>2,‡</sup> and O. P. Sushkov<sup>3,§</sup>

<sup>1</sup>*Budker Institute of Nuclear Physics, 630090, Novosibirsk, Russia*

<sup>2</sup>*Yale University, Department of Physics, P.O. Box 208120, New Haven, CT 06520-8120*

<sup>3</sup>*School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

In a recent paper, Wicht, Lämmerzahl, Lorek, and Dittus [Phys. Rev. **A** **78**, 013610 (2008)] come to the conclusion that a molecular rotational-vibrational quantum interferometer may possess the sensitivity necessary to detect gravitational waves. We do not agree with their results and demonstrate here that the true sensitivity of such interferometer is many orders of magnitude worse than that claimed in the mentioned paper. In the present comment we estimate the expected energy shifts and derive equations of motion for a quantum symmetric top (diatomic molecule or deformed nucleus) in the field of gravitational wave, and then estimate the sensitivity of possible experiments.

The direct detection of gravitational waves with an experimental apparatus located on or near the Earth is one of the forefront goals of modern science. Several large-scale detector projects, which are presently being either built or designed, promise the first glimpses of this form of radiation that most certainly must permeate the fabric of the Universe. Given the scale of these experiments, it is reasonable to ask whether gravitational waves might be more simply detected. In a recent paper, Wicht et al. [1] suggest that it is possible to detect such waves by measuring the energy shifts of rovibrational levels of a diatomic molecule. We show here that any such effect is many orders of magnitude smaller than predicted in [1].

Following Ref. [1] we start with the problem of the hydrogen atom in the field of a weak gravitational wave propagating along the  $z$  axis,  $h_{\mu\nu}(t-z)$ . The so-called TT gauge, where

$$h_{11} = -h_{22}, \quad h_{12} = h_{21}, \quad h_{0\mu} = h_{3\mu} = 0, \quad (1)$$

is convenient for the description of this wave. Wicht et al [1] base their analysis on the covariant Klein-Gordon equation. Expansion of this equation in powers of  $h_{ij}$  and  $1/c$  ( $c$  is speed of light) gives the following nonrelativistic hamiltonian for the electron of mass  $m$  and charge  $-e$

$$H = -\frac{\hbar^2}{2m}(\delta_{ij} - h_{ij})\partial_i\partial_j - eA_0, \quad (2)$$

and where the field of a Coulomb center with charge  $e$  in the presence of the gravitational wave is

$$A_0 = \frac{e}{r} \left( 1 - \frac{h_{ij}x_ix_j}{2r^2} \right). \quad (3)$$

This is the same hamiltonian used in [1], which we agree is valid.

Let us now calculate the energy shifts generated by the gravitational wave. The total system hamiltonian is

$$\begin{aligned} H &= H_0 + \delta H \\ H_0 &= \frac{p^2}{2m} - \frac{e^2}{r} \\ \delta H &= \frac{\hbar^2}{2m}h_{ij}\partial_i\partial_j + \frac{e^2h_{ij}x_ix_j}{2r^3}. \end{aligned} \quad (4)$$

Naively, the energy shift is of the order of the perturbation

$$\delta\epsilon \sim \langle \frac{\hbar^2}{2m}h_{ij}\partial_i\partial_j \rangle \sim \langle \frac{e^2h_{ij}x_ix_j}{2r^3} \rangle \sim h Ry, \quad (5)$$

where  $Ry$  stands for Rydberg, a typical atomic energy. However, we claim that the shift is much smaller, on the order of

$$\delta\epsilon \sim h \frac{(\hbar\omega)^2}{Ry}, \quad (6)$$

where  $\omega$  is the frequency of the gravitational wave, which is typically much lower than atomic or molecular frequencies of interest.

To prove this we perform the following transformation of coordinates  $x \rightarrow y$  in the hamiltonian (2),(3).

$$y_i = x_i + \frac{1}{2}h_{ij}x_j. \quad (7)$$

This is the transformation into the frame freely falling in the gravitational wave. After this transformation, the interaction with the gravitational wave is completely removed from Eqs. (2) and (3). However, this does not mean that the interaction with the gravitational wave vanishes completely. Consider the time-dependent Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \Big|_x \psi(t, x) = H\psi(t, x). \quad (8)$$

The partial time derivative is taken at constant  $x$ . Under the transformation in Eq.(7) the left hand side of Eq.(8) is changed as

$$\frac{\partial}{\partial t} \Big|_x = \frac{\partial}{\partial t} \Big|_y + \frac{\partial y_i}{\partial t} \frac{\partial}{\partial y_i} = \frac{\partial}{\partial t} \Big|_y + \frac{1}{2} \dot{h}_{ik} \frac{\partial}{\partial y_i}. \quad (9)$$

\*Electronic address: khriplovich@inp.nsk.su

†Electronic address: steve.lamoreaux@yale.edu

‡Electronic address: alex.sushkov@yale.edu

§Electronic address: sushkov@phys.unsw.edu.au

Therefore Schrödinger Eq. (8) is transformed to

$$i\hbar \frac{\partial}{\partial t} \psi(t, y) = \left[ H_0(y) + \frac{1}{2} \dot{h}_{ik} y_k p_i \right] \psi(t, y), \quad (10)$$

where the equation is written in the  $y$ -coordinates and the time derivative is necessarily taken at  $y = \text{constant}$ . Hence, the new hamiltonian is

$$H' = H_0 + \frac{1}{2} \dot{h}_{ik} y_k p_i = H_0 + \frac{1}{2} m \dot{h}_{ik} y_k \dot{y}_i, \quad (11)$$

where we have used  $p_i = m\dot{y}_i$ . The new perturbation is

$$\begin{aligned} \delta H' &= \frac{1}{2} m \dot{h}_{ik} y_k \dot{y}_i = \frac{d}{dt} \left( \frac{1}{4} m \dot{h}_{ik} y_k y_i \right) - \frac{1}{4} m \ddot{h}_{ik} y_k y_i \\ &\rightarrow -\frac{1}{4} m \ddot{h}_{ik} y_k y_i. \end{aligned} \quad (12)$$

In this equation we have thrown away the full time derivative because it influences neither the equations of motion nor the physical results. Equation (12) justifies our estimate (6) for the energy shift or for a typical matrix element, if we substitute  $\ddot{h}_{ik} \sim \omega^2 \hbar$  and  $y_i, y_k \sim a_B = \hbar^2/m_e^2$ , where  $a_B$  is the Bohr radius.

Obviously, the above statements are valid not only for a hydrogen atom. One can perform the transformation (7) for any quantum system in a gravitational wave; furthermore one can certainly work in terms of original  $x$ -coordinates as well. For example, for the problem of the gravitational radiation of atomic hydrogen, the equivalence of the two approaches has been demonstrated explicitly in Ref. [2]. However, in the case of low frequency gravitational waves, calculations in the  $x$ -coordinates (“ $x$ -frame” calculations) are “dangerous” because there is a dramatic exact cancelation (or compensation) between different large terms in the perturbation  $\delta H$  given by Eq. (4) and approximations can easily destroy this exact compensation. Because the result in [1] does not contain the frequency of the gravitational wave and parametrically coincides with the naive estimate (5), we believe that this compensation was destroyed in Ref. [1] due to approximations in the molecular calculations for the ionic molecule  $\text{HD}^+$ .

One can derive Eq. (12) in a much simpler way using a frame freely falling in the gravitational wave from the very beginning. In this frame the center of mass of the system remains at rest. The motion of any particle of this system with respect to the center of mass is described by the well-known covariant equation (see, e.g., [3, 4])

$$\frac{D^2 \eta^\mu}{Ds^2} + R^\mu{}_{\rho\nu\tau} u^\rho u^\tau \eta^\nu = 0. \quad (13)$$

In the general case, this equation for the so-called geodesic deviation  $\eta^\mu$  has the following meaning: Particles  $a$  and  $b$  move along two close geodesics  $x_a^\mu(s)$  and  $x_b^\mu(s)$ ; then, by definition,  $\eta^\mu(s) = x_a^\mu(s) - x_b^\mu(s)$ , and  $u^\mu = dx_b^\mu(s)/ds$ .

In our problem, let the index  $b$  label the coordinate of center of mass which is at rest in the chosen reference

frame. Then  $u^\mu = (1, 0, 0, 0)$ , and in the weak field of the gravitational wave  $h_{mn} = h_{mn}^{(0)} \exp[-i\omega(t-z)]$  we arrive at the following equation of motion for particle  $a$ :

$$\ddot{x}_m^a = -\frac{1}{2} \omega^2 h_{mn} x_n^a. \quad (14)$$

In the hydrogen atom (or hydrogen-like ion), where the coordinates of electron, center of mass, and proton (or nucleus),  $x^e$ ,  $x^c$ , and  $x^p$ , respectively, lie on a straight line, we arrive immediately at the following equation of motion for the relative coordinate  $x = x^e - x^p$ :

$$\ddot{x}_m = -\frac{1}{2} \omega^2 h_{mn} x_n. \quad (15)$$

The corresponding interaction hamiltonian for the hydrogen atom is therefore

$$H_{int} = \frac{1}{4} \mu^{ep} \omega^2 h_{mn} x_m x_n; \quad (16)$$

$$\mu^{ep} = \frac{m^e m^p}{m^e + m^p} \simeq m^e.$$

Let us note here that because  $h_{mn}$  depend on  $t-z$ , but not on  $x_{m,n}$ , the relations (15) and (16) are in exact correspondence. Eq. (16) coincides with Eq. (12).

The molecular ion  $\text{HD}^+$  considered in Ref. [1] consists of three constituents: an electron, a proton, and a deuteron. To simplify the description, we note that, with  $m^e \ll m^p, m^d$ , one can assume that the coordinates of proton, center of mass, and deuteron,  $x^p, x^c$ , and  $x^d$ , respectively, lie on a straight line, so that for the relative coordinate  $x = x^p - x^d$  we have the same equation, (15). Therefore, the interaction hamiltonian for the  $\text{HD}^+$  ion can be written as

$$H_{int} = \frac{1}{4} \mu^{dp} \omega^2 h_{mn} x_m x_n; \quad (17)$$

$$\mu^{dp} = \frac{m^d m^p}{m^d + m^p} \simeq \frac{2}{3} m^p.$$

We have neglected here the formally-arising term

$$\frac{1}{4} m^e \omega^2 h_{mn} x_m^e x_n^e,$$

because the assumption  $m^e \ll m^p$  has been made already for the derivation of hamiltonian (17); furthermore, the contribution of this term to the discussed effect is relatively suppressed as  $m^e/m^p$ .

The typical frequency shift for a  $\text{HD}^+$  ion, as estimated with hamiltonian in Eq. (17), is

$$\delta\nu \sim \frac{1}{\hbar} m^p \omega^2 h a_B^2. \quad (18)$$

This estimate is valid as long as the frequency  $\omega$  of the gravitational wave is much less than that of the molecular

vibrations and rotations. For the dimensionless amplitude  $h \sim 10^{-19}$  and frequency  $\omega \sim 10^4$  Hz, we estimate a frequency shift as  $\sim 10^{-23}$  Hz. This estimate is well below the corresponding estimate  $\sim 3 \times 10^{-5}$  Hz presented in Ref. [1]. Moreover, this frequency shift varies with time together with  $h(t-z)$ . With this time-dependence of the frequency shift, the observation of the effect seems even less realistic because high precision spin precession measurements generally require considerable averaging time (days to years).

Let us make some additional remarks about the behavior of a molecule (or a deformed nucleus) under the influence of the hamiltonian (17). This hamiltonian influences both the vibrational and rotational degrees of freedom of the diatomic molecule. The classical (non-quantum) vibrational dynamics in the gravitational wave are discussed in several textbooks, see e.g. Ref. [5], and quantum mechanical treatment does not bring significant differences in the dynamics. Here we derive the implications of Eq. (17) for rotational dynamics. The classical equation of motion for angular momentum  $\mathbf{L} = \mathbf{r} \times \mathbf{p}$  follows from Eq. (17)

$$\dot{L}_i = \frac{1}{2}\omega^2 \epsilon_{ikl} h_{ln} I_{kn} , \quad (19)$$

where is  $I_{kn}$  is the tensor of inertia of the molecule. Only the traceless part of  $I_{kn}$  contributes to Eq. (19), and the molecule is a symmetric top. As usual we denote by  $I_{||}$  the moment of inertia for rotations around axis of the top, and by  $I_{\perp}$  the moment of rotations around a perpendicular axis.<sup>1</sup> Then

$$I_{kn} = \frac{1}{3} (2I_{\perp} + I_{||}) \delta_{kn} + (I_{||} - I_{\perp}) \left( n_k n_n - \frac{1}{3} \delta_{kn} \right). \quad (20)$$

Clearly Eqs.(19) and (20) are valid for both the classical and quantum cases. Let us consider a molecule in a rotational quantum state with a given angular momentum  $J$ ,  $\mathbf{L} = \hbar \mathbf{J}$ ,  $\hat{\mathbf{J}}^2 = J(J+1)$ . Then (see e.g. [6])

$$I_{kn} \rightarrow \mathcal{I} \left( J_k J_n + J_n J_k - \frac{2}{3} J(J+1) \delta_{kn} \right) \\ \mathcal{I} = (I_{||} - I_{\perp}) \frac{3\Omega^2 - J(J+1)}{J(2J-1)(J+1)(2J+3)}. \quad (21)$$

Here we keep only the traceless part of the inertia tensor, and  $\Omega$  is projection of  $\mathbf{J}$  on axis of the top. Hence Eq. (19) is transformed to

$$\dot{J}_i = \frac{\mathcal{I}}{2\hbar} \omega^2 \epsilon_{ikl} h_{ln} \{ J_k J_n + J_n J_k \}. \quad (22)$$

The corresponding effective Hamiltonian is

$$H_{eff} = -\frac{\mathcal{I}}{2} \omega^2 h_{ln} J_l J_n. \quad (23)$$

Eqs. (21) and (23) are equally applicable to a diatomic molecule or to a deformed nucleus, and describe transitions or energy shifts between magnetic sublevels. For the ground state  $J = \Omega = S$ ,  $S$  is spin of the system. We assume that  $S > 1/2$ . Hence Eq. (22) is transformed to equation of spin precession in the field of gravitational wave

$$\dot{S}_i = -\frac{I_{\perp} - I_{||}}{2\hbar(S+1)(2S+3)} \omega^2 \epsilon_{ikl} h_{ln} (S_k S_n + S_n S_k). \quad (24)$$

This is a general equation for a free spin.

Given that a magnetic resonance frequency shift due to (24) is extremely small, we alternatively consider the case of a solid, where there is a possibility of a much larger nuclear electric quadrupole field shift. This shift is induced by the deformation of the lattice due to the gravitational wave. For a sample with size  $\sim 1$  m and for  $\omega \gtrsim 10^4$  Hz the frequency of the gravitational wave disappears from the equations because the frequency of the lowest acoustic standing wave  $\omega_0$  is lower than  $\omega$ . The cancellation of the  $\omega$ -dependence is similar to that for the LIGO experiment discussed below, see eqs. (27), (28). Hence the maximum possible nuclear quadrupole resonance frequency shift is

$$\dot{S} \sim 10^9 h \text{ Hz}. \quad (25)$$

for the case of deformed nuclei.

Unfortunately the frequency shift  $10^{-10}$  Hz given by Eq. (25) with  $h \sim 10^{-19}$  is still very small. The current state of the art nuclear spin precession frequency measurements accuracy (for a gas of  $^{199}\text{Hg}$  atoms used in a permanent electric dipole moment (EDM) search [7]) is barely at this level after several years of measurement and averaging, and this experimental accuracy far exceeds that of any other experiment performed to date. Of course, the possibility of a macroscopic sample with a large number of participating spins suggests that some increase in sensitivity might be possible, but there is a considerable loss in sensitivity due to large-sample measurement limitations. The only plausible measurement technique is the much less efficient monitoring of the sample's precessing magnetization with a SQUID or other magnetometer, compared to the direct optical detection of the spin state that can be used with gaseous samples.

Let us add a brief remark about gravity wave detectors based on strain measurements (such as LIGO). These are also described by Eq. (14). Suppose a mass (such as an interferometer mirror) is positioned in equilibrium at  $x = y = z = 0$ , and another mass (the second mirror) is positioned in equilibrium at  $x = L$ ,  $y = z = 0$ . The mass at the origin is unaffected by the gravity wave, the response of the mass at  $x = L$  is given by

$$\ddot{x} + \omega_0^2(x - L) = -\frac{1}{2}\omega^2 h_+(t)L, \quad (26)$$

where  $\omega_0$  is the natural oscillation frequency around the equilibrium point,  $h_+ = h_{11} = -h_{22}$ , and we neglect

<sup>1</sup> For a molecule or deformed nucleus one must set  $I_{||} = 0$ . Nevertheless, we keep  $I_{||}$  to stress that only traceless part of the inertia tensor contributes to the spin precession.

damping. For a wave  $h_+(t) = h_+^{(0)} e^{-i\omega t}$  the solution is

$$x = L + \frac{1}{2} \frac{\omega^2}{\omega^2 - \omega_0^2} h_+^{(0)} L e^{-i\omega t}. \quad (27)$$

LIGO is a “free-mass” detector, which means that  $\omega_0 \ll \omega$  ( $\omega_0 \approx 4$  Hz for Advanced LIGO [8]). This gives the strain

$$\Delta L/L = \frac{1}{2} h_+^{(0)} e^{-i\omega t}. \quad (28)$$

The strain in oscillator modes with frequencies much greater than the gravitational wave frequency is suppressed by  $\omega^2/\omega_0^2$ . Indeed, the more rigid the system,

the less sensitive it is to an external force. In a molecule, for example,  $\omega_0$  is on the order of THz, which gives an enormous strain suppression for gravitational wave frequencies  $\omega \lesssim 10^4$  Hz.

### Acknowledgements

The authors would like to acknowledge useful discussions with David DeMille, Dmitry Budker, and Max Zolotarev. The work by I.B. Khriplovich was supported by the Russian Foundation for Basic Research through Grant No. 08-02-00960-a.

- 
- [1] A. Wicht, C. Lämmerzahl, D. Lorek, and H. Dittus, *Phys. Rev. A* **78**, 013610 (2008).
  - [2] S. Boughn and T. Rothman, *Class. Quantum Grav.* **23**, 5839 (2006); gr-qc/0605052 v.2 (see, in particular, section 5 therein).
  - [3] J. Weber, *General Relativity and Gravitational Waves*, Interscience Publishers Inc., New York, 1961.
  - [4] A.P. Lightman, W.H. Press, R.H. Price, S.A. Teukolsky, *Problem Book in Relativity and Gravitation*, Princeton University Press, Princeton, New Jersey, 1975, problem 9.23.
  - [5] C. W. Misner, K. S. Thorne, J. A. Wheeler, *Gravitation*, San Francisco : W. H. Freeman, 1973, section 37.5.
  - [6] L.D. Landau and E.M. Lifshitz, *Quantum mechanics: non-relativistic theory*, Oxford, New York, Pergamon Press, 1965, section 119, problem 1.
  - [7] W. C. Griffith, M. D. Swallows, T. H. Loftus, M. V. Romalis, B. R. Heckel, and E. N. Fortson, *Phys. Rev. Lett.* **102**, 101601 (2009).
  - [8] Advanced LIGO Reference Design Document, LIGO M060056-08-M, 2007.